ANOVA a Misure Ripetute

Effetti Entro i Sggetti

	Correzione della Sfericità	Somma dei Quadrati	gdl	Media Quadratica	F	p	η^2	η^2_p
Training	Nessuna	0.9075	1	0.90749	227.9	<.001	0.347	0.962
	Greenhouse-Geisser	0.9075	1.00	0.90749	227.9	<.001	0.347	0.962
Residuo	Nessuna	0.0358	9	0.00398				
	Greenhouse-Geisser	0.0358	9.00	0.00398				
Blocco	Nessuna	0.5511	2	0.27553	55.1	<.001	0.211	0.860
	Greenhouse-Geisser	0.5511	1.54	0.35692	55.1	<.001	0.211	0.860
Residuo	Nessuna	0.0900	18	0.00500				
	Greenhouse-Geisser	0.0900	13.90	0.00648				
Training ✻ Blocco	Nessuna	0.1810	2	0.09052	10.4	<.001	0.069	0.537
	Greenhouse-Geisser	0.1810	1.12	0.16109	10.4	.008	0.069	0.537
Residuo	Nessuna	0.1561	18	0.00867				
	Greenhouse-Geisser	0.1561	10.11	0.01543				

Nota. Somma dei quadrati Tipo 3

Effetti Tra Soggetti

	Somma dei Quadrati	gdl	Media Quadratica	F	p	η^2	η^2_p
Residuo	0.694	9	0.0771				

Nota. Somma dei quadrati Tipo 3

Ipotesi

Test di Sfericità

	W di Mauchly	p	Greenhouse-Geisser ε	Huynh-Feldt ε
Training[a]	1.000	NaN	1.000	1.000
Blocco	0.705	.246	0.772	0.901
Training ✻ Blocco	0.220	.002	0.562	0.586

[a] Le misure ripetute hanno solo due livelli. L'assunzione di sfericità è sempre soddisfatta quando le misure ripetute hanno solo due livelli.

Test Post Hoc

Confronti Post Hoc - Training

Confronto											
Training		Training	Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
T1	-	T2	0.246	0.0163	9.00	15.1	<.001	<.001	<.001	<.001	<.001

Confronti Post Hoc - Blocco

Confronto										
Blocco		Blocco	Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
B1	-	B2	0.1889	0.0259	9.00	7.29	<.001	<.001	<.001	<.001	<.001
	-	B3	0.2151	0.0244	9.00	8.80	<.001	<.001	<.001	<.001	<.001
B2	-	B3	0.0262	0.0152	9.00	1.72	.119	.249	.277	.356	.119

Confronti Post Hoc - Training ✻ Blocco

Confronto													
Training	Blocco		Training	Blocco	Differenza Media	SE	gdl	t	p	p_{tukey}	$p_{scheffe}$	$p_{bonferroni}$	p_{holm}
T1	B1	-	T1	B2	0.2784	0.0497	9.00	5.606	<.001	.003	.009	.005	.003
		-	T1	B3	0.3469	0.0585	9.00	5.928	<.001	.002	.006	.003	.002
		-	T2	B1	0.3935	0.0530	9.00	7.419	<.001	<.001	.001	<.001	<.001
		-	T2	B2	0.4929	0.0486	9.00	10.143	<.001	<.001	<.001	<.001	<.001
		-	T2	B3	0.4768	0.0382	9.00	12.490	<.001	<.001	<.001	<.001	<.001
	B2	-	T1	B3	0.0686	0.0229	9.00	2.998	.015	.110	.210	.225	.060
		-	T2	B1	0.1151	0.0257	9.00	4.481	.002	.014	.034	.023	.009
		-	T2	B2	0.2145	0.0257	9.00	8.360	<.001	<.001	<.001	<.001	<.001
		-	T2	B3	0.1985	0.0297	9.00	6.691	<.001	<.001	.003	.001	<.001
	B3	-	T2	B1	0.0465	0.0177	9.00	2.630	.027	.184	.316	.410	.060
		-	T2	B2	0.1460	0.0256	9.00	5.701	<.001	.003	.008	.004	.003
		-	T2	B3	0.1299	0.0282	9.00	4.610	.001	.011	.029	.019	.009
T2	B1	-	T2	B2	0.0994	0.0259	9.00	3.845	.004	.033	.075	.059	.020
		-	T2	B3	0.0834	0.0283	9.00	2.946	.016	.119	.222	.245	.060
	B2	-	T2	B3	-0.0161	0.0179	9.00	-0.898	.393	.938	.971	1.000	.393