

Discrete Structures: Homework #2

Due on 30 June 2020

Professor Jensen Section 201

Brian Ton

Problem 1

Is the statement “ \forall real numbers x and y , $\sqrt{x+y} = \sqrt{x} + \sqrt{y}$ ” *True* or *False*? If it is false, provide a counterexample.

Solution

False

Consider the case $x = 4$, $y = 5$. In this case, $\sqrt{x+y} = \sqrt{4+5} = \sqrt{9} = 3$. However, $\sqrt{4} + \sqrt{5} = 2 + \sqrt{5} \neq 3$.

Problem 2

Is the statement “ \exists real numbers x and y such that $\sqrt{x+y} = \sqrt{x} + \sqrt{y}$ ” *True* or *False*? If it is true, provide an example.

Solution

True

Consider the case $x = 1$, $y = 0$. In this case, $\sqrt{x+y} = \sqrt{1+0} = \sqrt{1} = 1$ and $\sqrt{1} + \sqrt{0} = \sqrt{1} = 1$. More generally, this is true when at least one of x or y is 0 (which follows from the identity element axiom of addition for real numbers, as $\sqrt{0} = 0$ and $\sqrt{x+0} = \sqrt{x}$, meaning that $\sqrt{x+0} = \sqrt{x} + \sqrt{0}$).

Problem 3

Write a negation for each statement.

- There exists a real number x such that $x \leq -2$
- \forall computer programs P , if P compiles without error messages, then P is correct.
- \forall integers n , \exists a prime number p such that $n < p < 2n$.

Solution

- For all real numbers x , $x > -2$
- There exists a computer program P such that P compiles without error messages and P is not correct.
- There exists an integer n such that for all prime numbers p , $(p \leq x) \vee (p \geq 2x)$

Part A

Let $P(x)$ be the statement $x \leq -2$.

The statement can then be written as $\exists x P(x)$. By De Morgan's Law of Quantifiers, $\neg \exists x P(x) \equiv \forall x \neg P(x)$. Additionally, note that $\neg(x \leq -2) \equiv (x > -2)$. Thus, the negation of the original statement is $\forall x (x > -2)$, which in English is “For all real numbers x , $x > -2$ ”.

Part B

Let $E(x)$ be the statement “ x compiles without error messages.” and $C(x)$ be the statement “ x is correct.” The original statement can then be written as $\forall P (E(P) \rightarrow C(P))$. By De Morgan's Law of Quantifiers, $\neg \forall P (E(P) \rightarrow C(P)) \equiv \exists P \neg (E(P) \rightarrow C(P))$. By conditional disjunction, $\exists P \neg (E(P) \rightarrow C(P)) \equiv \exists P \neg (\neg E(P) \vee C(P))$, and by De Morgan's law of propositional logic, $\exists P \neg (\neg E(P) \vee C(P)) \equiv \exists P (E(P) \wedge \neg C(P))$, which in English would be “There exists a computer program P such that P compiles without error messages and P is not correct.”

Part C

Let $P(x, p)$ be the statement $x < p < 2x$. The original statement can then be written as $\forall n \exists p (P(n, p))$. Note that $\neg(x < p < 2x) \equiv (p \leq x) \vee (p \geq 2x)$. By De Morgan's Law of Quantifiers, $\neg \forall n \exists p (P(n, p)) \equiv \exists n \neg \exists p (P(n, p)) \equiv \exists n \forall p (\neg P(n, p))$. Then, taking note of the negation of $P(x)$, $\exists n \forall p (\neg P(n, p)) \equiv \exists n \forall p ((p \leq n) \vee (p \geq 2n))$. Written in English, the statement would be "There exists an integer n such that for all prime numbers p , $(p \leq n) \vee (p \geq 2n)$."

Problem 4

Show that $\exists x P(x) \wedge \exists x Q(x)$ and $\exists x (P(x) \wedge Q(x))$ are not logically equivalent.

Solution

Let $P(x)$ be the statement $x < 1$ and $Q(x)$ be the statement $x > 1$, where $x \in \mathbb{Z}$.

The first statement is then $\exists x (x < 1) \wedge \exists x (x > 1)$. This is a true statement, as there does exist an integer x that is less than 1 (e.g. 0) and an x that is greater than 1 (e.g. 2). However, the second statement is false, as there exists no integer x that is both greater than and less than 1. $\therefore \exists x P(x) \wedge \exists x Q(x) \not\equiv \exists x (P(x) \wedge Q(x))$

Problem 5

Show that $\forall x P(x) \vee \forall x Q(x)$ and $\forall x (P(x) \vee Q(x))$ are not logically equivalent.

Solution

Let $P(x)$ be the statement $x \geq 1$ and $Q(x)$ be the statement $x < 1$, where $x \in \mathbb{Z}$.

The first statement is false, since $\forall x P(x)$ is false, as not every integer is greater than or equal to 1, and $\forall x Q(x)$ is false, as not every integer is less than 1. However, the second statement is true, as every integer is greater than or equal to or less than 1. $\therefore \forall x P(x) \vee \forall x Q(x) \not\equiv \forall x (P(x) \vee Q(x))$.

Problem 6

Let $P(x, y)$ be the statement " $xy = 1$ ". If the domain for both variables is the set of nonzero real numbers, what are the truth values?

- a) $\exists y \forall x P(x, y)$
- b) $\forall x \exists y P(x, y)$

Solution

- a) False
- b) True