

# Resolution and accuracy in Congreve & Lamsdell matrices

*Martin R. Smith martin.smith@durham.ac.uk*


2019-07-20

## Contents


0.1	Summary . . . . .	1
0.2	Trees 1–10 . . . . .	2
0.3	Trees 11–20 . . . . .	5
0.4	Trees 21–30 . . . . .	8
0.5	Trees 31–40 . . . . .	11
0.6	Trees 41–50 . . . . .	14
0.7	Trees 51–60 . . . . .	17
0.8	Trees 61–70 . . . . .	20
0.9	Trees 71–80 . . . . .	23
0.10	Trees 81–90 . . . . .	26
0.11	Trees 91–100 . . . . .	29
	References . . . . .	31


This page depicts the analytical results of all 100 matrices generated by Congreve & Lamsdell [1] using a ternary plotting approach [2], with quartets and partitions used as distance metrics.

## 0.1 Summary






## 0.2 Trees 1–10











Dataset 5: CI=0.26





Dataset 6: CI=0.26









Dataset 7: CI=0.27




Dataset 8: CI=0.25









### 0.3 Trees 11–20













Dataset 15: CI=0.25





Dataset 16: CI=0.25









Dataset 17: CI=0.27




Dataset 18: CI=0.28








## 0.4 Trees 21–30











Dataset 25: CI=0.28




Dataset 26: CI=0.27




Dataset 27: CI=0.26






Dataset 28: CI=0.27








**Dataset 29: CI=0.29**




**Dataset 30: CI=0.25**





## 0.5 Trees 31–40









Dataset 35: CI=0.25




Dataset 36: CI=0.24




Dataset 37: CI=0.26




Dataset 38: CI=0.29








Dataset 39: CI=0.25





Dataset 40: CI=0.24







## 0.6 Trees 41–50









Dataset 45: CI=0.27




Dataset 46: CI=0.24




Dataset 47: CI=0.29






Dataset 48: CI=0.27








**Dataset 49: CI=0.28**






**Dataset 50: CI=0.25**






## 0.7 Trees 51–60






**Dataset 51: CI=0.3**





**Dataset 52: CI=0.25**




**Dataset 53: CI=0.25**





**Dataset 54: CI=0.28**











Dataset 55: CI=0.28




Dataset 56: CI=0.31




Dataset 57: CI=0.25





Dataset 58: CI=0.25











**Dataset 59: CI=0.26**




**Dataset 60: CI=0.24**






## 0.8 Trees 61–70











Dataset 65: CI=0.25





Dataset 66: CI=0.27









Dataset 67: CI=0.26




Dataset 68: CI=0.31








## 0.9 Trees 71–80











Dataset 75: CI=0.24




Dataset 76: CI=0.26




Dataset 77: CI=0.28






Dataset 78: CI=0.24








**Dataset 79: CI=0.24**




**Dataset 80: CI=0.27**






## 0.10 Trees 81–90











Dataset 85: CI=0.27





Dataset 86: CI=0.27









Dataset 87: CI=0.23




Dataset 88: CI=0.25










## 0.11 Trees 91–100









Dataset 95: CI=0.24




Dataset 96: CI=0.26




Dataset 97: CI=0.27



Dataset 98: CI=0.26





## References

1. Congreve CR, Lamsdell JC. 2016 Implied weighting and its utility in palaeontological datasets: a study using modelled phylogenetic matrices. *Palaeontology* **59**, 447–465. (doi:10.1111/pala.12236)
2. Smith MR. 2019 Bayesian and parsimony approaches reconstruct informative trees from simulated morphological datasets. *Biology Letters* **15**, 20180632. (doi:10.1098/rsbl.2018.0632)