


Statistics for Business and Economics

Mohammad Saifuddin

2026-09-04

Statistics for Business and Economics



Mohammad Saifuddin

MS in Applied Statistics,
University of Dhaka

Table of contents

Preface	4
1 Data and Statistics	5
1.1 Statistics	5
1.2 Types of statistics	5
1.3 Applications in Business and Economics	6
1.4 Data	6
1.5 Elements, Variables, and Observations	7
1.6 Scales of Measurement	8
1.7 Quantitative and Categorical and Data	8
1.8 Cross-Sectional and Time Series Data	9
1.9 Data vs. Information	9
1.10 Exercise	10
2 Descriptive statistic: Tabular and Graphical Presentations	11
2.1 Summarizing Categorical Data	11
2.2 Summarizing Quantitative Data	14
2.3 Data	19
2.4 Stem-and-leaf display	19
2.5 Solution-I	20
2.6 Solution-II	20
2.7 Exercises	21
2.8 Data	21
2.9 Histogram	22
2.10 Data	22
2.11 Histogram	23
2.12 Line Chart	23
2.13 Scatter diagram	23
2.14 Case Study: Lifestyle Indicators and Preferences	23
3 Descriptive statistics: Numerical Measures	26
3.1 Measures of location	26
3.2 Measures of variability	31
3.3 The mean and standard deviation of Grouped data	35
3.4 Measures of relative location: z-score	36
3.5 Five-Number summary	40
3.6 Box-plot	41
3.7 Measures of shape: Skewness and Kurtosis	45
3.8 Exercise	50
3.9 Data	53
3.10 Ordered data	53

4	Probability	54
4.1	Random experiment	54
4.2	Sample space	54
4.3	Event	54
4.4	Complement of an event	55
4.5	Mutually exclusive events	55
4.6	Collectively Exhaustive	55
4.7	Axiomatic definition of Probability	55
4.8	Probability of an event (Classical approach)	56
4.9	Probability of an event (Empirical approach)	57
4.10	Properties of Probability Laws	57
4.11	Conditional Probability	58
4.12	The Multiplication Rule	58
4.13	Independent events	59
4.14	Bivariate Probabilities: Joint and Marginal Probability	60
4.15	Independent Events in Joint probability table	61
4.16	Exercises 4.1	62
4.17	Probability Trees	64
4.18	Total Probability rule and Bayes' Theorem	65
4.19	Exercises 4.2	66
4.20	Random variables	68
5	Discrete Probability Distributions	70
5.1	Discrete random variable and Probability mass function (PMF)	70
5.2	Joint distribution of two discrete r.vs	74
5.3	Bernoulli distribution/r.v	76
5.4	Binomial r.v	77
5.5	Poisson r.v	79
5.6	Multinomial Probability Distribution	82
6	Continuous Probability Distributions	83
6.1	Probability density function (PDF)	83
6.2	Uniform probability distribution/r.v	86
6.3	Normal distribution/r.v	88
6.4	Exponential distribution	97
6.5	Joint distribution of two continuous r.vs	98
6.6	Bivariate Normal distribution	98
6.7	Some other important Probability Densities	99
7	Further topics on random variables	101
7.1	Linear functions of random variables	101
7.2	Functions of random variables: Transformations	102
7.3	Moments Moment-Generating Functions (MGF)	103
8	Sampling and Sampling distributions	105
8.1	Some preliminary idea (Anderson 2020a)	105
8.2	Sampling from a Finite Population	105
8.3	Sampling from an Infinite Population	105
8.4	Sampling distribution	108

8.5	Sampling distribution of $\bar{X}$	109
8.6	Sampling distribution of sample proportion, $\hat{p}$	112
8.7	Sampling Distribution of the Sample Variances	113
8.8	t -Distribution	116
8.9	F -Distribution	118
8.10	Other Sampling methods	118
9	Introduction to estimation	119
9.1	Point Estimation	119
9.2	Interval estimation	123
9.3	Sample size determination: Large Population	127
9.4	Interval estimation: Comparing TWO populations	129
10	Hypothesis test: Introduction and testing one population parameter	130
10.1	Definition	130
10.2	Types of hypothesis	130
10.3	Developing hypotheses	130
10.4	Types of test based on alternative hypothesis H_1	131
10.5	Types of error in hypothesis test and P-value	131
10.6	Hypothesis testing concerning population mean (μ)	132
10.7	Hypothesis test of a Population variance	136
10.8	Hypothesis test of a Population proportion	136
10.9	Normality test	136
11	Hypothesis test concerning TWO population parameters	138
11.1	Hypothesis test: Difference between two population means ($\mu_1 - \mu_2$)	138
11.2	Testing the Population Variances ($\sigma_1^2 = \sigma_2^2$)	140
11.3	Hypothesis test: Comparing TWO means when the samples are dependent/ matched/paired	143
12	Analysis of Variance	147
12.1	Sources of Variation for a One-Way Analysis of Variance	147
12.2	Assumptions for a One-Way Analysis of Variance	147
12.3	Hypotheses for a One-Way Analysis of Variance	147
12.4	Sample statistics used in one-way ANOVA	148
12.5	The Between-Sample Variability for One-Way Analysis of Variance	148
12.6	The Within-Sample Variability for One-Way Analysis of Variance	148
12.7	The F Statistic	148
12.8	Rejection rule	148
12.9	Example 12.1	148
12.10	Multiple comparison	151
12.11	Example 12.2	152
13	Chi-squared Test	155
13.1	Goodness of Fit Test	155
13.2	Test for Independence (Categorical Data)	160
14	Correlation and Simple Linear Regression	163
14.1	Scatter plot: Graphical method to explore correlation	163
14.2	Covariance	164

14.3 Coefficient of Correlation	166
14.4 Properties of coefficient of correlation	167
14.5 Rank correlation	170
14.6 Simple linear regression (SLR)	173
15 Summary	179
References	180

Preface

This book is specially for the undergrad students of Business and Economics program providing basic to advance statistical tools and techniques to handle data . The book is under development.

1 Data and Statistics

1.1 Statistics

Statistics is defined as the art and science of collecting, analyzing, presenting, and interpreting data.

Particularly in business and economics, the information provided by **collecting, analyzing, presenting, and interpreting data** gives managers and decision makers a better understanding of the business and economic environment and thus enables them to make more informed and better decisions.

1.2 Types of statistics

Descriptive Statistics

Most of the statistical information in newspapers, magazines, company reports, and other publications consists of data that are summarized and presented in a form that is easy for the reader to understand. Such summaries of data, which may be tabular, graphical, or numerical, are referred to as **descriptive statistics**.

Inferential statistics (Statistical Inference)

Many situations require information about a large group of elements (individuals, companies, voters, households, products, customers, and so on). But, because of time, cost, and other considerations, data can be collected from only a small portion of the group. The larger group of elements in a particular study is called the **population**, and the smaller group is called the **sample**. Formally, we use the following definitions.

- **Population** A population is the set of all elements of interest in a particular study.
- **Sample** A sample is a subset of the population.

The process of conducting a survey to collect data for the entire population is called a **census**.

The process of conducting a survey to collect data for a sample is called a **sample survey**.

As one of its major contributions, statistics uses data from a sample to make estimates and test hypotheses about the characteristics of a population through a process referred to as **statistical inference**.

1.3 Applications in Business and Economics

Business Applications

- **Marketing:** Market segmentation, customer surveys, A/B testing for marketing campaigns, and sales forecasting.
- **Finance:** Risk assessment, portfolio analysis, and stock price trend evaluation.
- **Operations:** Quality control using control charts, inventory management, and ongoing process improvement.
- **Human Resources:** Employee performance analysis and salary benchmarking.

Economics Applications

- **Macroeconomics:** GDP growth analysis, inflation tracking, and unemployment rate estimation.
- **Microeconomics:** Demand and supply estimation alongside price elasticity analysis.
- **Policy Analysis:** Evaluating the economic impacts of tax changes and minimum wage adjustments.
- **Econometrics:** Employing regression models to empirically test economic theories, such as the relationship between income and consumption.

Common Statistical Techniques Across Both Fields

- **Descriptive Statistics:** Utilizing metrics like the mean and variance to summarize and interpret baseline data.
- **Hypothesis Testing:** Testing specific assumptions, such as evaluating whether a new pricing strategy leads to a measurable increase in sales.
- **Regression Analysis:** Modeling relationships to predict key variables like demand, sales volume, or pricing.
- **Time Series Analysis:** Analyzing historical trends over time to forecast future metrics, such as revenue streams or stock prices.
- **Probability Distributions:** Quantifying uncertainty to model operational defect rates and financial risks.

1.4 Data

Data are the facts and figures collected, analyzed, and summarized for presentation and interpretation. All the data collected in a particular study are referred to as the data set for the study.

?@tbl-tab1__1 shows a data set containing information for 10 restaurant.

Table 1.1: Summary of 10 sampled local dining establishments

Restaurant Name	Cuisine Type	Average Meal Price (\$)	Customer Satisfaction (%)	Delivery Fee (\$)	Quality Rating
Bella Italia	Italian	22.50	88.5	3.50	4-Star
Taco Fiesta	Mexican	11.20	79.0	1.99	3-Star
Dragon Wok	Asian	14.80	92.4	2.50	5-Star
Burger Barn	American	13.50	81.0	2.00	3-Star
Green Leaf Cafe	Vegetarian	16.00	85.2	3.00	4-Star
Tokyo Ramen	Asian	15.75	94.0	2.50	5-Star
Mama Mia Pasta	Italian	24.00	76.8	4.00	3-Star
The Daily Bagel	Bakery	8.50	83.5	1.50	4-Star
Sizzling Steakhouse	American	32.00	90.1	4.50	5-Star
Curry Express	Indian	15.00	86.4	2.75	4-Star

1.5 Elements, Variables, and Observations

Elements are the entities on which data are collected.

For the data set in `?@tbl-tab1_1`, each individual restaurant is an element: the element names appear in the first column. With 10 restaurants, the data set contains 10 elements.

A **variable** is a *characteristic* of interest for the elements. The data set includes the following five variables:

- **Cuisine Type:** The culinary category or style of food served by the restaurant.
- **Average Meal Price (\$):** The average cost of a standard meal per customer in US dollars.
- **Customer Satisfaction (%):** The percentage score reflecting positive feedback from surveyed diners.
- **Delivery Fee (\$):** The flat fee charged for delivering an order to the customer.
- **Quality Rating:** The overall customer or inspection rating assigned to the establishment, ranging from 1-Star to 5-Stars.

An **observation** is the set of measurements obtained for a particular element.

The set of measurements obtained for a particular element is called an observation.

- Referring to `?@tbl-tab1_1`, we see that the set of measurements for the first observation (Bella Italia) is Italian, 22.50, 88.5, 3.50, and 4-Star.

1.6 Scales of Measurement

Data collection requires one of the following scales of measurement: *nominal*, *ordinal*, *interval*, or *ratio*.

- When the data for a variable consist of labels or names used to identify an attribute of the element, the scale of measurement is considered a *nominal* scale (**Example: Cuisine Type**).
- The scale of measurement for a variable is called an *ordinal* scale if the data exhibit the properties of nominal data and the order or rank of the data is meaningful (**Example: Quality Rating**).
- The scale of measurement for a variable is an *interval* scale if the data have all the properties of ordinal data and the interval between values is expressed in terms of a fixed unit of measure. Interval data are always numeric (**Example: Temperature**).
- The scale of measurement for a variable is a *ratio* scale if the data have all the properties of interval data and the ratio of two values is meaningful (**Example: distance, height, weight,time etc.**).

This scale requires that a zero value be included to indicate that nothing exists for the variable at the zero point.

1.7 Quantitative and Categorical and Data

Data can be classified as either *quantitative* or *categorical*.

Quantitative Data (Numerical Data)

- Data that represents numerical values.
- **Example:** Heights of people, temperatures, test scores.
- Subtypes:
 - **Discrete Data:** *Countable* values (e.g., number of students in a class).
 - **Continuous Data:** *Measurable* values that can take **any value** within a range (e.g., weight, time).

Qualitative Data (Categorical Data)

- Data that represents categories or labels.
- Example: Colors of cars, types of animals, survey responses (e.g., yes/no).
- Subtypes:
 - **Nominal Data:** Categories without a natural order (e.g., gender, blood type).
 - **Ordinal Data:** Categories with a meaningful order (e.g., rankings, education levels).

The statistical analysis appropriate for a particular variable depends upon whether the variable is categorical or quantitative.

1.8 Cross-Sectional and Time Series Data

For purposes of statistical analysis, distinguishing between cross-sectional data and time series data is important.

Cross-sectional data are data collected at the **same or approximately the same point in time**. The data in `?@tbl-tab1_1` are cross-sectional because they describe the five variables for the 25 mutual funds at the same point in time.

Time series data are data collected **over several time periods**. For example, the time series in Figure 1.1 shows the GDP growth (%) of Bangladesh 1990 and 2024.




Figure 1.1: Bangladesh GDP Growth Rate (1990–2024)

1.9 Data vs. Information

Data: The raw, unorganized facts, symbols, or measurements collected from a source. On its own, data is “context-poor” and typically lacks meaning.

- **Examples:** Individual test scores, a list of daily temperatures, or raw survey responses.

Information: Data that has been processed, organized, or structured to make it meaningful and useful. It provides context and answers specific questions.

- **Examples:** The average test score of a class, a trend line showing rising temperatures, or a summary report of survey findings.

1.10 Exercise

1. What is the **level of measurement / categorical (nominal, ordinal) or quantitative (discrete, continuous)** for each of the following variables?
 - a. Student IQ ratings.
 - b. Distance students travel to class.
 - c. The jersey numbers of a sorority soccer team.
 - d. A classification of students by state of birth.
 - e. A summary of students by academic class—that is, freshman, sophomore, junior, and senior.
 - f. Number of hours students study per week.
2. What is the **level of measurement / categorical (nominal, ordinal) or quantitative (discrete, continuous)** for these items related to the newspaper business?
 - a. The number of papers sold each Sunday during 2011.
 - b. The departments, such as editorial, advertising, sports, etc.
 - c. A summary of the number of papers sold by county.
 - d. The number of years with the paper for each employee.
3. What is the **level of measurement / categorical (nominal, ordinal) or quantitative (discrete, continuous)** for these following items?
 - a. Salary
 - b. Gender
 - c. Sales volume of MP3 players
 - d. Soft drink preference
 - e. Temperature
 - f. SAT scores
 - g. Student rank in class
 - h. Rating of a finance professor
 - i. Number of home computers
4. For each of the following, determine whether the group is a sample or a population.
 - a. The participants in a study of a new cholesterol drug.
 - b. The drivers who received a speeding ticket in Kansas City last month.
 - c. Those on welfare in Cook County (Chicago), Illinois.
 - d. The 30 stocks reported as a part of the Dow Jones Industrial Average.

2 Descriptive statistic: Tabular and Graphical Presentations

2.1 Summarizing Categorical Data

2.1.1 Frequency Distribution

A frequency distribution is a tabular summary of data showing the number (frequency) of items in each of several non overlapping classes.

Example 2.1 Consider the following data shown in ?@tbl-tbl2__1.

Table 2.1: Data from a sample of 50 soft drink purchases

Coke Classic	Coke Classic	Coke Classic
Diet Coke	Diet Coke	Coke Classic
Pepsi	Coke Classic	Pepsi
Diet Coke	Diet Coke	Dr. Pepper
Coke Classic	Coke Classic	Coke Classic
Coke Classic	Sprite	Diet Coke
Dr. Pepper	Pepsi	Pepsi
Diet Coke	Coke Classic	Pepsi
Pepsi	Coke Classic	Pepsi
Pepsi	Coke Classic	Pepsi
Coke Classic	Pepsi	Coke Classic
Dr. Pepper	Coke Classic	Dr. Pepper
Sprite	Sprite	Pepsi
Coke Classic	Dr. Pepper	Sprite
Diet Coke	Pepsi	Coke Classic
Coke Classic	Diet Coke	Sprite
Coke Classic	Pepsi	

Now we will construct a frequency distribution by simply counting each type of soft-drink.

Table 2.2: Frequency distribution of Soft Drink Purchases

Soft Drink	Frequency
Coke Classic	19
Diet Coke	8
Dr. Pepper	5
Pepsi	13

Table 2.2: Frequency distribution of Soft Drink Purchases

Soft Drink	Frequency
Sprite	5

Relative Frequency and Percent Frequency Distributions

- Relative Frequency = $\frac{\text{Frequency of the class}}{n}$
- The *percent frequency* of a class is the relative frequency multiplied by 100.

2.1.2 Bar Charts and Pie Charts

- **Bar chart:** A graphical device for depicting qualitative data that have been summarized in a frequency, relative frequency, or percent frequency distribution.
- **Pie chart:** A graphical device for presenting data summaries based on subdivision of a circle into sectors that correspond to the relative frequency for each class.

From the frequency table of soft drinks purchase, we will develop relative and percent frequency distribution (see Table 2.3) and will construct a **bar-chart** and **pie-chart**.

Table 2.3: Frequency, Relative And Percent Frequency Distributions Of Soft Drink Purchases

Soft Drink	Frequency (f)	Relative Frequency(Rf)	Percent Frequency (Pf)
Coke Classic	19	0.38	38
Diet Coke	8	0.16	16
Dr. Pepper	5	0.10	10
Pepsi	13	0.26	26
Sprite	5	0.10	10

Now we construct a bar chart and pie chart.

2.1.3 Cross-tabulation and its graphical presentation

Cross-tabulation (also known as crosstabs or contingency tables) is a fundamental data analysis technique used to examine the relationship between **two or more categorical variables**. For instance, consider the following cross-table between the **gender** and **fitness-level** of 30 individuals:

Table 2.4: Cross-Tabulation of Gender by Fitness Level

Gender	Excellent	Fair	Good	Poor	Grand Total
Male	2	7	3	3	15
Female	5	3	5	2	15
Grand Total	7	10	8	5	30




Figure 2.1: Bar chart of Soft drink purchases




Figure 2.2: Pie chart of Soft drink purchases

We can show the information of cross-tab using (a) Clustered column chart or (b) Stacked column chart (see Figure 2.3).




Figure 2.3: Visualization of relationship between two categorical variables

2.2 Summarizing Quantitative Data

2.2.1 Frequency Distribution of quantitative data

Consider the following data.

YEAR-END AUDIT TIMES (IN DAYS): 12, 14, 19, 18, 15, 15, 18, 17, 20, 27, 22, 23, 22, 21, 33, 28, 14, 18, 16, 13,

To construct a frequency distribution we have to

1. Determine the *number of non overlapping classes(k)*.

2. Determine the *width* of each class.
3. Determine the *class limits*.

2.2.2 Frequency Distribution of Audit time data

Here, $n = 20$, Smallest value=12, Largest value=33.

1. Determine number of classes, k as : $k = \sqrt{n} = \sqrt{20} = 4.47 \approx 5$. So 5 is the number of *classes*.
2. Class width w as: $w = \frac{\text{Largest}-\text{Smallest}}{k} = \frac{33-12}{5} = 4.2 \approx 5$
3. Class limits: Start from near *smallest* value (12) say from 10 we have the following classes (exclusive method-where upper bound of the class is excluded):

[10,15), [15,20), [20,25), [25,30), and [30,35)

Now count the data values in corresponding classes and thus we have the *frequency distribution*. Once we have the frequency distribution then we also can produce the *relative* and *percent frequency distribution* (Table 2.5).

Table 2.5: Frequency, relative frequency (rf) and percent frequency (pf) distribution for the audit time data (n=20)

Audit Time (days)	Frequency (f)	Relative frequency (rf)	Percent frequency(pf)
[10,15)	4	0.20	20
[15,20)	8	0.40	40
[20,25)	5	0.25	25
[25,30)	2	0.10	10
[30,35)	1	0.05	5

2.2.3 Histogram

A common graphical presentation of quantitative data is a *histogram*. This graphical summary can be prepared for data previously summarized in either a *frequency*, *relative frequency*, or *percent frequency* distribution.

Important Note: A **relative frequency/ percent frequency histogram** is ideal for comparing distributions across groups of different sizes, as it displays proportions instead of raw counts, allowing fair and meaningful comparisons.

Illustration (see Figure 2.5)

A dataset of **Marks** (out of 100) was collected from two student groups:

- **70 Female students**
- **40 Male students**




Figure 2.4: Histogram for the Audit Time data

The goal is to compare the distribution of marks between these two groups.

The **frequency histogram** shows how many students fall into each marks range (bin). However, since the number of female students (70) is greater than male students (40), their bars are naturally taller — even if the relative performance is similar. This makes direct comparison **unfair and misleading**.

A **relative frequency histogram** shows the **proportion** of students in each bin **within each group**. By dividing counts by the total number in the group:

- It **normalizes** the data,
- Allows for **fair comparisons** between groups of different sizes,
- Highlights **true differences in distribution**, not just differences in group size.

Example Observation

From the RF histogram:

- Around **40% of males** scored between 50–60.
- Around **39% of females** scored between 50–60.

This comparison is **valid** only because the histograms show **relative frequency**, not raw counts.

2.2.4 HISTOGRAM and shape of the distribution

See Figure 2.6 .



Figure 2.5: Comparison between Frequency histogram and relative frequency histogram




Figure 2.6: Histograms Showing Differing Levels of Skewness

2.2.5 Cumulative Distributions and Ogive

A variation of the frequency distribution that provides another tabular summary of quantitative data is the *cumulative frequency* distribution. Table 2.6 shows the **cumulative relative frequency** of Audit Time data.

Table 2.6: Frequency, relative frequency and Cumulative relative frequency distribution of Audit Time Data

Audit Time (days)	Frequency (f)	Relative frequency (rf)	Cumulative relative frequency(crf)
[10,15)	4	0.20	0.20
[15,20)	8	0.40	0.60
[20,25)	5	0.25	0.85
[25,30)	2	0.10	0.95
[30,35)	1	0.05	1.00

2.2.6 Ogive

Another way of presenting this information is the ogive, which is a graphical representation of the cumulative relative frequencies. Figure 2.7 is the drawn ogive for the cumulative relative frequency for Audit time data.




Figure 2.7: Ogive for Audit Time data

For instance, from both Table 2.6 and Figure 2.7 we can say (estimate) that 60% of audits took **less than 20 days** . Similarly, 95% of audits took **less than 30 days** and so on.

2.2.7 The Stem-and-Leaf Display

The techniques of **exploratory data analysis** consist of simple arithmetic and easy-to-draw graphs that can be used to summarize data quickly. One technique—referred to as a **stem-and-leaf display**—can be used to show both the rank order and shape of a data set simultaneously (Anderson and Sweeney 2011).

Steps to Construct a Stem-and-Leaf Diagram

- (1) Divide each number into two parts: a **stem**, consisting of one or more of the leading digits, and a **leaf**, consisting of the remaining digit.
- (2) List the stem values in a vertical column.
- (3) Record the leaf for each observation beside its stem.
- (4) Write the units for stems and leaves on the display.

Example 2.2 Here are the number of questions answered correctly on an aptitude test given to 50 individuals recently interviewed for a position at Haskens Manufacturing.

2.3 Data

112, 72, 69, 97, 107, 73, 92, 76, 86, 73, 126, 128, 118, 127, 124, 82, 104, 132, 134, 83, 92, 108, 96, 100, 92, 115, 76, 91, 102, 81, 95, 141, 81, 80, 106, 84, 119, 113, 98, 75, 68, 98, 115, 106, 95, 100, 85, 94, 106, 119

2.4 Stem-and-leaf display

The decimal point is 1 digit(s) to the right of the |

```
6 | 89
7 | 233566
8 | 01123456
9 | 12224556788
10 | 002466678
11 | 2355899
12 | 4678
13 | 24
14 | 1
```

Exception

In some data sets, providing more classes or stems may be desirable. One way to do this would be to modify the original stems as follows: For example, divide stem 5 into two new stems, 5L and 5U. Stem 5L has leaves 0, 1, 2, 3, and 4, and stem 5U has leaves 5, 6, 7, 8, and 9. This will double the

number of original stems. However, there may be various type of data in practical situations. So, we have to figure out the suitable stem-and-leaf plot.

Example 2.3: Construct a stem-and-leaf plot from the following data:

88.5, 98.8, 89.6, 92.2, 92.7, 88.4, 87.5, 90.9, 94.7, 88.3, 90.4, 83.4, 87.9, 92.6, 87.8, 89.9, 84.3, 90.4, 91.6, 91.0

2.5 Solution-I

The decimal point is 1 digit(s) to the right of the |

```
8 | 34
8 | 888889
9 | 0000112233
9 | 59
```

2.6 Solution-II

The decimal point is at the |

```
82 | 4
84 | 3
86 | 589
88 | 34569
90 | 44906
92 | 267
94 | 7
96 |
98 | 8
```

Example 2.4 (Another example): Construct a stem-and-leaf plot from the following data:

7,8,2,1,8,3,5,7,1,2,2,5,8,5,5,7,8,7,5,3

Solution:

The decimal point is at the |

```
1 | 00
2 | 000
3 | 00
4 |
5 | 00000
6 |
```

7 | 0000
8 | 0000

2.7 Exercises

2.1 A doctor's office staff studied the waiting times for patients who arrive at the office with a request for emergency service. The following data with waiting times in minutes were collected over a one-month period.

2, 5, 10, 12, 4, 4, 5, 17, 11, 8, 9, 8, 12, 21, 6, 8, 7, 13, 18, 3

Use class interval/width of 5 in the following (start your class limit from 0):

- Show the frequency distribution.
- Show the relative frequency distribution.
- Show the cumulative frequency distribution.
- Show the cumulative relative frequency distribution.
- What proportion of patients needing emergency service wait less than 10 minutes or less?

2.2 A shortage of candidates has required school districts to pay higher salaries and offer extras to attract and retain school district superintendents. The following data show the annual base salary (\$1000s) for superintendents in 20 districts in the greater Rochester, New York, area (The Rochester Democrat and Chronicle, February 10, 2008).

187, 184, 174, 185, 175, 172, 202, 197, 165, 208, 215, 164, 162, 172, 182, 156, 172, 175, 170, 183


Use appropriate number classes/ class width in the following.

- Show the frequency distribution.
- Show the percent frequency distribution.
- Show the cumulative percent frequency distribution.
- Develop a histogram for the annual base salary.
- Do the data appear to be skewed? Explain.
- Which salary range belongs to the highest percentage of superintendents ?

2.8 Data

187, 184, 174, 185, 175, 172, 202, 197, 165, 208, 215, 164, 162, 172, 182, 156, 172, 175, 170, 183

2.9 Histogram



2.3 NRF/BIG research provided results of a consumer holiday spending survey (USA Today, December 20, 2005). The following data provide the dollar amount of holiday spending for a sample of 25 consumers.

1200, 850, 740, 590, 340, 450, 890, 260, 610, 350, 1780, 180, 850, 2050, 770, 800, 1090, 510, 520, 220, 1450, 280, 1120, 200, 350


- What is the lowest holiday spending? The highest?
- Use a class width of \$250 to prepare a frequency distribution and a percent frequency distribution for the data.
- Prepare a histogram and comment on the shape of the distribution.
- What observations can you make about holiday spending?

2.10 Data

1200, 850, 740, 590, 340, 450, 890, 260, 610, 350, 1780, 180, 850, 2050, 770, 800, 1090, 510, 520, 220, 1450, 280, 1120, 200, 350

2.11 Histogram

Frequency histogram of holiday spending (USD)



2.4 Construct a stem-and-leaf display for the following data.

70, 72, 75, 64, 58, 83, 80, 82, 76, 75, 68, 65, 57, 78, 85, 72

2.5 Construct a stem-and-leaf display for the following data.

11.3, 9.6, 10.4, 7.5, 8.3, 10.5, 10.0, 9.3, 8.1, 7.7, 7.5, 8.4, 6.3, 8.8

2.6 A psychologist developed a new test of adult intelligence. The test was administered to 20 individuals, and the following data were obtained.

114, 99, 131, 124, 117, 102, 106, 127, 119, 115, 98, 104, 144, 151, 132, 106, 125, 122, 118, 118

Construct a stem-and-leaf display for the data.

2.12 Line Chart

2.13 Scatter diagram

2.14 Case Study: Lifestyle Indicators and Preferences

You're given data from a cross-sectional survey of 30 individuals. Variables include Age, Gender, Monthly Income, Hours of Exercise/Week, Fitness Level, Favorite Fruit and Temperature preference. This data-set enables practice with data visualization, scale classification, and relationship analysis using tools like bar charts, scatter plots, and histograms.

ID	Age	Gender	Monthly Income (USD)	Hours of Exer- cise/Week	Favorite Fruit	Fitness Level	Temper- ature Prefer- ence (°C)
1	25	Male	1520.75	2.5	Mango	Fair	23
2	32	Female	2280.50	1.2	Apple	Poor	22
3	20	Male	925.10	4.8	Banana	Good	19
4	29	Female	1980.90	3.3	Orange	Fair	22
5	40	Male	3055.45	0.7	Apple	Poor	21
6	23	Female	1425.00	5.9	Mango	Excellent	20
7	37	Male	3625.80	2.1	Banana	Fair	18
8	31	Female	2520.00	1.5	Apple	Poor	27
9	27	Male	1685.25	3.7	Orange	Good	22
10	22	Female	1190.00	6.2	Mango	Excellent	19
11	34	Male	3180.45	0.0	Banana	Poor	24
12	26	Female	1830.60	4.3	Orange	Good	19
13	39	Male	3999.00	1.4	Apple	Fair	22
14	24	Female	1335.75	5.6	Banana	Excellent	26
15	28	Male	2125.35	2.2	Mango	Fair	27
16	33	Female	2885.50	3.1	Orange	Good	22
17	21	Male	960.00	6.0	Apple	Excellent	18
18	30	Female	2190.00	1.7	Banana	Fair	27
19	36	Male	3425.90	2.3	Mango	Fair	27
20	29	Female	1920.40	4.5	Orange	Good	20
21	35	Male	3095.25	3.0	Apple	Good	23
22	22	Female	1125.00	5.1	Banana	Excellent	27
23	38	Male	4180.60	1.8	Mango	Fair	22
24	30	Female	2590.00	2.4	Orange	Fair	20
25	41	Male	3540.85	0.9	Apple	Poor	22
26	28	Female	1875.50	4.0	Banana	Good	24
27	24	Male	1275.00	6.7	Mango	Excellent	21
28	31	Female	2390.25	3.5	Orange	Good	25
29	36	Male	2965.80	2.6	Banana	Fair	22
30	21	Female	985.00	5.4	Apple	Excellent	25

Tasks

- 1) **Identify** the *scale of measurement* of each variable.
- 2) **Construct** suitable graph/chart for variables like *gender*, *fitness level* etc.
- 3) **Construct** relative frequency histogram for variables like age, Monthly Income (USD), etc. **Describe** what you have learned.
- 4) **Plot** Hours of Exercise/Week vs Temperature Preference (°C). What is your conclusion.
- 5) **Draw** the scatter plot of Monthly Income vs Hours of Exercise/Week. Is there any relation?
- 6) **Cross-tabulation**

- a) Make a cross-tabulation between Gender and Favorite Fruit. Show the result in a stacked bar-chart.
- b) Make a cross-tabulation between Favorite Fruit and Fitness Level. Show the result in a stacked bar-chart.

3 Descriptive statistics: Numerical Measures

Numerical measures of location, dispersion, shape, and association are introduced. If the measures are computed for data from a sample, they are called **sample statistics**. If the measures are computed for data from a population, they are called **population parameters**. In statistical inference, a sample statistic is referred to as the **point estimator** of the corresponding population parameter (Anderson and Sweeney 2011).

3.1 Measures of location

In statistics, measures of location, also known as measures of central tendency, are used to describe the central value or position of a distribution. They provide information about where the “center” of the distribution lies. Common measures of location include:

a) Mean b) Median c) Mode d) Percentiles e) Quartiles

3.1.1 Mean

- **Sample mean:** Suppose n observation of a variable X is drawn from a population. Then the sample mean is denoted by $\bar{x}$ and

$$\bar{x} = \frac{\sum x}{n}$$

The sample mean $\bar{x}$ is a sample statistic.

- **Population mean:** Suppose in a population there are N values of variable X . Then the population mean is denoted by μ and

$$\mu = \frac{\sum x}{N}$$

The $\bar{x}$ is a point estimator of the population mean μ .

3.1.2 Weighted mean

The weighted mean is a special case of the arithmetic mean. It occurs when there are several observations of the same value.

- **Weighted mean:**

$$\bar{x} = \frac{\sum w_i x_i}{\sum w_i}$$

where, w_i = weight for observation i

Example 3.1 (Lind, Marchal, and Wathen 2012): The Carter Construction Company pays its hourly employees \$16.50, \$19.00, or \$25.00 per hour. There are 26 hourly employees, 14 of which are paid at the \$16.50 rate, 10 at the \$19.00 rate, and 2 at the \$25.00 rate. What is the mean hourly rate paid the 26 employees?

Solution:

Hourly wage (\$), x_i	Weight (w_i)	$w_i x_i$
16.50	14	231
19.00	10	190
25.00	2	50

Here, $\sum w_i x_i = 471$ and $\sum w_i = 26$

Hence, $\bar{x} = \frac{\sum w_i x_i}{\sum w_i} = \frac{471}{26} = 18.1154$

So, the weighted mean hourly wage is rounded to \$18.12.

Example 3.2 (Anderson 2020a) : The grade point average for college students is based on a weighted mean computation. For most colleges, the grades are given the following data values: A (4), B (3), C (2), D (1), and F (0). After 60 credit hours of course work, a student at State University earned 9 credit hours of A, 15 credit hours of B, 33 credit hours of C, and 3 credit hours of D.

- Compute the student's grade point average.
- Students at State University must maintain a 2.5 grade point average for their first 60 credit hours of course work in order to be admitted to the business college. Will this student be admitted?

Example 3.3 (Lind, Marchal, and Wathen 2012): Springers sold 95 Antonelli men's suits for the regular price of \$400. For the spring sale, the suits were reduced to \$200 and 126 were sold. At the final clearance, the price was reduced to \$100 and the remaining 79 suits were sold.

- What was the weighted mean price of an Antonelli suit?
- Springers paid \$200 a suit for the 300 suits. Comment on the store's profit per suit if a salesperson receives a \$25 commission for each one sold.

Ans: (a) \$237 (b) \$12

3.1.3 Median

The **median** is another measure of central location. The median is the value in the middle when the data are arranged in ascending order (smallest value to largest value).

- For an *odd* number of observations, median is the middle value
- For an *even* number of observations, median is the average of the two middle values

Example 3.4 (n is odd): Let us consider the following class size data for a sample of five college classes.

46, 54, 42, 46, 32

Arranged data: 32, 42, 46, 46, 54.

Because $n = 5$ is odd, the median is the middle value. Thus the median class size is **46** students.

Example 3.5 (n is even): Let us consider the following class size data for a sample of five college classes.

46, 54, 42, 46, 32, 40

Arranged data: 32, 40, 42, 46, 46, 54.

Because $n = 6$ is even, the

$$\text{Median} = \frac{42 + 46}{2} = 44$$

i Note

Although the mean is the more commonly used measure of central location, in some situations the median is preferred. For example, the median is the measure of location most often reported for annual income and property value data because a few extremely large incomes or property values can inflate the mean. In such cases, the median is the preferred measure of central location.

3.1.4 Mode

The **mode** is the value that occurs with greatest frequency.

i Note

Situations can arise for which the greatest frequency occurs at two or more different values. In these instances, more than one mode exists. If the data contain exactly two modes, we say that the data are **bimodal**. If data contain more than two modes, we say that the data are multimodal. In **multimodal** cases the mode is almost never reported because listing three or more modes would not be particularly helpful in describing a location for the data.

3.1.5 Percentiles

A **percentile** provides information about how the data are spread over the interval from the smallest value to the largest value.

- The p^{th} percentile is a value such that *at least* p percent of the observations are less than or equal to this value and *at least* $(100 - p)$ percent of the observations are greater than or equal to this value.
- **Formula:**

$$p^{th} \text{ percentile} = (p \times \frac{n+1}{100})^{th} \text{ value}$$

Example 3.6: Here is the monthly starting salary (\$) of 12 graduates:

3450 ,3550 ,3650 ,3480 ,3355, 3310 ,3490 ,3730, 3540 ,3925, 3520 ,3480

Let us determine the 85th percentile for the starting salary data.

Solution:

Arranged data: 3310, 3355, 3450, 3480, 3480, 3490, 3520, 3540, 3550, 3650, 3730, 3925

Now,

$$\begin{aligned} P_{85} &= (85 \times \frac{12+1}{100})^{th} \text{ value} \\ &= 11.05^{th} \text{ value} = 11^{th} \text{ value} + 0.05(12^{th} - 11^{th}) \\ &= 3730 + 0.05(3925 - 3730) \\ &= 3739.75 \text{ dollars} \end{aligned}$$

Interpretation: Here, 85th percentile = 3739.75 implies that at least 85% of the total observations (salaries) are less or equal to 3739.75 dollars.

3.1.6 Quartiles

It is often desirable to divide data into four parts, with each part containing approximately one-fourth, or 25% of the observations. The division points are referred to as the quartiles and are defined as

Q_1 = first quartile, or 25th percentile

Q_2 = second quartile, or 50th percentile (also the median)

Q_3 = third quartile, or 75th percentile.

Example 3.7: Here is the monthly starting salary (\$) of 12 graduates:

3450, 3550, 3650, 3480, 3355, 3310, 3490, 3730, 3540, 3925, 3520, 3480

Compute Q_1 and Q_3 of the above data (*Will be solved in class*).

3.1.7 Geometric mean

The **geometric mean** is useful in finding the average change of percentages, ratios, indexes, or growth rates over time. It has a wide application in business and economics because we are often interested in finding the percentage changes in sales, salaries, or economic figures, such as the Gross Domestic Product, which compound or build on each other.

- **Geometric mean (GM):** GM is the n^{th} root of the product of n values.

$$GM = \sqrt[n]{(x_1)(x_2) \cdots (x_n)} = [(x_1)(x_2) \cdots (x_n)]^{1/n}$$

Example 3.8: Compute the geometric mean of the following percent increases: 8, 12, 14, 26, and 5.

Solution: Here, $n = 5$. The geometric mean is:

$$GM = [8 \cdot 12 \cdot 14 \cdot 26 \cdot 5]^{1/5} = [174720]^{1/5} \approx 11.18$$

Exercise 3.9 (Lind, Marchal, and Wathen 2012): The percent increase in sales for the last 4 years at Combs Cosmetics were: 4.91, 5.75, 8.12, and 21.60.

- Find the geometric mean percent increase.
- Find the arithmetic mean percent increase.
- Is the arithmetic mean equal to or greater than the geometric mean?

Example 3.10: Listed below is the percent increase in sales for the MG Corporation over the last 5 years. **Determine** the geometric mean percent increase in sales over the period.

9.4, 13.8, 11.7, 11.9, 14.7

3.2 Measures of variability

Variability in data means lack of uniformity. It is also referred to as spread, scatter, or dispersion. We turn now to a discussion of some commonly used measures of variability.

3.2.1 Range

Range = Largest value – Smallest value

- The simplest one, but is highly influenced by extreme values.

3.2.2 Interquartile Range (IQR)

$$IQR = Q_3 - Q_1$$

- The interquartile range is the range for the middle 50% of the data.

3.2.3 Variance

The variance is a measure of variability that utilizes all the data. The variance is based on the difference between the value of each observation (x_i) and the mean. The difference between each x_i and the mean ($\bar{x}$ for a sample, μ for a population) is called a deviation about the mean.

- **Population variance**

$$\sigma^2 = \frac{(x_1 - \mu)^2 + (x_2 - \mu)^2 + \cdots + (x_N - \mu)^2}{N} = \frac{\sum (x_i - \mu)^2}{N}$$

- **Sample variance**

$$s^2 = \frac{(x_1 - \bar{x})^2 + (x_2 - \bar{x})^2 + \cdots + (x_n - \bar{x})^2}{n - 1} = \frac{\sum (x_i - \bar{x})^2}{n - 1}$$

The sample variance s^2 is the estimator of the population variance σ^2 .

- An alternative formula for the computation of the sample variance is:

$$s^2 = \frac{\sum x_i^2 - n\bar{x}^2}{n - 1}$$

where, $\sum x_i^2 = x_1^2 + x_2^2 + \dots + x_n^2$

i Derivation

Since, $\bar{x} = \frac{\sum x}{n}$ so, $\sum x = n \cdot \bar{x}$
Now, $\sum_{i=1}^n (x_i - \bar{x})^2$
 $= \sum_{i=1}^n (x_i^2 - 2 \cdot x_i \cdot \bar{x} + \bar{x}^2)$

$$\begin{aligned}
&= \sum_{i=1}^n x_i^2 - 2\bar{x} \sum_{i=1}^n x_i + \sum_{i=1}^n \bar{x}^2 \\
&= \sum_{i=1}^n x_i^2 - 2\bar{x} \cdot n\bar{x} + n\bar{x}^2 \\
&= \sum_{i=1}^n x_i^2 - 2n\bar{x}^2 + n\bar{x}^2 = \sum_{i=1}^n x_i^2 - n\bar{x}^2
\end{aligned}$$

Example 3.11: Here is the monthly starting salary (\$) of 6 graduates:

3450 ,3550 ,3650 ,3480 ,3355, 3545.

Compute sample variance (s^2).

Solution:

Here, **sample size**, $n = 6$.

The sample mean salary,

$$\bar{x} = \frac{\sum x}{n} = \frac{3450 + \dots + 3545}{6} = 3505 \text{ dollars}$$

Table 3.2: Computation of the sample variance for the starting Salary data

Salary (x_i)	Sample mean, $\bar{x}$	$(x_i - \bar{x})$	$(x_i - \bar{x})^2$
3450	3505	-55	3025
3550	3505	45	2025
3650	3505	145	21025
3480	3505	-25	625
3355	3505	-150	22500
3545	3505	40	1600
		$\sum(x_i - \bar{x}) = 0$	$\sum(x_i - \bar{x})^2 = 50800$

Hence, the sample variance is:

$$s^2 = \frac{\sum(x_i - \bar{x})^2}{n - 1} = \frac{50800}{6 - 1} = 10160 \text{ (dollars)}^2$$

3.2.4 Standard deviation

The **standard deviation** is defined to be the positive square root of the variance

- **Sample standard deviation** $= s = \sqrt{s^2}$
- **Population standard deviation** $= \sigma = \sqrt{\sigma^2}$

The sample standard deviation s is the estimator of population standard deviation σ .

Example 3.12: The standard deviation of the previous example is :

$$s = \sqrt{10160} = 100.7968 \approx 100.80 \text{ dollars}$$

i Note:

The standard deviation is easier to interpret than the variance because the standard deviation is measured in the same units as the data.

For example, the sample variance for the starting salary data of business school graduates is $s^2 = 10160$ (dollars)².

Because the standard deviation is the square root of the variance, the units of the variance, dollars squared, are converted to dollars in the standard deviation.

Thus, the standard deviation of the starting salary data is 100.80 dollar. In other words, the standard deviation is measured in the same units as the original data. For this reason the standard deviation is more easily compared to the mean and other statistics that are measured in the same units as the original data.

i Properties of variance

1. **Non-negativity:** $Var(X) \geq 0$.
2. For any constant say $X = c$, $Var(c) = 0$.
3. Variance is affected by outliers.
4. Variance is NOT affected by **change origin**; but affected by **change of scale** that is:

$$Var(aX + b) = a^2 Var(X)$$

Here, a and b are both constants.

Proof: For a population data $X = \{x_1, x_2, \dots, x_N\}$ the population mean of X is

$$\mu_X = \frac{\sum_{i=1}^N x_i}{N} \text{ and variance of } X \text{ is}$$

$$Var(X) = \frac{\sum_{i=1}^N (x_i - \mu_X)^2}{N}$$

Now let, $Y = aX + b$

So, the population mean of Y is

$$\begin{aligned} \mu_Y &= \frac{\sum_{i=1}^N y_i}{N} = \frac{\sum_{i=1}^N (ax_i + b)}{N} \\ &= \frac{\sum_{i=1}^N (ax_i) + \sum_{i=1}^N b}{N} = a \frac{\sum_{i=1}^N x_i}{N} + \frac{Nb}{N} = a \cdot \mu_X + b \end{aligned}$$

Hence,

$$\begin{aligned} Var(Y) &= \frac{\sum_{i=1}^N (y_i - \mu_Y)^2}{N} = \frac{\sum_{i=1}^N (a \cdot x_i + b - a \cdot \mu_X - b)^2}{N} \\ &= \frac{\sum_{i=1}^N (a \cdot x_i - a \cdot \mu_X)^2}{N} \\ &= a^2 \frac{\sum_{i=1}^N (x_i - \mu_X)^2}{N} = a^2 Var(X) \end{aligned}$$

$$\therefore Var(Y) = Var(aX + b) = a^2 Var(X).$$

3.2.5 Coefficient of variation (CV)

In some situations we may be interested in a descriptive statistic that indicates how large the standard deviation is relative to the mean. This measure is called the **coefficient of variation** and is usually expressed as a percentage.

Coefficient variation,

$$CV = \frac{\text{Standard deviation}}{\text{Mean}}$$

- The coefficient of variation is a relative measure of variability; it measures the standard deviation relative to the mean.
- In general, the coefficient of variation is a useful statistic for comparing the variability of variables that have different standard deviations and different means.

Example 3.13: The table at the left shows the population heights (in inches) and weights (in pounds) of the members of a basketball team. Find the coefficient of variation for the heights and the weights. Then compare the results.

3.2.5.1 Data

Heights (inches)	Weights (pounds)
72	180
74	168
68	225
76	201
74	189
69	192
72	197
79	162
70	174
69	171
77	185
73	210

3.2.5.2 Coefficient of variation

The mean height $\mu = \frac{\sum x}{N} = \frac{72+74+\dots+73}{12} \approx 72.8$ inches with a standard deviation $\sigma = \sqrt{\frac{\sum x^2}{N} - \mu^2} = 3.3$ inches.

The coefficient of variation for the heights is

$$CV_{\text{height}} = \frac{\sigma}{\mu} \cdot 100\% = \frac{3.3}{72.8} \cdot 100\% \approx 4.5\%.$$

Similarly,

the mean weight $\mu \approx 187.8$ pounds with a standard deviation $\sigma = 17.7$ pounds.

The coefficient of variation for the weights is

$$CV_{weight} = \frac{\sigma}{\mu} \cdot 100\% = \frac{17.7}{187.8} \cdot 100\% \approx 9.4\%$$

Interpretation: The weights (9.4%) are more variable than the heights (4.5%).

3.3 The mean and standard deviation of Grouped data

In most cases, measures of location and variability are computed by using the individual data values. Sometimes, however, data are available only in a grouped or frequency distribution form. In the following discussion, we show how the weighted mean formula can be used to obtain approximations of the mean, variance, and standard deviation for **grouped data**.

3.3.1 Sample mean for grouped data

$$\bar{x} = \frac{\sum f_i M_i}{n}$$

where,

$$M_i = \frac{\text{Lower limit} + \text{Upper limit}}{2} = \text{the midpoint for class } i$$

f_i = the frequency for class i

n = the sample size = $\sum f_i$

3.3.2 Sample variance for grouped data

$$s^2 = \frac{\sum f_i (M_i - \bar{x})^2}{n - 1} = \frac{\sum M_i^2 f_i - n \cdot \bar{x}^2}{n - 1}$$

Eventually the **standard deviation** is $\sqrt{s^2}$

Example 3.14 The frequency distribution of audit times is given below:

Table 3.4: Frequency distribution of audit times

Audit Time (days)	Frequency
10-14	4
15-19	8
20-24	5
25-29	2
30-34	1
Total	20

Compute sample mean and standard deviation of Audit time (days) from the above frequency distribution / grouped data.

Solution:

Table 3.5: Computation of the sample mean audit time for grouped data

Audit Time (days)	Mid point(M_i)	Frequency (f_i)	$f_i M_i$	$f_i M_i^2$
10-14	12	4	48	576
15-19	17	8	136	2312
20-24	22	5	110	2420
25-29	27	2	54	1458
30-34	32	1	32	1024
Total		20	380	7790

Sample mean,

$$\bar{x} = \frac{\sum f_i M_i}{n} = \frac{380}{20} = 19 \text{ days}$$

Sample variance,

$$s^2 = \frac{\sum f_i M_i^2 - n \cdot \bar{x}^2}{n - 1} = \frac{7790 - 20 \cdot 19^2}{20 - 1} = 30 \text{ (days)}^2$$

Hence the **standard deviation** is:

$$s = \sqrt{30} \text{ days} = 5.48 \text{ days}$$

3.4 Measures of relative location: z-score

In addition to measures of location, variability, and shape, we are also interested in the relative location of values within a data set. Measures of relative location help us determine how *far a particular value* is from the **mean**.

3.4.1 z-score

The **z-score** provide how far an observation or value is from the mean or average.

z-score

Let, $X = \{x_1, x_2, \dots, x_n\}$ has the *sample mean* $\bar{x}$ and the *sample standard deviation* s . Then the **z-score** for x_i is :

$$z_i = \frac{x_i - \bar{x}}{s}$$

- The z-score is often called the *standardized value*. The z-score, z_i , can be interpreted as the *number of standard deviations* x_i is from the mean $\bar{x}$. For example, $z_1 = 1.2$ would indicate that x_1 is 1.2 standard deviations greater than the sample mean. Similarly, $z_2 = -0.5$ would indicate that x_2 is 0.5, or $1/2$, standard deviation less than the sample mean.

- A z-score greater than zero occurs for observations with a value greater than the mean, and a z-score less than zero occurs for observations with a value less than the mean. A z-score of zero indicates that the value of the observation is equal to the mean.
- The z-score for any observation can be interpreted as a measure of the relative location of the observation in a data set. Thus, observations in two different data sets with the same z-score can be said to have the same relative location in terms of being the same number of standard deviations from the mean.

Example 3.15 Suppose $X = \{46, 54, 42, 46, 32\}$. Here X is the number students in each class.

- Compute *sample mean* and *standard deviation* of X
- Compute *z-scores*
- Interpret the *z-scores* for 54 and 32
- Compute the mean and variance of *z-scores*

Example 3.16 Consider a very large number of students taking a college entrance exam such as the SAT. And suppose the mean score on the mathematics section of the SAT is 570 with a standard deviation of 40.

- Find the z-score for a student who scored 600.
- A student is told that his z-score on this test is -1.5. What was his actual SAT math score?

3.4.2 Chebyshev's Theorem

Regardless of the shape of a distribution **Chebyshev's Theorem** provides lower bound of proportion of observations lie within a certain interval.

i Note

Chebyshev's Theorem

At least $(1 - \frac{1}{z^2})$ of the data values must be within z standard deviations of the mean, where $z > 1$.

Mathematically,

$$P(\bar{x} - z \cdot s < X < \bar{x} + z \cdot s) \geq (1 - \frac{1}{z^2})$$

Example 3.17 Suppose that the midterm test scores for 100 students in a college business statistics course had a mean of 70 and a standard deviation of 5.

- How many students (in %) had test scores between 60 and 80?
- How many students (in %) had test scores between 58 and 82?

Solution:

Here, $\bar{x} = 70$; $s = 5$

i) For $x = 60$; $z = \frac{60-70}{5} = -2$

For, $x = 80$; $z = \frac{80-70}{5} = +2$

Applying Chebyshev's theorem with $z = 2$, we have

$$(1 - \frac{1}{z^2}) = (1 - \frac{1}{2^2}) = 0.75$$

So, at least 75% of the students must have test scores between 60 and 80.

ii) DIY

Example 3.18 Suppose that the midterm test scores for 100 students in a college business statistics course had a mean of 70 and a standard deviation of 5. **Find** the interval in which at least 80% data values lie.

3.4.3 Empirical Rule

If a **distribution is approximately bell-shaped/symmetric/normal** then

- Approximately 68% of the data values will be within **one** standard deviation of the mean.
- Approximately 95% of the data values will be within **two** standard deviations of the mean.
- Almost all of the data values (99%) will be within **three** standard deviations of the mean.




Figure 3.1: Empirical rule

Example GMAT Exam Scores. The Graduate Management Admission Test (GMAT) is a standardized exam used by many universities as part of the assessment for admission to graduate study in business. The average GMAT score is 547 (Magoosh website). Assume that GMAT scores are bell-shaped with a standard deviation of 100.

- What percentage of GMAT scores are 647 or higher?
 - What percentage of GMAT scores are 747 or higher?
 - What percentage of GMAT scores are between 447 and 547?
 - What percentage of GMAT scores are between 347 and 647?
- (Hints: Use empirical rule)

3.4.4 Detecting Outliers

Sometimes a data set will have one or more observations with unusually large or unusually small values. These extreme values are called **outliers**.

i Using z-score to detect outlier

An observation say x_i is treated as *outlier* if its corresponding **z-score** is *less than -3* or *greater than +3*.

Equivalently, if an observation x_i falls outside the interval $[\bar{x} - 3 \cdot s, \bar{x} + 3 \cdot s]$.


Note: Z-score rule works well for **large, symmetric, bell-shaped datasets**.

Example 3.19 Consider the following data:

35, 68, 52, 80, 45, 60, 30, 75, 48, 53, 70, 42, 66, 55, 40, 58, 78, 36, 50, 62, 33, 71, 49, 44, 64, 38, 85, 57, 39, 61, 73, 46, 67, 32, 56, 65, 41, 59, 51, 54, 10, 100, 110, 77, 63, 69, 82, 37, 47, 60

- Plot the data (histogram). Is it approximately bell-shaped/symmetric?

Solution (i):



- Now identify outlier(s) using z-score rule.

Solution (ii):

i Using $1.5(IQR)$ rule to detect outlier

We define two limits as follows:

Lower limits, $LL = Q_1 - 1.5(IQR)$

Upper limits, $UL = Q_3 + 1.5(IQR)$

Any data value or observation falls outside the interval $[LL, UL]$ will be treated as outlier.

Note: IQR rule is **better for skewed or small datasets** .

Example 3.20 The following data shows the annual salaries (in thousand USD) of 10 employees in a company:

50, 45, 52, 37, 42, 70, 40, 35, 47, 17

Identify the **outlier(s)** in the dataset using $1.5(IQR)$ rule.

3.5 Five-Number summary

In a **five-number summary**, five numbers are used to summarize the data:

1. Smallest value
2. First quartile (Q_1)
3. Median (Q_2)
4. Third quartile (Q_3)
5. Largest value

Example 3.21 (Anderson 2020a, 158): **Household Incomes**. The following data represent a sample of 14 household incomes(\$1000s). Answer the following questions based on this sample.

49.4, 52.4, 53.4, 51.3, 52.1, 48.7, 52.1, 52.2, 64.5, 51.6, 46.5, 52.9, 52.5, 51.2

- a. What is the median household income for these sample data?
- b. According to a previous survey, the median annual household income five years ago was \$55,000. Based on the sample data above, estimate the percentage change in the median household income from five years ago to today.
- c. Compute the first and third quartiles.
- d. Provide a five-number summary.
- e. Using the z-score approach, do the data contain any outliers? Does the approach that uses the values of the first and third quartiles and the interquartile range to detect outliers provide the same results?

3.6 Box-plot

The **box plot** is a graphical display that simultaneously describes several important features of a data set, such as **center**, **spread**, a **departure from symmetry**, and identification of unusual observations or **outliers**.

A key to the development of a boxplot is the computation of the interquartile range, $IQR = Q_3 - Q_1$. Figure 3.2 shows a boxplot for the monthly starting salary data. The steps used to construct the boxplot follow (Anderson 2020b).




Figure 3.2: Boxplot of the Monthly Starting Salary Data with Lines Showing the Lower and Upper Limits


Salary data (assending order)

5710, 5755, 5850, 5880, 5880, 5890, 5920, 5940, 5950, 6050, 6130, 6325

1. A box is drawn with the ends of the box located at the first and third quartiles. For the salary data, $Q_1 = 5857.5$ and $Q_3 = 6025$. This box contains the middle 50% of the data.
2. A vertical line is drawn in the box at the location of the **median** (5905 for the salary data).
3. By using the interquartile range, $IQR = Q_3 - Q_1$, *limits* are located at $1.5(IQR)$ below Q_1 and $1.5(IQR)$ above Q_3 . For the salary data, $IQR = Q_3 - Q_1 = 6025 - 5857.5 = 167.5$. Thus, the *limits* are $LL = 5857.5 - 1.5(167.5) = 5606.25$ and $UL = 6025 + 1.5(167.5) = 6276.25$. Data outside these limits are considered outliers.
4. The horizontal lines extending from each end of the box in Figure 3.2 called *whiskers*. The whiskers are drawn from the ends of the box to the smallest and largest values *inside* the *limits* computed in step 3. Thus, the whiskers end at salary values of 5710 and 6130.

Here is the computer generated boxplot of salary data using R programming language (R Core Team 2024).

Boxplot of the Monthly Starting Salary Data




3.6.1 Boxplot and skewness of the data

When we discuss the frequency histogram we also learned about shape of the distribution. By visual inspection of boxplot we can also tell about the distribution shape of a variable. The following boxplots are the typical examples of skewness of the data.


Symmetric (approximately)



Positively skewed



Negatively skewed



3.6.2 Comparative box-plot

An example: Cell Phone Companies Customer Satisfaction. Consumer Reports provides overall customer satisfaction scores for AT&T, Sprint, T-Mobile, and Verizon cell-phone services in major metropolitan areas throughout the United States. The **rating** for each service reflects the overall customer satisfaction considering a variety of factors such as cost, connectivity problems, dropped

calls, static interference, and customer support. A satisfaction scale from 0 to 100 is used with 0 indicating completely dissatisfied and 100 indicating completely satisfied. Suppose that the ratings for the four cell-phone services in 20 metropolitan areas are as shown below (Anderson 2020a).

Table 3.6: Cell Phone Companies Customer Satisfaction

Metropolitan Area	AT&T	Sprint	T-Mobile	Verizon
Atlanta	70	66	71	79
Boston	69	64	74	76
Chicago	71	65	70	77
Dallas	75	65	74	78
Denver	71	67	73	77
Detroit	73	65	77	79
Jacksonville	73	64	75	81
Las Vegas	72	68	74	81
Los Angeles	66	65	68	78
Miami	68	69	73	80
Minneapolis	68	66	75	77
Philadelphia	72	66	71	78
Phoenix	68	66	76	81
San Antonio	75	65	75	80
San Diego	69	68	72	79
San Francisco	66	69	73	75
Seattle	68	67	74	77
St. Louis	74	66	74	79
Tampa	73	63	73	79
Washington	72	68	71	76

We can easily compare the **ratings** for 4- cell-phone services using **comparative/parallel box-plots**

- Now, **discuss** what a comparison of the boxplots tells about the four services.
- Which service does *Consumer Reports* recommend as being best in terms of overall customer satisfaction?

Problem 3.22 Consider a sample with data values of 27, 25, 20, 15, 30, 34, 28, and 25. Provide the five-number summary for the data. Also construct a boxplot.

Problem 3.23 A sample of 28 time shares in the Orlando, Florida, area revealed the following daily charges (in USD dollars) for a one-bedroom suite. For convenience, the data are ordered from smallest to largest.

116, 121, 157, 192, 207, 209, 209, 229, 232, 236, 236, 239, 243, 246,
260,264, 276, 281, 283, 289, 296, 307, 309, 312, 317, 324, 341, 353

- Compute the lower and upper limits and check for outlier(s).
- Then construct a boxplot of the daily charges and show the outlier(s) if any in the boxplot.
- Comment on the distribution of the daily charges.




Figure 3.3: Comparative Boxplots of Rating by customer for cell-phone services

Problem 3.24 Amateur Golfer Scores. Scores turned in by an amateur golfer at the Bonita Fairways Golf Course in Bonita Springs, Florida, during 2017 and 2018 are as follows:

2017 Season: 74, 78, 79, 77, 75, 73, 75, 77

2018 Season: 71, 70, 75, 77, 85, 80, 71, 79

- Construct a side-by-side comparative boxplot of scores for two seasons. Show outlier(s) if any.
- Describe your findings.


Problem 3.25 Automobile Fuel Efficiencies. In automobile mileage and gasoline-consumption testing, 13 automobiles were road tested for 300 miles in both city and highway driving conditions. The following data were recorded for miles-per-gallon performance.

City: 16.2, 16.7, 15.9, 14.4, 13.2, 15.3, 16.8, 16.0, 16.1, 15.3, 15.2, 15.3, 16.2

Highway: 19.4, 20.6, 18.3, 18.6, 19.2, 17.4, 17.2, 18.6, 19.0, 21.1, 19.4, 18.5, 18.7

- Construct a side-by-side comparative boxplot of mileage between city and highway. Show outlier(s) if any.
- Describe your findings.

Answer (a):



3.7 Measures of shape: Skewness and Kurtosis

Measures of shape are tools that can be used to describe the shape of a distribution of data. In this section, we examine two measures of shape, **skewness** and **kurtosis**.

3.7.1 Skewness

Skewness refers to lack of symmetry or departure from symmetry. There are three types of skewness based on the histogram or density plot of data.

- a) **Positive skewness**/ Skewed right- where $mean > median > mode$
- b) **Symmetrical distribution**- in a perfect symmetrical distribution $mean = median = mode$
- c) **Negative skewness**/ Skewed left- where $mean < median < mode$

The typical example of skewness is exhibited in Figure 3.4 with the relative position of mean, median and mode.




Figure 3.4: Types of skewness and relative position of mean, median, mode

3.7.2 Kurtosis

Kurtosis describes the amount of peakedness of a distribution.

- Distributions that are high and thin are referred to as **leptokurtic** distributions.
- Distributions that are flat and spread out are referred to as **platykurtic** distributions.
- Between these two types are distributions that are more “normal” in shape, referred to as **mesokurtic** distributions.

These three types of kurtosis are illustrated in Figure 3.5.




Figure 3.5: Different types of kurtosis

3.7.3 Measures of skewness and kurtosis using Moments

Moments:

Suppose a sample of size n of variable X with observations $x_1, x_2, \dots, x_n$.

The r^{th} sample **raw moment** is

$$m'_r = \frac{\sum_{i=1}^n x_i^r}{n} \quad (3.1)$$

Hence the first 4 raw moments are:

$$m'_1 = \frac{\sum_{i=1}^n x_i}{n} = \bar{x}$$

$$m'_2 = \frac{\sum_{i=1}^n x_i^2}{n}$$

$$m'_3 = \frac{\sum_{i=1}^n x_i^3}{n}$$

$$m'_4 = \frac{\sum_{i=1}^n x_i^4}{n}$$

The r^{th} sample **central moment** (about mean) is:

$$m_r = \frac{\sum_{i=1}^n (x_i - \bar{x})^r}{n} \quad (3.2)$$

Hence the first 4 central moments are:

$$m_1 = \frac{\sum (x - \bar{x})}{n} = \frac{0}{n} = 0$$

$$m_2 = \frac{\sum (x - \bar{x})^2}{n}$$

$$m_3 = \frac{\sum (x - \bar{x})^3}{n}$$

$$m_4 = \frac{\sum (x - \bar{x})^4}{n}$$

Relation between raw moments and central moments

a) $m_1 = 0$

b) $m_2 = m'_2 - m_1'^2$

c) $m_3 = m'_3 - 3m_1' m'_2 + 2m_1'^3$

d) $m_4 = m'_4 - 4m_1' m'_3 + 6m_1'^2 m'_2 - 3m_1'^4$

Example 3.18

Calculate the first 4 central moments from the following sample data.

3, 5, 6, 9, 12

Solution:

Here sample mean, $\bar{x} = \frac{\sum x}{n} = \frac{3+5+\dots+12}{5} = 7$

x	$(x - \bar{x})$	$(x - \bar{x})^2$	$(x - \bar{x})^3$	$(x - \bar{x})^4$
3	-4	16	-64	256
5	-2	4	-8	16
6	-1	1	-1	1
9	2	4	8	16
12	5	25	125	625
$\sum x = 35$	$\sum (x - \bar{x}) = 0$	$\sum (x - \bar{x})^2 = 50$	$\sum (x - \bar{x})^3 = 60$	$\sum (x - \bar{x})^4 = 914$

The central moments are:

$$m_1 = 0$$

$$m_2 = \frac{50}{5} = 10$$

$$m_3 = \frac{60}{5} = 12$$

$$m_4 = \frac{914}{5} = 182.8$$

Coefficient of skewness and kurtosis based on moments

The **coefficient skewness** say γ_1 is defined as

$$\gamma_1 = \frac{m_3}{\sqrt{m_2^3}}$$

Decision:

- a) $\gamma_1 > 0$ indicates positive skewness;
- b) $\gamma_1 \approx 0$ indicates symmetrical distribution;
- c) $\gamma_1 < 0$ indicates negative skewness.

The **coefficient of kurtosis** say β_2 is defined as:

$$\beta_2 = \frac{m_4}{m_2^2}$$

Decision:

- a) $\beta_2 > 3$ indicates *leptokurtic* distribution;
- b) $\beta_2 \approx 3$ indicates *mesokurtic/normal* distribution;
- c) $\beta_2 < 3$ indicates *platykurtic* distribution.

Example 3.18 (continued)

From the calculated moments in **Example 3.18** now we compute coefficients of skewness and kurtosis.

Now,

$$\gamma_1 = \frac{m_3}{\sqrt{m_2^3}} = \frac{12}{\sqrt{10^3}} = 0.38 > 0$$

So the distribution is **positively skewed**.

Again,

$$\beta_2 = \frac{m_4}{m_2^2} = \frac{182}{10^2} = 1.82 < 3$$

So the distribution is **platykurtic**.

Example 3.19 A sample of 7 data entry clerks employed in the Harry County Tax Office revised the following *number of tax records* last hour: 100, 75, 70, 65, 80, 68 and 50.

Comment about *skewness* and *kurtosis* of the *number of tax records*.

NOTE:

Some authors (Newbold, Carlson, and Thorne 2013) use the following formulas to measure skewness and kurtosis without mentioning the moments.

$$Skewness = \frac{1}{n} \left[\sum \left(\frac{X_i - \bar{X}}{s} \right)^3 \right] = \frac{1}{n} \sum z_i^3$$

For calculation purpose the above formula is written as

$$Skewness = \frac{1}{n} \frac{\sum (X - \bar{X})^3}{s^3}$$

and

$$Kurtosis = \frac{1}{n} \left[\sum \left(\frac{X_i - \bar{X}}{s} \right)^4 \right] = \frac{1}{n} \sum z_i^4$$

For calculation purpose the above formula is written as

$$Kurtosis = \frac{1}{n} \frac{\sum (X - \bar{X})^4}{s^4}$$

3.7.4 Some other formulas to calculate skewness

1) Karl Pearson's Coefficient of Skewness,

$S_k = \frac{Mean - Mode}{SD}$. When mean *Mode* is not available use median:

$$S_k = \frac{3(Mean - Median)}{SD}$$

2) Bowley's Skewness (Based on quartiles):

$$S_k = \frac{Q_3 + Q_1 - 2Median}{Q_3 - Q_1}$$

3) Kelly's Skewness (Uses 10th and 90th percentiles):

$$S_k = \frac{P_{90} + P_{10} - 2P_{50}}{P_{90} - P_{10}}$$

Home work A sample of 7 data entry clerks employed in the Harry County Tax Office revised the following *number of tax records* last hour: 100, 75, 70, 65, 80, 68 and 50.

Compute coefficient of *skewness* using *Pearson*, *Bowley* and *Kelly's* methods. Compare the results.

Home work

1) **Compute** coefficient of *skewness* and *kurtosis* and comment for the following data : 20, 21, 5, 9, 14, 6, 19, 16.

2) Suppose the following data are the ages of Internet users obtained from a sample.


41, 15, 31, 25, 24, 23, 21, 22, 22, 18, 30, 20, 19, 19, 16, 23, 27, 38, 34, 24, 19, 20, 29, 17, 23.

From sample data we have following statistics: $\sum (X - \bar{X})^2 = 1062$, $\sum (X - \bar{X})^3 = 7020$ and $\sum (X - \bar{X})^4 = 153198$.

a) **Plot** a histogram. What is your observation about skewness? Is it possible to have an idea about form the histogram?

b) Now compute the coefficient of skewness and kurtosis of the given data and comment.

c) Does your observation in part (a) match with the result in part (b)?



3.8 Exercise

3.1 What are the common measures of central tendency/ location?

3.2 When median is preferable to mean?

3.3 Discuss the nature of unimodal, bimodal and multimodal data/ distribution.

3.4 What are the common measures of dispersion/ variation?

3.5 (a) Compute the mean of the following sample values: 5, 9, 4, 10 (b) Show that $\sum(x - \bar{x}) = 0$.

3.6 (a) Compute the mean of the following sample values: 1.3, 7.0, 3.6, 4.1, 5.0 (b) Show that $\sum(x - \bar{x}) = 0$.

3.7 Show that variance is affected by change of scale; but not by origin.

3.8 The monthly starting salary (\$) of 12 graduates:

3450 ,3550 ,3650 ,3480 ,3355, 3310 ,3490 ,3730, 3540 ,3925, 3520 ,3480

i) **Compute** sample mean and standard deviation.

ii) **Compute** sample median and IQR.

iii) To be in top 10% earners what should be the starting salary of a graduate?

3.9 Automobile Fuel Efficiencies. In automobile mileage and gasoline-consumption testing, 13 automobiles were road tested for 300 miles in both city and highway driving conditions. The following data were recorded for miles-per-gallon performance.

City: 16.2, 16.7, 15.9, 14.4, 13.2, 15.3, 16.8, 16.0, 16.1, 15.3, 15.2, 15.3, 16.2

Highway: 19.4, 20.6, 18.3, 18.6, 19.2, 17.4, 17.2, 18.6, 19.0, 21.1, 19.4, 18.5, 18.7

Use the mean, median, and mode to make a statement about the difference in performance for city and highway driving.

3.10 Air Quality Index. The Los Angeles Times regularly reports the air quality index for various areas of Southern California. A sample of air quality index values for Pomona provided the following data: 28, 42, 58, 48, 45, 55, 60, 49, and 50.

- i) Compute the range and interquartile range.
- ii) Compute the sample variance and sample standard deviation.
- iii) A sample of air quality index readings for Anaheim provided a sample mean of 48.5, a sample variance of 136, and a sample standard deviation of 11.66. What comparisons can you make between the air quality in Pomona and that in Anaheim on the basis of these descriptive statistics?

3.11 Reliability of Delivery Service. The following data were the number of days required to fill orders for Dawson Supply, Inc., and J.C. Clark Distributors.

Dawson Supply Days for Delivery: 11, 10, 9, 10, 11, 11, 10, 11, 10, 10

Clark Distributors Days for Delivery: 8, 10, 13, 7, 10, 11, 10, 7, 15, 12

Which company is more *consistent* to fill orders?

Hints: Compute and compare standard deviation (SD) of the number days for each company. The less SD would indicate more consistency.

3.12 Amateur Golfer Scores. Scores turned in by an amateur golfer at the Bonita Fairways Golf Course in Bonita Springs, Florida, during 2017 and 2018 are as follows:

2017 Season: 74, 78, 79, 77, 75, 73, 75, 77

2018 Season: 71, 70, 75, 77, 85, 80, 71, 79

- i) Use the mean and standard deviation to evaluate the golfer's performance over the two-year period.
- ii) What is the primary difference in performance between 2017 and 2018? What improvement, if any, can be seen in the 2018 scores?

3.13 Consistency of Running Times. The following times were recorded by the quarter-mile and mile runners of a university track team (times are in minutes).

Quarter-Mile Times: 0.92, 0.98, 1.04, 0.90, 0.99

Mile Times: 4.52, 4.35, 4.60, 4.70, 4.50

After viewing this sample of running times, one of the coaches commented that the quarter-milers turned in the more consistent times.

- i) Use the standard deviation and the coefficient of variation to summarize the variability in the data.
- ii) Does the use of the coefficient of variation indicate that the coach's statement should be qualified?

3.14 Automobiles traveling on a road with a posted speed limit of 55 miles per hour are checked for speed by a state police radar system. Following is a frequency distribution of speeds.

Speed (miles per hour)	Frequency
45-49	10
50-54	40
55-59	150
60-64	175
65-69	75
70-74	15
75-79	10
Total	475

- i) What is the *mean speed* of the automobiles traveling on this road?
- ii) **Compute** the *variance* and the *standard deviation*.

3.15 Consider a sample with a mean of 500 and a standard deviation of 100. What are the z-scores for the following data values: 520, 650, 500, 450, and 280?

3.16 Consider a sample with a mean of 30 and a standard deviation of 5. Use Chebyshev's theorem to determine the percentage of the data within each of the following ranges:

- a. 20 to 40
- b. 15 to 45

3.17 The results of a national survey showed that on average, adults sleep 6.9 hours per night. Suppose that the standard deviation is 1.2 hours.

- a. Use Chebyshev's theorem to calculate the percentage of individuals who sleep between 4.5 and 9.3 hours.
- b. Use Chebyshev's theorem to calculate the percentage of individuals who sleep between 3.9 and 9.9 hours.
- c. Assume that the number of hours of sleep follows a bell-shaped distribution. Use the empirical rule to calculate the percentage of individuals who sleep between 4.5 and 9.3 hours per day. How does this result compare to the value that you obtained using Chebyshev's theorem in part (a)?

3.18 The high costs in the California real estate market have caused families who cannot afford to buy bigger homes to consider backyard sheds as an alternative form of housing expansion. Many are using the backyard structures for home offices, art studios, and hobby areas as well as for additional storage. The mean price of a customized wooden, shingled backyard structure is \$3100 (Newsweek, September 29, 2003). Assume that the standard deviation is \$1200. a. What is the z-score for a backyard structure costing \$2300?

- b. What is the z-score for a backyard structure costing \$4900?
- c. Interpret the z-scores in parts (a) and (b). Comment on whether either should be considered an outlier.

- d. The Newsweek article described a backyard shed-office combination built in Albany, California, for \$13,000. Should this structure be considered an outlier? Explain.

3.20 A data set has a first quartile of 42 and a third quartile of 50. Compute the lower and upper limits for the corresponding box plot. Should a data value of 65 be considered an outlier?

3.22 Suppose a consumer group asked 18 consumers to keep a yearly log of their shopping practices and that the following data represent the number of coupons used by each consumer over the yearly period.

3.9 Data

81, 68, 70, 100, 94, 47, 66, 70, 82, 110, 105, 60, 21, 70, 66, 90, 78, 85

3.10 Ordered data

21, 47, 60, 66, 66, 68, 70, 70, 70, 78, 81, 82, 85, 90, 94, 100, 105, 110

- a) Use the data to construct a box-and-whisker plot.
- b) Discuss the skewness of the distribution of these data and point out any outliers.

4 Probability

A probability is the chance, or likelihood, that a particular event will occur. These are examples of events representing typical probability-type questions:

- How many customers will arrive in a super shop in next 30 minutes?
- What is probability that a stock price will rise or fall?

To answer these kind of questions in the face of uncertainty we need to study probability. To answer these type of questions which are raised in real life; at first we have to learn some basic concepts of probability.

4.1 Random experiment

A **random experiment** is a process leading to two or more possible outcomes, without knowing exactly which outcome will occur (Newbold, Carlson, and Thorne 2013).

Example 4.1: Tossing a coin, throwing a dice, change in the stock prices etc.

4.2 Sample space

A **sample space** is the collection of all outcomes of a random experiment. The sample space is usually denoted by S or Greek letter Ω (omega).

Example 4.2:

- If we toss a coin then the sample space is: $S = \{H, T\}$
- If we toss 2 coins then the sample space is: $S = \{HH, HT, TH, TT\}$

4.3 Event

An **event** is a *subset* of a *sample space*.

For example suppose, $S = \{HH, HT, TH, TT\}$ and $A = \{HH, TT\}$ is an event which a subset of sample space S .

4.4 Complement of an event

The complement of an event A with respect to Ω is the subset of all elements of Ω that are not in A . We denote the complement of A by the symbol A^C .

Example 4.3: Consider the sample space:

$$\Omega = \{1, 2, 3, 4, 5, 6\}$$

Let, $A = \{1, 3, 5\}$. Then the complement of A is $A^C = \Omega - A = \{2, 4, 6\}$

4.5 Mutually exclusive events

The occurrence of one event means that none of the other events can occur at the same time.

Example 4.4:

- The variable “Employment status” presents mutually exclusive outcomes, *employed* and *unemployed*. An employee selected at random is either male or female but cannot be both.
- A manufactured part is acceptable or unacceptable. The part cannot be both acceptable and unacceptable at the same time.

4.6 Collectively Exhaustive

Given the K events $E_1, E_2, \dots, E_K$ in the sample space, S , if $E_1 \cup E_2 \cup \dots \cup E_K = S$, these K events are said to be collectively exhaustive.

4.7 Axiomatic definition of Probability

The **probability** of an event A is the sum of the weights of all sample points in A . Therefore,

(a) $0 \leq P(A) \leq 1$

(b) If $A_1, A_2, A_3, \dots$ is a sequence of mutually exclusive events, then

$$P(A_1 \cup A_2 \cup A_3 \cup \dots) = P(A_1) + P(A_2) + P(A_3) + \dots$$

(c) $P(\Omega) = 1$

4.8 Probability of an event (Classical approach)

Suppose an event A is defined in the sample space S . Then the probability of event A is defined as :

$$P(A) = \frac{n(A)}{n(S)};$$

Here,

$n(A)$ = number of outcomes favorable to event A ;

$n(S)$ = total number of outcomes in the sample space S .

Example 4.5 Consider a random experiment of throwing two six-sided fair dices. Then the sample space is:

	Dice2					
Dice1	(1,1)	(1,2)	(1,3)	(1,4)	(1,5)	(1,6)
	(2,1)	(2,2)	(2,3)	(2,4)	(2,5)	(2,6)
	(3,1)	(3,2)	(3,3)	(3,4)	(3,5)	(3,6)
	(4,1)	(4,2)	(4,3)	(4,4)	(4,5)	(4,6)
	(5,1)	(5,2)	(5,3)	(5,4)	(5,5)	(5,6)
	(6,1)	(6,2)	(6,3)	(6,4)	(6,5)	(6,6)

Now **compute** the following probabilities:

a) probability of same number in both dices;

b) probability that sum of the numbers of two dices are equal to 5.

Solution: Here $n(\Omega) = 36$

a) Let, $A = \{\text{same number in both dices}\} = \{(1,1), (2,2), (3,3), (4,4), (5,5), (6,6)\}$.

Hence, $n(A) = 6$. So, $P(A) = \frac{n(A)}{n(\Omega)} = \frac{6}{36} = \frac{1}{6}$.

b) DIY (do it yourself).

Example 4.6 A box/ an urn contains 6 black balls and 4 white balls. If two balls are selected at random (at a time) what is the probability that the

i) both balls will be black?

ii) both balls will be white?

Solution-i) Here, 2 balls can be selected in total $\binom{10}{2} = 45$ ways. So, $n(\Omega) = 45$.

Suppose, $B = \{2 \text{ black balls selected}\}$. Two black balls can be selected in $\binom{6}{2} = 15$ ways. So, $n(B) = 15$.

$\therefore P(B) = \frac{n(B)}{n(\Omega)} = \frac{15}{45} = \frac{1}{3}$.

Solution-i) DIY.

4.9 Probability of an event (Empirical approach)

Empirical Probability is a type of probability that is calculated based on actual observations, experiments, or historical data rather than theoretical assumptions. It measures the likelihood of an event occurring by analyzing past occurrences or experimental results.

Formula for Empirical Probability:

$$P(E) = \frac{\text{Number of times the event occurs}}{\text{Total number of trials}}$$

Where:

- $P(E)$ is the probability of the event E ,
- The numerator is the count of occurrences of the event, and
- The denominator is the total number of trials or observations.

Example 4.7: Suppose in a class there are 30 students; 20 are male and 10 are females. If a student is selected at random what is the probability that he is a male?

Solution: Let, E_1 = set of male students and E_2 = set of female students. And, S = set of all students

So, probability that a male student is selected is:

$$P(E_1) = \frac{n(E_1)}{n(S)} = \frac{20}{30} = 0.66667 \approx 0.67$$

Interpretation There is almost 67% chance that the selected student will be male.

4.10 Properties of Probability Laws

Probability laws have a number of properties, which can be deduced from the axioms. Some of them are summarized below.

- $P(A^C) = 1 - P(A)$ [**complement rule**]
- $P(A \cap B^C) = P(A) - P(A \cap B)$ [**only A happens**]
- $P(A \cup B) = P(A) + P(B) - P(A \cap B)$ [**additive rule**]
- $P(A^C \cap B^C) = P(A \cup B)^C = 1 - P(A \cup B)$. [**neither A NOR B happens**]
- $P(\text{only } A \text{ or only } B) = P(A \cap B^C) + P(A^C \cap B)$
 $= P(A) + P(B) - 2P(A \cap B)$

Example 4.8: In a class 65% students prefer tea and 35% students prefer coffee. While 15% students prefer both tea and coffee. If a student is selected at random from the class **find** the probability that

- he/she prefers only coffee

- ii) he/she prefers tea or coffee
- iii) he/she prefers none (neither tea nor coffee)

Example 4.9 (Lind, Marchal, and Wathen 2012, 166) A local bank reports that 80 percent of its customers maintain a checking account, 60 percent have a savings account, and 50 percent have both. If a customer is chosen at random, what is the probability the customer has either a checking or a savings account? What is the probability the customer does not have either a checking or a savings account?

Example 4.10 (Lind, Marchal, and Wathen 2012, 166) All Seasons Plumbing has two service trucks that frequently need repair. If the probability the first truck is available is .75, the probability the second truck is available is .50, and the probability that both trucks are available is .30, what is the probability neither truck is available?

4.11 Conditional Probability

The conditional probability of an event A , *given* an event B with $P(B) > 0$, is defined by,

$$P(A|B) = \frac{P(A \cap B)}{P(B)} = \frac{n(A \cap B)}{n(B)}$$

Example 4.11 The probability that a regularly scheduled flight departs on time is $P(D) = 0.83$; the probability that it arrives on time is $P(A) = 0.82$; and the probability that it departs and arrives on time is $P(D \cap A) = 0.78$. Find the probability that a plane:

- (a) arrives on time, given that it departed on time, and
- (b) departed on time, given that it has arrived on time.

4.12 The Multiplication Rule

The multiplication rule is used to calculate the joint probability of two events.

The joint probability of any two events A and B is

$$P(A \cap B) = P(B).P(A|B) \quad [\text{Considering } B \text{ as prior}]$$

or, altering the notation,

$$P(A \cap B) = P(A).P(B|A) \quad [\text{Considering } A \text{ as prior}]$$

Example 4.12: Suppose a box contains 10 balls; 4 are black and 6 are white. If 2 balls are drawn at random successively without replacement, what is the probability that both balls are white?

Solution:

Let, W_1 = 1st ball is white ; W_2 = 2nd ball is also white.

According to question,

$$P(\text{both balls are white}) = P(W_1 \cap W_2) = P(W_1) \cdot P(W_2|W_1)$$

$$= \frac{6}{10} \cdot \frac{5}{9} = \frac{1}{3}$$

Example 4.13 Suppose $P(A) = 0.40$ and $P(B|A) = 0.30$. What is the joint probability of A and B ?

Example 4.14 Suppose $P(X_1) = 0.75$ and $P(Y_2|X_1) = 0.30$. What is the joint probability of X_1 and Y_2 ?

4.13 Independent events

If two events A and B are independent, the probability that both of them occur is equal to the product of their individual probabilities i.e.

$$P(A \cap B) = P(A)P(B)$$

- **Corollary:** If A and B are independent events then their complement events also be independent that is,

$$P(A^C \cap B^C) = P(A^C)P(B^C)$$

- **Independence Rule for Multiple events:**

$$P(A \cap B \cap C) = P(A)P(B)P(C)$$

Example 4.15 (Lind, Marchal, and Wathen 2012, 182) You take a trip by air that involves three independent flights. If there is an 80 percent chance each specific leg of the trip is done on time, what is the probability all three flights arrive on time?

Example 4.16 (Lind, Marchal, and Wathen 2012, 182) The probability a HP network server is down is .05. If you have three independent servers, what is the probability that at least one of them is operational?

Solution:

Given, $P(\text{server is down}) = 0.05$.

So, $P(\text{server is operational}) = 0.95$

Now, let $O_i = \{i^{\text{th}} \text{ server is operational}\}$

So,

$$\begin{aligned} &P(\text{at least one of them is operational}) \\ &= P(O_1 \cup O_2 \cup O_3) = 1 - P(O_1^C \cap O_2^C \cap O_3^C) \end{aligned}$$

$$\begin{aligned}
&= 1 - P(O_1^C) \cdot P(O_2^C) \cdot P(O_3^C) \\
&= 1 - (0.05)(0.05)(0.05) = 0.9999875.
\end{aligned}$$

Example 4.17 (Lind, Marchal, and Wathen 2012, 182) Twenty-two percent of all liquid crystal displays (LCDs) are manufactured by Samsung. What is the probability that in a collection of three independent LCD purchases, at least one is a Samsung?

4.14 Bivariate Probabilities: Joint and Marginal Probability

The **Intersection** of events A and B is the event that occurs when both A and B occur.

It is denoted as A and B or $(A \cap B)$.

The probability of the intersection is called the joint probability that is $P(A \cap B)$.

Example 4.18: Suppose that our sample space S is the population of 900 adults in a small town who have completed the requirements for a college degree. We shall categorize them according to gender and employment status. The data are given in Table 4.2 (also referred as *joint frequency table* or *cross-table*)

Table 4.2: Categorization of the Adults in a Small Town

	Employed	Unemployed
Male	460	40
Female	140	260

Question i: Construct a joint probability table

Solution i: Let,

A_1 = Male adults

A_2 = Female adults

B_1 = Employed adults

B_2 = Unemployed adults

Here $n(S) = 900$. Now divide all cell frequency by 900 and round to 2 decimal points, hence we get joint probability table below(see Table 4.3):

Table 4.3: Joint probability table

	B_1	B_2
A_1	0.51	0.04
A_2	0.16	0.29

Joint probability: In Table 4.3 the joint probabilities are:

i) $P(A_1 \cap B_1) = 0.51$

- ii) $P(A_1 \cap B_2) = 0.04$
- iii) $P(A_2 \cap B_1) = 0.16$ and
- iv) $P(A_2 \cap B_2) = 0.29$

Marginal probability: In Table 4.3 the marginal probabilities are:

- i) $P(A_1) = 0.51 + 0.04 = 0.55$
- ii) $P(A_2) = 0.16 + 0.29 = 0.45$
- iii) $P(B_1) = 0.51 + 0.16 = 0.67$
- iv) $P(B_2) = 0.04 + 0.29 = 0.33$

From a joint probability table we can also compute **conditional probabilities**. For example,

$$P(A_1|B_1) = \frac{P(A_1 \cap B_1)}{P(B_1)} = \frac{0.51}{0.67} \approx 0.7612$$

4.15 Independent Events in Joint probability table

Let A and B be a pair of events, each broken into mutually exclusive and collectively exhaustive event categories denoted by labels $A_1, A_2, \dots, A_H$ and $B_1, B_2, \dots, B_K$. If every event A_i is statistically independent of every event B_j , then A and B are independent events (Newbold, Carlson, and Thorne 2013).

Example 4.19 Students in a business statistics class were asked what grade they expected in the course and whether they worked on additional problems beyond those assigned by the instructor. The following table gives proportions of students in each of eight joint classifications (Newbold, Carlson, and Thorne 2013, exercise 3.68).

Worked Problems	Expected Grade			
	A	B	C	Below C
Yes	0.12	0.06	0.12	0.02
No	0.13	0.21	0.26	0.08

- a. **Find** the probability that a randomly chosen student from this class worked on additional problems.
- b. **Find** the probability that a randomly chosen student from this class expects an A.
- c. **Find** the probability that a randomly chosen student expects an A given that he/she worked on additional problems .
- d. **Find** the probability that a randomly chosen student worked on additional problems given that he/she expects an A .
- e. Are “worked additional problems” and “expected grade” statistically independent?

Solution:

Let, $Y = \{\text{Yes}\}$ and $N = \{\text{No}\}$.

The joint probability table with marginal probability table is given below:

	A	B	C	D (Below C)	Row total
Y	0.12	0.06	0.12	0.02	0.32
N	0.13	0.21	0.26	0.08	0.68
Column total	0.25	0.27	0.38	0.10	1.00

Solution of (e): To show whether “worked additional problems” and “expected grade” statistically independent we have to verify whether “Y, N” and “A”, “B”, “C” are independent events.

Now from joint probability table, $P(Y \cap A) = 0.12$.

And $P(Y \cap A) = P(Y) \cdot P(A) = 0.32 \times 0.25 = 0.08$.

Since $P(Y \cap A) \neq P(Y) \cdot P(A)$ so, Y and A are not independent. Hence, we do not need to test other combinations.

In conclusion we can say that “worked additional problems” and “expected grade” are not statistically independent.

4.16 Exercises 4.1

4.1) (Anderson and Sweeney 2011) Suppose that we have two events, A and B , with $P(A) = 0.50$, $P(B) = 0.60$, and $P(A \cap B) = 0.40$.

i) Find $P(A|B)$.

ii) Find $P(B|A)$.

iii) Are A and B independent? Why or why not?

4.2) Suppose $P(A)=0.40$ and $P(B|A)=0.30$. What is the joint probability of A and B ?

4.3) A local bank reports that 80 percent of its customers maintain a checking account, 60 percent have a savings account, and 50 percent have both. If a customer is chosen at random, what is the probability the customer has either a checking or a savings account? What is the probability the customer does not have either a checking or a savings account?

4.4) (Keller 2014) Suppose we have the following joint probabilities.

	A ₁	A ₂	A ₃
B ₁	0.15	0.20	0.10
B ₂	0.25	0.25	0.05

Compute the marginal probabilities.

4.5) Refer to Exercise 4.

- a. Compute $P(A_2|B_2)$. Also compute $P(\bar{A}_2|B_2)$
- b. Compute $P(B_1|A_2)$.

4.6) Refer to Exercise 2.

- a. Compute $P(A_1 \text{ or } A_2)$.
- b. Compute $P(A_2 \text{ or } B_2)$.

4.7) A survey was conducted among 100 college students about their preferred study method and performance in exams.

	Low Score	Medium Score	High Score
Group Study	10	15	5
Self Study	5	25	40

- a. **Construct** a joint probability table.
- b. **Compute** the probability that a randomly selected student did group study and scored medium?
- c. **Find** the probability that a student prefers Self Study, regardless of score.
- d. **Determine** the probability that a student either got a High Score or prefers Group Study.
- e. Given that a student scored Low, what is the probability they did group study?
- f. A student scored high. **Find** the probability that he did self study.
- g. Are preferred study method and performance in exam independent?

4.8) A company surveys 600 customers about their income level and purchasing behavior of a premium product.

	Did Not Buy	Bought Once	Bought Multiple Times
Low Income	150	60	30
High Income	60	90	210

- a. What is the probability that a customer is high income and bought multiple times?
- b. What is the probability that a customer is from the low-income group?
- c. What is the probability that a customer is either high income or bought multiple times?
- d. What is the probability that a customer is high income, given that they bought multiple times?
- e. What is the probability that a customer bought once, given that they are low income?
- f. Are income level and purchasing behavior independent?

4.9) A firm has classified its customers in two ways: (1) according to whether the account is overdue and (2) whether the account is new (less than 12 months) or old. An analysis of the firm's records provided the input for the following table of joint probabilities.


	Overdue	Not overdue
New	0.06	0.13
Old	0.52	0.29

One account is randomly selected.

- If the account is overdue, what is the probability that it is new?
- If the account is new, what is the probability that it is overdue?
- Is the age of the account related to whether it is overdue? Explain.

4.17 Probability Trees

Consider a sequential experiment where in the **first stage** either A_1 or A_2 can be happened with some probabilities . And in the **second stage** event B can be happened. If B^C is the complement of B then this experiment can be shown in the following **tree diagram**.




Example 4.20:

Two balls are drawn in succession, without replacement, from a box containing 3 blue and 2 white balls .

- What is the probability that both balls will be white ?

Solution: Here, two balls are drawn in succession (one by one) without replacement. This experiment can be shown in the following tree:



The probability of drawing a white ball on the first draw and a white ball on the second draw (both are white) is:

$$P(W_1 \cap W_2) = P(W_1)P(W_2|W_1) = \left(\frac{2}{5}\right)\left(\frac{1}{4}\right) = \frac{1}{10}$$

ii) What is the probability that the second ball is white?

Solution:

$$\begin{aligned}
 P(W_2) &= P(W_1 \cap W_2) + P(B_1 \cap W_2) \\
 &= P(W_1)P(W_2|W_1) + P(B_1)P(W_2|B_1) \\
 &= \left(\frac{2}{5}\right)\left(\frac{1}{4}\right) + \left(\frac{3}{5}\right)\left(\frac{2}{4}\right) = \frac{1}{10} + \frac{3}{10} = \frac{4}{10} = \frac{2}{5}.
 \end{aligned}$$

4.18 Total Probability rule and Bayes' Theorem

Suppose A_1 , A_2 , and A_3 are mutually exclusive and exhaustive events, that is:

$$P(A_i \cap A_j) = 0 \text{ for } i \neq j = 1, 2, 3;$$

and

$$P(A_1 \cup A_2 \cup A_3) = 1$$

- The *prior* probabilities are $P(A_1)$, $P(A_2)$ and $P(A_3)$.
- The *likelihood/conditional* probabilities are $P(B|A_1)$, $P(B|A_2)$ and $P(B|A_3)$ (see Figure 4.1).




Figure 4.1

The total probability rule

$$P(B) = P(A_1).P(B|A_1) + P(A_2).P(B|A_2) + P(A_3).P(B|A_3)$$

Bayes' theorem

The Bayes' theorem is used to find the *posterior/revised/update* probabilities of *prior* probabilities.

$$P(A_1|B) = \frac{P(A_1 \cap B)}{P(B)} = \frac{P(A_1).P(B|A_1)}{P(A_1).P(B|A_1) + P(A_2).P(B|A_2) + P(A_3).P(B|A_3)}$$

In the same way we can compute $P(A_2|B)$ and $P(A_3|B)$.

Example 4.21: A marketing company is studying customer behavior by analyzing how individuals respond to different types of advertisements and whether those interactions lead to a purchase. Each customer is exposed to one of three types of ads: email with a probability of 0.3, social media with a probability of 0.5, and television with a probability of 0.2. The effectiveness of these ads varies: 20% of customers who receive an email make a purchase, while 40% respond positively to social media ads, and 50% of those shown a TV ad proceed to buy.

- Construct** a probability tree from the above information.
- Find** the probability that a customer will make a purchase (regardless which type of ad he was exposed).
- Suppose a customer made a purchase. What is the probability that he saw the ad in social media?

4.19 Exercises 4.2

4.9) (Anderson and Sweeney 2011) The prior probabilities for events A_1 and A_2 are $P(A_1) = .40$ and $P(A_2) = .60$. It is also known that $P(A_1 \cap A_2) = 0$. Suppose $P(B|A_1) = .20$ and $P(B|A_2) = .05$.

- Are A_1 and A_2 mutually exclusive? Explain.
- Compute $P(A_1 \cap B)$ and $P(A_2 \cap B)$.
- Compute $P(B)$.

d. Apply Bayes' theorem to compute $P(A_1|B)$ and $P(A_2|B)$.

4.10) (Lind, Marchal, and Wathen 2012, 171) The Ludlow Wildcats baseball team, a minor league team in the Cleveland Indians organization, plays 70 percent of their games at night and 30 percent during the day. The team wins 50 percent of their night games and 90 percent of their day games. According to today's newspaper, they won yesterday. What is the probability the game was played at night?

4.11) (Lind, Marchal, and Wathen 2012, 171) Dr. Stallter has been teaching basic statistics for many years. She knows that 80 percent of the students will complete the assigned problems. She has also determined that among those who do their assignments, 90 percent will pass the course. Among those students who do not do their homework, 60 percent will pass. Mike Fishbaugh took statistics last semester from Dr. Stallter and received a passing grade. What is the probability that he completed the assignments?

4.12) (Anderson and Sweeney 2011) A local bank reviewed its credit card policy with the intention of recalling some of its credit cards. In the past approximately 5% of cardholders defaulted, leaving the bank unable to collect the outstanding balance. Hence, management established a prior probability of .05 that any particular cardholder will default. The bank also found that the probability of missing a monthly payment is .20 for customers who do not default. Of course, the probability of missing a monthly payment for those who default is 1.

- a. Given that a customer missed one or more monthly payments, compute the posterior probability that the customer will default.
- b. The bank would like to recall its card if the probability that a customer will default is greater than .20. Should the bank recall its card if the customer misses a monthly payment? Why or why not?

4.13) (Black 2012) In a manufacturing plant, machine A produces 10% of a certain product, machine B produces 40% of this product, and machine C produces 50% of this product. Five percent of machine A products are defective, 12% of machine B products are defective, and 8% of machine C products are defective. The company inspector has just sampled a product from this plant and has found it to be defective. Determine the revised probabilities that the sampled product was produced by machine A, machine B, or machine C.

4.14) (Black 2012) Suppose 70% of all companies are classified as small companies and the rest as large companies. Suppose further, 82% of large companies provide training to employees, but only 18% of small companies provide training. A company is randomly selected without knowing if it is a large or small company; however, it is determined that the company provides training to employees. What are the prior probabilities that the company is a large company or a small company? What are the revised probabilities that the company is large or small? Based on your analysis, what is the overall percentage of companies that offer training?


4.15) (Black 2012) Alex, Alicia, and Juan fill orders in a fast-food restaurant. Alex incorrectly fills 20% of the orders he takes. Alicia incorrectly fills 12% of the orders she takes. Juan incorrectly fills 5% of the orders he takes. Alex fills 30% of all orders, Alicia fills 45% of all orders, and Juan fills 25% of all orders. An order has just been filled.

- a. What is the probability that Alicia filled the order?
- b. If the order was filled by Juan, what is the probability that it was filled correctly?

- c. Who filled the order is unknown, but the order was filled incorrectly. What are the revised probabilities that Alex, Alicia, or Juan filled the order?
- d. Who filled the order is unknown, but the order was filled correctly. What are the revised probabilities that Alex, Alicia, or Juan filled the order?

i Review exercises

1) One urn has 3 blue and 2 white balls; a second urn has 1 blue and 3 white balls. A single fair die is rolled and if 1 or 2 comes up, a ball is drawn out of the first urn; otherwise, a ball is drawn out of the second urn. If the drawn ball is blue, what is the probability that it came out of the first urn? Out of the second urn?



2) One bag contains 4 white balls and 3 black balls, and a second bag contains 3 white balls and 5 black balls. One ball is drawn from the first bag and placed unseen in the second bag. What is the probability that a ball now drawn from the second bag is black? (**Hints:** Apply probability tree)

3) From 4 red, 5 green, and 6 yellow apples, **find** the probability that, out of 9 randomly selected apples, exactly 3 of each color are selected.

4) Pollution of the rivers in the United States has been a problem for many years. Consider the following events:

A: the river is polluted,

B : a sample of water tested detects pollution,

C : fishing is permitted.

Assume $P(A) = 0.3$, $P(B|A) = 0.75$, $P(B|A') = 0.20$, $P(C|A \cap B) = 0.20$, $P(C|A' \cap B) = 0.15$, $P(C|A \cap B') = 0.80$, and $P(C|A' \cap B') = 0.90$.

(a) Find $P(A \cap B \cap C)$.

(b) Find $P(B' \cap C)$.

(c) Find $P(C)$.

(d) Find the probability that the river is polluted, given that fishing is permitted and the sample tested did not detect pollution.

4.20 Random variables

A **variable** is said to be **random** if its values are determined by a random experiment. In other word, **random variable** is a numerical description of the outcome of an experiment.

- A random variable often denoted with an uppercase letter (say X)
- The value of a random variable is denoted with a lowercase letter (say x)

Illustration Consider a random experiment of tossing a coin (fair/unfair) 2 times. Then the sample space is

$$S = \{HH, HT, TH, TT\}$$

Now let, $X = \text{number of heads occur}$

From the sample space we can see that X can take following values:

Sample point	x
HH	2
HT	1
TH	1
TT	0

Since the values of X completely determined by the outcomes of the random experiment, so X is a random variable (discrete).

4.20.1 Types of random variable

There are two types of random variables, **discrete** and **continuous**.

A **discrete random variable** can assume only a certain number of separated values. A discrete random variable is usually the result of *counting* something. For example, number of customers arrive, number of calls receive etc.

A **continuous random variable** is one whose values are uncountable or which can take any value in a given interval. Generally a continuous random variable is usually the result of *measuring* something.

Task Which of these variables are discrete and which are continuous random variables?

- The number of new accounts established by a salesperson in a year.
- The time between customer arrivals to a bank ATM.
- The number of customers in Big Nick's barber shop.
- The amount of fuel in your car's gas tank.
- The number of minorities on a jury.
- The outside temperature today.

5 Discrete Probability Distributions

5.1 Discrete random variable and Probability mass function (PMF)

Suppose X is a discrete random variable. The **probability mass function (PMF)** of X can be denoted as $f(x)$ where

$$f(x) = P(X = x)$$

For each possible outcome x ; $f(x)$ must satisfies:

1.

$$f(x) \geq 0$$

2.

$$\sum_x f(x) = 1$$

The **PMF** $f(x)$ is also called probability distribution of the discrete random variable X .

Example 5.1 John Ragsdale sells new cars for Pelican Ford. John usually sells the largest number of cars on Saturday. He has developed the following probability distribution for the number of cars he expects to sell on a particular Saturday.

Number of cars sold, x	Probability, $f(x)$
0	0.10
1	0.20
2	0.30
3	0.30
4	0.10

Compute (i) $P(X = 2)$; (ii) $P(X < 2)$; (iii) $P(X \geq 3)$

5.1.1 Expectation (Mean) of discrete random variable

Let X be a discrete random variable with probability mass function $f(x) = P(X = x)$.

The **expected value of X** the mean of X is denoted by $E(X)$ and defined by:

$$E(X) = \sum_x x.f(x)$$

The expected value of X is sometimes called the population mean of X that is $\mu = E(X)$.

Example 5.2 John Ragsdale sells new cars for Pelican Ford. John usually sells the largest number of cars on Saturday. He has developed the following probability distribution for the number of cars he expects to sell on a particular Saturday.

Number of cars sold, x	Probability, $f(x)$
0	0.10
1	0.20
2	0.30
3	0.30
4	0.10

On a typical Saturday, how many cars does John expect to sell?

Solution:

Table 5.3: Calculation of the Expected Value for the Number of Cars Sold

x	$f(x)$	$x \cdot f(x)$
0	0.10	0.00
1	0.20	0.20
2	0.30	0.60
3	0.30	0.90
4	0.10	0.40
Total	$\sum f(x) = 1$	$\mu = \sum x \cdot f(x) = 2.10$

Alternative: The mean number of cars is:

$$\begin{aligned}\mu &= E[X] = \sum_{x=0}^4 x \cdot f(x) \\ &= 0(0.10) + 1(0.20) + 2(0.30) + 3(0.30) + 4(0.10) = 2.1\end{aligned}$$

So on a typical Saturday, John Ragsdale expects to sell a mean of 2.1 cars a day.

5.1.2 Variance of discrete random variable

Let X be a discrete random variable with probability distribution $f(x)$ and mean μ . The variance of X is

$$\text{var}(X) = \sigma^2 = E[(X - \mu)^2] = \sum_x (x - \mu)^2 f(x)$$

Alternative:

$$\text{var}(X) = E(X^2) - \mu^2$$

Where,

$$E(X^2) = \sum_x x^2 \cdot f(x)$$

Example 5.3: From Example 5.2 **compute** *variance* and *standard deviation* of X .

Solution: From Example 5.2 we have $\mu = 2.1$.

Table 5.4: Calculation of the Variance for the Number of Cars Sold

x	$f(x)$	$x - \mu$	$(x - \mu)^2$	$(x - \mu)^2 f(x)$
0	0.10	-2.1	4.41	0.441
1	0.20	-1.1	1.21	0.242
2	0.30	-0.1	0.01	0.003
3	0.30	0.9	0.81	0.243
4	0.10	1.9	3.61	0.361
Total	$\sum f(x) = 1$			$\sigma^2 = 1.290$

Alternative: Here,

$$E(X^2) = \sum_{x=0}^4 x^2 \cdot f(x)$$

$$= 0^2(0.10) + 1^2(0.20) + 2^2(0.30) + 3^2(0.30) + 4^2(0.10)$$

$$= 5.70$$

$$\text{Hence, } \text{var}(X) = \sigma^2 = E(X^2) - \mu^2 = 5.70 - (2.10)^2 = 1.29$$

- The variance is, $\sigma^2 = 1.29$ and
- The standard deviation is, $\sigma = \sqrt{1.29} = 1.136$

i Properties of $E(\cdot)$ and $\text{var}(\cdot)$

If a and b are constants, then

- $E(b) = b$
- $E(aX + b) = aE(X) + b$
- $\text{var}(b) = 0$
- $\text{var}(aX + b) = a^2 \text{var}(X)$

5.1.3 Exercise: Discrete random variable

5.1) Compute the mean and variance of the following probability distribution.

x	$f(x)$
5	0.10
10	0.30
15	0.20
20	0.40

5.2) The information below is the number of daily emergency service calls made by the volunteer ambulance service of Walterboro, South Carolina, for the last 50 days. To explain, there were 22 days on which there were 2 emergency calls, and 9 days on which there were 3 emergency calls.

Number of calls	Frequency
0	8
1	10
2	22
3	9
4	1
Total	50

- Convert this information on the number of calls to a probability distribution.
 - Is this an example of a discrete or continuous probability distribution?
 - What is the mean number of emergency calls per day?
 - What is the standard deviation of the number of calls made daily?
- 5.3) Consider the following probability distribution of random variable X :

x	1	3	5	7
$f(x)$	k	2k	2k	3k

- Find the value of k .
- Find the probability of the value of X exactly 4.
- Find the probability of the value of X between 3 and 7 (inclusive).
- Estimate expected value and standard deviation of X .

5.4) Suppose that an antique jewelry dealer is interested in purchasing a gold necklace for which the probabilities are 0.22, 0.36, 0.28, and 0.14, respectively, that she will be able to sell it for a profit of \$250, sell it for a profit of \$150, break even, or sell it for a loss of \$150. What is her expected profit?

5.5) The monthly sales at a computer store have a mean of \$25,000 and a standard deviation of \$4,000. Profits are calculated by multiplying sales by 30% and subtracting fixed costs of \$6,000. **Find** the mean and standard deviation of monthly profits.

5.6) When parking a car in a downtown parking lot, drivers pay according to the number of hours or parts thereof. The probability distribution of the number of hours cars are parked has been estimated as follows.

x	1	2	3	4	5	6	7	8
$f(x)$	0.24	0.18	0.13	0.10	0.07	0.04	0.04	0.20

- (a) **Find** the mean and standard deviation of the number of hours cars are parked in the lot.
- (b) The cost of parking is \$2.50 per hour. **Calculate** the mean and standard deviation of the amount of revenue each car generates.

5.2 Joint distribution of two discrete r.vs

The function $f(x, y)$ is a **joint probability distribution** or **probability mass function** of the discrete random variables X and Y if

1. $f(x, y) \geq 0$ for all (x, y) ,
2. $\sum_x \sum_y f(x, y) = 1$,
3. $P(X = x, Y = y) = f(x, y)$

5.2.1 Marginal distribution X and Y (discrete)

The marginal distributions of X alone and of Y alone are

1. $f_X(x) = \sum_y f(x, y)$
2. $f_Y(y) = \sum_x f(x, y)$

5.2.2 Stochastic independence of Jointly Distributed Random Variables

If $f(x, y) = f_X(x) \times f_Y(y)$ for all x and y then the random variables X and Y will said to be independent.

5.2.3 Covariance and correlation between X and Y

i Covariance

$$Cov(X, Y) = \sigma_{XY} = E[(X - \mu_X)(Y - \mu_Y)]$$

In other way,

$$Cov(X, Y) = \sigma_{XY} = E(XY) - \mu_X \mu_Y$$

i Correlation coefficient

$$\rho = \frac{\sigma_{XY}}{\sigma_X \sigma_Y} \quad ; -1 \leq \rho \leq +1$$

5.2.4 Laws of Expected Value and Variance of the Linear combination of Two Variables

Suppose a new random variable is Z as follows:

$$Z = aX + bY$$

Where a and b are both constants.

1. $E(Z) = E(aX + bY) = aE(X) + bE(Y)$,
2. $Var(Z) = Var(aX + bY) = a^2Var(X) + b^2Var(Y) + 2ab Cov(X, Y)$

N.B: If X and Y are independent, $Cov(X, Y) = 0$.

5.2.5 Some problems on discrete joint distribution

Problem 1 The joint probability distribution of X and Y is shown in the following table.

		y	
		1	2
x	1	0.3	0.2
	2	0.1	0.4

- a. Determine the marginal probability distribution of X and Y .
- b. Find $P(Y = 1|X = 2)$.
- c. Compute $E(X)$, $E(Y)$ and $Var(X)$, $Var(Y)$.
- d. Verify whether X and Y are independent or not.
- e. Compute $E(XY)$.
- f. Compute $Cov(X, Y)$ and correlation coefficient between X and Y .
- g. Derive the probability distribution of $Z = X + Y$.

Problem 7.5.1 The joint probability distribution of X and Y is shown in the following table.

y	x		
	1	2	3
1	.42	.12	.06
2	.28	.08	.04

- a. Determine the marginal distributions of X and Y .
- b. Compute the covariance and coefficient of correlation between X and Y .
- c. Develop the probability distribution of $X + Y$.
- d. Find $P(X + Y \leq 3)$.

Problem 7.5.2 After analyzing several months of sales data, the owner of an appliance store produced the following joint probability distribution of the number of refrigerators and stoves sold daily.

Stoves	Refrigerators		
	0	1	2
0	.08	.14	.12
1	.09	.17	.13
2	.05	.18	.04

- Find the marginal probability distribution of the number of refrigerators sold daily.
- Find the marginal probability distribution of the number of stoves sold daily.
- Compute the mean and variance of the number of refrigerators sold daily.
- Compute the mean and variance of the number of stoves sold daily.
- Compute the covariance and the coefficient of correlation.

In the following sections we will discuss some commonly used discrete probability distributions which are used to predict number of success in finite number of random trials, or number of occurrence in a given interval or space and so on.

5.3 Bernoulli distribution/r.v

Bernoulli r.v comes from **Bernoulli trial**-a trial which has **TWO** possible outcomes (*success* or *failure*).

Consider the toss of a **biased coin**, which comes up a head with probability p , and a tail with probability $1 - p$. The **Bernoulli random variable** takes the two values 1 and 0, depending on whether the outcome is a head or a tail:

$$X = \begin{cases} 1, & \text{if HEAD appears,} \\ 0, & \text{if TAIL appears} \end{cases}$$

PMF: $P(X = x) = f(x) = p^x(1 - p)^{1-x}; \quad x = 0, 1$

Mean: $E(X) = p$

Variance: $Var(X) = p(1 - p)$

For all its simplicity, the Bernoulli random variable is very important. In practice, it is used to model generic probabilistic situations with just two outcomes, such as:

- The state of a telephone at a given time that can be either free or busy.
- A person who can be either healthy or sick with a certain disease.

- (c) The preference of a person who can be either for or against a certain political candidate.

Furthermore, by combining multiple Bernoulli random variables, one can construct more complicated random variables.

i Note

Derivation of Mean and Variance of Bernoulli r.v

Mean:

$$E(X) = \sum_{x=0}^1 x \cdot f(x) = (0)f(0) + (1)f(1) = 0 + 1 \cdot p = p$$

Variance:

$$Var(X) = E(X^2) - [E(X)]^2 = p - p^2 = p(1 - p)$$

5.4 Binomial r.v

In a Binomial experiment, the **Bernoulli trial** is repeated n times with the following conditions:

- a) The trials are independent
- b) In each trial $P(\text{success}) = p$ remains constant

Suppose $X = \text{number of successes in } n \text{ trials}$. Then X is called a **Binomial r.v** or follows **Binomial distribution**.

PMF:

$$P(X = x) = f(x) = \binom{n}{x} p^x (1 - p)^{n-x}; x = 0, 1, 2, \dots, n$$

Mean: $E(X) = np$

Variance: $Var(X) = np(1 - p)$

We write $X \sim \text{Bin}(n, p)$

i Note

If $Y = \text{number of failures in } n \text{ trials}$ then
 $Y \sim \text{Bin}(n, 1 - p)$

i Relation between Bernoulli r.v and Binomial r.v

A Binomial Random Variable Is a Sum of Bernoulli Random Variables

Let, Y_i is a Bernoulli r.v appeared in i^{th} Bernoulli trial. If we conduct n independent Bernoulli trials then we have n independent Bernoulli r.v.s such as $Y_1, Y_2, \dots, Y_n$. Each Y_i has values of either 1 or 0.

Now if X is a Binomial r.v then,

$$X = Y_1 + Y_2 + \dots + Y_n = \sum_{i=1}^n Y_i$$

i Note

Derivation of Mean and Variance of Binomial r.v

From previous note, we know if Y_i is a Bernoulli r.v then

$$E(Y_i) = p \text{ and } Var(Y_i) = p(1 - p)$$

So, the mean of Binomial r.v that is

$$E(X) = E(Y_1 + Y_2 + \dots + Y_n)$$

$$= E(Y_1) + E(Y_2) + \dots + E(Y_n)$$

$$= p + p + \dots + p = np$$


Now, the variance of X is:

$$Var(X) = Var(Y_1 + Y_2 + \dots + Y_n)$$

$$= Var(Y_1) + Var(Y_2) + \dots + Var(Y_n)$$

$$= p(1 - p) + p(1 - p) + \dots + p(1 - p) = np(1 - p)$$

Probability plot of binomial r.v for different values of p and shape characteristics



Example 5.4 Consider a binomial experiment with $n = 10$ and $p = 0.30$.

- Compute $P(X = 0)$;
- Compute $P(X = 2)$;
- Compute $P(X \leq 1)$;
- Compute $P(X \geq 2)$;

e) Compute $E(X)$; f) Compute $Var(X)$ and σ .

Example 5.5 A manufacturer of window frames knows from long experience that 30 percent of the production will have some type of minor defect that will require an adjustment. What is the probability that in a sample of 20 window frames:

- a) none will need adjustment?
- b) at most two will need adjustment?
- c) at least two will need adjustment?
- d) Estimate the mean and standard deviation of number of adjustment.

Example 5.6 A certain type of tomato seed germinates 90% of the time. A backyard farmer planted 25 seeds.

- a) What is the probability that exactly 20 germinate?
- b) What is the probability that 23 or more germinate?
- c) What is the probability that 24 or fewer germinate?
- d) What is the expected number of seeds that germinate?

Example 5.7 A shoe store's records show that 30% of customers making a purchase use a credit card to pay. This morning, 10 customers purchased shoes from the store. Answer the following:

- a) Find the probability that at least 8 of the customers used a credit card.
- b) What is the probability that at least three customers, but not more than five, used a credit card?
- c) What is the expected number of customers who used a credit card? What is the standard deviation?
- *d) Find the probability that exactly 5 customers *did not use* a credit card.
- *e) Find the probability that at least nine customers *did not use* a credit card

5.5 Poisson r.v

In this section we consider a discrete random variable that is often useful in estimating the number of occurrences over a specified interval of time or space. *For example*, the random variable of interest might be

- the *number of arrivals* at a car wash in one hour,
- the *number of repairs* needed in 10 miles of highway, or
- the *number of leaks* in 100 miles of pipeline.

PROPERTIES OF A POISSON EXPERIMENT

1. The probability of an occurrence is the same for any two intervals of equal length.
2. The occurrence or nonoccurrence in any interval is independent of the occurrence or nonoccurrence in any other interval.

Suppose X be the number occurrences in a given **interval**. Then,

PMF:

$$P(X = x) = f(x) = \frac{\mu^x e^{-\mu}}{x!} \quad ; \quad x = 0, 1, 2, \dots, \infty$$

Where, μ is the expected value or mean number of occurrences in an interval.

Mean: $E(X) = \mu$

Variance: $Var(X) = \mu$

We write, $X \sim Pois(\mu)$

5.5.1 Finding probability of Poisson r.v

Let, X be a Poisson r.v with $\mu = 2.5$. Find the following probabilities using PMF:

- i) $P(X = 2)$
- ii) $P(X \leq 1)$
- iii) $P(X > 3)$

5.5.2 The recursion formula for Poisson PMF

If $X \sim Pois(\mu)$ then

$$P(X = k) = \frac{e^{-\mu} \mu^k}{k!} \text{ and}$$

$$P(X = k - 1) = \frac{e^{-\mu} \mu^{k-1}}{(k-1)!}$$

Hence we have

$$\frac{P(X = k)}{P(X = k - 1)} = \frac{\mu}{k}$$

$$\text{Hence, } P(X = k) = \frac{\mu}{k} P(X = k - 1)$$

So, for $k = 1$, $P(X = 1) = \frac{\mu}{1} P(X = 0)$ Or,

$$f(1) = \frac{\mu}{1} f(0)$$

$$f(2) = \frac{\mu}{2} f(1)$$

...

and so on.

Example 911 Calls. Emergency 911 calls to a small municipality in Idaho come in at the rate of one every 2 minutes. (Anderson 2020a, page no. 261)

- a. What is the expected number of 911 calls in one hour?
- b. What is the probability of three 911 calls in five minutes?
- c. What is the probability of no 911 calls in a five-minute period?

Example Airport Passenger-Screening Facility. Airline passengers arrive randomly and independently at the passenger-screening facility at a major international airport. The mean arrival rate is 10 passengers per minute. (Anderson 2020a, page no. 261)

- a. Compute the probability of no arrivals in a one-minute period.
- b. Compute the probability that three or fewer passengers arrive in a one-minute period.
- c. Compute the probability of no arrivals in a 15-second period.
- d. Compute the probability of at least one arrival in a 15-second period.

Example Customers arrive at a busy checkout counter at an average rate of 3 per minute. If the distribution of arrivals is Poisson, find the probability that in any given minute there will be 2 or fewer arrivals.

Example The number of accidents in a production facility has a Poisson distribution with a mean of 2.6 per month.

- a. For a given month what is the probability there will be fewer than 2 accidents?
- b. For a given month what is the probability there will be more than 3 accidents?

5.5.3 Poisson Approximation to the Binomial Distribution

When,

- $p \rightarrow 0$ (*Success rate is very low*);
- $n \rightarrow \infty$ (*Number of trials is very large*);

Then **Binomial distribution** can be *approximated* by **Poisson distribution**.

- Mathematically, $\text{Bin}(x; n, p) \approx \text{Pois}(x; \mu)$; where $\mu = np$.

i Note

In practical situation if $n > 20$ and $np \leq 7$; then the approximation is close enough to use the Poisson distribution for binomial problems(Black 2012).

Example A college has 250 personal computers. The probability that any 1 of them will require repair in a given week is 0.01. Find the probability that fewer than 3 of the personal computers will require repair in a particular week. Use the Poisson approximation to the binomial distribution.

Example It is estimated that 0.5 percent of the callers to the Customer Service department of Dell Inc. will receive a busy signal. What is the probability that of today's 1,200 callers at least 3 received a busy signal?

Example Ms. Bergen is a loan officer at Coast Bank and Trust. From her years of experience, she estimates that the probability is .025 that an applicant will not be able to repay his or her installment loan. Last month she made 40 loans.

- a. What is the probability that 3 loans will be defaulted?
- b. What is the probability that at least 3 loans will be defaulted?

5.6 Multinomial Probability Distribution

Suppose that a random experiment consists of a sequence of n trials. Assume that

- (1) The result of each trial is classified into one of k categories.
- (2) The probability of a trial generating a result in category 1, category 2, ... , category k is constant over the trials and equal to $p_1, p_2, \dots, p_k$, respectively.
- (3) The trials are independent.

The random variables $X_1, X_2, \dots, X_k$ that denote the number of trials that result in class 1, class 2, ... , class k , respectively, have a multinomial distribution and the joint probability mass function is

$$P(X_1 = x_1, X_2 = x_2, \dots, X_k = x_k) = \frac{n!}{x_1!x_2!\dots x_k!} p_1^{x_1} p_2^{x_2} \dots p_k^{x_k}$$

for $x_1 + x_2 + \dots + x_k = n$ and $p_1 + p_2 + \dots + p_k = 1$.

Example The probabilities are 0.4, 0.2, 0.3, and 0.1, respectively, that a delegate to a certain convention arrived by air, bus, automobile, or train. What is the probability that among 9 delegates randomly selected at this convention, 3 arrived by air, 3 arrived by bus, 1 arrived by automobile, and 2 arrived by train?

Example According to a genetics theory, a certain cross of guinea pigs will result in red, black, and white offspring in the ratio 8:4:4. **Find** the probability that among 8 offspring, 5 will be red, 2 black, and 1 white.

6 Continuous Probability Distributions

6.1 Probability density function (PDF)

Definition: A continuous r.v X must have a probability density function (PDF) $f(x)$ such that

1) $f(x) \geq 0$ [Non-negativity]

2) $\int_{x \in \mathbb{R}} f(x)dx = 1$ [Total AREA under the curve $f(x)$ always 1]

6.1.1 Illustration with an example

Given $f(x) = \frac{1}{2}x$; $0 \leq x \leq 2$

a) Show/plot the graph of $f(x)$.


b) Is $f(x)$ a PDF?

c) Find $P(X < 1.0)$.

d) Find $P(X = 1.0)$

Solution:

(a)




b) Here, $f(x) \geq 0$ for all values of x in the interval $0 \leq x \leq 2$.

Now, **total area under curve** $f(x)$ from $x = 0$ to $x = 2$ is

$$\int_0^2 f(x)dx$$

= AREA of the SHADED Triangle



$$= \frac{1}{2} \times \text{base} \times \text{height}$$

$$= \frac{1}{2} \times 2 \times 1 = 1$$


So, total area under curve $f(x)$ is 1 that is $\int_0^2 f(x)dx = 1$.

Hence, $f(x)$ is a PDF.

c) Here,

$P(X < 1) = \text{Area under the curve from } x = 0 \text{ to } x = 1$

$= \text{Area of the SHADED Triangle}$



$$= \frac{1}{2} \times 1 \times f(1) = \frac{1}{2} \times 1 \times 0.5 = 0.25$$

Therefore $P(X < 1) = 0.25$

d) $P(X = 1.0) = 0$ [Because there is no area at $x = 1.0$]

i Note

We always remember that **Probability in an interval of X is actually the AREA under the pdf $f(x)$.**

Problem 6.2.1 A random variable has the following density function.

$$f(x) = 1 - 0.5x \quad ; \quad 0 < x < 2$$

- a) Graph the density function.
- b) Verify that $f(x)$ is a density function.
- c) Find $P(X > 1)$.
- d) Find $P(X < 0.5)$.
- e) Find $P(X = 1.5)$.

Problem 6.2.2 The following function is the density function for the random variable X :

$$f(x) = \frac{x-1}{8} \quad ; \quad 1 < x < 5$$

- a) Graph the density function.
- b) Find the probability that X lies between 2 and 4.
- c) What is the probability that X is less than 3?

6.1.2 Expectation and variance of continuous r.v

If X is a continuous r.v with PDF $f(x)$ then

Expected value of X is

$$\mu = E(X) = \int_{x \in \mathbb{R}} x \cdot f(x) dx$$

Variance of X is

$$Var(X) = E(X^2) - \mu^2 = \int_{x \in \mathbb{R}} x^2 \cdot f(x) dx - \mu^2$$




Figure 6.1: Graph of $f(x)$

6.2 Uniform probability distribution/r.v

A continuous r.v X is said to be uniform r.v ranges between a to b if it has the following PDF

$$f(x) = \frac{1}{b-a} \quad ; \quad a < x < b \quad (6.1)$$

with

Mean: $\mu = E(X) = \frac{a+b}{2}$

Variance: $\sigma^2 = \frac{(b-a)^2}{12}$

We write, $X \sim U(a, b)$

6.2.1 Finding probability for uniform r.v

If $X \sim U(a, b)$ then the $P(x_1 < X < x_2)$ is actually the **area of the shaded rectangle**.




Figure 6.2: Computing area for an interval of Uniform distribution

That is,

$$P(x_1 < X < x_2) = \text{Base} \times \text{Height} = (x_2 - x_1) \times \frac{1}{b-a}$$

Problem 6.4.1 The random variable X is known to be uniformly distributed between 10 and 30.

- a) Show the graph of the probability density function.
- b) Compute $P(X < 15)$.
- c) Compute $P(X \geq 22)$.
- d) Compute $P(13 \leq X < 23)$.
- e) Compute $P(X = 29)$.
- f) Compute $E(X)$.
- g) Compute $Var(X)$ and $SD(X)$.

Problem 6.4.2 (Keller 2014, 263) The amount of gasoline sold daily at a service station is uniformly distributed with a minimum of 2,000 gallons and a maximum of 5,000 gallons.

- a. Find the probability that daily sales will fall between 2,500 and 3,000 gallons.
- b. What is the probability that the service station will sell at least 4,000 gallons?
- c. What is the probability that the station will sell exactly 2,500 gallons?
- d. What is the **mean** and **standard deviation** of amount of daily gasoline sold? (*)

Problem 6.4.3 (Keller 2014, 265) The weekly output of a steel mill is a uniformly distributed random variable that lies between 110 and 175 metric tons.

- a. Compute the probability that the steel mill will produce more than 150 metric tons next week.
- b. Determine the probability that the steel mill will produce between 120 and 160 metric tons next week.
- c. The operations manager labels any week that is in the bottom 20% of production a “bad week.” How many metric tons should be used to define a bad week? (*)

Problem 6.4.4 (Keller 2014, 265) The amount of time it takes for a student to complete a statistics quiz is uniformly distributed between 30 and 60 minutes. One student is selected at random. Find the probability of the following events.

- a. The student requires more than 55 minutes to complete the quiz.
- b. The student completes the quiz in a time between 30 and 40 minutes.
- c. The student completes the quiz in exactly 37.23 minutes.

Problem 6.4.5 (Keller 2014, 265) Refer to previous problem.

- a. The professor wants to reward (with bonus marks) students who are in the lowest quarter of completion times. What completion time should he use for the cutoff for awarding bonus marks? (*)
- b. The professor would like to track (and possibly help) students who are in the top 10% of completion times. What completion time should he use? (*)

6.3 Normal distribution/r.v

The normal distribution is arguably the most popular and commonly used distribution. It is compatible with a wide range of human attributes, including height, weight, length, speed, IQ, academic success, and years of life expectancy.

A large number of business and industrial variables are also normally distributed. Several variables, such as the annual cost of household insurance, the cost per square foot of warehouse space rental, and managers' happiness with ownership support on a five-point scale, could result in data that are normally distributed. Also, most things that are manufactured or filled by machines are normally distributed.

Due to its numerous uses, the normal distribution is a very significant distribution. In addition to the several variables that are normally distributed that have been described, **statistical inference**, **statistical process control** rely heavily on the normal distribution. No matter the form of the underlying distribution from which they are derived, many statistics are normally distributed when sufficiently large sample sizes are obtained (Black 2012).

6.3.1 Definition

A continuous r.v X is said to be normal r.v if it has the following **PDF**:

$$f(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2} ; -\infty < x < \infty \quad (6.2)$$

The graph of $f(x)$ is called **normal curve** (Figure 6.3).




Figure 6.3: Normal Curve

Mean: $E(X) = \mu$

Variance: $Var(X) = \sigma^2$

We write: $X \sim N(\mu, \sigma^2)$

Properties of normal distribution

- The **total area** under the normal curve $f(x)$ is 1 that is

$$\int_{-\infty}^{\infty} f(x)dx = 1$$

- Normal distribution is symmetric about mean, μ
- Mean, median and mode is identical in normal distribution that is $Mean = Median = Mode = \mu$
- Almost 99% observations of X lie within **3 standard deviation of mean** that is

$$P(\mu - 3\sigma < X < \mu + 3\sigma) \approx 0.99$$

- Almost 95% observations of X lie within **2 standard deviation of mean** that is

$$P(\mu - 2\sigma < X < \mu + 2\sigma) \approx 0.95$$

- Almost 68% observations of X lie within **1 standard deviation of mean** that is

$$P(\mu - \sigma < X < \mu + \sigma) \approx 0.68$$

6.3.2 Standard normal r.v

Suppose $X \sim N(\mu, \sigma^2)$. Then the variable $Z = \frac{X-\mu}{\sigma}$ is said to be **standard normal variable** with **PDF**

$$f(z) = \frac{1}{\sqrt{2\pi}} e^{-\frac{z^2}{2}} \quad ; -\infty < z < \infty \quad (6.3)$$

Mean: $E(Z) = 0$

Variance: $Var(Z) = 1$

We write: $Z \sim N(0, 1)$

6.3.3 Computing probability(area) under standard normal curve

To compute area (probability) under the standard normal curve for a given interval of z we use **standard Normal Distribution table** which provides cumulative probabilities.

RULE-I: Suppose we want to find $P(Z < 1.25)$.

From TABLE 1 (Appendix B) in Anderson (2020a) we have

$$P(Z < 1.25) = 0.8944$$




Figure 6.4: Standard normal curve

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990

The probability $P(Z < 1.25)$ is shown in Figure 6.5.

RULE-II: Now we find $P(Z > 1.36)$

So, due to symmetry we can write $P(Z > 1.36) = P(Z < -1.36) = 0.0869$

RULE-III: Let us evaluate $P(-1.96 < Z < 2.58)$.

We can write

$$\begin{aligned}
 &= P(-1.96 < Z < 2.58) \\
 &= P(Z < 2.58) - P(Z < -1.96)
 \end{aligned}$$




Figure 6.5: Area under standard normal curve for $Z < 1.25$




Figure 6.6: Area under standard normal curve for $Z > 1.36$

$$= 0.9951 - 0.0250 = 0.9701$$




Figure 6.7: Area under standard normal curve for $-1.96 < Z < 2.58$

6.3.4 Finding quantiles (percentiles, quartiles, deciles etc) of Z

What is the 90th percentile of Z ? To answer this question, let k is the 90th percentile of Z . So we can write

$$P(Z < k) = 0.90 \quad \dots (1)$$

From TABLE 1 (Appendix-B) (Anderson 2020a) we have

$$P(Z < 1.28) = 0.90 \quad \dots (2)$$

Comparing eq.(1) with eq.(2) we have $k = 1.28$. So the 90th percentile of Z is 1.28.

Problem 1 Find c such that $P(Z > c) = 0.05$.

Problem 2 Find c such that $P(-c < Z < c) = 0.95$.

6.3.5 Computing probability(area) under normal curve:

Suppose $X \sim N(30, 5^2)$. Then find the following:

- $P(X < 22)$
- $P(X > 44)$
- $P(20 < X < 35)$
- If $P(X < x) = 0.25$ then find the value of x .

Solution:

Here, $\mu = 30$ and $\sigma = 5$

$$\text{a) } P(X < 22) = P\left(\frac{X-\mu}{\sigma} < \frac{22-30}{5}\right) = P(Z < -1.60) = 0.0548.$$

$$\begin{aligned}\text{b) } P(X > 44) &= P\left(\frac{X-\mu}{\sigma} > \frac{44-30}{5}\right) \\ &= P(Z > 2.80) = P(Z < -2.80) = 0.0026\end{aligned}$$

$$\begin{aligned}\text{c) } P(20 < X < 35) &= P\left(\frac{20-30}{5} < \frac{X-\mu}{\sigma} < \frac{35-30}{5}\right) \\ &= P(-2 < Z < 1) = P(Z < 1) - P(Z < -2) \\ &= 0.8413 - 0.0228 = 0.8185\end{aligned}$$

d) To find the value of x we proceed this way.

$$P(X < x) = 0.25$$

$$\Rightarrow P\left(\frac{X-\mu}{\sigma} < \frac{x-30}{5}\right) = 0.25$$

$$\Rightarrow P\left(Z < \frac{x-30}{5}\right) = 0.25 \quad \dots (1)$$

From TABLE (Appendix B) we have

$$P(Z < -0.67) = 0.25 \quad \dots (2)$$

Comparing (1) with (2) we can write

$$\frac{x-30}{5} = -0.67$$

$$\Rightarrow x = 30 + (-0.67) \times 5$$

$$\therefore x = 26.65$$

i Note

If $P(X < x) = p$ and
 $P(Z < z) = p$ then

$$x = \mu + z\sigma$$

6.3.6 Applications

In this section we will discuss about some problems which are connected to the normal distribution.

Problem 6.5.1 (Anderson 2020a, 298) Automobile repair costs continue to rise with an average 2015 cost of \$367 per repair (U.S. News & World Report website). Assume that the cost for an automobile repair is normally distributed with a standard deviation of \$88. Answer the following questions about the cost of automobile repairs.

- What is the probability that the cost will be more than \$450?
- What is the probability that the cost will be less than \$250?
- What is the probability that the cost will be between \$250 and \$450?
- If the cost for your car repair is in the lower 5% of automobile repair charges, what is your cost?

Problem 6.5.2 (Anderson 2020a, 298) Labor Day Travel Costs. The American Automobile Association (AAA) reported that families planning to travel over the Labor Day weekend spend an average of \$749. Assume that the amount spent is normally distributed with a standard deviation of \$225.

- What is the probability of family expenses for the weekend being less than \$400?
- What is the probability of family expenses for the weekend being \$800 or more?
- What is the probability that family expenses for the weekend will be between \$500 and \$1000?
- What would the Labor Day weekend expenses have to be for the 5% of the families with the most expensive travel plans?

Problem 6.5.3 (Keller 2014, 280) A new gas-electric hybrid car has recently hit the market. The distance traveled on 1 gallon of fuel is normally distributed with a mean of 65 miles and a standard deviation of 4 miles. Find the probability of the following events.

- The car travels more than 70 miles per gallon.
- The car travels less than 60 miles per gallon.
- The car travels between 55 and 70 miles per gallon.

Problem 6.5.4 (Anderson 2020a, 298) Mensa Membership. A person must score in the upper 2% of the population on an IQ test to qualify for membership in Mensa, the international high-IQ society. If IQ scores are normally distributed with a mean of 100 and a standard deviation of 15, what score must a person have to qualify for Mensa?

Problem 6.5.5 (Keller 2014 , 282) The lifetimes of televisions produced by the Hishobi Company are normally distributed with a mean of 75 months and a standard deviation of 8 months. If the manufacturer wants to have to replace only 1% of its televisions, what should its warranty be?

Problem 6.5.6 (Newbold, Carlson, and Thorne 2013, 218) I am considering two alternative investments. In both cases I am unsure about the percentage return but believe that my uncertainty can be represented by normal distributions with the means and standard deviations shown in the accompanying table. I want to make the investment that is more likely to produce a return of at least 10%. Which investment should I choose?

	Mean	Standard deviation
Investment A	10.4	1.2
Investment B	11.0	4.0

6.3.7 Normal Approximation to the Binomial Distribution

Let $X \sim \text{Bin}(n, p)$. When n is large so that both $np \geq 5$ and $n(1 - p) \geq 5$. We can use the normal distribution to get an approximate answer. Remember to use **continuity correction**.

$X \sim N(\mu = np, \sigma^2 = np(1 - p))$, approx.

Problem 7.8.1 A car-rental company has determined that the probability a car will need service work in any given month is 0.2. The company has 900 cars (Newbold, Carlson, and Thorne 2013).

- (a) What is the probability that more than 200 cars will require service work in a particular month?
- (b) What is the probability that fewer than 175 cars will need service work in a given month?

Problem 7.8.2 The tread life of Stone Soup tires can be modeled by a normal distribution with a mean of 35,000 miles and a standard deviation of 4,000 miles. A sample of 100 of these tires is taken. What is the probability that more than 25 of them have tread lives of more than 38,000 miles? (Newbold, Carlson, and Thorne 2013)

6.3.8 Normal Approximation to the Poisson Distribution

Let $X \sim \text{Poisson}(\mu)$. When μ is large ($\mu > 5$) then the Normal distribution can be used to approximate the Poisson distribution.

$X \sim N(\mu, \mu)$ approx.

Problem 7.8.3 Hits to a high-volume Web site are assumed to follow a Poisson distribution with a mean of 10,000 per day. Approximate each of the following: (Montgomery and Runger 2014)

- (a) Probability of more than 20,000 hits in a day,
- (b) Probability of less than 9900 hits in a day .

6.4 Exponential distribution

Consider an arrival process where events arrive in a constant rate say λ per unit time.

Let, T be a continuous r.v expresses the **time between arrivals**. T will follow **Exponential distribution** with the PDF:

$$f(t) = \lambda e^{-\lambda t} \quad ; \quad t > 0 \quad (6.4)$$

Where,

- Mean of T : $E(T) = \frac{1}{\lambda}$
- Variance of T : $Var(T) = \frac{1}{\lambda^2}$
- We write, $T \sim Exp(\lambda)$

6.4.1 Probability calculation of Exponential distribution

- Probability (arrival **within** t time) $= P(T < t) = 1 - e^{-\lambda t}$
- Probability (arrival **after** t time) $= P(T > t) = e^{-\lambda t}$

6.4.2 The k^{th} percentile of T

If P_k is the k^{th} percentile of T then

$$P_k = \frac{\ln(1 - \frac{k}{100})}{-\lambda} \quad (6.5)$$

Problem 7.8.4 Waiting times to receive food after placing an order at the local Subway sandwich shop follow an exponential distribution with a mean of 60 seconds. **Calculate** the probability a customer waits:

- Less than 30 seconds.
- More than 120 seconds.
- Between 45 and 75 seconds
- Find** the mean and median waiting time

Problem 7.8.5 The lifetime of plasma and LCD TV sets follows an exponential distribution with a mean of 100,000 hours. **Compute** the probability a television set:

- Fails in less than 10,000 hours.
- Lasts more than 120,000 hours.
- Fails between 60,000 and 100,000 hours of use.
- Find** the 90th percentile. So 10 percent of the TV sets last more than what length of time?

Problem 7.8.6 A checkout counter at a supermarket completes the process according to an exponential distribution with a service rate of 6 per hour. A customer arrives at the checkout counter. Find the probability of the following events.

- The service is completed in fewer than 5 minutes.
- The customer leaves the checkout counter more than 10 minutes after arriving.
- Find the mean and median service time (in minutes)

6.5 Joint distribution of two continuous r.vs

The function $f(x, y)$ is a joint density function of the continuous random variables X and Y if

- $f(x, y) \geq 0$,
- $\int_{-\infty}^{\infty} \int_{-\infty}^{\infty} f(x, y) \, dx \, dy = 1$.

6.5.1 Marginal distribution X and Y (continuous)

The marginal distributions of X alone and of Y alone are

- $f_X(x) = \int_{-\infty}^{\infty} f(x, y) \, dy$,
- $f_Y(y) = \int_{-\infty}^{\infty} f(x, y) \, dx$

6.6 Bivariate Normal distribution

Two random variables X and Y are said to have a **bivariate normal distribution** with parameters $\mu_X, \sigma_X^2, \mu_Y, \sigma_Y^2$, and ρ , if their joint PDF is given by

$$f_{XY}(x, y) = \frac{1}{2\pi\sigma_X\sigma_Y\sqrt{1-\rho^2}} \exp \left\{ -\frac{1}{2(1-\rho^2)} \left[\left(\frac{x-\mu_X}{\sigma_X} \right)^2 + \left(\frac{y-\mu_Y}{\sigma_Y} \right)^2 - 2\rho \frac{(x-\mu_X)(y-\mu_Y)}{\sigma_X\sigma_Y} \right] \right\} \quad (6.6)$$

where $\mu_X, \mu_Y \in \mathbb{R}$, $\sigma_X, \sigma_Y > 0$ and $\rho \in (-1, 1)$ are all constants.

i Theorem

Suppose $X \sim N(\mu_X, \sigma_X^2)$, $Y \sim N(\mu_Y, \sigma_Y^2)$ and the Pearson correlation coefficient is ρ . Consider a new variable $W = aX \pm bY$.

Then the variable W also follows normal distribution with

- Mean, $E(W) = a\mu_X + b\mu_Y$
- Variance, $Var(W) = a^2Var(X) + b^2Var(Y) \pm 2abCov(X, Y)$

Problem 7.1 A random variable X is normally distributed with a mean of 100 and a variance of 100, and a random variable Y is normally distributed with a mean of 200 and a variance of 400. The

random variables have a correlation coefficient equal to -0.5. Find the mean and variance of the random variable:

$$W = 5X + 4Y$$

Problem 7.2 A random variable X is normally distributed with a mean of 500 and a variance of 100, and a random variable Y is normally distributed with a mean of 200 and a variance of 400. The random variables have a correlation coefficient equal to 0.5. Find the mean and variance of the random variable:

$$W = 5X - 4Y$$

Problem 7.3 The nation of Olecarl, located in the South Pacific, has asked you to analyze international trade patterns. You first discover that each year it exports 10 units and imports 10 units of wonderful stuff. The price of exports is a random variable with a mean of 100 and a variance of 100. The price of imports is a random variable with a mean of 90 and a variance of 400. In addition, you discover that the prices of imports and exports have a correlation of $\rho = -0.40$. The prices of both exports and imports follow a normal probability density function. Define the balance of trade as the difference between the total revenue from exports and the total cost of imports.

- What are the mean and variance of the balance of trade?
- What is the probability that the balance of trade is negative?

Problem 7.4 The nation of Waipo has recently created an economic development plan that includes expanded exports and imports. It has completed a series of extensive studies of the world economy and Waipo's economic capability, following Waipo's extensive 10-year educational-enhancement program. The resulting model indicates that in the next year exports will be normally distributed with a mean of 100 and a variance of 900 (in billions of Waipo yuan). In addition, imports are expected to be normally distributed with a mean of 105 and a variance of 625 in the same units. The correlation between exports and imports is expected to be +0.70. Define the trade balance as exports minus imports.

- Determine the mean and variance of the trade balance (exports minus imports) if the model parameters given above are true.
- What is the probability that the trade balance will be positive?

6.7 Some other important Probability Densities

6.7.1 Chi-square (χ^2) distribution

A continuous random variable X is said to follow chi-square distribution if it has the following PDF:

$$f(x) = \frac{1}{2^{\nu/2}\Gamma(\nu/2)} x^{\frac{\nu-2}{2}} e^{-\frac{x}{2}} \quad ; \quad x > 0$$

- $E(X) = \nu$
- $Var(X) = 2\nu$

6.7.2 t-Distribution

6.7.3 F-Distribution

In next chapter we will learn how to derive χ^2 , Student- t and F -distributions using **transformation**

7 Further topics on random variables

7.1 Linear functions of random variables

Given random variables $X_1, X_2, \dots, X_n$ and constants $a_1, a_2, \dots, a_n \in \mathbb{R}$

$$Y = a_1X_1 + a_2X_2 + \dots + a_nX_n$$

is a **linear combination** of $X_1, X_2, \dots, X_n$.

- Mean of Linear function

$$E(Y) = a_1E(X_1) + a_2E(X_2) + \dots + a_nE(X_n)$$

- Variance of Linear function

$$V(Y) = a_1^2V(X_1) + a_2^2V(X_2) + \dots + a_n^2V(X_n) + 2 \sum_{i < j} a_i a_j \text{Cov}(X_i, X_j)$$

- If $X_1, X_2, \dots, X_n$ are **independent** then,

$$V(Y) = a_1^2V(X_1) + a_2^2V(X_2) + \dots + a_n^2V(X_n)$$

If n random variables X_i have common mean μ and common variance σ^2 then,

- $E(Y) = (a_1 + a_2 + \dots + a_n)\mu$
- $\text{Var}(Y) = (a_1^2 + a_2^2 + \dots + a_n^2)\sigma^2$

Average of Independent Random Variables:

$X_1, X_2, \dots, X_n$ are n independent random variables

- $\bar{X} = \frac{X_1 + X_2 + \dots + X_n}{n}$
- $E[\bar{X}] = \frac{1}{n}[E(X_1) + E(X_2) + \dots + E(X_n)]$
- $\text{Var}[\bar{X}] = \frac{1}{n^2}[\text{Var}(X_1) + \text{Var}(X_2) + \dots + \text{Var}(X_n)]$

If n random variables X_i have common mean μ and common variance σ^2 then,

- $E[\bar{X}] = \mu$
- $\text{Var}[\bar{X}] = \frac{\sigma^2}{n}$

7.2 Functions of random variables: Transformations

7.2.1 Transformation of Discrete random variable

7.2.2 Transformation Continuous random variable

- **CDF Method:**

We know $P(X \leq x) = F_X(x)$.

Let, $Y = g(X)$. If $g(X)$ is a **one-to-one** function we will have the inverse function solving for X we will have $X = w(Y)$

By definition

$$F_Y(y) = P(Y \leq y) = P[g(X) \leq y] = P[X \leq w(y)] = F_X[w(y)]$$

$$\therefore f_Y(y) = \frac{d}{dy} F_Y(y) = f_X[w(y)] \cdot \frac{d}{dy} w(y)$$

Example 7.1 Let X be a continuous r.v with the following PDF

$$f(x) = \begin{cases} \frac{x}{12}, & 1 < x < 5, \\ 0, & \text{elsewhere.} \end{cases}$$

Find the probability distribution of the random variable $Y = 3X - 4$.

Here, $Y = 3X - 4$. Solving for X we have $X = \frac{Y+4}{3}$

Now,

$$f_Y(y) = \frac{d}{dy} F_X\left(\frac{y+4}{3}\right) = f_X\left(\frac{y+4}{3}\right) \cdot \frac{1}{3}$$

$$f_Y(y) = \frac{\frac{y+4}{3}}{12} \cdot \frac{1}{3} = \frac{y+4}{108}$$

Hence the PDF of Y is:

$$f_Y(y) = \begin{cases} \frac{y+4}{108}, & -1 < y < 11, \\ 0, & \text{elsewhere.} \end{cases}$$

- **Jacobian Method:**

If $Y = u(X)$ then we will have $x = w(y)$

Then the **Jacobian** $J = \frac{d}{dy}w(y)$. Finally

$$f_Y(y) = f_X[w(y)]|J|$$

Example 7.2 From previous example we have

$$x = \frac{y+4}{3} = w(y).$$

$$\text{So, } J = \frac{d}{dy}w(y) = \frac{1}{3}$$

$$\text{Finally, } f_Y(y) = f_X[w(y)]|J| = \frac{\frac{y+4}{3}}{12} \cdot \frac{1}{3} = \frac{y+4}{108}$$

Example 7.3 If $X \sim N(\mu, \sigma^2)$ then derive the distribution of Z where $Z = \frac{X-\mu}{\sigma}$.

Solution:

We know

$$f_X(x) = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}\left(\frac{x-\mu}{\sigma}\right)^2}$$

Given

$$z = \frac{x - \mu}{\sigma}$$

$$\Rightarrow x = \mu + z\sigma$$

$$\text{Jacobian, } J = \frac{dx}{dz} = \frac{d}{dz}(\mu + z\sigma) = \sigma$$

For $-\infty < x < \infty$; $-\infty < z < \infty$

$$\therefore f_Z(z) = f_X(z) \cdot |J| = \frac{1}{\sigma\sqrt{2\pi}} e^{-\frac{1}{2}z^2} \cdot \sigma = \frac{1}{\sqrt{2\pi}} e^{-\frac{1}{2}z^2}$$

NOTE: CDF and Jacobian are almost same for uni-variate case; but **Jacobian** is more robust when we deal with multiple random variables.

7.3 Moments Moment-Generating Functions (MGF)

7.3.1 Moments

The r^{th} raw moment defined as:

$$\mu'_r = E(X^r) = \begin{cases} \sum x^r f(x), & \text{if } X \text{ is discrete,} \\ \int_{-\infty}^{\infty} x^r f(x), & \text{if } X \text{ is continuous.} \end{cases}$$

For example $\mu = \mu'_1 = E(X)$ and $\mu'_2 = E(X^2)$. So $\sigma^2 = \mu'_2 - \mu^2$.

7.3.2 MGF

$$M_X(t) = E(e^{tX})$$

7.3.3 Finding moments using MGF

$$\left. \frac{d^r M_X(t)}{dt^r} \right|_{t=0} = \mu'_r.$$

MGF of Binomial distribution

$$M_X(t) = E(e^{tX}) = \sum_{x=0}^n e^{tx} \binom{n}{x} p^x q^{n-x} = \sum_{x=0}^n \binom{n}{x} (pe^t)^x q^{n-x} = (pe^t + q)^n$$

Now

$$\frac{dM_X(t)}{dt} = n(pe^t + q)^{n-1} pe^t = np(pe^t + q)^{n-1} e^t$$

Setting $t = 0$ we have $\mu'_1 = np(p + q)^{n-1} \cdot 1 = np \cdot 1 \cdot 1 = np$

MGF of Poisson distribution

Let $X \sim \text{Poisson}(\lambda)$. The PMF is $P(X = x) = \frac{e^{-\lambda} \lambda^x}{x!}$ for $x = 0, 1, \dots$

The MGF is derived as follows:

$$\begin{aligned} M_X(t) &= E[e^{tX}] = \sum_{x=0}^{\infty} e^{tx} \frac{e^{-\lambda} \lambda^x}{x!} \\ &= e^{-\lambda} \sum_{x=0}^{\infty} \frac{(\lambda e^t)^x}{x!} \\ &= e^{-\lambda} e^{\lambda e^t} \quad (\text{using } \sum \frac{z^k}{k!} = e^z) \\ &= e^{\lambda(e^t - 1)}. \end{aligned}$$

MGF of Normal Distribution

MGF of Standard Normal Distribution

MGF of Chi-square Distribution

i Theorem

If Z_i 's are independent standard normal random variables then $U = \sum_{i=1}^k Z_i^2$ has a chi-square distribution with k degrees of freedom.

Proof:

8 Sampling and Sampling distributions

8.1 Some preliminary idea (Anderson 2020a)

- An **element** is the entry on which data are collected.
- A **population** is the collection of all the elements of interest.
- A **sample** is a subset of the population.
- A **sampling frame** is the *list* of all the elements in the population of interest.
- A numerical measure (e.g., mean, median, standard deviation, proportion) computed from population is referred as **population parameter**.
- A numerical measure (e.g., mean, median, standard deviation, proportion) computed from sample is referred as **sample statistic**.

8.2 Sampling from a Finite Population

8.2.1 Simple random sample (Finite population)

A **simple random sample (SRS)** of size n from a *finite* population of size N is a sample selected such that each possible sample of size n has the same probability of being selected.

- Sampling can be *with* replacement (**SWR**).
- Sampling can be *without* replacement (**SWOR**).
- In both case, the probability of a particular element being selected is n/N .

8.3 Sampling from an Infinite Population

In the case of infinite population, it is not possible to develop a sampling from. In that case statisticians recommend selecting a random sample

8.3.1 Random sample (Infinite population)

A **random sample** of size n from an *infinite population* is a sample selected such that the following conditions are satisfied.

1. Each element selected comes from the same population.
2. Each element is selected independently.

Notes and Comments

1) A sample selected randomly from a population (finite or infinite) is referred as a *random sample*. The procedure of selecting a sample randomly is known as *probability sampling*.

2) The *number* of ways simple random samples by **SWR** of size n that can be selected from a finite population of size N is

$$N^n$$

3) The *number* of ways **different** simple random samples by **SWOR** of size n that can be selected from a finite population of size N is

$$\binom{N}{n} = \frac{N!}{n!(N-n)!}$$

4) Some other probability sampling methods are *stratified random sampling*, *cluster sampling*, and *systematic sampling*. We will discuss these methods later.

5) We use the term “simple” in simple random sampling to clarify that this is the probability sampling method that assures each sample of size n has the same probability of being selected.

8.3.2 Selecting simple random sample using R

Suppose we have a variable X and let it contains $N = 4$ elements as follows:

$X = \{3, 5, 7, 9\}$

Using R we can draw several simple random samples of size $n = 2$.

Now we draw a random sample (by using both SWR and SWOR)using `sample()` function in base R.

Sampling without replacement (SWOR)

```
set.seed(2103) # To keep reproducibility
X=c(3,5,7,9) # The elements in X variable

# Drawing a random sample without replacement
sample(X,2, replace = FALSE)
```

```
[1] 5 9
```

The all possible that is $\binom{N}{n} = \binom{4}{2} = 6$ samples (by *SWOR*) are :

Table 8.1: Samples of size n=2 Without replacement

Sample #	Sample	Probability
1	3,5	1/6
2	3,7	1/6
3	3,9	1/6
4	5,7	1/6
5	5,9	1/6
6	7,9	1/6

Sampling with replacement (SWR)

The all possible that is $N^n = 4^2 = 16$ samples (by *SWR*) are :

```
# Drawing a random sample with replacement
sample(X,2, replace = TRUE)
```

```
[1] 9 9
```

```
sampls_swr=expand.grid(X, X)
```

Table 8.2: Samples of size n=2 with replacement

Sample #	Sample	Probability
1	3,3	1/16
2	3,5	1/16
3	3,7	1/16
4	3,9	1/16
5	5,3	1/16
6	5,5	1/16
7	5,7	1/16
8	5,9	1/16
9	7,3	1/16
10	7,5	1/16
11	7,7	1/16
12	7,9	1/16
13	9,3	1/16
14	9,5	1/16
15	9,7	1/16
16	9,9	1/16

8.4 Sampling distribution

The probability distribution of a **sample statistic** is called a **sampling distribution**.

For example, due to sampling variability the **sample mean** $\bar{X}$ has a sampling distribution.

Illustration Consider a population of variable X: 1,3,5,7,9.

Task-1: Compute population mean μ .

Solution:

$$\text{Here, } \mu = \frac{\sum x}{N} = \frac{1+3+\dots+9}{5} = 5$$

Task-2: Draw all possible samples (without replacement) of size $n = 2$ from this population. Then compute the means of all samples.

Solution:

Table 8.3: All Samples of Size 2 and Their Means

Sample	Sample mean, $\bar{x}$
1,3	2
1,5	3
1,7	4
1,9	5
3,5	4
3,7	5
3,9	6
5,7	6
5,9	7
7,9	8

Task-3: Construct a probability distribution of sample means, $\bar{X}$ (discrete) and **plot** it.

Solution:

Table 8.4: Sampling distribution of $\bar{x}$

$\bar{x}$	$f(\bar{x})$
2	$\frac{1}{10}$
3	$\frac{1}{10}$
4	$\frac{2}{10}$
5	$\frac{2}{10}$
6	$\frac{2}{10}$
7	$\frac{1}{10}$
8	$\frac{1}{10}$

Task-4: Find the $E(\bar{X})$. Does $E(\bar{X})$ same as population mean μ ?

Solution:




Figure 8.1: Sampling distribution of $\bar{x}$

$$E(\bar{X}) = \sum \bar{x} \cdot f(\bar{x})$$

$$= 2(1/10) + 3(1/10) + 4(2/10) + \dots + 8(1/10) = 5$$

We can see that $E(\bar{X}) = 5$ is same as $\mu = 5$.

NOTE: This phenomenon is known as the **unbiasedness** of a sample statistic or an estimator. We will discuss it briefly in next chapter.

Home work: Consider a population of variable X: 3,6,9,12,15.

- i) **Compute** population mean μ .
- ii) **Draw** all possible samples of size $n = 3$ from this population without replacement. Then compute the means of all samples.
- iii) **Construct** a probability distribution of sample mean, $\bar{x}$ (discrete) and **plot** it.
- iv) **Find** the $E(\bar{X})$. Does $E(\bar{X})$ same as population mean μ ?

8.5 Sampling distribution of $\bar{X}$

The sampling distribution of $\bar{X}$ is the probability distribution of all possible values of the sample mean $\bar{X}$.

- Expected value of $\bar{X}$:

$$E(\bar{X}) = \mu_{\bar{X}} = \mu$$

- Standard deviation of $\bar{X}$:

CASE I: Sampling without replacement (**SWR**) or if N is **infinite** (or finite but $\frac{n}{N} \leq 0.05$)

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}}$$

CASE II: Sampling without replacement (**SWOR**) and N is **finite**:

$$\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}} \sqrt{\frac{N-n}{N-1}}$$

Standard deviation of $\bar{X}$ is also known as **Standard error of $\bar{X}$** or $s.e(\bar{X})$.

But what is the **form** of the sampling distribution of $\bar{X}$?

8.5.1 Central limit theorem (CLT)

The sampling distribution of the mean of a random sample drawn from any population is approximately normal for a sufficiently large sample size. The larger the sample size, the more closely the sampling distribution of $\bar{X}$ will resemble a normal distribution.

i The Central Limit Theorem

Let, $X_1, X_2, \dots$ be a sequence of independently and identically distributed random variables with common mean μ and common variance σ^2 . We define

$$Z = \frac{\bar{X} - \mu_{\bar{X}}}{\sigma_{\bar{X}}}$$

Then the Z will be approximately normally distributed as the sample size $n \rightarrow \infty$.

The definition of “sufficiently large” depends on the extent of non-normality of X . Some authors consider a sample will be sufficiently large if $n \geq 30$ (Walpole et al. 2017).


8.5.2 Central Limit Theorem through simulation

In this section we illustrates how sampling distributions of sample means approximate to normal or bell shaped distribution as we increase the sample size .

At first, we consider a population data regarding **gdp per capita (USD)**, 2023 of 218 countries. We can see that the distribution of **gdp per capita** is highly skewed to the right (see Figure 8.2).


Now we draw 1000 random samples (without replacement) of different sample sizes and then plot the histogram of samples means.

From Figure 8.3 we can see that as the sample size increases, the sampling distribution of **sample mean** tends to bell-shaped or normal though the population data was very skewed to the right. This simulation clearly demonstrate the fact of Central Limit Theorem (CLT).



Source: World Bank, 2023

Figure 8.2: Frequency histogram of GDP percapita of N=218 countries



- (a) Sampling distribution of sample mean for sample size $n=10$ (b) Sampling distribution of sample mean for sample size $n=30$ (c) Sampling distribution of sample mean for sample size $n=100$

Figure 8.3: Demonstration of Central Limit Theorem through simulation

For more interactive simulation of **CLT** please [click here to visit the ShinyApp for Central Limit Theorem Simulation](#).

Problem 8.1 The foreman of a bottling plant has observed that the amount of soda in each 32-ounce bottle is actually a normally distributed random variable, with a mean of 32.2 ounces and a standard deviation of .3 ounce (Keller 2014, 308).

- a. If a customer buys one bottle, what is the probability that the bottle will contain more than 32 ounces?
- b. If a customer buys a carton of four bottles, what is the probability that the mean amount of the four bottles will be greater than 32 ounces?

Problem 8.2 Suppose a subdivision on the southwest side of Denver, Colorado, contains 2215 houses. The subdivision was built in 1983. A sample of 100 houses is selected randomly and evaluated by an appraiser. If the mean appraised value of a house in this subdivision for all houses is \$177,000, with a standard deviation of \$8,500, what is the probability that the sample average is greater than \$185,000? (Black 2012, 243 (population size is changed))

Problem 8.3 A scientist is studying the heights of men in Australia. The true population mean μ is unknown but the true population standard deviation is assumed to be 2.5 inches. Suppose the scientist randomly samples 100 men. **Find** the probability that the difference between the sample mean and the true population mean is less than 0.5 inches.

Problem 8.4 In rainy season, it tends to rain a lot in Dhaka. Suppose that the amount of rain that falls on any given winter day in Dhaka is normally distributed with a mean of 2.3mm and a variance of 1.21 mm^2 . Assume rainfalls are independent from day to day.

- (a) **Find** the probability that between 1.9 and 3.4 mm of rain fell today.
- (b) **Find** the probability that the total amount of rain that falls over the next 20 days is between 54.3 and 57.1 mm.

Problem 8.5 Suppose, we load on a plane 100 packages whose weights are independent random variables that are uniformly distributed between 5 and 50 pounds. What is the probability that the total weight will exceed 3000 pounds?

8.6 Sampling distribution of sample proportion, $\hat{p}$

The sample proportion $\hat{p}$ is the point estimator of the population proportion p . The formula for computing the sample proportion is

$$\hat{p} = \frac{x}{n}$$

Where,

x = number of *successes* in the sample of size n .

Case-I (small sample): If $X \sim \text{Bin}(n, p)$ then $\hat{p}$ also follows **binomial distribution** with

$$\text{Mean : } E(\hat{p}) = p$$

$$\text{Variance : } \sigma_{\hat{p}}^2 = \frac{p(1-p)}{n}$$

Case-II (large sample): When the sample size is large enough so that np and $n(1-p)$ are greater than or equal to 5 then $\hat{p}$ will be approximately normally distributed with

$$\text{Mean : } E(\hat{p}) = p$$

$$\text{Variance : } \sigma_{\hat{p}}^2 = \frac{p(1-p)}{n}$$

Problem 8.6 According to the Internal Revenue Service, 75% of all tax returns lead to a refund. A random sample of 100 tax returns is taken.

- What is the mean of the distribution of the sample proportion of returns leading to refunds?
- What is the variance of the sample proportion?
- What is the standard error of the sample proportion?
- What is the probability that the sample proportion exceeds 0.8?

Problem 8.7 A random sample of 270 homes was taken from a large population of older homes to estimate the proportion of homes with unsafe wiring. If, in fact, 20% of the homes have unsafe wiring, what is the probability that the sample proportion will be between 16% and 24%?

8.7 Sampling Distribution of the Sample Variances

Like as sample mean, **sample variance** is also considered as a random variable due to sampling variability.

If we have a random sample $\{X_1, X_2, \dots, X_n\}$ of size n then the quantity

$$S^2 = \frac{1}{n-1} \sum_{i=1}^n (X_i - \bar{X})^2$$

is called the **sample variance**.

i Sampling Distribution of the Sample Variances

If S^2 is the variance of a random sample of size n taken from a normal population having the variance σ^2 , then the statistic

$$\chi^2 = \frac{(n-1)S^2}{\sigma^2} = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{\sigma^2}$$

has a **Chi-squared distribution** with $\nu = n - 1$ degrees of freedom.

Mean of S^2 : $E(S^2) = \sigma^2$

Variance of S^2 : $Var(S^2) = \frac{2\sigma^4}{(n-1)}$

How to determine the area under the curve of χ^2 distribution?

In every statistics textbook area under the χ^2 distribution can be determined for a given **degrees of freedom**. The distribution is defined for only positive values, since variances are all positive values. For a given probability or area say α and degrees of freedom ν we can determine the value of χ^2 to the upper tail such that:

$$P(\chi^2 > \chi_\alpha^2) = \alpha$$




Figure 8.4: The chi-squared distribution

For example, when $\alpha = 0.05$ and $\nu = 10$ the value of χ_α^2 is 18.307.

Problem 8.8 A random sample of size $n = 18$ is obtained from a normally distributed population with a population mean of $\mu = 46$ and a variance of $\sigma^2 = 50$.

- What is the probability that the sample mean is greater than 50?
- What is the value of the sample variance such that 5% of the sample variances would be less than this value?
- What is the value of the sample variance such that 5% of the sample variances would be greater than this value?

Problem 8.9 A process produces batches of a chemical whose impurity concentrations follow a normal distribution with a variance of 1.75. A random sample of 20 of these batches is chosen. Find the probability that the sample variance exceeds 3.10.

Solution:

Let, X be the impurity concentration

Given, $\sigma^2 = 1.75$; $n = 20$. We have to compute

Degrees of Freedom	Area in Upper Tail									
	.995	.99	.975	.95	.90	.10	.05	.025	.01	.005
1	.000	.000	.001	.004	.016	2.706	3.841	5.024	6.635	7.879
2	.010	.020	.051	.103	.211	4.605	5.991	7.378	9.210	10.597
3	.072	.115	.216	.352	.584	6.251	7.815	9.348	11.345	12.838
4	.207	.297	.484	.711	1.064	7.779	9.488	11.143	13.277	14.860
5	.412	.554	.831	1.145	1.610	9.236	11.070	12.832	15.086	16.750
6	.676	.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.647	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.300
13	3.565	4.107	5.009	5.892	7.041	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.041	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.558
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.878	14.573	16.151	18.114	36.741	40.113	43.195	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.994
29	13.121	14.256	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.335

Figure 8.5: Chi-square distribution. Source: Appendix B, TABLE 3 (Anderson 2020)

$$P[s^2 > 3.10] = P\left[\frac{(n-1)s^2}{\sigma^2} > \frac{(20-1)(3.10)}{1.75}\right]$$

$$= P[\chi^2 > 33.657] \approx 0.01$$

So there is approximately 1% chance that the sample variance exceeds 3.10.

8.8 t-Distribution

Let $Z \sim N(0, 1)$ and $V \sim \chi_\nu^2$. If Z and V are independent then the random variable

$$T = \frac{Z}{\sqrt{V/\nu}}$$

said to have a *Student-t distribution with ν degrees of freedom*. The PDF of T is

$$f(t) = \frac{\Gamma[(\nu+1)/2]}{\sqrt{\pi\nu} \Gamma(\nu/2)} \left(1 + \frac{t^2}{\nu}\right)^{-(\nu+1)/2}; -\infty < t < \infty.$$

i Theorem

Given a random sample of n observations, with sample mean $\bar{X}$ and sample standard deviation S , from a normally distributed population with mean μ , the random variable T follows the *Student's t distribution* with $\nu = (n-1)$ degrees of freedom and is given by

$$T = \frac{\bar{X} - \mu}{S/\sqrt{n}}$$

Properties:

- 1) Symmetry: t -distribution is symmetric about mean (zero). So
if $P(T > t_\nu) = \alpha$ then $P(T < -t_\nu) = \alpha$.
- 2) Convergence to Normal: As $n \rightarrow \infty$ then the distribution of T_ν approaches the standard Normal distribution.
- 3) Cauchy as special case: The T_1 distribution is the same as the Cauchy distribution.

How to determine the area under the curve of t - distribution?


From t -distribution table we can determine the value of t for a given *area* and *degrees of freedom*.

For example, with $n = 11$ and area in upper tail 0.05 the the value of t is 1.812. That is

$$P(T > 1.812) = 0.05$$

Due to symmetry

$$P(T < -1.812) = 0.05$$



Degrees of Freedom	Area in Upper Tail					
	.20	.10	.05	.025	.01	.005
1	1.376	3.078	6.314	12.706	31.821	63.656
2	1.061	1.886	2.920	4.303	6.965	9.925
3	.978	1.638	2.353	3.182	4.541	5.841
4	.941	1.533	2.132	2.776	3.747	4.604
5	.920	1.476	2.015	2.571	3.365	4.032
6	.906	1.440	1.943	2.447	3.143	3.707
7	.896	1.415	1.895	2.365	2.998	3.499
8	.889	1.397	1.860	2.306	2.896	3.355
9	.883	1.383	1.833	2.262	2.821	3.250
10	.879	1.372	1.812	2.228	2.764	3.169
11	.876	1.363	1.796	2.201	2.718	3.106
12	.873	1.356	1.782	2.179	2.681	3.055
13	.870	1.350	1.771	2.160	2.650	3.012
14	.868	1.345	1.761	2.145	2.624	2.977
15	.866	1.341	1.753	2.131	2.602	2.947
16	.865	1.337	1.746	2.120	2.583	2.921
17	.863	1.333	1.740	2.110	2.567	2.898
18	.862	1.330	1.734	2.101	2.552	2.878
19	.861	1.328	1.729	2.093	2.539	2.861

Figure 8.6: t-distribution. Source: Appendix B, TABLE 3 (Anderson 2020)

8.9 F -Distribution

8.9.1 The F -Distribution with Two Sample Variances

If S_1^2 and S_2^2 are the variances of independent random samples of size n_1 and n_2 taken from normal populations with variances σ_1^2 and σ_2^2 , respectively, then the random variable

$$F = \frac{S_1^2/\sigma_1^2}{S_2^2/\sigma_2^2}$$

has an F distribution with numerator degrees of freedom $(n_1 - 1)$ and denominator degrees of freedom $(n_2 - 1)$.

8.10 Other Sampling methods

8.10.1 Probability sampling

Stratified Random Sampling

Cluster Sampling

Systematic Sampling

8.10.2 Non-probability sampling

9 Introduction to estimation

In previous chapter we discuss the sampling properties of the sample mean and variance. In this chapter we discuss about the **parameter estimation**. It falls under the branch of **Statistical Inference**. The process of estimation involves determining the approximate value of a population parameter on the basis of sample data. There are two types **parameter estimation**-(i) **point estimation** and (ii) **interval estimation**.

9.1 Point Estimation

- To estimate the value of a **population parameter**, we compute a corresponding characteristic of the sample, referred to as a **sample statistic**.
- By making the preceding computations, we perform the statistical procedure called **point estimation**. For instance, we refer to the sample mean $\bar{x}$ as the **point estimator** of the population mean μ .
- The numerical value obtained for , is called the **point estimate**.

Table 9.1: Some common population parameters and their estimators

Population parameter	Symbol	Point estimator
Population mean	μ	Sample mean, $\bar{X} = \frac{\sum x}{n}$
Population standard deviation	σ	Sample standard deviation, $S = \sqrt{\frac{\sum (x - \bar{x})^2}{n-1}} = \sqrt{\frac{\sum x^2 - n \cdot \bar{x}^2}{n-1}}$
Population proportion	p	Sample proportion, $\hat{p} = \frac{\# \text{ of outcomes of interest}}{n}$

9.1.1 Properties of Point Estimators

Suppose

θ be the population parameter of interest

$\hat{\theta}$ be the sample statistic or point estimator of θ

A “good” estimator has some desirable properties like **unbiasedness, efficiency, consistency**.

i Unbiasedness

A sample statistic $\hat{\theta}$ is said to be unbiased estimator of the population parameter θ if

$$E(\hat{\theta}) = \theta$$

Problem 9.1 Show that the function of sample mean $\bar{X}$ is the unbiased estimator of population mean μ .

Solution:

Here X is the variable of interest and let $X_1, X_2, \dots, X_n$ is a sequence of random sample provided $E(X_i) = \mu$. The sample mean is $\bar{X} = \frac{\sum_{i=1}^n X_i}{n}$.

Now

$$\begin{aligned} E(\bar{X}) &= E\left(\frac{\sum_{i=1}^n X_i}{n}\right) = \frac{1}{n} E\left(\sum_{i=1}^n X_i\right) \\ &= \frac{1}{n} \sum_{i=1}^n E(X_i) = \frac{1}{n} \sum_{i=1}^n \mu = \frac{1}{n} \cdot n\mu = \mu \\ \therefore E(\bar{X}) &= \mu \end{aligned}$$

So, $\bar{X}$ is an unbiased estimator of μ .

Problem 9.2 Let, $X_1, X_2, \dots, X_n$ be a random sample from a population with mean μ and variance σ^2 . That is X_i 's are independent and each X_i has $E(X_i) = \mu$ and $Var(X_i) = \sigma^2$.

Show that the function of sample variance $S^2 = \frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}$ is an unbiased estimator of population variance σ^2 .

Solution: See Newbold, Carlson, and Thorne (2013), page 283, or we can proof it as follows:

We know $E(\bar{X}) = \mu_{\bar{X}} = \mu$.

So,

$$Var(X), \sigma^2 = E(X^2) - \mu^2$$

$$\text{Or, } E(X^2) = \mu^2 + \sigma^2$$

and

$$Var(\bar{X}), \sigma_{\bar{X}}^2 = E(\bar{X}^2) - \mu_{\bar{X}}^2$$

$$\text{Or, } E(\bar{X}^2) = \mu_{\bar{X}}^2 + \sigma_{\bar{X}}^2 = \mu^2 + \frac{\sigma^2}{n}$$

Now,

$$\begin{aligned} E(S^2) &= E\left[\frac{\sum_{i=1}^n (X_i - \bar{X})^2}{n-1}\right] = \frac{1}{n-1} E\sum_{i=1}^n (X_i - \bar{X})^2 = \frac{1}{n-1} E\sum_{i=1}^n (X_i^2 + \bar{X}^2 - 2X_i\bar{X}) \\ &= \frac{1}{n-1} E\left(\sum_{i=1}^n X_i^2 - n\bar{X}^2\right) = \frac{1}{n-1} \left[\sum_{i=1}^n E(X_i^2) - nE(\bar{X}^2)\right] \end{aligned}$$

$$\begin{aligned}
&= \frac{1}{n-1} \left[\sum_{i=1}^n (\mu^2 + \sigma^2) - n(\mu^2 + \frac{\sigma^2}{n}) \right] = \frac{1}{n-1} [n\mu^2 + n\sigma^2 - n\mu^2 - \sigma^2] \\
&= \frac{1}{n-1} (n-1)\sigma^2 = \sigma^2 \\
&\therefore E(S^2) = \sigma^2.
\end{aligned}$$

Hence S^2 is an unbiased estimator of σ^2 .

Important NOTE: This proof only valid when the actual sample size, n , is a small proportion of the population size N or for small population size samples are taken with replacement (which me).

i Efficiency

Suppose $\hat{\theta}_1$ and $\hat{\theta}_2$ be two unbiased estimators of population parameter θ .
 $\hat{\theta}_1$ is said to be more efficient than $\hat{\theta}_2$ if $Var(\hat{\theta}_1) < Var(\hat{\theta}_2)$.

i Consistency

A point estimator $\hat{\theta}$ is said to be consistent estimator of the parameter θ if the variance of the estimator becomes smaller as sample size increases.

In other word,

$$\lim_{n \rightarrow \infty} Var(\hat{\theta}) = 0$$

Problem 9.3 (Anderson 2020a) A simple random sample of 30 managers and the corresponding data on annual salary and management training program participation are as shown in **Table 9.2**

Table 9.2: Annual Salary and Training Program Status for a Simple Random Sample of 30 EAI Managers

Annual Salary (\$000)	Management Training Program
49.09	Yes
53.26	Yes
49.64	Yes
49.89	Yes
47.62	No
45.92	Yes
49.09	Yes
51.40	Yes
50.96	Yes
45.11	Yes
45.92	No
57.27	Yes
55.69	No
51.56	No
56.19	No

Annual Salary (\$000)	Management Training Program
51.77	Yes
52.54	No
44.98	Yes
51.93	Yes
52.97	Yes
45.12	Yes
51.75	Yes
54.39	No
50.16	No
52.97	Yes
50.24	Yes
52.79	Yes
50.98	No
55.86	Yes
57.31	No

a) **Compute** *sample mean* and *standard deviation* of annual salary (\$) of a random sample of 30 EAI managers.

Solution:

Let μ be the population mean of annual salary of all EAI managers.

If X is the annual salary in '000 USD, then the to estimate μ we use **sample mean** $\bar{x}$ as follows:

$$\bar{x} = \frac{\sum_{i=1}^n x_i}{n} = \frac{49.09 + 53.26 + \dots + 57.31}{30} \approx 51.1457$$

So the **sample mean** is \$51145.7.

Similarly let σ be the population standard deviation of annual salary of all EAI managers.

The estimate of σ is the **sample standard deviation** s as follows:

$$s = \sqrt{\frac{\sum_{i=1}^n x_i^2 - n \cdot (\bar{x})^2}{n - 1}}$$

$$= \sqrt{\frac{(49.09^2 + 53.26^2 + \dots + 57.31^2) - 30 \cdot (51.1457)^2}{30 - 1}}$$

$$\approx 3.5408$$

So the **sample standard deviation** is \$3540.8.

b) Also, **estimate** the *proportion* of managers in the population who completed the management training program.

Solution: Here, $n = 30$

Let, p be the population proportion of managers who completed the training

The estimate of p is:

$$\hat{p} = \frac{\# \text{ of yes}}{n} = \frac{20}{30} = 0.6667 \approx 66.67\%$$

Problem 9.4 (Anderson 2020a) Many drugs used to treat cancer are expensive. Business Week reported on the cost per treatment of Herceptin, a drug used to treat breast cancer (Business Week, January 30, 2006). Typical treatment costs (in dollars) for Herceptin are provided by a simple random sample of 10 patients.

4376, 5578, 2717, 4920, 4495, 4798, 6446, 4119, 4237, 3814

- Develop** a point estimate of the mean cost per treatment with Herceptin.
- Develop** a point estimate of the standard deviation of the cost per treatment with Herceptin.

9.2 Interval estimation

Instead of estimating a population parameter by a single value (point estimator) it is more reasonable to estimate with an **interval** with some confidence (probability) that our **parameter** value will be in the **interval**.

i Interval Estimator

An **interval estimator** is a rule for determining (based on sample information) an interval that is likely to include the parameter. The general form of an interval estimate is as follows:

$$\text{Point estimate} \pm \text{margin of error}$$

Due to sampling variability, **interval estimator** is also random.

9.2.1 Interval estimate of a population mean: σ known

The $(1 - \alpha)100\%$ confidence interval for μ is :

$$\bar{x} \pm z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \quad (9.1)$$

Or,

$$\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}}, \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

We can express this confidence interval in a probabilistic way:

$$P \left(\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}} \right) = 1 - \alpha$$

NOTE:

1) Here, $z_{\alpha/2}$ is the z value providing an area of $\alpha/2$ in the upper tail of the standard normal distribution that is $P(Z > z_{\alpha/2}) = \alpha/2$.


2) $z_{\alpha/2} \cdot \frac{\sigma}{\sqrt{n}}$ is often called **margin of error (ME)**.

9.2.2 Interpretation of confidence interval

The probabilistic equation of confidence interval says that, if we repeatedly construct confidence intervals in this manner, we will expect $(1 - \alpha)100\%$ of them contain μ .

9.2.3 Understanding confidence interval through Simulation

Suppose $X \sim N(50, 5^2)$. Now consider a population data of size $N = 10000$ and the histogram of X is:



Now we draw a random sample of size $n = 50$ from this population and construct a 95% confidence interval (CI) for μ . The CI may or may not include the $\mu = 50$!!!

Sample data : 52.60842 55.16664 59.23435 44.2092 50.94234 43.34063 44.65922 53.81687 47.60075 4

Sample mean: 49.3

95% CI:

[Lower ,Upper]

[47.91 , 50.68]




Figure 9.1: Simulation of 95% confidence intervals for μ

Luckily our 95% CI contains the true population mean $\mu = 50$.

Lets simulate 100 samples each of size $n = 50$ and construct all 95% CIs.

We can see that out of 100 CIs , 95 of them contain true population mean $\mu = 50$ and the rest 5 do not.

Table 9.3: Four Commonly Used Confidence Levels and $z_{\alpha/2}$

$1 - \alpha$	α	$z_{\alpha/2}$
0.90	0.10	1.645
0.95	0.05	1.96
0.98	0.02	2.33
0.99	0.01	2.575

9.2.4 Interval estimate of a population mean: σ unknown

The $(1 - \alpha)100\%$ confidence interval for μ is :

$$\bar{x} \pm t_{\alpha/2} \frac{s}{\sqrt{n}} \quad (9.2)$$

Or,

$$\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}}, \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}$$

We can express this confidence interval in a probabilistic way:

$$P\left(\bar{x} - t_{\alpha/2} \frac{s}{\sqrt{n}} < \mu < \bar{x} + t_{\alpha/2} \frac{s}{\sqrt{n}}\right) = 1 - \alpha$$

Here, $t_{\alpha/2}$ is the t value providing an area of $\alpha/2$ in the upper tail of the t distribution with $(n - 1)$ degrees of freedom that is $P(T > t_{\alpha/2, n-1}) = \alpha/2$.

Problem 9.5 (Keller 2014, 341) In a survey conducted to determine, among other things, the cost of vacations, 64 individuals were randomly sampled. Each person was asked to compute the cost of her or his most recent vacation and the sample mean was \$1810.16. Assuming that the standard deviation is \$400, **estimate** with 95% confidence interval for the average cost of all vacations.

Solution: Let, X be the vacation cost in \$. Also let μ be the mean of all vacations.

Here sample size is, $n = 64$ with the sample mean, $\bar{x} = \$1810.16$.

The assumed (population) standard deviation, $\sigma = \$400$. Since the sample size is sufficiently large so according to CLT, $\bar{X} \sim N(\mu, \sigma_{\bar{X}}^2)$ *approximately*.

Where $\sigma_{\bar{X}} = \frac{\sigma}{\sqrt{n}} = \frac{400}{\sqrt{64}} = 50$

For 99% confidence interval (CI), $z_{\alpha/2} = 2.575$.

So the 99% CI for the μ is:

$$\bar{x} \pm z_{\alpha/2} \times \frac{\sigma}{\sqrt{n}}$$

$$\text{Or, } 1810.16 \pm 2.575 \times 50$$

$$\text{Or, } 1810.16 \pm 128.75$$

$$[1681.41, 1938.91]$$

$$\therefore P(1681.41 < \mu < 1938.91) = 99\%.$$

Problem 9.6 (Keller 2014, 340) It is known that the amount of time needed to change the oil on a car is normally distributed with a standard deviation of 5 minutes. The amount of time to complete a random sample of 10 oil changes was recorded and listed here. **Compute** the 99% confidence interval estimate of the mean of the population.

11, 10, 16, 15, 18, 12, 25, 20, 18, 24

Problem 9.7 (Newbold, Carlson, and Thorne 2013, 302) How much do students pay, on the average, for textbooks during the first semester of college? From a random sample of 400 students the mean cost was found to be \$357.75, and the sample standard deviation was \$37.89. Assuming that the population is normally distributed, find the 95% confidence interval for the population mean.

Problem 9.8 (Newbold, Carlson, and Thorne 2013, 302) Twenty people in one large metropolitan area were asked to record the time (in minutes) that it takes them to drive to work. These times were as follows:

30, 42, 35, 40, 45, 22, 32, 15, 41, 45, 28, 32, 45, 27, 47, 50, 30, 25, 46, 25

Assuming that the population is normally distributed find the 99% confidence interval for the population mean of time it takes to drive to work.

9.2.5 Interval estimation for a population proportion : Large sample

From previous chapter we know if np and $np(1-p)$ is equal or greater than 5 then the sample proportion $\hat{p}$ will approximately follow **normal distribution** with **mean** p and **variance** $\frac{p(1-p)}{n}$.

Mathematically,

$$Z = \frac{\hat{p} - p}{\sqrt{\frac{p(1-p)}{n}}} \approx \text{follows } N(0, 1)$$

Since p is unknown we estimate $\text{var}(\hat{p})$ as $\frac{\hat{p}(1-\hat{p})}{n}$.

Confidence Interval for Population Proportion, p (Large Samples)

$$\hat{p} \pm z_{\alpha/2} \sqrt{\frac{\hat{p}(1-\hat{p})}{n}} \quad (9.3)$$

which is valid provided that $n\hat{p}$ and $n(1-\hat{p})$ are greater than 5.

Problem 9.9 (Newbold, Carlson, and Thorne 2013, 305) In a random sample of 95 manufacturing firms, 67 indicated that their company attained ISO certification within the last two years. Find a 99% confidence interval for the population proportion of companies that have been certified within the last 2 years.

Problem 9.10 (Newbold, Carlson, and Thorne 2013, 305) From a random sample of 400 registered voters in one city, 320 indicated that they would vote in favor of a proposed policy in an upcoming election. Find a 98% confidence interval for the population proportion in favor of this policy.

9.3 Sample size determination: Large Population

9.3.1 Sample Size for Population Mean

If E be the *margin of error* (E), σ be the population standard deviation (assume known) then the required **sample size** (n) to estimate a population mean μ with $(1-\alpha) \times 100\%$ confidence level is:

$$n = \left(\frac{z_{\alpha/2} \sigma}{E} \right)^2 \quad (9.4)$$

i Sample size determination: Derivation

From the interval estimation regarding μ we know

$$P(\bar{x} - z_{\alpha/2} \frac{\sigma}{\sqrt{n}} < \mu < \bar{x} + z_{\alpha/2} \frac{\sigma}{\sqrt{n}}) = 1 - \alpha$$

That is,

$$P(\bar{x} - E < \mu < \bar{x} + E) = 1 - \alpha$$

Hence we can write,

$$E = z_{\alpha/2} \frac{\sigma}{\sqrt{n}}$$

Solving for n we have

$$n = \left(\frac{z_{\alpha/2} * \sigma}{E} \right)^2$$

Problem 9.11 (Keller 2014)

- a) Determine the sample size required to estimate a population mean to within 10 units given that the population standard deviation is 50. A confidence level of 90% is judged to be appropriate.
- b) Repeat part (a) changing the standard deviation to 100.
- c) Re-do part (a) using a 95% confidence level.
- d) Repeat part (a) wherein we wish to estimate the population mean to within 20 units.

Problem 9.12 (Keller 2014) The operations manager of a large production plant would like to estimate the average amount of time workers take to assemble a new electronic component. After observing a number of workers assembling similar devices, she guesses that the standard deviation is 6 minutes. How large a sample of workers should she take if she wishes to estimate the mean assembly time to within 20 seconds? Assume that the confidence level is to be 99%.

9.3.2 Sample Size for Population Proportion

If E be the *margin of error* (E), p^* be the assumed population proportion or planning value of $\hat{p}$. Then the required **sample size** (n) to estimate a population proportion p with $(1 - \alpha) \times 100\%$ confidence level is:

$$n = \frac{z_{\alpha/2}^2 \times p^*(1 - p^*)}{E^2} \quad (9.5)$$

N.B: From Equation 9.5 it can be proved that the sample size will be maximum for $p^* = 0.5$. So to obtain the maximum sample size we can take $p^*(1 - p^*) = 0.5 \times 0.5 = 0.25$ in Equation 9.5.

Problem 9.13 (Anderson 2020a) At 95% confidence, how large a sample should be taken to obtain a margin of error of 0.03 for the estimation of a population proportion? Assume that past data are not available for developing a planning value for population proportion p ?

Problem 9.14 (Anderson 2020a) **Stay-at-Home Parenting.** In June 2014, Pew Research reported that in 16% of all homes with a stay-at-home parent, the father is the stay-at-home parent. An independent research firm has been charged with conducting a sample survey to obtain more current information.

- a. What sample size is needed if the research firm's goal is to estimate the current proportion of homes with a stay-at-home parent in which the father is the stay-at-home parent with a margin of error of 0.03? Use a 95% confidence level.
- b. Repeat part (a) using a 99% confidence level.

9.4 Interval estimation: Comparing TWO populations

10 Hypothesis test: Introduction and testing one population parameter

10.1 Definition

A statistical hypothesis is a *statement* about the *parameters* of one or more populations.

Example 1: A manufacturer claims that the mean life of a smartphone is more than 1.5 years.

Example 2: A local courier service claims that they deliver a ordered product within 30 minutes on average.

Example 3: A sports drink maker claims that the mean calorie content of its beverages is 72 calories per serving.

10.2 Types of hypothesis

Statistical hypothesis are stated in two forms- (i) Null hypothesis (H_0) and (ii) Alternative hypothesis (H_1).

Both null and alternative hypothesis are the written about the parameter of interest based on the claim.

- We will always state the null hypothesis as an **equality claim**.
- However, when the alternative hypothesis is stated with the “<” sign, the implicit claim in the null hypothesis can be taken as “ ” or “=” sign.
- When the alternative hypothesis is stated with the “>” sign, the implicit claim in the null hypothesis can be taken as “ ” or “=” sign.

10.3 Developing hypotheses

To develop or state null and alternative hypothesis, at first we have to clearly identify the “**claim**” about population parameter. Now we will see some examples.

Example 1: A manufacturer claims that the mean life of a smartphone is more than 1.5 years.

Hypothesis:

$$H_0 : \mu = 1.5$$

$$H_1 : \mu > 1.5 \text{ (claim)}$$

Example 2: A local courier service claims that they deliver a ordered product within 30 minutes on average.

Hypothesis:

$$H_0 : \mu = 30$$

$$H_1 : \mu < 30 \text{ (claim)}$$

Example 3: A sports drink maker claims that the mean calorie content of its beverages is 72 calories per serving.

Hypothesis:

$$H_0 : \mu = 72 \text{ (claim)}$$

$$H_1 : \mu \neq 72$$

10.4 Types of test based on alternative hypothesis H_1

- $H_1 : \mu < \mu_0$ (Lower tailed)
- $H_1 : \mu > \mu_0$ (Upper tailed)
- $H_1 : \mu \neq \mu_0$ (Two-tailed)

10.5 Types of error in hypothesis test and P-value

While testing a statistical hypothesis concerning population parameter we commit two types of errors.

- **Type I error** occurs when we **reject** a **TRUE** H_0
- **Type II error** occurs when we **FAIL to reject** a **FALSE** H_0
- The **Level of significance** is the probability of comiting **Type I error**. It is denoted by α .

$$\alpha = P(\text{Type I error})$$

- The probability of committing a **Type II error**, denoted by β .

$$\beta = P(\text{Type II error})$$

! Note

Type I error is more serious than **Type II error**. Because rejecting a TRUE statement is more devastating than FAIL to reject a FALSE statement. So, we always try to keep our probability of Type I error as small as possible (1% or at most 5%). For more detail see (Keller 2014).

- **P-value:** If the null hypothesis is true, then a P-value (or probability value) of a hypothesis test is the probability of obtaining a sample statistic with a value as extreme or more extreme than the one determined from the sample data.

The smaller the P-value of the test, the more evidence there is to reject the null hypothesis. A very small P-value indicates an unusual event

! P-value Explanation

A hypothesis test functions like a **legal trial**. You start by assuming the **null hypothesis** (H_0) is true, just as a jury assumes *a defendant is innocent until proven guilty*.

The data you collect acts as the evidence. By comparing this evidence to your initial assumption, you calculate a **p-value** (ranging from 0 to 1), which quantifies how well the sample data “fits” with H_0 .

- **A high p-value** suggests the evidence is consistent with the null hypothesis.
- **A low p-value** indicates a sharp disagreement between the data and the null hypothesis.

When the p-value is small enough, it provides “beyond a reasonable doubt” proof that the initial assumption is unlikely. At this point, you **reject the null hypothesis** and shift your belief to the **alternative hypothesis** (H_1).

So, how these hypotheses will be tested?

To test a hypothesis we have to determine

- a **test-statistic**; and
- **Critical/Rejection region** based on the sampling distribution of test-statistic for a given α ;
- If the value of test-statistic **falls in Critical/Rejection region**, then we reject Null (H_0) hypothesis; otherwise not. Or,
- If **P-value** $\leq \alpha$ then we reject Null (H_0) hypothesis.

10.6 Hypothesis testing concerning population mean (μ)

The following two hypotheses tests are used concerning population mean (μ):

1. One sample z-test (with known σ)
2. One sample t-test (with unknown σ)

10.6.1 One sample z-test

When sampling is from a **normally distributed population** or **sample size is sufficiently large** and **the population variance is known**, the test statistic for testing $H_0 : \mu = \mu_0$ at α is

$$z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}}$$

Decision (Critical value approach): If calculated z falls in rejection region (CR) , then reject H_0 . Otherwise, do not reject H_0 .

- For lower tailed test, reject H_0 if $z_0 < -z_\alpha$;
- For upper tailed test, reject H_0 if $z_0 > z_\alpha$;
- For two-tailed test, reject H_0 if $z_0 < -z_{\alpha/2}$ or $z_0 > z_{\alpha/2}$.

Decision (P-value approach)

Alternative Hypothesis	P-value
$H_1 : \mu < \mu_0$ (Lower-tailed)	$P(Z < z_0)$
$H_1 : \mu > \mu_0$ (Upper-tailed)	$P(Z > z_0)$
$H_1 : \mu \neq \mu_0$ (Two-tailed)	$P(Z < - z_0) + P(Z > z_0)$ $= 2P(Z < - z_0)$ $= 2P(Z > z_0)$

i Exercises 10.1

Methods

1) Consider the following hypothesis test:

$$H_0 : \mu \geq 20$$

$$H_a : \mu < 20$$

A sample of 50 provided a sample mean of 19.4. The population standard deviation is 2.

- Compute the value of the test statistic.
- What is the p -value?
- Using $\alpha = 0.05$, what is your conclusion?
- What is the rejection rule using the critical value? What is your conclusion?

Solution:

- Test statistic, $z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} = \frac{19.4 - 20}{2/\sqrt{50}} = -2.12132 \approx -2.12$
- For lower-tail test, **P-value** $= P(Z < z_0) = P(Z < -2.12) = 0.0170$
- Since **P-value** $< \alpha$; so reject H_0 .
- For $\alpha = 0.05$, critical value is $-z_\alpha = -1.645$. Since $z_0 < -z_\alpha$ that is z_0 falls in **Critical region (CR)** so we reject the H_0 .

2) Consider the following hypothesis test:

$$H_0 : \mu \leq 25$$

$$H_a : \mu > 25$$

A sample of 40 provided a sample mean of 26.4. The population standard deviation is 6.

- Compute** the value of the test statistic.
- What is the p -value?
- At $\alpha = 0.01$, what is your conclusion?
- What is the rejection rule using the critical value? What is your conclusion?

Solution: Do it yourself.

3) **Consider the following hypothesis test:**

$$H_0 : \mu = 15$$

$$H_a : \mu \neq 15$$

A sample of 50 provided a sample mean of 14.15. The population standard deviation is 3.

- Compute the value of the test statistic.
- What is the p -value?
- At $\alpha = 0.05$, what is your conclusion?
- What is the rejection rule using the critical value? What is your conclusion?

Solution:

- Test statistic, $z_0 = \frac{\bar{x} - \mu_0}{\sigma/\sqrt{n}} = \frac{14.15 - 15}{3/\sqrt{50}} = -2.003 \approx -2.00$.
- For TWO-tailed test, $P\text{-value} = P(Z < -2.00) + P(Z > 2.00) = 0.0228 + 0.0228 = 0.0456$.
- Since **P-value** $< \alpha$; so reject H_0 .
- For $\alpha = 0.05$, critical value is $\pm z_{\alpha/2} = \pm 1.96$. Since $z_0 < -z_{\alpha/2}$ that is z_0 falls in **Critical region (CR)** so we reject the H_0 .

Problem 10.1 The waiting time for customers at MacMillan Restaurants follows a normal distribution with a mean of 3 minutes and a standard deviation of 1 minute. At the Mirpur Road MacMillan, the quality-assurance department sampled 50 customers and found that the mean waiting time was 2.75 minutes.

- At the 0.05 significance level, can we conclude that the mean waiting time is less than 3 minutes?
- Compute** P-value and conclude whether the mean waiting time is less than 3 minutes

Problem 10.2 At the time she was hired as a server at the Grumney Family Restaurant, Beth Brigden was told, “You can average \$80 a day in tips.” Assume the population of daily tips is normally distributed with a standard deviation of \$3.24. Over the first 35 days she was employed at the restaurant, the mean daily amount of her tips was \$84.85. At the 0.01 significance level, can Ms. Brigden conclude that her daily tips average more than \$80?

Problem 10.3 The manufacturer of the X-15 steel-belted radial truck tire claims that the mean mileage the tire can be driven before the tread wears out is 60,000 miles. Assume the mileage wear follows the normal distribution and the standard deviation of the distribution is 5,000 miles. Crosset

Truck Company bought 48 tires and found that the mean mileage for its trucks is 59,500 miles. Is Crosset's experience different from that claimed by the manufacturer at the 0.05 significance level?

10.6.2 One sample t-test

When sampling is from a **normally distributed population** or **sample size is sufficiently large** and **the population variance is unknown**, the test statistic for testing $H_0 : \mu = \mu_0$ at α is

$$t_0 = \frac{\bar{x} - \mu_0}{s/\sqrt{n}}$$

Test statistic t follows a Student's t distribution with $(n - 1)$ degrees of freedom.

Decision (Critical value approach): If calculated t falls in rejection region (CR) , then reject H_0 . Otherwise, do not reject H_0 .

- For lower tailed test, reject H_0 if $t_0 < -t_\alpha$;
- For upper tailed test, reject H_0 if $t_0 > t_\alpha$;
- For two-tailed test, reject H_0 if $t_0 < -t_{\alpha/2}$ or $t_0 > t_{\alpha/2}$.

P-value calculation for t-statistic:

Exact P-value calculation needs using of calculus applied to PDF of t-distribution. But there are a number of tools like Excel, Spreadsheet, R, etc. which provide P-value.

- a) In R , `pt(q, df, lower.tail=TRUE)` is used to compute **left-tail area** of a given value of q .
- b) In Excel, `=T.DIST.RT(x,deg_freedom)` is used to compute **right-tail area** of a given value of x .

Problem 10.4 Annual per capita consumption of milk is 21.6 gallons (*Statistical Abstract of the United States: 2006*). Being from the Midwest, you believe milk consumption is higher there and wish to support your opinion. A sample of 16 individuals from the Midwestern town of Webster City showed a sample mean annual consumption of 24.1 gallons with a standard deviation of $s = 4.8$.

- a) **Develop** a hypothesis test that can be used to determine whether the mean annual consumption in Webster City is higher than the national mean.
- b) **Test** the hypothesis at $\alpha = 0.05$.
- c) **Draw** a conclusion.

Problem 10.5 The mean length of a small counterbalance bar is 43 millimeters. The production supervisor is concerned that the adjustments of the machine producing the bars have changed. He asks the Engineering Department to investigate. Engineer selects a random sample of 10 bars and measures each. The results are reported below in millimeters.

42, 39, 42, 45, 43, 40, 39, 41, 40, 42

Is it reasonable to conclude that there has been a change in the mean length of the bars?

10.7 Hypothesis test of a Population variance

When sampling is from a **normally distributed population** or **sample size is sufficiently large** the test statistic to test the $H_0 : \sigma^2 = \sigma_0^2$ is at α level of significance:

$$\chi_0^2 = \frac{(n-1)s^2}{\sigma_0^2}$$

Where χ_0^2 follows the χ^2 -distribution with degrees of freedom $\nu = n - 1$.

Decision rule for rejecting H_0 :

Alternative hypothesis	Rejection rule
$H_1 : \sigma^2 < \sigma_0^2$	$\chi_0^2 < \chi_{1-\alpha}^2$
$H_1 : \sigma^2 > \sigma_0^2$	$\chi_0^2 > \chi_{\alpha}^2$
$H_1 : \sigma^2 \neq \sigma_0^2$	$\chi_0^2 < \chi_{1-\alpha/2}^2$ OR $\chi_0^2 > \chi_{\alpha/2}^2$

Problem : A manufacturer of car batteries claims that the life of the company's batteries is approximately normally distributed with a standard deviation equal to 0.9 year. If a random sample of 10 of these batteries has a standard deviation of 1.2 years, do you think that $\sigma > 0.9$ year? Use a 0.05 level of significance.

Problem : The content of containers of a particular lubricant is known to be normally distributed with a variance of 0.03 liter. A random sample of 10 containers are 10.2, 9.7, 10.1, 10.3, 10.1, 9.8, 9.9, 10.4, 10.3, and 9.8 liters. Test the hypothesis that $\sigma^2 = 0.03$ against the alternative that $\sigma^2 \neq 0.03$.

10.8 Hypothesis test of a Population proportion

10.9 Normality test

In parametric (distribution based) hypothesis test the checking normality assumption of study variable is a common practice especially when the sample size is small ($n < 30$). For large samples, the **Central Limit Theorem (CLT)** often makes this test robust to non-normality.

The normality assumption is checked in two ways:

- a) Graphically
- b) Numerically using some normality tests
- a) **Graphical procedure to check normality**

We often plot the data (i.e., histogram, density plot, boxplot) to explore so called bell-shaped of the data. But the most popular and effective way to check normality is **Q-Q plot (Quantile-Quantile plot)**.

b) Normality test

A number of normality tests are available; of them a common test is **Shapiro-Wilk** test of normality suitable for small to medium sample size (3 to 5000) (Shapiro and Wilk 1965; Royston 1982).

Shapiro-Wilk Test Statistic W

$$W = \frac{(\sum_{i=1}^n a_i x_{(i)})^2}{\sum_{i=1}^n (x_i - \bar{x})^2}$$

Where,

- $x_{(i)}$: the i^{th} **order statistic** (i.e., the i -th smallest value in the sample)
- $\bar{x}$: the **sample mean**
- a_i : constants calculated based on the **expected values and variances** of order statistics from a **standard normal distribution** (Tabulated in Shapiro Wilk Table)
- n : sample size

Hypotheses:

- **Null Hypothesis H_0** : The data are **normally distributed**.
- **Alternative H_1** : The data are **not normally distributed**.

We reject H_0 if the **p-value** is less than our significance level (e.g., 0.05).

Almost all statistical software and package routinely provide the **Shapiro-Wilk** test.

In **R** Shapiro-Wilk test is available as `shapiro.test` .

Shapiro-Wilk normality test

```
data: uniform.data
W = 0.93903, p-value = 0.0001683
```

The p-value<0.05 implies (reject H_0) that the data is not normally distributed.

Shapiro-Wilk normality test

```
data: normal.data
W = 0.99212, p-value = 0.83
```

The p-value > 0.05 implies (do not reject H_0) that the data is normally distributed.

11 Hypothesis test concerning TWO population parameters

In order to test difference between two population parameters we need to draw samples from two populations. Using sample statistics and appropriate test statistics we can test whether there is any significant difference between the parameters of interest. In this chapter we will discuss:

- 1) testing difference between two population means;
- 2) testing difference between two population variances;
- 3) testing difference between two means from matched pairs experiment;
- 4) testing difference between two population proportions.

11.1 Hypothesis test: Difference between two population means ($\mu_1 - \mu_2$)

Assumptions:

- The two samples must be **independent** of each other.
- The populations from which the samples are drawn should be **normally distributed**, especially when the sample size is small (typically $n < 30$).
- With larger samples, the **Central Limit Theorem** justifies the use of normal approximation even if the population is not normal.

11.1.1 Case-I: When σ_1^2 and σ_2^2 are known

To test $H_0 : \mu_1 - \mu_2 = D_0$ the test statistic is:

$$z_0 = \frac{(\bar{x}_1 - \bar{x}_2) - D_0}{s.e(\bar{x}_1 - \bar{x}_2)} = \frac{(\bar{x}_1 - \bar{x}_2) - D_0}{\sqrt{\frac{\sigma_1^2}{n_1} + \frac{\sigma_2^2}{n_2}}} \quad (11.1)$$

The test-statistic z_0 follows standard normal distribution. The rejection rule of H_0 is same as **one-sample z-test**.

Problem 12.1 Consider the following information:

Sample 1	Sample 2
$n_1 = 80$	$n_1 = 70$

Sample 1	Sample 2
$\bar{x}_1 = 104$	$\bar{x}_2 = 106$
$\sigma_1 = 8.4$	$\sigma_2 = 7.6$

Now test the following hypothesis:

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

Problem 12.2 (Larson and Farber 2015) A credit card watchdog group claims that there is a difference in the mean credit card debts of households in California and Illinois. The results of a random survey of 250 households from each state are shown at the bottom. The two samples are independent. Assume that $\sigma_1 = \$1045$ for California and $\sigma_2 = \$1350$ for Illinois. Do the results support the group's claim? Use $\alpha = 0.05$.

Table 11.2: Sample statistics for Credit Card Debt

California	Illinois
$\bar{x}_1 = \$4777$	$\bar{x}_2 = \$4866$
$n_1 = 250$	$n_2 = 250$

Problem 12.3 (Larson and Farber 2015) A travel agency claims that the average daily cost of meals and lodging for vacationing in Alaska is greater than the average daily cost in Colorado. The table at the bottom shows the results of a random survey of vacationers in each state. The two samples are independent. Assume that $\sigma_1 = \$24$ for Alaska and $\sigma_2 = \$19$ for Colorado, and that both populations are normally distributed. At $\alpha = 0.05$, is there enough evidence to support the claim?

Table 11.3: Sample Statistics for Daily Cost of Meals and Lodging for Two Adults

	Alaska	Colorado
Sample mean	\$ 296	\$ 293
Sample size	15	20

11.1.2 Case-II(A): σ_1^2 and σ_2^2 are unknown but equal ($\sigma_1^2 = \sigma_2^2$)

To test $H_0 : \mu_1 - \mu_2 = D_0$ the test statistic is:

$$t_0 = \frac{(\bar{x}_1 - \bar{x}_2) - D_0}{\sqrt{s_p^2 \left(\frac{1}{n_1} + \frac{1}{n_2} \right)}} \quad (11.2)$$

Here, s_p^2 is the pooled variance and

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2}$$

The test statistic **t** follows the **Student t distribution** with $df = n_1 + n_2 - 2$. The rejection rule of H_0 is same as **one-sample t-test**.

11.1.3 Case-II(B): σ_1^2 and σ_2^2 are unknown and NOT equal ($\sigma_1^2 \neq \sigma_2^2$)

To test $H_0 : \mu_1 - \mu_2 = D_0$ the test statistic is:

$$t_0 = \frac{(\bar{x}_1 - \bar{x}_2) - D_0}{\sqrt{\left(\frac{s_1^2}{n_1} + \frac{s_2^2}{n_2}\right)}} \quad (11.3)$$

There are two procedure of obtaining degrees of freedom of t-distribution (Moore 2010).

- (i) Use **Satterthwaite approximation** to compute $df = \frac{(s_1^2/n_1 + s_2^2/n_2)^2}{\frac{(s_1^2/n_1)^2}{n_1 - 1} + \frac{(s_2^2/n_2)^2}{n_2 - 1}}$
- (ii) Use degrees of freedom equal to the smaller of $n_1 - 1$ and $n_2 - 1$.

11.2 Testing the Population Variances ($\sigma_1^2 = \sigma_2^2$)

The hypothesis to test the equality of two population variances is:

$$H_0 : \sigma_1^2 = \sigma_2^2$$

$$H_1 : \sigma_1^2 \neq \sigma_2^2$$

Test statistic

$$F_0 = \frac{s_1^2}{s_2^2} \quad (11.4)$$

The F_0 -statistic is F -distributed with degrees of freedom $\nu_1 = n_1 - 1$ and $\nu_2 = n_2 - 1$.

Assuming $s_1^2 > s_2^2$, **we can reject the H_0 if $F_0 > F_{\alpha/2}$** (Here $\alpha/2$ is the area in the upper tail).

NOTE: We refer to the population providing the larger sample variance as *population 1* .

Problem 12.4 The following data were collected from two population- population A and population B.

	Population A	Population B
Sample size	35	40

	Population A	Population B
Sample mean	13.6	10.1
Sample variance	5.2	8.5

a) *Test the equality of variances of the two populations at $\alpha = 5\%$.*

Solution (a):

Since the sample variance of population B is greater than that of population A; we consider population B as population 1.

So, $s_1^2 = 8.5$, $s_2^2 = 5.2$ and $n_1 = 40$; $n_2 = 35$

We have to test the following hypothesis :

$$H_0 : \sigma_1^2 = \sigma_2^2$$

$$H_1 : \sigma_1^2 \neq \sigma_2^2$$

Test statistic:

$$F_0 = \frac{s_1^2}{s_2^2} = \frac{8.5}{5.2} = 1.635$$

Critical value: For $\alpha/2 = 0.025$

$$F_{0.025} \approx 1.93$$

Decision: Since $F_0 < F_{\alpha/2}$; so we cannot reject the H_0 . So the the equality assumption of the two population variances are met that is $\sigma_1^2 = \sigma_2^2$.

b) *Then use **appropriate test statistic** to test equality of two population means at $\alpha = 5\%$.*

Solution (b):

We have to test the following hypothesis:

$$H_0 : \mu_1 - \mu_2 = 0$$

$$H_1 : \mu_1 - \mu_2 \neq 0$$

Since equality of population variance is fulfilled so the appropriate **test statistic** is :

$$t_0 = \frac{(\bar{x}_1 - \bar{x}_2)}{\sqrt{s_p^2(\frac{1}{n_1} + \frac{1}{n_2})}}$$

Where,

$$s_p^2 = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2}{n_1 + n_2 - 2} = 6.963014$$

$$\text{So, } t_0 = \frac{10.1 - 13.6}{\sqrt{6.963014(\frac{1}{40} + \frac{1}{35})}} = -5.731$$

Critical value: At $\alpha/2 = 0.025$, with $df = 40 + 35 - 2 = 73$

$$-t_{\alpha/2} \approx -1.992$$

Decision: Since the value of t_0 falls in the rejection region (RR) we can reject the H_0 .

Conclusion: With 95% confidence we can conclude that the population means are not equal; rather mean of population 1 (B) is significantly lower than the mean of population 2 (A).

Problem 12.5 An accounting firm is interested in providing opportunities for its auditors to gain more expertise in statistical sampling methods. They wish to compare traditional classroom instruction with online self-paced tutorials. Auditors were assigned at random to one type of instruction, and the auditors were then given an exam. The table shows how the two groups performed.

Program	n	Sample mean score	Sample SD
Traditional	30	76.74	11.2
Online	30	72.33	12.3

a) **Test** equality of variances of two methods of learning.

b) **Test** whether there is any significant difference in mean scores between Traditional and Online method.

Problem 12.6 The bank randomly assigns **160 credit card customers** to each of two marketing strategies: **Proposal A** (waive the annual fee if yearly spending is at least \$3600) and **Proposal B** (offer a small cash rebate based on total yearly spending). After one year, the **total amount charged by each customer** is recorded and compared to determine which proposal leads to higher credit card spending. Here are the summary statistics:

Group	n	$\bar{x}$	s
A	160	\$3285	\$492
B	160	\$3124	\$415

(a) **Test** whether the population variances are equal or not.

(b) Now **examine** whether the data show a significant difference between the mean amounts charged by customers offered the two plans.

- Give the null and alternative hypotheses, and
- Calculate the two-sample t-statistic
- State your practical conclusions.

11.3 Hypothesis test: Comparing TWO means when the samples are dependent/ matched/paired

The **paired t-test** is used to compare the means of two related groups (e.g., before-and-after measurements on the **same subjects**). To validly apply the test, the following assumptions must be met:

Assumptions of the Paired t-test:

- 1) Paired observations: Each subject or entity provides a pair of observations (e.g., pre-treatment and post-treatment). The test assumes that data are dependent.
- 2) Continuous or interval scale: The differences between paired observations should be measured on a continuous (interval or ratio) scale.
- 3) Normality of the differences:

The distribution of the differences (not the raw values) between the paired observations should be approximately normally distributed.

- For small samples (typically $n < 30$), this assumption is critical.
- For larger samples, the Central Limit Theorem provides robustness.

- 4) No significant outliers in the differences:

Extreme outliers in the differences can heavily influence the result and violate the normality assumption.

Test procedure

Suppose n subjects are measured in two different occasion/conditions regarding the measurement variable X .

Let, μ_A be the population mean of X under condition A and

μ_B be the population mean of X under condition B

So the difference between the means is $\mu_d = \mu_A - \mu_B$

To test the **hypotheses**:

$$H_0 : \mu_d = D_0$$

$$H_1 : \mu_d \neq 0$$

the **test statistic** is:

$$t_0 = \frac{\bar{d} - D_0}{s_d / \sqrt{n}}$$

which is Student t distributed with $n - 1$ degrees of freedom, provided that the differences are normally distributed.

The rejection rule is similar as one-sample t-test.

Here,

- n = sample size
- d_i = difference in measurements between two conditions from i^{th} subject

$$\bar{d} = \frac{\sum_{i=1}^n d_i}{n}$$

$$s_d = \sqrt{\frac{\sum_{i=1}^n (d_i - \bar{d})^2}{n - 1}}$$

Problem 12.6 (Black 2012, 10.27)

Eleven employees were put under the care of the company nurse because of high cholesterol readings. The nurse lectured them on the dangers of this condition and put them on a new diet. Shown are the cholesterol readings of the 11 employees both before the new diet and one month after use of the diet began.

Employee	Before	After
1	255	197
2	230	225
3	290	215
4	242	215
5	300	240
6	250	235
7	215	190
8	230	240
9	225	200
10	219	203
11	236	223

Use $\alpha = 0.05$ to test for a significant difference between population means for the before and after cholesterol readings.

Solution:

Let, μ_B be the population of cholesterol readings **before** nurse's lecture and

μ_A be the population of cholesterol readings **after** nurse's lecture.

The difference is : $\mu_A - \mu_B = \mu_D$

Hypotheses:

$$H_0 : \mu_D = 0$$

$$H_1 : \mu_D \neq 0$$

Required calculation for test of hypotheses

Employee	Before	After	Difference, $d = \text{After} - \text{Before}$
1	255	197	-58
2	230	225	-5
3	290	215	-75
4	242	215	-27
5	300	240	-60
6	250	235	-15
7	215	190	-25
8	230	240	10
9	225	200	-25
10	219	203	-16
11	236	223	-13

Using calculator;

$$\sum d = -259$$

$$\bar{d} = \frac{\sum d}{n} = \frac{-259}{11} = -28.09$$

$$s_D = \sqrt{\frac{\sum d^2 - n(\bar{d})^2}{n-1}} = 25.81$$

Test statistic

$$t_0 = \frac{\bar{d} - D_0}{s_D/\sqrt{n}} = \frac{-28.09 - 0}{25.81/\sqrt{11}} = -3.609$$

Critical value

For $\alpha = 0.05$ and $df = n - 1 = 11 - 1 = 10$;

$$-t_{\alpha/2} = -t_{0.025} = -2.228$$

Since t_0 falls in CR ($t_0 < -t_{\alpha/2}$), so reject H_0 .

Hence we can conclude with 95% confidence that the cholesterol reading before and after nurse's lecture are not same; actually after lecture mean cholesterol reading was decreased.

Problem 12.7 (Lind, Marchal, and Wathen 2012) Advertisements by Sylph Fitness Center claim that completing its course will result in losing weight. A random sample of eight recent participants showed the following weights before and after completing the course. At the 0.01 significance level, can we conclude the students lost weight?

Name	Before	After
Hunter	155	154

Name	Before	After
Cashman	228	207
Mervine	141	147
Massa	162	157
Creola	211	196
Peterson	164	150
Redding	184	170
Poust	172	165

Solution:

Let, μ_B = pop. mean of weight **before** completing the course

μ_A = pop. mean of weight **after** completing the course

According to the **claim** ; $\mu_A < \mu_B$ that is $\mu_A - \mu_B < 0 \implies \mu_D < 0$. So the

Hypotheses

$$H_0 : \mu_D \geq 0$$

$$H_1 : \mu_D < 0$$

Do it yourself the rest of the calculation.....

[Ans: $\bar{d} = -8.875$; $s_D = 8.774$; $t = -2.861$, $-t_\alpha = -1.895$. Reject H_0 .]

Problem 12.8 (Lind, Marchal, and Wathen 2012) The management of Discount Furniture, a chain of discount furniture stores in the Northeast, designed an incentive plan for salespeople. To evaluate this innovative plan, 12 salespeople were selected at random, and their weekly incomes (\$) before and after the plan were recorded.

Salesperson	Before	After
Sid Mahone	320	340
Carol Quick	290	285
Tom Jackson	421	475
Andy Jones	510	510
Jean Sloan	210	210
Jack Walker	402	500
Peg Mancuso	625	631
Anita Loma	560	560
John Cuso	360	365
Carl Utz	431	431
A. S. Kushner	506	525
Fern Lawton	505	619

Was there a significant increase in the typical salesperson's weekly income due to the innovative incentive plan? Use the 0.05 significance level.

12 Analysis of Variance

The one-way analysis of variance, in particular, is used to test whether or not the averages from several **different situations** (AKA *Treatments*) are significantly different from one another.

This is the simplest kind of analysis of variance. Although more complex situations require more complicated calculations, the general ANOVA idea remains the same: to test significance by comparing one source of variability (the one being tested) against another source of variability (the underlying randomness of the situation).

The F test for the one-way analysis of variance will tell you whether the averages of several independent samples are significantly different from one another.

12.1 Sources of Variation for a One-Way Analysis of Variance

- Between-sample variability (from one sample to another).
- Within-sample variability (inside each sample).

12.2 Assumptions for a One-Way Analysis of Variance

- For each population, the response variable is normally distributed.
- The variance of the response variable, denoted σ^2 , is the same for all of the populations.
- The observations must be independent.

12.3 Hypotheses for a One-Way Analysis of Variance

Hypothesis

$H_0 : \mu_1 = \mu_2 = \dots = \mu_k$ (*The population means are equal*)

$H_a : \text{At least TWO population means are NOT equal}$

12.4 Sample statistics used in one-way ANOVA

- Total sample size, $n_T = n_1 + n_2 + \dots + n_k$
- Overall sample mean,

$$\bar{\bar{x}} = \frac{n_1\bar{x}_1 + n_2\bar{x}_2 + \dots + n_k\bar{x}_k}{n_T}$$

If, $n_1 = n_2 = \dots = n_k = n$ (say) then

$$\bar{\bar{x}} = \frac{\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_k}{k} \quad (\text{why?})$$

12.5 The Between-Sample Variability for One-Way Analysis of Variance

Between-sample variability AKA **MSTR** (mean square due to treatments):

$$MSTR = \frac{n_1(\bar{x}_1 - \bar{\bar{x}})^2 + n_2(\bar{x}_2 - \bar{\bar{x}})^2 + \dots + n_k(\bar{x}_k - \bar{\bar{x}})^2}{k - 1}$$

12.6 The Within-Sample Variability for One-Way Analysis of Variance

Within-sample variability AKA **MSE** (mean square due to error):

$$MSE = \frac{(n_1 - 1)s_1^2 + (n_2 - 1)s_2^2 + \dots + (n_k - 1)s_k^2}{n_T - k}$$

12.7 The F Statistic

$$F = \frac{\text{Between sample variability}}{\text{Within sample variability}} = \frac{MSTR}{MSE}$$

- The test statistic follows an F -distribution with $df_1 = k - 1$ degrees of freedom in the **numerator** and $df_2 = n_T - k$ degrees of freedom in the **denominator**.

12.8 Rejection rule

- Reject H_0 if $F \geq F_\alpha$.

12.9 Example 12.1


The following data are from a completely randomized design.

Treatment		
A	B	C
162	142	126
142	156	122
165	124	138
145	142	140
148	136	150
174	152	128

Sample data in *long/tidy* format

Treatment	Value
A	162
A	142
A	165
A	145
A	148
A	174
B	142
B	156
B	124
B	142
B	136
B	152
C	126
C	122
C	138
C	140
C	150
C	128

Now we see how the “*Values*” are distributed for 3 different *Treatments*.



Question

At the $\alpha = 0.05$ level of significance, test whether the means for the three treatments are equal.

Solution

- Hypothesis*

$H_0 : \mu_1 = \mu_2 = \mu_3$ (The population means are equal)

$H_a : \text{At least TWO population means are NOT equal}$

- Treatment-wise sample mean and sample variance

Treatment	Sample size	Sample mean	Sample variance
A	6	156	164.4
B	6	142	131.2
C	6	134	110.4

- Overall mean:

$$\bar{\bar{x}} = \frac{\bar{x}_1 + \bar{x}_2 + \dots + \bar{x}_k}{k} = \frac{156 + 142 + 134}{3} = 144$$

- MSTR: Since, $n_1 = n_2 = n_3 = n = 6$ so,

$$\begin{aligned} MSTR &= \frac{n[(\bar{x}_1 - \bar{\bar{x}})^2 + (\bar{x}_2 - \bar{\bar{x}})^2 + (\bar{x}_3 - \bar{\bar{x}})^2]}{k - 1} \\ &= \frac{6[(156 - 144)^2 + (142 - 144)^2 + (134 - 144)^2]}{3 - 1} = 744 \end{aligned}$$

- MSE: Since, $n_1 = n_2 = n_3 = n = 6$ so,

$$MSE = \frac{(n-1)s_1^2 + (n-1)s_2^2 + (n-1)s_3^2}{n_T - k}$$

$$= \frac{(n-1)[s_1^2 + s_2^2 + s_3^2]}{n_T - k} = \frac{(6-1)[164.4 + 131.2 + 110.4]}{18 - 3} = 135.33$$

- F -statistic:

$$F = \frac{MSTR}{MSE} = \frac{744}{135.33} = 5.4978 \approx 5.50$$

- Critical value

At $\alpha = 0.05$ and for $df_1 = 2$ and $df_2 = 15$, $F_\alpha = 3.68$

- Decision

Since $F > F_\alpha$ so reject H_0 .

- Conclusion

So, the equality of 3 means claim is rejected. Hence, at least TWO of the means are not equal.

12.10 Multiple comparison

Rejection of the null hypothesis (k population means are equal) in one-way ANOVA suggests that at least 2 population means are not equal. To investigate further which means are significantly differs we conduct multiple comparison test. In this section we will introduce Fisher's Least significant difference (LSD) method, then we discuss Bonferroni Adjustment to LSD Method.

- **Fisher's LSD method**

We define the LSD as

$$LSD = t_{\alpha/2} \sqrt{MSE \left(\frac{1}{n_i} + \frac{1}{n_j} \right)} ; i \neq j = 1, 2, \dots, k$$

We conclude that $\mu_i \neq \mu_j$ if $|\bar{x}_i - \bar{x}_j| > LSD$.

- **Bonferroni Adjustment to LSD Method**

To control the **Type I** error rate, we adjust the α as follows:

$$\alpha = \frac{\alpha_E}{C}$$

Where, α_E is the experiment-wise **Type I error** (that is given default)

$C = \binom{k}{2}$, the number of pairs to be compared.

Based on update α we take $t_{\alpha/2}$ and compute **LSD**.

12.11 Example 12.2

12.11.1 One-way ANOVA in R

ANOVA table

Source of variation	df	SSTR	MSTR	F-statistic	p-value
Treatment	2	1488	744.000	5.498	0.016
Residuals	15	2030	135.333	NA	NA

- Since $p\text{-value} < \alpha$ so, reject H_0 .
- Therefore the equality of 3 means claim is rejected. Hence, at least TWO of the means are not equal.

Homework

- 1) “Does the height of the shelf affect daily sales of dog food?”. To answer this question daily sales data were collected where dog foods were randomly allocated in three different height of shelves in 8 days.

Shelf Height		
Knee Level	Waist Level	Eye Level
77	88	85
82	94	85
86	93	87
78	90	81
81	91	80
86	94	79
77	90	87
81	87	93

- (a) Based on the data, is there a significant difference in the average daily sales of this dog food based on shelf height? Use a 0.01 level of significance.
- (b) If null hypothesis is rejected in (a) then conduct a multiple comparison test using Fisher LSD.

- 2) Many college and university students obtain summer jobs. A statistics professor wanted to determine whether students in different degree programs earn different amounts. A random sample of 5 students in the BA, BSc, and BBA programs were asked to report what they earned the previous summer. The results (in \$1,000s) are listed here.

Can the professor infer at the 5% significance level that students in different degree programs differ in their summer earnings?

B.A.	B.Sc.	B.B.A.
3.3	3.9	4.0
2.5	5.1	6.2
4.6	3.9	6.3
5.4	6.2	5.9
3.9	4.8	6.4

- 3) Perform a one-way ANOVA to determine whether there is a significant difference in the mean ages of the workers at the three plants. Use $\alpha = 0.01$ and note that the sample sizes are equal.

Plant	Age					
Plant 1	29	27	30	27	28	
Plant 2	32	33	31	34	30	
Plant 3	25	24	24	25	26	

4. A corporation is trying to decide which of three makes of automobile to order for its fleet—domestic, Japanese, or European. Five cars of each type were ordered, and, after 10,000 miles of driving, the operating cost per mile of each was assessed. The accompanying results in cents per mile were obtained.

Domestic	Japanese	European
18.0	20.1	19.3
15.6	15.6	15.4
15.4	16.1	15.1
19.1	15.3	18.6
16.9	15.4	16.1

- Prepare the analysis of variance table for these data.
- Test the null hypothesis that the population mean operating costs per mile are the same for these three types of car. Use $\alpha = 0.01$.
- Conduct Fisher's LSD pairwise comparison if applicable.

13 Chi-squared Test

13.1 Goodness of Fit Test

In this section we use a chi-square test to determine whether a population being sampled has a specific probability distribution.

13.1.1 A Multinomial Population

Multinomial Experiment

A multinomial experiment is one that possesses the following properties.

1. The experiment consists of a fixed number n of trials.
2. The outcome of each trial can be classified into one of k categories, called *cells*.
3. The probability p_i that the outcome will fall into cell i remains constant for each trial. Moreover, $p_1 + p_2 + \dots + p_k = 1$
4. Each trial of the experiment is independent of the other trials.

Testing Market Shares

Company A has recently conducted aggressive advertising campaigns to maintain and possibly increase its share of the market (currently 45%) for fabric softener. Its main competitor, company B, has 40% of the market, and a number of other competitors account for the remaining 15%.

To determine whether the market shares changed after the advertising campaign, the marketing manager for company A solicited the preferences of a random sample of 200 customers of fabric softener.

Of the **200** customers, *102 indicated a preference for company A's product, 82 preferred company B's fabric softener, and the remaining 16 preferred the products of one of the competitors.* Can the analyst infer at the 5% significance level that customer preferences have changed from their levels before the advertising campaigns were launched?

We recognize this experiment as a multinomial experiment, and we identify the technique as the chi-squared goodness-of-fit test. Because we want to know whether the market shares have changed, we specify those precampaign market shares in the null hypothesis.

$$H_0 : p_1 = 0.45; \quad p_2 = 0.40; \quad p_3 = 0.15$$

The alternative hypothesis attempts to answer our question, Have the proportions changed? Thus,

H_1 : At least one p_i is not equal to its specified value

Chi-Squared Goodness-of-Fit Test Statistic

$$\chi^2 = \sum_{i=1}^k \frac{(f_i - e_i)^2}{e_i}$$

Where, f_i = observed frequency and e_i = expected frequency

Note that, $e_i = n * p_i$

The sampling distribution of the test statistic is approximately chi-squared distributed with $k - 1$ degrees of freedom, provided that the sample size is large.

Test Statistic calculation

Company	Observed frequency, f_i	Expected frequency, e_i	$(f_i - e_i)$	$\frac{(f_i - e_i)^2}{e_i}$
A	102	90	12	1.60
B	82	80	2	0.05
Other	16	30	-14	6.53
Total	200	200		$\chi^2 = 8.18$

Critical value

At $\alpha = 0.05$ and for $df = 3 - 1 = 2$, $\chi_\alpha^2 = 5.99$.

Decision

Since $\chi^2 > \chi_\alpha^2$ so reject null hypothesis.

Interpretation/ Conclusion

There is sufficient evidence at the 5% significance level to infer that the proportions have changed since the advertising campaigns were implemented.

Problem 14.1 Test the following hypotheses by using the χ^2 goodness of fit test.

$$H_0 : p_A = 0.40; \quad p_B = 0.40; \quad p_C = 0.20$$

$$H_a : \text{At least one } p \text{ is not equal to } H_0 \text{ value}$$

A sample of size 200 yielded 60 in category A, 120 in category B, and 20 in category C. Use $\alpha = 0.01$ and test to see whether the proportions are as stated in H_0 .

Problem 14.2 Television Audiences Across Networks. During the first 13 weeks of the television season, the Saturday evening 8 p.m. to 9 p.m. audience proportions were recorded as ABC 29%, CBS 28%, NBC 25%, and independents 18%. A sample of 300 homes two weeks after a Saturday night schedule revision yielded the following viewing audience data: ABC 95 homes, CBS 70 homes,

NBC 89 homes, and independents 46 homes. **Test** with $\alpha = .05$ to determine whether the viewing audience proportions changed.

Problem 14.3 M&M Candy Colors. Mars, Inc. manufactures M&M's, one of the most popular candy treats in the world. The milk chocolate candies come in a variety of colors including blue, brown, green, orange, red, and yellow. The overall proportions for the colors are .24 blue, .13 brown, .20 green, .16 orange, .13 red, and .14 yellow. In a sampling study, several bags of M&M milk chocolates were opened and the following color counts were obtained.

Color	Blue	Brown	Green	Orange	Red	Yellow
Count	105	72	89	84	70	80

Use a 0.05 level of significance and the sample data to test the hypothesis that the overall proportions for the colors are as stated above. What is your conclusion?

Problem 14.4 Traffic Accidents by Day of Week. The National Highway Traffic Safety Administration reported the percentage of traffic accidents occurring each day of the week. Assume that a sample of 420 accidents provided the following data.

Day	Sun	Mon	Tue	Wed	Thur	Fri	Sat
No. of accidents	66	50	53	47	55	69	80

Conduct a hypothesis test to determine if the proportion of traffic accidents is the same for each day of the week. Using a .05 level of significance, what is your conclusion?

13.1.2 Normal population (continuous)

To test whether a variable follows **normal distribution** with mean μ and variance σ^2 we will illustrate the following example.

Problem 14.5 A random sample of 500 car batteries was taken and the life of each battery was measured. Letting X denote battery life in years, suppose that the sample revealed the following distribution of battery life:

Life (in years)	Frequency
$X < 1$	12
$1 < X \leq 2$	94
$2 < X \leq 3$	170
$3 < X \leq 4$	188
$4 < X \leq 5$	28
$5 < X$	8
Total	500

Based on this data, test whether battery life follows a normal distribution with $\mu = 2.8$ and $\sigma^2 = 1.1^2$. Clearly state your hypotheses and use a significance level of $\alpha = 5\%$.

Solution:

Hypotheses


H_0 : The battery life follows a normal distribution with $\mu = 2.8$ and $\sigma^2 = 1.1^2$.

H_a : The battery life does not follow a normal distribution.

Test Statistic calculation

Life (in years)	Probability	$e_i = np_i$	f_i	$\frac{(f_i - e_i)^2}{e_i}$
$X < 1$	$P(X < 1)$ $= P(Z < -1.64)$ $= 0.0505$	25.25	12	6.9530
$1 < X \leq 2$	$P(1 < X \leq 2)$ $= P(-1.64 < Z \leq -0.73)$ $= 0.1826$	91.30	94	0.0798
$2 < X \leq 3$	0.3386	169.30	170	0.0029
$3 < X \leq 4$	0.2902	145.10	188	12.6837
$4 < X \leq 5$	0.1149	57.45	28	15.0966
$5 < X$	0.0228	11.40	8	1.0140
				$\chi^2 = 35.83$

Frequency histogram of Battery life (in years)



Critical value

At $\alpha = 0.05$, and for $df=6-1=5$, $\chi_{\alpha,5}^2 = 11.1$

Decision

Since $\chi^2 > \chi_{\alpha,5}^2$ so we can reject the null hypothesis.

13.1.3 Uniform distribution (continuous)

To test whether a variable follows **uniform distribution** between a to b we will illustrate the following example.

Problem 14.6 Suppose X , the amount of time a person stays in bed after their alarm goes off, is uniformly distributed between 5 and b minutes. Over 100 days, how long they slept past their alarm (X) were recorded (in minutes). However, only the number of days for which X within certain ranges was reported in the table below:

Time stays in bed	No. of days
$5 < X < 7$	40
$7 < X < 8$	22
$8 < X < 10$	38
Total	100

Based on the data given above, test whether b is equal to 10. Clearly state your hypotheses and use a significance level of $\alpha = 5\%$.

Solution:

If the X , the amount of time a person stays in bed after their alarm goes off, is uniformly distributed between 5 and b minutes then the data will fit the **uniform distribution** with parameter 5 to $b=10$ minutes. So following hypotheses can be formed:

Hypothesis

H_0 : The amount of time a person stays in bed after their alarm goes off, is uniformly distributed between 5 and $b = 10$ minutes

H_a : The amount of time a person stays in bed after their alarm goes off, is NOT uniformly distributed between 5 and $b = 10$ minutes.

Test statistic calculation

If $X \sim U(5, 10)$ then PDF,

$$f(x) = \frac{1}{10 - 5} = \frac{1}{5}; \quad 5 < x < 10$$

So, $P(5 < X < 7) = (7 - 5) * \frac{1}{5} = \frac{2}{5}$ and so on

Bed time (in mins)	f_i	p_i	$e_i = np_i$	$\frac{(f_i - e_i)^2}{e_i}$
$5 < X < 7$	40	$\frac{2}{5}$	40	0.00
$7 < X < 8$	22	$\frac{1}{5}$	20	0.20
$8 < X < 10$	38	$\frac{3}{5}$	40	0.10
Totals	100			$\chi^2 = 0.3$

Critical value At $\alpha = 5\%$ and $df = 3 - 1 = 2$, $\chi_{\alpha, 2}^2 = 5.99$.

Decision Since, $\chi^2 < \chi^2_{\alpha,2}$ so we cannot reject null hypothesis. Hence the amount of time a person stays in bed after their alarm goes off, is uniformly distributed between 5 and $b = 10$ minutes.

Problem 14.7 Suppose Y , the number of minutes they are late to work is uniformly distributed between 0 and b minutes. Over 100 days, how late they were to work (Y) were recorded (in minutes). However, only the number of days for which Y fell within certain ranges was reported in the table below:

Time late to work	No. of days
$0 < Y < 2$	39
$2 < Y < 3$	25
$3 < Y < 5$	36
Total	100

Based on the data given above, test whether b is equal to 5. Clearly state your hypotheses and use a significance level of $\alpha = 5\%$.

13.2 Test for Independence (Categorical Data)

To determine whether two categorical variables are independent summarized in a contingency table we use **Chi-squared test of association /independence**. That is, to determine whether the distribution of one categorical variable is the same across all categories of the other categorical variable.

Consider the following example:

In an experiment to study the dependence of hypertension on smoking habits, the following data were taken on 180 individuals:

	Non-smokers	Moderate Smokers	Heavy Smokers
Hypertension	21	36	30
No hypertension	48	26	19

Test the hypothesis that the presence or absence of *hypertension* is independent of *smoking habits*. Use a 0.05 level of significance.

Solution: We have to test the following hypothesis:

H_0 : The column variable is independent of the row variable

H_a : The column variable is not independent of the row variable

Test statistic

$$\chi^2 = \sum_{i=1}^r \sum_{j=1}^c \frac{(f_{ij} - e_{ij})^2}{e_{ij}}$$

The sampling distribution of the test statistic is approximately chi-squared distributed with $(r - 1) \times (c - 1)$ degrees of freedom, provided that the sample size is large.

i Note

f_{ij} = Observed frequency of $(i, j)^{th}$ cell;
 e_{ij} = Expected frequency of $(i, j)^{th}$ cell = $\frac{\text{Row } i \text{ total} \times \text{Column } j \text{ total}}{\text{Sample size}(n)}$

Table: Contingency table with Row total and Column total

	Non-smokers	Moderate Smokers	Heavy Smokers	Row Total
Hypertension	21	36	30	87
No hypertension	48	26	19	93
Column Total	69	62	49	180

For example,

$$e_{11} = \frac{87 \times 69}{180}$$

Chi-square Statistic calculation

Observed, f_i	Expected, e_i	$\frac{(f_i - e_i)^2}{e_i}$
21	33.35	4.57
36	29.97	1.21
30	23.68	1.68
48	35.65	4.28
26	32.03	1.14
19	25.32	1.14
		$\chi^2 = 14.46$

Critical value At $\alpha = 0.01$ and with $df = (2 - 1) * (3 - 1) = 2$, $\chi^2_{\alpha} = 9.21$

Decision Since $\chi^2 > \chi^2_{\alpha}$ so reject the null hypothesis.

Interpretation/conclusion There is sufficient evidence at the 5% significance level to infer that the *smoking habits (column variable)* is not independent of the *the presence or absence of hypertension (row variable)*, rather the two variables are associated.

Problem 14.8 Suppose X, the amount of time a person stays in bed after their alarm goes off, is uniformly distributed between 5 and 10 minutes. Also, suppose Y, the number of minutes they are late to work is uniformly distributed between 0 and 5 minutes. Over 100 days, how long they slept past their alarm (X) and how late they were to work (Y) were recorded (in minutes). However, only the number of days for which X and Y fell within certain ranges was reported in the table below:

	$5 < X < 7$	$7 < X < 8$	$8 < X < 10$
$0 < Y < 2$	18	9	12
$2 < Y < 3$	9	4	12
$3 < Y < 5$	13	9	14

Based on the data above, **test** whether X and Y are independent (that is, whether the rows and columns are independent). Clearly state your hypotheses and use a significance level of $\alpha = 5\%$.

Problem 14.9 The contingency table shows the results of a random sample of students by the location of school and the number of those students achieving basic skill levels in three subjects. At $\alpha = 0.01$, **test** the hypothesis that the variables (Subject vs. Location) are independent.

	Reading	Math	Science
Urban	43	42	38
Suburban	63	66	65

Problem 14.10 The Athlete Injury Data by Stretching Status is given below:

	Athlete has: Stretched	Athlete has: Not stretched
Injury	18	22
No injury	211	189

Do these data suggest that the result of the athlete's activity (Injury vs. No injury) is statistically independent of whether the athlete stretched prior to the activity (Stretched vs. Not stretched)?

14 Correlation and Simple Linear Regression

In real world we often observe that a change in one variable is associated with the change in another variable.

In statistics, **correlation** refers to *degree* and *direction* of **linear relationship** between two quantitative (interval or ratio scale) variables. For example-

- As *income* increases *expense* also increases (positive correlation);
- As *resistance* increases *current flow* decreases (negative correlation) etc.

To understand the nature and to measure the **linear relationship (correlation)** between two quantitative variable we use some techniques. In the following section we will discuss about that.

14.1 Scatter plot: Graphical method to explore correlation

A **scatter plot** shows the relationship between two quantitative variables measured for the same individuals. The values of one variable appear on the horizontal axis, and the values of the other variable appear on the vertical axis. Each individual in the data appears as a point on the graph.

- A scatterplot *displays* the strength, direction, and form of the relationship between two quantitative variables (*see* Figure 14.1).

Lets draw a scatter plot for the following data.

Table 14.1: Data

	1	2	3	4	5	6	7	8	9	10
x	2	5	1	3	4	1	5	3	4	2
y	50	57	41	54	54	38	63	48	59	46

In fact, the scatter lot suggests that a straight line could be used as an approximation of the relationship. In the following discussion, we introduce **covariance, coefficient of correlation, and coefficient of determination** as descriptive measures which provide *direction* and *strength* of the linear relationship between two variables.




Figure 14.1: Types of correlations that can be represented using a scatter plot

14.2 Covariance

The covariance between X and Y is defined as follows

Population covariance

$$\sigma_{XY} = \frac{\sum_{i=1}^N (x_i - \mu_X)(y_i - \mu_Y)}{N}$$

Sample covariance

$$s_{XY} = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{n - 1}$$

Shortcut for Sample Covariance

$$s_{XY} = \frac{\sum xy - n\bar{x}\bar{y}}{n - 1}$$

Or,

$$s_{XY} = \frac{1}{n - 1} \left[\sum xy - \frac{\sum x \sum y}{n} \right] \quad (14.1)$$

Example: Compute sample covariance between X and Y from Table 14.1.




Figure 14.2: Scatter plot between X and Y

Solution:

Here, $n = 10$; $\sum x = 30$; $\sum y = 510$.

$$\sum xy = x_1y_1 + x_2y_2 + \cdots + x_ny_n = 1629$$

Hence the sample covariance,

$$\begin{aligned} s_{XY} &= \frac{1}{n-1} \left[\sum xy - \frac{\sum x \sum y}{n} \right] \\ &= \frac{1}{10-1} \left[1629 - \frac{30 \times 510}{10} \right] \end{aligned}$$

$$\therefore s_{XY} = 11$$

Since, $s_{XY} > 0$ so there exists positive correlation between X and Y .

Drawback of covariance

According to Keller (2014) , “Unfortunately, the magnitude may be difficult to judge. For example, if you’re told that the covariance between two variables is 500, does this mean that there is a strong linear relationship? The answer is that it is impossible to judge without additional statistics. Fortunately, we can improve on the information provided by this statistic by creating another one.”

14.3 Coefficient of Correlation

The **Pearson product-moment coefficient of correlation** is defined as the covariance divided by the standard deviations of the variables.

Population correlation coefficient

$$\rho = \frac{\sigma_{XY}}{\sigma_X \times \sigma_Y} \quad ; \quad -1 \leq \rho \leq +1$$

Sample correlation coefficient

$$r = \frac{s_{XY}}{s_X \times s_Y} \quad ; \quad -1 \leq r \leq +1 \quad (14.2)$$

Where

s_{XY} is sample covariance between X and Y s_X is sample standard deviation of X and s_Y is sample standard deviation of Y .

Note: The sample correlation coefficient can be expressed in other forms:

(a)

$$r = \frac{\sum (x_i - \bar{x})(y_i - \bar{y})}{\sqrt{\sum (x_i - \bar{x})^2} \sqrt{\sum (y_i - \bar{y})^2}}$$

(b)

$$r = \frac{n \sum xy - (\sum x)(\sum y)}{\sqrt{[n \sum x^2 - (\sum x)^2]} \sqrt{[n \sum y^2 - (\sum y)^2]}}$$

(c)

$$r = \frac{1}{n-1} \sum_{i=1}^n z_{x_i} z_{y_i}$$

Where, z_x and z_y are the z-score of X and Y respectively.

14.4 Properties of coefficient of correlation

1. The value of ρ is always between -1 and 1 inclusive. That is $-1 \leq \rho \leq +1$.
2. If all values of either variable are converted to a different scale, the value of ρ does not change (NOT affected by change of *origin* and *scale*).
3. The value of ρ is not affected by the choice of X or Y . Interchange all x values and y values, and the value of ρ will not change (ρ is a symmetric measure).
4. ρ measures the strength of a *linear relationship*. It is not designed to measure the strength of a relationship that is not linear.
5. ρ is very sensitive to outliers in the sense that a single outlier could dramatically affect its value.

i Proof of $-1 \leq \rho \leq +1$

From Cauchy-Schwarz Inequality we know that,

$$(E[XY])^2 \leq E[X^2]E[Y^2]$$

Now center the variables X and Y with respect to their mean we have,

$$(E[(X - \mu_X)(Y - \mu_Y)])^2 \leq E[(X - \mu_X)^2]E[(Y - \mu_Y)^2]$$

$$\Rightarrow [Cov(X, Y)]^2 \leq Var(X) \cdot Var(Y)$$

$$\Rightarrow (\sigma_{XY})^2 \leq \sigma_X^2 \sigma_Y^2$$

$$\Rightarrow |\sigma_{XY}| \leq \sigma_X \sigma_Y$$

$$\Rightarrow \left| \frac{\sigma_{XY}}{\sigma_X \sigma_Y} \right| \leq 1$$

$$\Rightarrow |\rho| \leq 1$$

Which implies that $-1 \leq \rho \leq +1$.

i Effect of change of origin and scale on ρ

Let, $U = \frac{X-a}{b}$ and $V = \frac{Y-c}{d}$. Here a, b, c, d are any arbitrary real numbers.

We have to prove that, $\rho_{XY} = \rho_{UV}$.

Proof:

So we have,

$X = a + bU$; and $\bar{X} = a + b\bar{U}$. Similarly

$Y = c + dV$; and $\bar{Y} = c + d\bar{V}$

Now,

$$\sigma_{XY} = \frac{\sum(X-\bar{X})(Y-\bar{Y})}{N} = \frac{bd\sum(U-\bar{U})(V-\bar{V})}{N} = bd \times \sigma_{UV}$$

$$\sigma_X^2 = Var(X) = Var(a + bU) = b^2 Var(U) = b^2 \sigma_U^2$$

Hence, $\sigma_X = b\sigma_U$ (standard deviation is always ≥ 0).

Similarly, $\sigma_Y = d\sigma_V$

Finally we have,

$$\rho_{XY} = \frac{\sigma_{XY}}{\sigma_X \sigma_Y} = \frac{bd \times \sigma_{UV}}{b\sigma_U \times d\sigma_V} = \frac{\sigma_{UV}}{\sigma_U \sigma_V} = \rho_{UV} \quad (PROVED.)$$

14.4.1 Interpretation of correlation coefficient

- (a) $r = -1$ implies *perfect negative* correlation,
- (b) $r = +1$ implies *perfect positive* correlation,
- (c) $r \approx 0$ implies no correlation or very weak correlation,
- (d) As r close to -1 , the degree of *negative* correlation becomes stronger,
- (e) As r close to $+1$, the degree of *positive* correlation becomes stronger.

14.4.2 Computing the Coefficient of Correlation

Let us compute sample correlation coefficient between X and Y from Table 14.1.

Here, $n = 10$; $\sum x = 30$; $\sum y = 510$.

$$\sum xy = x_1y_1 + x_2y_2 + \dots + x_ny_n = 1629$$

$$\sum x^2 = 2^2 + 5^2 + \dots + 2^2 = 110$$

$$\sum y^2 = 50^2 + 57^2 + \dots + 46^2 = 26576$$

$$\bar{x} = 3$$

$$\bar{y} = 51$$

So,

$$s_X = \sqrt{\frac{\sum x^2 - n \bar{x}^2}{n - 1}} = 1.490712$$

$$s_Y = \sqrt{\frac{\sum y^2 - n \bar{y}^2}{n-1}} = 7.930252$$

$$s_{XY} = \frac{\sum xy - n \bar{x} \bar{y}}{n-1} = 11$$

Hence,

$$r = \frac{s_{XY}}{s_X \times s_Y} = \frac{11}{1.490712 \times 7.930252} = 0.9305$$

which is close to 1. So the correlation between X and Y is *strong* and *positive*.

14.4.3 Exercises: Constructing a Scatter Plot and Determining Correlation

In Exercises 1–4, (a) display the data in a scatter plot, (b) calculate the sample correlation coefficient r , and (c) describe the type of correlation and interpret the correlation in the context of the data.

1. **Age and Blood Pressure.** The ages (in years) of 10 men and their systolic blood pressures (in millimeters of mercury) (Larson and Farber 2015, 482)

Age, x	16	25	39	45	49	64	70	29	57	22
Systolic blood pressure, y	109	122	143	132	199	185	199	130	175	118

2. **Driving Speed and Fuel Efficiency.** A department of transportation's study on driving speed and miles per gallon for midsize automobiles resulted in the following data (Anderson 2020a, 149):

Speed (Miles per Hour)	30	50	40	55	30	25	60	25	50	55
Miles per Gallon	28	25	25	23	30	32	21	35	26	25

3. Are the marks one receives in a course related to the amount of time spent studying the subject? To analyze this mysterious possibility, a student took a random sample of 10 students who had enrolled in an accounting class last semester. He asked each to report his or her mark in the course and the total number of hours spent studying accounting. These data are listed here (Keller 2014).

Study time	40	42	37	47	25	44	41	48	35	28
Marks	77	63	79	86	51	78	83	90	65	47

4. The owner of a paint store was attempting to analyse the relationship between advertising and sales, and recorded the monthly advertising budget (\$'000) and the sales (\$m) for a sample of 12 months. The data are listed here

Advertising	23	46	60	54	28	33	25	31	36	88	95	99
Sales	8	11	13	13	8.9	10.7	9	10.4	11	14	14.4	15.9

14.4.4 Coefficient of determination

The coefficient of determination measures the amount of variation in the dependent variable that is explained by the variation in the independent variable.

For example, if $r = 0.8764$ between X and Y then **coefficient of determination** $r^2 = (0.8764)^2 \approx 0.7681$.

Interpretation: The $r^2 = 0.7681$ tells us that 76.81% variation in Y (dependent variable) can be explained by X (independent variable).

14.4.5 Correlation vs. causation

Correlation does not always imply *causation*. For example,

- A study (Messerli 2012) found that there was a significant ($r = 0.791$) positive correlation between *chocolate consumption per capita* and *number of Nobel laureates per 10 million persons*. This does not necessarily implies that more a country consumes chocolate, more the chance of getting a Nobel prize. Rather differences in socioeconomic status from country to country and geographic and climatic factors may play some role to win a Nobel prize.
- We might find that there is a positive correlation between the time spent driving on road and the number of accidents but this does not mean that spending more time on road causes accident. Because in that case, in order to avoid accidents one may drive fast so that time spent on road is less (Selvamuthu and Das 2024).

14.4.6 Effect of outlier on correlation coefficient

The correlation coefficient is heavily affected by outlier (see Figure 14.3). It changes the magnitude of the correlation coefficient.

Even sometime the outlier(s) can change the sign of the correlation coefficient (see Figure 14.4).

14.5 Rank correlation

To measure the association between two ordinal or rank-ordered data we use the **Spearman Rank-correlation coefficient (RCC)**. Even if in presence of outliers in interval or ratio scale data we can use RCC. The sample Spearman RCC is computed as follows:

$$r_s = 1 - \frac{6 \sum_{i=1}^n d_i^2}{n(n^2 - 1)} \quad (14.3)$$




Figure 14.3: Effect of outlier on the magnitude of correlation coefficient (r)




Figure 14.4: Effect of outlier on the sign of correlation coefficient (r)

Where,

n = the number of observations in the sample

x_i = the rank of observation i with respect to the first variable

y_i = the rank of observation i with respect to the second variable

$$d_i = x_i - y_i$$

The interpretation is as usual as Pearson correlation coefficient.

Problem 15.1 Two sportswriters have ranked 10 Olympic sports according to how interesting they are to watch in person, with the results shown here. **Calculate** and **interpret** the Spearman coefficient of rank correlation.

Olympic Sport	Bob's Ranking	Tom's Ranking
Track & field	1	2
Basketball	2	4
Volleyball	3	5
Swimming	4	1
Boxing	5	8
Ice skating	6	3
Weight lifting	7	6
Wrestling	8	7
Judo	9	10
Equestrian	10	9

Problem 15.2 Quality of Teaching Assessments. A student organization surveyed both current students and recent graduates to obtain information on the quality of teaching at a particular university. An analysis of the responses provided the following teaching-ability rankings. Do the rankings given by the current students agree with the rankings given by the recent graduates ? (Anderson 2020b)

Professor	Current Students	Recent Graduates
1	4	6
2	6	8
3	8	5
4	3	1
5	1	2
6	2	3
7	5	7
8	10	9
9	7	4
10	9	10

14.6 Simple linear regression (SLR)

In regression analysis we try to estimate or predict the **outcome/ response** of one variable (dependent variable) on the basis of other variables (independent variable). For example, sales of certain product depends of price, advertising cost, quality of the product, brand etc.

It involves developing a mathematical model or equation that describes the relationship between the **dependent variable** and the **independent variables**.

In the following section we will discuss about the **Simple linear regression (SLR)** where a independent variable and a dependent variable are involved.

14.6.1 Population regression function (PRF)

$$y_i = \beta_0 + \beta_1 x_i + \epsilon_i \quad ; i = 1, 2, 3, \dots, N \quad (14.4)$$

Where,

y = dependent variable

x = independent variable

β_0 = y-intercept

β_1 = slope of the line/ regression coefficient

ϵ = error variable

Assumptions

- i) The errors are *independently, identically* normally distributed with **constant variance** σ_ϵ^2 that is $\epsilon_i \stackrel{\text{i.i.d.}}{\sim} \mathcal{N}(0, \sigma_\epsilon^2)$.
- ii) The independent variable should not be correlated with the error term (no endogeneity)

Taking conditional expectation of of **PRF** for a given x_i we have

$$E(y_i/x_i) = \beta_0 + \beta_1 x_i \quad (14.5)$$

From sample data we have to estimate $E(y_i/x_i)$ which is equivalent to estimate β_0 and β_1 .

14.6.2 Ordinary least square (OLS) estimate of $E(y_i/x_i)$

Let we have n pairs of sample data: $\{X, Y\} = \{(x_1, y_1), (x_2, y_2), \dots, (x_n, y_n)\}$. From the sample data suppose the **estimated regression line of $E(y_i/x_i)$ is**

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i \quad ; i = 1, 2, \dots, n$$

So,

$$y_i = \hat{y}_i + e_i \quad ; i = 1, 2, \dots, n$$

Where, e_i is the **estimated error or residual**.

Now, the **residuals sum of square (RSS)** is given as:

$$RSS = \sum_{i=1}^n e_i^2 = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = \sum_{i=1}^n (y_i - \hat{\beta}_0 - \hat{\beta}_1 x_i)^2 \quad (14.6)$$

In **OLS** method we *minimize* the **RSS** with respect to $\hat{\beta}_0$ and $\hat{\beta}_1$. To minimize **RSS** we have to set two equations using *partial derivative*:

$$\frac{\partial (RSS)}{\partial \hat{\beta}_0} = 0 \quad (14.7)$$

$$\frac{\partial (RSS)}{\partial \hat{\beta}_1} = 0 \quad (14.8)$$

By solving the two equations we will have the **OLS estimates** of β_0 and β_1 as follows:

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x}$$

and

$$\hat{\beta}_1 = \frac{\sum_{i=1}^n (x_i - \bar{x})(y_i - \bar{y})}{\sum_{i=1}^n (x_i - \bar{x})^2} = \frac{s_{xy}}{s_x^2}$$

Or,

$$\hat{\beta}_1 = r_{xy} \left(\frac{s_y}{s_x} \right)$$

Thus, the **estimated regression line** is given by

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i$$

14.6.3 Point prediction of y for a given x

Given $x = x_g$. Then the estimated y for given x is

$$\hat{y}_g = \hat{\beta}_0 + \hat{\beta}_1 \times x_g$$

14.6.4 Partition of sum squares

i) Total sum of square,

$$SS(Total) = \sum_{i=1}^n (y_i - \bar{y})^2 = (n-1)s_y^2$$

ii) Regression sum of square,

$$SSR = \sum_{i=1}^n (\hat{y}_i - \bar{y})^2$$

In the case of **SLR**, SSR can be written as

$$SSR = \hat{\beta}_1^2 \sum_{i=1}^n (x_i - \bar{x})^2 = \hat{\beta}_1^2 (n-1)s_x^2$$

iii) Sum of square of error,

$$SSE = \sum_{i=1}^n (y_i - \hat{y}_i)^2 = SS(Total) - SSR$$

14.6.5 Coefficient of determination (Goodness of fit)

$$R^2 = \frac{SSR}{SS(Total)} \quad ; \quad 0 \leq R^2 \leq 1$$

Interpretation: The R^2 explains the amount of variation in Y (dependent variable) by the estimated model.

In the case of **SLR**,

$$R^2 = r_{xy}^2 = \frac{s_{xy}^2}{s_x^2 \times s_y^2}$$

We can also write R^2 as:

$$R^2 = \frac{\hat{\beta}_1^2 \sum (x - \bar{x})^2}{\sum (y - \bar{y})^2}$$

14.6.6 Some problems on SLR

Problem 15.3 : The owner of a paint store was attempting to analyse the relationship between advertising and sales, and recorded the monthly advertising budget (\$'000) and the sales (\$m) for a sample of 12 months. The data are listed here:

Advertising	23	46	60	54	28	33	25	31	36	88	95	99
Sales	8	11	13	13	8.9	10.7	9	10.4	11	14	14.4	15.9

- i) **Identify** the dependent and independent variable.
- ii) **Plot** the data. Is it appear to be linear?
- iii) Now, **fit/ estimate** a linear regression line.
- iv) **Interpret** the regression/slope coefficient.
- v) **Predict** the sales for the advertising cost \$70, 000.
- vi) **Comment** about goodness of fit of the estimated model.

Problem 15.4 Age and Blood Pressure. The ages (in years) of 10 men and their systolic blood pressures (in millimeters of mercury).

Age	16	25	39	45	49	64	70	29	57	22
Systolic blood pressure (SBP)	109	128	143	145	180	185	185	199	175	118

- i) **Identify** the dependent and independent variable.
- ii) **Plot** the data. Is it appear to be linear?
- iii) Now, **fit/ estimate** a linear regression line.
- iv) **Interpret** the regression/slope coefficient.
- v) **Predict** the SBP for the age of a person is 40 years .
- vi) **Comment** about goodness of fit of the estimated model.

Solution:

Let, Y =Systolic blood pressure, SBP and X = age (in years)

i) Here SBP is dependent variable and age is independent variable

ii) **Scatter plot of Age versus SBP**

The scatter plot appears to be linear.

iii) Let the estimated regression line is :

$$\hat{y}_i = \hat{\beta}_0 + \hat{\beta}_1 x_i \quad (1)$$




Figure 14.5: Sacter plot of Age and SBP

From sample data;

$$n = 10, \sum x = 416, \sum y = 1512, \sum x^2 = 20398,$$

$$\sum y^2 = 237414, \sum xy = 67977$$

$$\bar{x} = 41.6, \quad \bar{y} = 151.2$$

So,

$$s_{xy} = \frac{\sum xy - n \cdot \bar{x} \cdot \bar{y}}{n - 1} = \frac{67977 - 10 \times 41.6 \times 151.2}{10 - 1} = 564.2$$

$$s_x^2 = \frac{\sum x^2 - n \times \bar{x}^2}{n - 1} = 343.6$$

$$s_y^2 = \frac{\sum y^2 - n \times \bar{y}^2}{n - 1} = 977.7333$$

Hence,

$$\hat{\beta}_1 = \frac{s_{xy}}{s_x^2} = \frac{564.2}{343.6} = 1.642026 \approx 1.642$$

and,

$$\hat{\beta}_0 = \bar{y} - \hat{\beta}_1 \bar{x} = 151.2 - 1.642026 \times 41.6 = 82.89172 \approx 82.8917$$

$$\therefore \hat{y}_i = 82.8917 + 1.642 x_i$$

iv) **Interpretation of the regression/slope coefficient:**

Here $\hat{\beta}_1 = 1.642$ implies that as one year increases, the SBP will increase by 1.642 Hg m on average.

v) **Predict the SBP for the age of a person is 40 years:**

For $x = 40$ years, the predicted SBP (in Hg m) is

$$\hat{y}_g = 82.8917 + 1.642 \times 40 = 148.5728 \text{ Hg m}$$

vi) **Comment about goodness of fit of the estimated model:**

The goodness of fit measure is

$$R^2 = \frac{s_{xy}^2}{s_x^2 \cdot s_y^2} = \frac{(564.2)^2}{(343.6)(977.7333)} = 0.9475$$

Hence, $R^2 = 0.9475$ implies that 94.75% variation in “SBP” can be explained by the estimated model.

15 Summary

References

- Anderson, David R. 2020a. *Statistics for Business & Economics*. 14e ed. Boston, MA: Cengage.
- . 2020b. *Statistics for Business & Economics*. 14e ed. Boston, MA: Cengage.
- Anderson, David R., and Dennis J. Sweeney. 2011. *Statistics for Business and Economics*. 11e [ed.]. Australia ; Mason, Ohio: South-Western Cengage Learning.
- Black, Ken. 2012. *Business statistics: for contemporary decision making*. 7th ed. Hoboken, NJ: Wiley.
- Keller, Gerald. 2014. *Statistics for Management and Economics*. 10e ed. Stamford, CT, USA : Cengage Learning.
- Larson, Ron, and Betsy Farber. 2015. *Elementary statistics: picturing the world*. 6. ed., global ed. Always Learning. Boston, Mass.: Pearson.
- Lind, Douglas A., William G. Marchal, and Samuel Adam Wathen. 2012. *Statistical Techniques in Business & Economics*. 15th ed. New York, NY: McGraw-Hill/Irwin.
- Messerli, Franz H. 2012. “Chocolate Consumption, Cognitive Function, and Nobel Laureates.” *New England Journal of Medicine* 367 (16): 1562–64. <https://doi.org/10.1056/nejmon1211064>.
- Montgomery, Douglas C., and George C. Runger. 2014. *Applied Statistics and Probability for Engineers*. Sixth edition. Hoboken, NJ: John Wiley; Sons, Inc.
- Moore, David S. 2010. *The Practice of Statistics for Business and Economics , Third Edition*. 3rd Revised edition. W. H. Freeman; libgen.li/file.php?md5=35437b5e2724b5b9af628d6bc2c86d97.
- Newbold, Paul, William L. Carlson, and Betty M. Thorne. 2013. *Statistics for business and economics*. 8. ed., global ed. Always learning. Boston, Mass. Munich: Pearson.
- R Core Team. 2024. *R: A Language and Environment for Statistical Computing*. Vienna, Austria: R Foundation for Statistical Computing. <https://www.R-project.org/>.
- Royston, J. P. 1982. “Algorithm AS 181: The w Test for Normality.” *Journal of the Royal Statistical Society. Series C (Applied Statistics)* 31 (2): 176–80. <http://www.jstor.org/stable/2347986>.
- Selvamuthu, Dharmaraja, and Dipayan Das. 2024. *Introduction to Probability, Statistical Methods, Design of Experiments and Statistical Quality Control*. Springer Nature Singapore. <https://doi.org/10.1007/978-981-99-9363-5>.
- Shapiro, Samuel Sanford, and Martin B Wilk. 1965. “An Analysis of Variance Test for Normality (Complete Samples).” *Biometrika* 52 (3-4): 591–611.
- Walpole, Ronald E., Raymond H. Myers, Sharon L. Myers, and Keying Ye. 2017. *Probability & statistics for engineers & scientists: MyStatLab update*. Ninth edition. Boston: Pearson.