

---

---

# DISCRETE MATHEMATICS

## MATH 381

---

---

BASIC CONCEPTS AND EXAMPLES EXPLAINING THE FUNDAMENTALS OF  
DISCRETE MATHEMATICS.

By  
MANSI SAKARVADIA  
*The University of North Carolina*  
*Chapel Hill*

## Contents

<b>1 February 17, 2020</b>	<b>1</b>
1.1 Restriction of Domain . . . . .	1
1.2 Restriction of Codomain . . . . .	2
1.3 Arithmetic of functions . . . . .	2

## 1 February 17, 2020

Recall:

- Functions  $f: A \rightarrow B$
- Image  $Im(f) = f(A)$

If  $f(a) = b$ , say "a is a preimage of b"

- $Gr(f) = \{(a, b) | f(a) = b\} \subseteq A \times B$
- $Gr(f) = \{(a, f(a)) | a \in A\}$

Graph  $Gr(f)$  is a relation between A and B

Which binary relations (subsets of  $A \times B$ ) are graphs of functions?

- A subset  $s \subseteq A \times B$  is the graph of a function if for every element  $a \in A$ , there is a unique element  $b \in B$  such that  $(a, b) \in S$ .
- key Can't have  $(a, b_1)$  and  $(a, b_2) \in S$  where  $b_1 \neq b_2$  and expect S to be a graph
- (abstraction of "straight line test" about graphs  $f: \mathbb{R} \rightarrow \mathbb{R}$ )

### 1.1 Restriction of Domain

suppose  $f: A \rightarrow B$

Consider  $A' \subseteq A$

DEFINITION: the restriction of f to A' is  $f|_{A'}: A' \rightarrow B$  defined by  $(f|_{A'})(a) = f(a), a \in A'$

KEY POINT: What does it mean for 2 functions  $f: A \rightarrow B$  and  $g: C \rightarrow D$  to be equal?

- NEED:  $A=C$ ,  $B=D$ , and  $f(a) = g(a) \forall a \in A$

## 1.2 Restriction of Codomain

If  $B'$  is a set with  $Im(f) \subseteq B' \subseteq B$ , then we consider:

$$f' : A \rightarrow B'$$

defined by  $f'(a) = f(a)$  for all  $a \in A$

$$\text{EX. } Im(f|_{A'}) = f(A')$$

- $Im(f|_{A'}) \leftarrow$  image of the restriction of  $f$  to  $A'$
- $f(A') \leftarrow$  image of the subset  $A' \subseteq A$  under  $f$

## 1.3 Arithmetic of functions

- A function is called real-valued if its codomain is  $\in \mathbb{R}$
- A function is called inter-valued if its codomain is  $\in \mathbb{Z}$
- DEFINITION: Suppose that  $f_1$  and  $f_2$  are two real-valued functions both w/ domain  $A$ . Then we have  $f_1 + f_2$  and  $f_1 f_2$  (the sum and product), two real-valued functions on  $A$ , defined by:

$$- f_1 + f_2(x) = f_1(x) + f_2(x)$$

$$- f_1 f_2(x) = f_1(x) * f_2(x)$$