

**CBSE Sample Paper-06**  
**Mathematics**  
**Class – XII**

Time allowed: 3 hours

**ANSWERS**

Maximum Marks: 100

**Section A**

1. Solution:

$$[2.1] = 2, [2.3] = 2, \text{ thus it is not one-one.}$$

Since it takes only integral values, hence it is not onto also.

2. Solution:

$$\text{Consider } (AB - BA)' = (AB)' - (BA)' = B'A' - A'B' = BA - AB = -(AB - BA)$$

Thus, AB-BA is skew symmetric.

3. Solution:

The operation \* is not a binary operation as  $2*3 = 2-3 = -1 \notin \mathbb{Z}^+$ .

4. Solution:

$$(AB)^{-1} = B^{-1}A^{-1} = \begin{bmatrix} 3 & 0 \\ -1 & 6 \end{bmatrix} \begin{bmatrix} 1 & 3 \\ -1 & 2 \end{bmatrix} = \begin{bmatrix} 3 & 9 \\ -7 & 9 \end{bmatrix}$$

5. Solution:

$$\vec{a} = 3i + 4j \text{ and } \vec{b} = 4i + 3j$$

$$\cos \theta = \frac{\vec{a} \cdot \vec{b}}{|\vec{a}| |\vec{b}|} = \frac{12+12}{5.5} = \frac{24}{25} \Rightarrow \theta = \cos^{-1} \left( \frac{24}{25} \right)$$

6. Solution:

$$\begin{vmatrix} 1 & a & b+c \\ 1 & b & a+c \\ 1 & c & a+b \end{vmatrix} = \begin{vmatrix} 1 & a & a+b+c \\ 1 & b & b+a+c \\ 1 & c & c+a+b \end{vmatrix} \quad (C_3 \rightarrow C_2 + C_3)$$

$$= (a+b+c) \begin{vmatrix} 1 & a & 1 \\ 1 & b & 1 \\ 1 & c & 1 \end{vmatrix} = 0 \quad (\because C_3 = C_1)$$

**Section B**

7. Solution:

$$\text{Let } x = \cos 2\theta$$

$$\theta = \frac{1}{2} \cos^{-1} x$$

$$\therefore \tan^{-1} \left( \frac{\sqrt{1+x} - \sqrt{1-x}}{\sqrt{1+x} + \sqrt{1-x}} \right) = \tan^{-1} \left( \frac{\sqrt{1+\cos 2\theta} - \sqrt{1-\cos 2\theta}}{\sqrt{1+\cos 2\theta} + \sqrt{1-\cos 2\theta}} \right)$$

$$\text{Now, } \frac{\sqrt{1+\cos 2\theta} - \sqrt{1-\cos 2\theta}}{\sqrt{1+\cos 2\theta} + \sqrt{1-\cos 2\theta}} = \frac{\sqrt{2\cos^2 \theta} - \sqrt{2\sin^2 \theta}}{\sqrt{2\cos^2 \theta} + \sqrt{2\sin^2 \theta}} = \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta}$$

$$\tan^{-1} \left( \frac{\cos \theta - \sin \theta}{\cos \theta + \sin \theta} \right) = \tan^{-1} \left( \frac{1 - \tan \theta}{1 + \tan \theta} \right) = \tan^{-1} \left( \tan \left( \frac{\pi}{4} - \theta \right) \right) = \frac{\pi}{4} - \theta$$

$$= \frac{\pi}{4} - \frac{1}{2} \cos^{-1} x$$

8. Solution:

$$\vec{a} = 5i + 2j - 4k, \vec{N} = 2i + 3j - k$$

The equation of a plane is given by  $(\vec{r} - \vec{a}) \cdot \vec{N} = 0$

$$\Rightarrow [\vec{r} - (5i + 2j - 4k)] \cdot (2i + 3j - k) = 0$$

Transforming into Cartesian form we get,  $[(x-5)i + (y-2)j + (z+4)k] \cdot (2i + 3j - k) = 0$ , i.e.  $2x + 3y - z = 20$ .

9. Solution:

(A) Let A denote the event that problem is solved by A and let B denote the event that problem is solved by B.

$$\therefore P(A) = 1/2, P(B) = 1/3, P(\bar{A}) = 1 - 1/2 = 1/2, P(\bar{B}) = 2/3$$

$$P(\text{Problem is solved}) = 1 - P(\text{Problem is not solved}) = 1 - P(\bar{A}\bar{B}) = 1 - (1/2)(2/3) = 2/3$$

$$(B) P(\text{exactly one of them solves the problem}) = P(\bar{A}B \text{ or } A\bar{B}) = (1/2)(2/3) + (1/2)(1/3) = 1/2$$

10. Solution:

Since f is constant for  $x < 2$ ,  $x > 10$ , f(x) is continuous for  $x < 2$ ,  $x > 10$ .

At  $x=2$ ,

$$\lim_{x \rightarrow 2^-} (f(x)) = \lim_{x \rightarrow 2^-} (5) = 5$$

$$\lim_{x \rightarrow 2^+} (f(x)) = \lim_{x \rightarrow 2^+} (ax + b) = 2a + b$$

$$\therefore 2a + b = 5$$

At  $x=10$ ,

$$\lim_{x \rightarrow 10^-} (f(x)) = \lim_{x \rightarrow 10^-} (ax + b) = 10a + b$$

$$\lim_{x \rightarrow 10^+} (f(x)) = \lim_{x \rightarrow 10^+} (21) = 21$$

$$\therefore 10a + b = 21$$

Solving the above two equations we get  $a=2$ ,  $b=1$ .

11. Solution:

$$F(x) = \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix}, F(y) = \begin{bmatrix} \cos y & -\sin y & 0 \\ \sin y & \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$$\begin{aligned}
 \therefore F(x)F(y) &= \begin{bmatrix} \cos x & -\sin x & 0 \\ \sin x & \cos x & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} \cos y & -\sin y & 0 \\ \sin y & \cos y & 0 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \cos x \cos y - \sin x \sin y & -\sin x \cos y - \cos x \sin y & 0 \\ \sin x \cos y + \cos x \sin y & \cos x \cos y - \sin x \sin y & 0 \\ 0 & 0 & 1 \end{bmatrix} \\
 &= \begin{bmatrix} \cos(x+y) & -\sin(x+y) & 0 \\ \sin(x+y) & \cos(x+y) & 0 \\ 0 & 0 & 1 \end{bmatrix}
 \end{aligned}$$

12. Solution:

$$\begin{aligned}
 \frac{d}{dx} \left( \cos^{-1} \sqrt{\frac{1+x}{2}} \right) &= \frac{-1}{\sqrt{1 - \left( \sqrt{\frac{1+x}{2}} \right)^2}} \cdot \frac{d}{dx} \sqrt{\frac{1+x}{2}} = \frac{-1}{\sqrt{1-x}} \left( \frac{1}{2} \right) \left( \frac{1+x}{2} \right)^{-1/2} \cdot \frac{1}{2} \\
 &= \frac{-1}{2\sqrt{(1-x)(1+x)}} = \frac{-1}{2\sqrt{1-x^2}}
 \end{aligned}$$

13. Solution:

$$\begin{aligned}
 f \circ g(x) &= f \left( \frac{x+3}{2} \right) = 2 \left( \frac{x+3}{2} \right) - 3 = x = I_R(x) \Rightarrow f \circ g = I_R \\
 g \circ f(x) &= g(2x-3) = \frac{2x-3+3}{2} = x = I_R(x) \Rightarrow g \circ f = I_R
 \end{aligned}$$

14. Solution:

$$y = x^3 - 11x + 5$$

$$\Rightarrow \frac{dy}{dx} = 3x^2 - 11$$

Also, from equation of tangent we get  $dy/dx=1$

$$\therefore 3x^2 - 11 = 1 \Rightarrow x = \pm 2.$$

Solving  $y=x-11$  for  $y$  we get the possible points are  $(2,-9), (-2,-13)$ .

But  $(-2,-13)$  does not lie on the curve, hence required point is  $(2,-9)$ .

15. Solution:

$$\frac{dT}{dt} = -c(T-S)$$

$$\int \frac{dT}{T-S} = \int -cdt$$

$$\therefore \log(T-S) = -ct + k$$

$$\Rightarrow e^{-ct+k} = T-S$$

Putting the condition  $T(0)=40$ , we get  $(40-S)e^{-ct} = T-S$ .

Measures to control global warming:

(1) Planting more trees

(2) Car pools to prevent emission of carbon dioxide which in turn causes global warming.

16. Solution:

$$\begin{aligned}
 \because \vec{a} \times \vec{b} &= \vec{c} \text{ and } \vec{b} \times \vec{c} = \vec{a} \Rightarrow \vec{a} \perp \vec{c}, \vec{b} \perp \vec{c}, \vec{a} \perp \vec{b}, \vec{a} \perp \vec{c} \Rightarrow \vec{a}, \vec{b}, \vec{c} \text{ are mutually perpendicular.} \\
 |\vec{a} \times \vec{b}| &= |\vec{c}|, |\vec{b} \times \vec{c}| = |\vec{a}| \Rightarrow |\vec{a}| |\vec{b}| \sin 90 = |\vec{c}| \text{ and } |\vec{b}| |\vec{c}| \sin 90 = |\vec{a}| (\because \vec{a}, \vec{b}, \vec{c} \text{ are mutually perpendicular}) \\
 \Rightarrow |\vec{a}| |\vec{b}| &= |\vec{c}| \text{ and } |\vec{b}| |\vec{c}| = |\vec{a}| \\
 \Rightarrow |\vec{b}| |\vec{c}| |\vec{b}| &= |\vec{c}| \Rightarrow |\vec{b}|^2 |\vec{c}| = |\vec{c}| \Rightarrow |\vec{b}| = 1 \Rightarrow |\vec{a}| = |\vec{c}|
 \end{aligned}$$

17. Solution:

$$\begin{aligned}
 \int e^x \left( \frac{1}{x} - \frac{1}{x^2} \right) dx &= \int e^x \left( \frac{1}{x} \right) dx - \int e^x \left( \frac{1}{x^2} \right) dx \\
 &= \frac{1}{x} \int e^x dx - \int \left( \frac{d}{dx} \left( \frac{1}{x} \right) \right) \int e^x dx - \int e^x \left( \frac{1}{x^2} \right) dx \\
 &= \frac{1}{x} e^x - \int \frac{-1}{x^2} e^x - \int e^x \left( \frac{1}{x^2} \right) dx \quad \text{(using integration by parts)} \\
 &= \frac{1}{x} e^x + \int \frac{1}{x^2} e^x - \int e^x \left( \frac{1}{x^2} \right) dx \\
 &= \frac{1}{x} e^x
 \end{aligned}$$

18. Solution:

Projection vector of  $\vec{a}$  along  $\vec{b}$  is given by  $\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b}$ .

$$\vec{a} = 2\vec{i} + 3\vec{j} - 3\vec{k}, \vec{b} = 5\vec{j} - \vec{k}$$

$$\left( \frac{\vec{a} \cdot \vec{b}}{|\vec{b}|^2} \right) \vec{b} = \left( \frac{(2\vec{i} + 3\vec{j} - 3\vec{k}) \cdot (5\vec{j} - \vec{k})}{(\sqrt{26})^2} \right) (5\vec{j} - \vec{k}) = \frac{9}{13} (5\vec{j} - \vec{k})$$

19. Solution:

$$x_1 = -3, y_1 = 1, z_1 = 5, x_2 = -1, y_2 = 2, z_2 = 5$$

$$a_1 = -3, b_1 = 1, c_1 = 5, a_2 = -1, b_2 = 2, c_2 = 5,$$

$$\text{The lines are coplanar iff } \begin{vmatrix} x_2 - x_1 & y_2 - y_1 & z_2 - z_1 \\ a_1 & b_1 & c_1 \\ a_2 & b_2 & c_2 \end{vmatrix} = 0$$

Substituting the values, we get the value of determinant as 0.

Hence, the lines are co-planar.

## Section C

20. Solution:

Let  $B_1$  denote the event that the item is produced by A.  $P(B_1) = 60/100 = .6$

Let  $B_2$  denote the event that the item is produced by B.  $P(B_2) = 40/100 = .4$

Let E denote the event that the item is defective.

$P(B_2/E) = ?$ ,  $P(E/B_1) = 0.02$ ,  $P(E/B_2) = 0.01$

By Baye's theorem,

$$\begin{aligned}
 P(B_2 / E) &= \frac{P(E / B_2)P(B_2)}{P(E / B_1)P(B_1) + P(E / B_2)P(B_2)} \\
 &= \frac{(1/100)(40/100)}{(2/100)(60/100) + (1/100)(40/100)} = \frac{1}{4}
 \end{aligned}$$

21. Solution:

$$\text{Let } y = \tan^{-1} \frac{2\sqrt{x}}{1-x}$$

$$\text{Let } x = \tan^2 \theta$$

$$\therefore y = \tan^{-1} \frac{2 \tan \theta}{1 - \tan^2 \theta} = \tan^{-1}(\tan 2\theta) = 2\theta = 2 \tan^{-1} \sqrt{x}$$

$$\frac{dy}{dx} = 2 \frac{1}{1 + (\sqrt{x})^2} \frac{1}{2\sqrt{x}} = \frac{1}{(1+x)\sqrt{x}}$$

$$\text{Let } z = \sin^{-1} \frac{2\sqrt{x}}{1+x}$$

$$\text{Let } x = \tan^2 \theta$$

$$\therefore z = \sin^{-1} \frac{2 \tan \theta}{1 + \tan^2 \theta} = \sin^{-1}(\sin 2\theta) = 2\theta = 2 \tan^{-1} \sqrt{x}$$

$$\frac{dz}{dx} = \frac{1}{(1+x)\sqrt{x}}$$

$$\frac{dy}{dz} = \frac{dy}{dx} \div \frac{dz}{dx} = \frac{1}{(1+x)\sqrt{x}} \div \frac{1}{(1+x)\sqrt{x}} = 1$$

22. Solution :

$$I = \int_0^{\pi} \frac{x dx}{4 \cos^2 x + 9 \sin^2 x} = \int_0^{\pi} \frac{(\pi - x) dx}{4 \cos^2 x + 9 \sin^2 x}$$

$$\therefore 2I = \pi \int_0^{\pi} \frac{dx}{4 \cos^2 x + 9 \sin^2 x} = 2\pi \int_0^{\frac{\pi}{2}} \frac{dx}{4 \cos^2 x + 9 \sin^2 x}$$

$$= 2\pi \left[ \int_0^{\frac{\pi}{4}} \frac{dx}{4\cos^2 x + 9\sin^2 x} + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{dx}{4\cos^2 x + 9\sin^2 x} \right]$$


$$= 2\pi \left[ \int_0^{\frac{\pi}{4}} \frac{\sec^2 x dx}{4 + 9\tan^2 x} + \int_{\frac{\pi}{4}}^{\frac{\pi}{2}} \frac{\csc^2 x dx}{4\cot^2 x + 9} \right]$$

Putting  $\tan x = t$  and  $\cot x = u$ , we get

$$2I = 2\pi \left[ \int_0^1 \frac{dt}{4 + 9t^2} - \int_1^0 \frac{du}{4u^2 + 9} \right] = 2\pi \left[ \frac{1}{9} \left( \frac{3}{2} \right) \tan^{-1} \frac{t}{2/3} \Big|_0^1 - \frac{1}{4} \left( \frac{2}{3} \right) \tan^{-1} \frac{u}{3/2} \Big|_1^0 \right]$$

23.  $= 2\pi \left[ \frac{1}{6} \tan^{-1} \left( \frac{3}{2} \right) + \frac{1}{6} \tan^{-1} \left( \frac{2}{3} \right) \right] = \frac{2\pi}{6} \left( \frac{\pi}{2} \right) = \frac{\pi^2}{6}$  Solution:

$$\therefore I = \frac{\pi^2}{12}$$



Let  $l, h, r, \alpha$  denote the slant height, height, radius and semi-vertical angle of the cone respectively.

$$\sin \alpha = \frac{r}{l} \Rightarrow r = l \sin \alpha, \cos \alpha = \frac{h}{l} \Rightarrow h = l \cos \alpha$$

$$\text{Volume} = V = \frac{\pi}{3} r^2 h = \frac{\pi}{3} l^3 \sin^2 \alpha \cos \alpha$$

$$\therefore \frac{dV}{d\alpha} = \frac{\pi}{3} l^3 [2 \sin \alpha \cos^2 \alpha + \sin^2 \alpha (-\sin \alpha)] = \frac{\pi}{3} l^3 [\sin \alpha (2 \cos^2 \alpha - \sin^2 \alpha)]$$

$$\frac{dV}{d\alpha} = 0 \Rightarrow \alpha = \tan^{-1}(\sqrt{2})$$

$$\frac{d^2V}{d\alpha^2} = \frac{\pi}{3} l^3 [\cos \alpha (2 \cos^2 \alpha - \sin^2 \alpha) + \sin \alpha (4 \cos \alpha (-\sin \alpha) - 2 \sin \alpha \cos \alpha)] = \frac{\pi}{3 \cos^3 \alpha} l^3 [2 - 7 \tan^2 \alpha]$$

$$\text{If } \tan \alpha = \sqrt{2} \Rightarrow \frac{d^2V}{d\alpha^2} < 0$$

Thus, for maximum volume  $\tan \alpha = \sqrt{2}$ .

24. Solution:

Let the three numbers be  $x, y, z$ . We can formulate the above as the mathematical problem:

$$x + y + z = 6$$

$$y + 3z = 11$$

$$x - 2y + z = 0$$

$$\text{Let } A = \begin{bmatrix} 1 & 1 & 1 \\ 0 & 1 & 3 \\ 1 & -2 & 1 \end{bmatrix}, b = \begin{bmatrix} 6 \\ 11 \\ 0 \end{bmatrix}$$

$$|A| = 9, A^{-1} = \frac{1}{9} \begin{bmatrix} 7 & -3 & 2 \\ 3 & 0 & -3 \\ -1 & 3 & 1 \end{bmatrix}, X = A^{-1}b = \frac{1}{9} \begin{bmatrix} 7 & -3 & 2 \\ 3 & 0 & -3 \\ -1 & 3 & 1 \end{bmatrix} \begin{bmatrix} 6 \\ 11 \\ 0 \end{bmatrix} = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

$$\therefore x = 1, y = 2, z = 3$$

25. Solution:

Suppose the factory produces  $x$  units of machine A and  $y$  units of machine B.

Then, Profit  $Z = 10,500x + 9000y$


The mathematical formulation of the problem is as follows:

$$\text{Max } Z = 10,500x + 9000y$$

$$\text{s.t. } 10x + 20y \leq 480, x + 2y \leq 48 \text{ (metal constraint)}$$

$$15x + 10y \leq 400, 3x + 2y \leq 80 \text{ (painting constraint)}$$

$$x \geq 0, y \geq 0$$



We graph the above inequalities. The feasible region is as shown in the figure. We observe the feasible region is bounded and the corner points are O, B, E and C. The co-ordinates of the corner points are  $(0,0)$ ,  $(0,24)$ ,  $(16,16)$ ,  $(80/3,0)$ .

Corner Point	$Z = 10,500x + 9000y$
$(0,0)$	0
$(0,24)$	2,16,000
$(16,16)$	<b>3,12,000</b>
$(80/3,0)$	2,80,000

Thus profit is maximized by producing 16 units each of machine A and B.

26. Solution:


The point of intersection of the two curves:

$$x^2 + y^2 = 16, x^2 = 6y$$

$$\Rightarrow y^2 + 6y - 16 = 0$$

$$\Rightarrow y = 2, -8$$

Rejecting  $y = -8$ , we get  $x = \pm 2\sqrt{3}$ .



Shaded area = Required area =  $\text{Ar}(\text{OAB}) + \text{Ar}(\text{OBC}) = 2 \text{Ar}(\text{OAB})$

$$\begin{aligned}
 \text{Area} &= 2 \int_0^{2\sqrt{3}} (y_1 - y_2) dx = 2 \int_0^{2\sqrt{3}} \left( \sqrt{16 - x^2} - \frac{x^2}{6} \right) dx \\
 &= 2 \left[ \frac{x}{2} \sqrt{16 - x^2} + \frac{16}{2} \sin^{-1} \left( \frac{x}{4} \right) - \frac{1}{6} \left( \frac{x^3}{3} \right) \right]_0^{2\sqrt{3}} \\
 &= 2 \left[ 2\sqrt{3} + \frac{8\pi}{3} - \frac{4\sqrt{3}}{3} \right] = \frac{4}{3} [4\pi + \sqrt{3}]
 \end{aligned}$$