

Table S2. LPR error relative to the LPG (m) across pyruvate concentrations

	P_{min} (μM)	Relative LPR error^a at P_{min} (%)	P_{max} (μM)	Relative LPR error^a at P_{max} (%)	LPR range
Unstratified	64 (41, 89)	66.8% (20.2%, 190.6%)	172 (139, 225)	23.3% (7.1%, 55.6%)	14.4 (9.2, 27.6)
Type P _b	61 (38, 88)	103.4% (39.1%, 254.7%)	171 (137, 225)	36.0% (16.8%, 74.5%)	16.0 (9.5, 34.7)
Type N _b	75 (51, 106)	-19.9% (-38.0%, -8.6%)	170 (140, 232)	-11.5% (-17.4%, -4.9%)	10.9 (7.5, 17.9)

Values are cohort medians (IQR) of patient-level metrics. Patient metrics: segment-weighted means (weights = segment size) of relative LPR error at segment P_{min} and P_{max}; patient medians of segment-level P_{min}, P_{max}, and LPR range. Cohort sizes: Unstratified = 506 patients, 1,339 segments; Type P_b = 476 patients, 967 segments; Type N_b = 276 patients, 372 segments.

^aRelative LPR error = $100 \times (\text{LPR} - \text{LPG}) / \text{LPG}$.

b, linear intercept; **IQR**, interquartile range; **LPG (m)**, lactate–pyruvate gradient; **LPR**, lactate/pyruvate ratio; **N_b**, negative intercept ($b < 0$); **P_b**, positive intercept ($b > 0$); **P_{max}**, segment-maximum pyruvate; **P_{min}**, segment-minimum pyruvate.