

Table S5. LPR error relative to the LPG (*m*) across pyruvate concentrations in the Uppsala cohort

	P_{min} (μM)	Relative LPR error^a at P_{min} (%)	P_{max} (μM)	Relative LPR error^a at P_{max} (%)	LPR range
Unstratified	101 (78, 136)	19.1% (-0.4%, 74.9%)	196 (153, 248)	10.1% (-0.5%, 34.8%)	10.4 (7.4, 14.3)
Type P _b	94 (72, 137)	54.9% (30.6%, 122.0%)	195 (155, 258)	28.8% (14.5%, 65.7%)	10.3 (7.3, 16.5)
Type N _b	108 (86, 136)	-22.3% (-35.2%, -8.4%)	202 (160, 249)	-13.6% (-21.2%, -4.9%)	9.5 (6.1, 11.7)

Values are cohort medians (IQR) of patient-level metrics. Patient metrics: segment-weighted means (weights = segment size) of relative LPR error at segment P_{min} and P_{max}; patient medians of segment-level P_{min}, P_{max}, and LPR range. Cohort sizes: Unstratified = 81 patients, 215 segments; Type P_b = 72 patients, 132 segments; Type N_b = 61 patients, 83 segments.

^aRelative LPR error = 100 × (LPR – LPG) / LPG.

b, linear intercept; **IQR**, interquartile range; **LPG (*m*)**, lactate–pyruvate gradient; **LPR**, lactate/pyruvate ratio; **N_b**, negative intercept (*b* < 0); **P_b**, positive intercept (*b* > 0); **P_{max}**, segment-maximum pyruvate; **P_{min}**, segment-minimum pyruvate.