
This copy is for your personal, non-commercial use only.

If you wish to distribute this article to others, you can order high-quality copies for your colleagues, clients, or customers by [clicking here](#).

Permission to republish or repurpose articles or portions of articles can be obtained by following the guidelines [here](#).

The following resources related to this article are available online at www.sciencemag.org (this information is current as of March 21, 2011):

Updated information and services, including high-resolution figures, can be found in the online version of this article at:

<http://www.sciencemag.org/content/325/5947/1509.full.html>

This article appears in the following **subject collections**:

Geochemistry, Geophysics

http://www.sciencemag.org/cgi/collection/geochem_phys

models that are used to describe high-temperature superconductors.

References and Notes

1. G. -B. Jo *et al.*, *Science* **325**, 1521 (2009).
2. F. Bloch, *Z. Phys.* **57**, 545 (1929).
3. For a detailed discussion of direct and itinerant exchange, see the articles in (4, 5).
4. C. Herring, in *Magnetism*, Vol. IIB, G. T. Radu, H. Suhl, Eds. (Academic Press, New York, 1966), pp. 1–181.
5. C. Herring, in *Magnetism*, Vol. IV, G. T. Radu, H. Suhl, Eds. (Academic Press, New York, 1966).
6. F. H. Zong, C. Lin, D. M. Ceperley, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **66**, 036703 (2002).
7. E. C. Stoner, *Philos. Mag.* **15**, 1018 (1933).
8. As a student in Cambridge in 1924, Stoner was in fact very close to stating the exclusion principle before Pauli; see (9).
9. I. Duck, E. C. G. Sudarshan, *Pauli and the Spin-Statistics Theorem* (World Scientific, Singapore, 1997).
10. W. Ketterle, M. W. Zwiernlein, in *Proceedings of the International School of Physics "Enrico Fermi," Course CLXIV, Varenna, 20–30 June 2006*, M. Inguscio, W. Ketterle, C. Salomon, Eds. (IOS Press, Amsterdam, 2008).
11. I. Bloch, J. Dalibard, W. Zwerger, *Rev. Mod. Phys.* **80**, 885 (2008).
12. D. S. Petrov, *Phys. Rev. A* **67**, 010703(R) (2003).
13. S. Trotzky *et al.*, *Science* **319**, 295 (2008); published online 20 December 2007 (10.1126/science.1150841).
14. H. Von Löhneysen, A. Rosch, M. Vojta, P. Wölfle, *Rev. Mod. Phys.* **79**, 1015 (2007).

10.1126/science.1179767

GEOPHYSICS

How River Beds Move

Philippe Frey¹ and Michael Church²

Insights into sediment transport at river beds can come from experiments in granular physics.


The transport of sediment through river channels has major consequences for public safety, management of water resources, and environmental sustainability. Most studies of sediment transport in rivers have focused on mass flux and its relation to water flow. Yet, after more than a century of work, there is no satisfactory theory for bedload, the component of the sediment load transported in contact with the stream bed. Bedload transport formulae often overpredict the actual rate by orders of magnitude (1). It is therefore difficult to predict, for example, the impact of disturbances such as extreme floods on the channel. Better insight may come from viewing bedload as a granular phenomenon.

Bedload transport can be divided into two stages (2): partial mobility of local bed surface material when part of the bed remains static but exposed grains may eventually move, and full mobility, when all grains move to a depth of several grain diameters. Grain-grain interactions over short time and length scales bear importantly on the predictability of both stages.

No single constitutive law reproduces the diversity of behaviors of cohesionless granular materials (3). Granular flows are often classified into three states: a gaseous state, in which flow is very rapid and dilute, and the particles interact by collision; an intermediate state, in which the material is dense but still flows like a liquid, the particles interact-

ing both by collision and friction; and a dense, quasistatic state, in which the deformations are very slow and the particles interact by frictional contacts. All three states might be found in free surface flows and in bedload.

Probably the most important phenomenon relevant to bedload is size segregation by shearing in free surface flows. Two distinct size segregation phenomena occur. When the coarsest fractions of the bed do not move and the smallest fractions are sufficiently fine, spontaneous percolation occurs. But when the bed is moving, kinematic sieving of the finer particles take place even if the size ratio is close to unity (4, 5). The usual result is a downward flux of smaller particles and an upward flux of larger particles, resulting in the segregation observed in river deposits (see the figure, panel A) and in a substantial reduction in transport.



Segregation through motion. (A) This vertical profile in a gravel river bar (in Vedder River, British Columbia, Canada) shows size sorting with an armored surface and finer material below. (B) In a quasi-two-dimensional experiment (13), kinematic sieving leads to the formation of layers of smaller transparent beads under larger moving black beads. This panel was redrawn from a video snapshot.

¹Cemagref, Unité de recherche Erosion Torrentielle, Neige et Avalanches, BP 76, 38402 Saint-Martin-d'Hères, France. E-mail: philippe.frey@cemagref.fr ²Department of Geography, University of British Columbia, Vancouver, BC, V6T 1Z2, Canada.

into a bedload flow initially formed only of larger moving beads (13). After a while, a quasi-continuous layer of small particles developed beneath larger moving beads and above quasi-immobile larger beads (see the figure, panel B).

In the regime of partial mobility, processes are restricted to the surface of the bed, and all particles experience long periods of rest. This condition is characteristic of gravel transport. The propensity for grains of similar size to block each other leads to accumulations of similar-sized grains in restricted areas of the channel bed. Two phenomena command attention. Mobile materials collect in patches of similar size in the streambed, a phenomenon that mediates the overall sediment flux (14), while the largest stones in streambeds—usually only marginally mobile—congregate into clusters, chains, and cell-like arrangements that dramatically increase the overall stability of the bed (15).

The second case is particularly interesting from the granular perspective, because the stone structures represent a natural case of force chains that have been studied in the laboratory for more than a decade (16). In the extreme case of steep mountain channels containing relatively large stones, stone lines become channel-spanning force chains,

forming a distinctive step-and-pool morphology that maintains a stable channel in situations when any unconstrained stone would be swept away.

Heuristic models have been constructed for the development of surface structures, but the mechanisms that promote patch development and bed surface structures require additional experimental study before physically sound models may be developed. Stone lines and cells on the surface are relatively long-lived because, during most flows, their ultimate strength is not tested. This allows time for additional mechanisms to strengthen them further, beyond the state achieved by force chains in continuously deforming media. Hence, failure mechanisms are of particular interest. When extreme flows do break the stability of steep channels, life-threatening debris flows result.

Granular physics provides a good basis for improving our understanding of bedload transport at relatively high rates. However, surface phenomena that would simulate partial bedload transport remain essentially uninvestigated in granular physics. While imparting insight into the bedload problem, experiments on these phenomena would also open a new perspective in granular physics.

References and Notes

1. J. C. Bathurst, *J. Hydraul. Eng.* **133**, 1192 (2007).
2. P. R. Wilcock, B. W. Mcardell, *Water Resour. Res.* **33**, 235 (1997).
3. Y. Forterre, O. Pouliquen, *Annu. Rev. Fluid Mech.* **40**, 1 (2008).
4. S. B. Savage, C. K. K. Lun, *J. Fluid Mech.* **189**, 311 (1988).
5. N. Thomas, *Phys. Rev. E Stat. Phys. Plasmas Fluids Relat. Interdiscip. Topics* **62**, 961 (2000).
6. N. Jain, J. M. Ottino, R. M. Lueptow, *J. Fluid Mech.* **508**, 23 (2004).
7. I. Goldhirsch, *Annu. Rev. Fluid Mech.* **35**, 267 (2003).
8. K. M. Hill, J. Zhang, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **77**, 061303 (2008).
9. B. M. Sumer, A. Kozakiewicz, J. Fredsoe, R. Deigaard, *J. Hydraul. Eng.* **122**, 549 (1996).
10. A. Armanini, H. Capart, L. Fraccarollo, M. Larcher, *J. Fluid Mech.* **532**, 269 (2005).
11. D. Berzi, J. T. Jenkins, *Phys. Rev. E Stat. Nonlin. Soft Matter Phys.* **78**, 011304 (2008).
12. J. K. Wooster *et al.*, *Water Resour. Res.* **44**, W03424 (2008).
13. T. Böhm *et al.*, *Exp. Fluids* **41**, 1 (2006).
14. C. Paola, R. Seal, *Water Resour. Res.* **31**, 1395 (1995).
15. M. Church, M. A. Hassan, J. F. Wolcott, *Water Resour. Res.* **34**, 3169 (1998).
16. M. E. Cates, J. P. Wittmer, J. P. Bouchaud, P. Claudin, *Phys. Rev. Lett.* **81**, 1841 (1998).
17. P.F. acknowledges the support of Cemagref and the University of British Columbia, Department of Geography, where this Perspective was written while on sabbatical leave. This work was partly funded by the French Institut National des Sciences de l'Univers and Agence Nationale de la Recherche grant ANR-05-ECCO-015. M.C. is supported by the Natural Sciences and Engineering Research Council of Canada.

10.1126/science.1178516

PHYSICS

The Super of Superradiance

Marlan O. Scully^{1,2} and Anatoly A. Svidzinsky¹

In 1954, Robert Dicke introduced the concept of superradiance in describing the cooperative, spontaneous emission of photons from a collection of atoms. The concept of superradiance can be understood by picturing each atom as a tiny antenna emitting electromagnetic waves. Thermally excited atoms emit light randomly, and the emitted intensity is a function of the number of atoms, N . However, when the atomic “antennas” are coherently radiating in phase with each other, the net electromagnetic field is proportional to N , and therefore, the emitted intensity goes as N^2 . As a result, the atoms radiate their energy N times faster than for incoherent emission. It is this anomalous radiance that Dicke dubbed “superradiance” (1–3).

An even more interesting kind of radiation speedup can occur when a single photon is stored uniformly in a cloud of N atoms (see the figure, panel A). Suppose you have one atom that decays with a rate γ . Then suppose there are N such atoms close together in an atom cloud with only one of the atoms excited (but we don't know which one). Because there is only one atom excited, you might expect the decay rate to be γ . But if the atoms are symmetrically organized within the cloud, the decay rate is actually $N\gamma$ (1). This enhanced single-photon emission rate is “the greatest radiation anomaly” inherent in superradiance. Single-photon superradiance has become a subject of current interest (4–12), and promises to yield new tools for storing quantum information and deeper insight into the physics of virtual processes.

Dicke's point is that the N atoms act like one big atom and decay collectively. This is intuitive when the atoms are close together

Cooperative single-photon emission from an atom ensemble will provide insights into quantum electrodynamics and applications in quantum communication.

compared to the wavelength of radiation λ . When the same symmetric state is formed but the atomic cloud size is larger than λ , there is no longer constructive cooperation in radiation emission. The atoms will trap the light, decreasing the emission rate.

Nevertheless, it is possible to produce a state such that the large cloud also emits radiation with an enhanced rate proportional to $N\gamma$ (4, 5). This is important because in quantum optics the sample is usually large compared to λ .

However, things are a bit trickier here. More subtle and interesting physics come into play, extending from quantum information and a new kind of cavity quantum electrodynamics (QED) (13), to new insights into quantum field theory (9–12).

The essential new physics is the transition from the coherent antenna array to the single-photon state in which cooperative emission is due to N entangled atoms (not N coherent

¹Texas A&M University, College Station, TX 77843, USA.

²Princeton University, Princeton, NJ 08544, USA. E-mail: scully@tamu.edu