
homework sheet 03

Marco Seravalli
03626387
marco.seravalli@tum.de

Nikola Tchipev
03625168
n.tchipev@tum.de

1 Basic Probability

Problem 1

Probability to be a terrorist $P(x = T) = \frac{1}{100}$

Probability to be a citizen $P(x = C) = \frac{99}{100}$

Probability to recognize a terrorist as such $P(T|x = T) = \frac{95}{100}$

Probability to recognize a citizen as terrorist $P(T|x = C) = 1 - P(C|x = C) = \frac{5}{100}$

Probability that my neighbor is actually a terrorist

$$P(x = T|T) = \frac{P(x = T)P(T|x = T)}{P(x = T)P(T|x = T) + P(x = C)P(T|x = C)}$$

$$P(x = T|T) = \frac{\frac{1}{100} \frac{95}{100}}{\frac{1}{100} \frac{95}{100} + \frac{99}{100} \frac{5}{100}} = \frac{19}{118} \approx 0.16$$

Problem 2

Notation $P(2d = RR)$ probability to draw two red balls, $P(RR)$ probability that the box contains two red balls.

Probability of having 2 red balls in the box $P(RR) = \frac{1}{4}$

Probability of having one red and one white ball in the box $P(WR) = \frac{1}{2}$

Probability of having 2 white balls in the box $P(WW) = \frac{1}{4}$

Probability that all the balls in the box are red given that 3 red balls were extracted in 3 different draws:

$$\begin{aligned} P(RR|3d = RRR) &= \\ &= \frac{P(1d = R|RR)^3 P(RR)}{P(1d = R|RR)^3 P(RR) + P(1d = R|WR)^3 P(WR) + P(1d = R|WW)^3 P(WW)} \end{aligned}$$

We know that

$$P(1d = R|WW) = 0$$

$$P(1d = R|WR) = \frac{1}{2}$$

$$P(1d = R|RR) = 1$$

Hence

$$P(RR|3d = RRR) = \frac{P(RR)}{P(RR) + P(1d = R|WR)^3 P(WR)} = \frac{\frac{1}{4}}{\frac{1}{4} + \frac{1}{8} \frac{1}{2}} = \frac{4}{5}$$

Problem 3

Expected value:

$$\begin{aligned} E[X] &= \int_{-\infty}^{+\infty} xf(x)dx = \int_{-\infty}^0 xf(x)dx + \int_0^1 xf(x)dx + \int_1^{+\infty} xf(x)dx \\ E[X] &= 0 + \int_0^1 xdx + 0 = \left| \frac{x^2}{2} \right|_0^1 = \frac{1}{2} \end{aligned}$$

Variance:

$$\begin{aligned} V[X] &= E[(X - E[X])^2] = \int_0^1 \left(x - \frac{1}{2}\right)^2 dx \\ V[X] &= \int_0^1 x^2 - x + \frac{1}{4} dx = \left| \frac{x^3}{3} - \frac{x^2}{2} + \frac{x}{4} \right|_0^1 = \frac{1}{3} - \frac{1}{2} + \frac{1}{4} = \frac{1}{12} \end{aligned}$$

Problem 4**Part A**

Prove

$$E[X] = E_Y[E_{X|Y}[X]]$$

We know that

$$\begin{aligned} E[X] &= \sum_x xp_X(x) \\ E[X] &= \sum_x x \sum_y p_{XY}(x, y) \\ E[X] &= \sum_x x \sum_y \frac{p_{XY}(x, y)}{p_Y(y)} p_Y(y) \\ E[X] &= \sum_y \sum_x x \frac{p_{XY}(x, y)}{p_Y(y)} p_Y(y) \end{aligned}$$

Use Bayes rule

$$\begin{aligned} E[X] &= \sum_y \sum_x xp_{X|Y}(x|y)p_Y(y) \\ E[X] &= \sum_y E_{X|Y}[X] p_Y(y) \\ E[X] &= E_y[E_{X|Y}[X]] \end{aligned}$$

Part B

Prove

$$Var_X[X] = E_Y[Var_{X|Y}[X]] + Var_Y[E_{X|Y}[X]]$$

The R.H.S. can be also rewritten as

$$\begin{aligned} &E_Y[Var_{X|Y}[X]] + Var_Y[E_{X|Y}[X]] = \\ &= E_Y[E_{X|Y}[X^2] - E_{X|Y}[X]^2] + E_Y[E_{X|Y}[X]^2] - E_Y[E_{X|Y}[X]]^2 \end{aligned}$$

By exploiting the linearity of E we get

$$\begin{aligned} &E_Y[Var_{X|Y}[X]] + Var_Y[E_{X|Y}[X]] = \\ &= E_Y[E_{X|Y}[X^2]] - E_Y[E_{X|Y}[X]^2] + E_Y[E_{X|Y}[X]^2] - E_Y[E_{X|Y}[X]]^2 \\ &= E_Y[E_{X|Y}[X^2]] - E_Y[E_{X|Y}[X]]^2 \end{aligned}$$

If the result of the Part A is reused we obtain

$$E_Y[Var_{X|Y}[X]] + Var_Y[E_{X|Y}[X]] = E_X[X^2] - E_X[X]^2 = Var_X[X]$$

2 Probability Inequalities

2.1 Markov Inequality

Let $I_{X>c}$ be the indicator random variable of the event $X > c$:

$$I_{X>c} = \begin{cases} 1 & \text{if } X > c, \\ 0 & \text{otherwise.} \end{cases}$$

We then have

$$cI_{X>c} = \begin{cases} c & \text{if } X > c, \\ 0 & \text{otherwise,} \end{cases}$$

and hence

$$cI_{X>c} \leq X,$$

since X is non-negative. Now, by the monotonicity of the expectation operator we have

$$E[cI_{X>c}] \leq E[X].$$

Taking linearity of E , into account, we get

$$cE[I_{X>c}] \leq E[X].$$

Note that we can compute $E[I_{X>c}]$:

$$\begin{aligned} E[I_{X>c}] &= \sum_x p(x)I_{X>c} \\ &= \sum_{x \leq c} p(x)I_{X>c} + \sum_{x > c} p(x)I_{X>c} \\ &= \sum_{x \leq c} p(x)0 + \sum_{x > c} p(x)1 \\ &= \sum_{x > c} p(x) \\ &= P(X > c). \end{aligned}$$

Thus, we arrive at

$$cP(X > c) = cE[I_{X>c}] \leq E[X],$$

and hence

$$P(X > c) \leq \frac{E[X]}{c}.$$

For the second part, let X be the random variable, describing the number of “heads” out of n coin tosses. Since we are throwing a fair coin, we clearly have

$$E[X] = \frac{n}{2}.$$

Now we have $c = \frac{3}{4}n$ and by the Markov inequality we obtain

$$P\left(X > \frac{3}{4}n\right) \leq \frac{\frac{n}{2}}{\frac{3}{4}n} = \frac{2}{3}$$

and

$$P\left(X > \frac{3}{4}n\right) \leq \frac{2}{3}.$$

2.2 Chebyshev Inequality

Similarly, let $I_{(X-E[X])^2 > c^2}$ be the indicator random variable of the event $(X - E[X])^2 > c^2$:

$$I_{(X-E[X])^2 > c^2} = \begin{cases} 1 & \text{if } (X - E[X])^2 > c^2, \\ 0 & \text{otherwise.} \end{cases}$$

We then have

$$c^2 I_{(X-E[X])^2 > c^2} = \begin{cases} c^2 & \text{if } (X - E[X])^2 > c^2, \\ 0 & \text{otherwise,} \end{cases}$$

and hence

$$c^2 I_{(X-E[X])^2 > c^2} \leq (X - E[X])^2.$$

Now, by the monotonicity of the expectation operator we have

$$E[c^2 I_{(X-E[X])^2 > c^2}] \leq E[(X - E[X])^2] = \text{Var}[E].$$

Taking linearity of E , into account, we get

$$c^2 E[I_{(X-E[X])^2 > c^2}] \leq \text{Var}[E].$$

Note that we can compute $E[I_{(X-E[X])^2 > c^2}]$:

$$\begin{aligned} E[I_{(X-E[X])^2 > c^2}] &= \sum_x p(x) I_{(X-E[X])^2 > c^2} \\ &= P((X - E[X])^2 > c^2) \\ &= P(|X - E[X]| > c) \end{aligned}$$

Thus, we arrive at

$$P(|X - E[X]| > c) \leq \frac{\text{Var}[X]}{c^2}.$$

For the second part, let again X be the random variable, describing the number of “heads” out of n coin tosses. Now, due to the symmetry of the problem and the fact that $E[X] = \frac{n}{2}$, we make the following observation:

$$P\left(X > \frac{3}{4}n\right) = P\left(X < \frac{1}{4}n\right).$$

Thus,

$$P\left(X > \frac{3}{4}n\right) = \frac{2P\left(X > \frac{3}{4}n\right)}{2} \tag{1}$$

$$= \frac{P\left(X > \frac{3}{4}n\right) + P\left(X < \frac{1}{4}n\right)}{2} \tag{2}$$

$$= \frac{P\left(\{X > \frac{3}{4}n\} \cup \{X < \frac{1}{4}n\}\right)}{2} \tag{3}$$

$$= \frac{P\left(|X - E[X]| > \frac{1}{4}n\right)}{2}. \tag{4}$$

Now, after applying Chebyshev’s inequality, we obtain:

$$P\left(X > \frac{3}{4}n\right) \leq \frac{\text{Var}[X]}{2\left(\frac{1}{4}n\right)^2} = \frac{8\text{Var}[X]}{n^2}.$$

Now we just need the variance:

$$\begin{aligned} \text{Var}[X] &= E[(X - E[X])^2] \\ &= \sum_x p(x) \left(x - \frac{n}{2}\right)^2 \\ &= \sum_{k=0}^n \frac{\binom{n}{k}}{2^n} \left(k - \frac{n}{2}\right)^2. \end{aligned}$$

2.3 Jensen's Inequality

The statement is obviously true for $n = 1$:

$$f(1 \cdot x_1) \leq 1 \cdot f(x_1)$$

and for $n = 2$ since f is convex:

$$f(\lambda x_1 + (1 - \lambda)x_2) \leq \lambda f(x_1) + (1 - \lambda)f(x_2),$$

which gives us the base step of the induction.

Now suppose it is true for $n = k$ and consider $n = k + 1$:

$$f\left(\sum_{i=1}^{k+1} \lambda_i x_i\right) = f\left(\sum_{i=1}^k \lambda_i x_i + \lambda_{k+1} x_{k+1}\right) \quad (5)$$

$$= f\left((1 - \lambda_{k+1})\left(\frac{\sum_{i=1}^k \lambda_i x_i}{1 - \lambda_{k+1}}\right) + \lambda_{k+1} x_{k+1}\right) \quad (6)$$

$$\leq (1 - \lambda_{k+1})f\left(\sum_{i=1}^k \frac{\lambda_i x_i}{1 - \lambda_{k+1}}\right) + \lambda_{k+1}f(x_{k+1}) \quad (7)$$

$$\leq (1 - \lambda_{k+1})f\left(\sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} x_i\right) + \lambda_{k+1}f(x_{k+1}) \quad (8)$$

$$\leq (1 - \lambda_{k+1})\left(\sum_{i=1}^k \frac{\lambda_i}{1 - \lambda_{k+1}} f(x_i)\right) + \lambda_{k+1}f(x_{k+1}) \quad (9)$$

$$= \sum_{i=1}^k \lambda_i f(x_i) + \lambda_{k+1}f(x_{k+1}) \quad (10)$$

$$= \sum_{i=1}^{k+1} \lambda_i f(x_i). \quad (11)$$

In going from (6) to (7) we used the definition of convexity of f . In going from (7) to (8) we use the fact that

$$\sum_{i=1}^k \lambda_i = 1 - \lambda_{k+1}$$

and apply the inductive hypothesis for $n = k$.