Part 2: Descriptive Statistics

307312 Applied Statistics — Masters Level

307312 Applied Statistics — Masters Level

University of Petra

What is Statistics?

Statistics is a mathematical science that includes methods for collecting, organizing, analyzing and visualizing data to extract useful information that can be used to make decisions.

Statistics help us solve problems, answer questions and make decisions based on data (Data Driven Decisions).

Statistics Helps to in Reduce Uncertainty.

Statistics is the science of making decisions under uncertainty (Savage, The Foundations of Statistics, 1954).



Statistics Branches

Statistics is generally divided into two main branches:

1. Descriptive statistics.

Descriptive statistics refers to the summarizing and explaining data using numbers, tables and charts.

2- Inferential statistics.

Inferential statistics refers to drawing conclusions about a population based on a sample data.

A population refers to all members of a specified group (not necessarily people), whereas a sample is a subset of that population.

Population: the collection of all items we are interested in (people, things, plants, stars, trees…etc.), for the purpose of making our decision.

Example of Useful Descriptive Statistics

1. Business Sales:

- A store owner calculates the average daily sales to get a sense about his income.
- The owner also computes the standard deviation of daily sales to understand the variability or consistency in income.

2. Classroom Grades:

- A teacher calculates the average grade of a test to understand the overall performance of the class.
- The teacher also determines the highest and lowest scores to understand the spread of the grades.

3. Survey Results:

- After a customer satisfaction survey, a company finds that 70% of respondents are "satisfied" or "very satisfied."
- The survey also indicates that the mode response to the question, "What's the most important feature?" is "product durability."

Examples of Useful Inferential Statistics

1. Medicine:

- A research group tests a new drug on 100 patients and finds that 60 of them show significant improvement.
- Using inferential statistics, they can make conclusions about the drug's effectiveness on the entire population.

2. Agriculture:

- A farmer tests a new fertilizer on a sample plot of land to see if it increases crop yield.
- Inferential statistics can help predict the impact of using the fertilizer on all their plots.

3. Marketing:

- A company launches a new ad campaign in a small geographic area.
- They use inferential statistics to determine if rolling out the campaign nationwide would likely produce similar results.

Statistical Process Life Cycle

1. Define Problem – Question
2. Collect Data
3. Prepare Data
4. Analyze Data
5. Find Insights
6. Make Decisions

The process returns to the question as needed.

Types of Data & Levels of Measurements

Categorical Data vs Numerical Data


1. **Qualitative** or **categorical** (non-numerical) data:

Consists of descriptive information, such as colors, tastes, textures, or any other characteristics that can be counted.

2. **Quantitative** or **numerical** data:

Includes data that can be measured and can be represented numerically, such as height, weight, or temperature.

Types of data



Categorical data represents groups or categories.


Examples:

1. Car brands: Audi, BMW and Mercedes.
2. Answers to yes/no questions: yes and no

Numerical data represents numbers. It is divided into two groups: discrete and continuous. Discrete data can be usually counted in a finite matter, while continuous is infinite and impossible to count.

Examples:
Discrete: # children you want to have, SAT score
Continuous: weight, height

Levels of measurement



There are two qualitative levels: nominal and ordinal. The nominal level represents categories that cannot be put in any order, while ordinal represents categories that can be ordered.

Examples:
Nominal: four seasons (winter, spring, summer, autumn)
Ordinal: rating your meal (disgusting, unappetizing, neutral, tasty, and delicious)

There are two quantitative levels: interval and ratio. They both represent "numbers", however, ratios have a true zero, while intervals don't.

Examples:
Interval: degrees Celsius and Fahrenheit
Ratio: degrees Kelvin, length

Why Types of Data and Levels of Measurements Matter?

- Data Visualization: e.g. Categorical data might be best represented in a bar chart, while continuous data might be better suited for a histogram or scatter plot.
- Data Transformation: e.g. Ordinal data might be converted into interval data under certain conditions, or continuous data might be categorized into ordinal data.
- Data Quality: e.g. If you expect a variable to be discrete and find continuous values, this could indicate a data quality issue.
- Interpretation of Results: e.g. If you have ordinal data, you can make statements about the order of values but not the difference between values.
- Appropriate Analysis: e.g. Nominal data can be analyzed using a Chi-square test, while interval data can be analyzed using a t-test or ANOVA. Using the wrong test can lead to incorrect conclusions.

Types of attributes and scales of measurement

Scale of measurement	Type of attribute	Examples	Notes	Operations
Nominal scale	Qualitative / Categorical	Gender, blood group, rhesus factor	Lowest level	A = B A ≠ B
Ordinal scale	Qualitative / Categorical	Grades, medical scores	Order is defined	A = B A ≠ B A < B A > B
Interval scale	Quantitative / Metric	Temperature in °C, intelligence quotient	Arbitrary zero, distance is defined	A = B A ≠ B A < B A > B d = A − B
Ratio scale	Quantitative / Metric	Temperature in Kelvin, body height	Highest level, natural zero, ratio is defined	A = B A ≠ B A < B A > B d = A − B r = A:B

Figure 2.2: Types of attributes and scales of measurement.

Curtesy of: Introduction to statistical analysis with R, Matheus Khol

Data collection and survey quality

Before summarizing data, ask how they were obtained.

- **Source:** Are suitable data already available, or must we collect them?
- **Selection:** Who or what should be included?
- **Measurement:** How will we obtain comparable, trustworthy answers?

By the end of this section, you should be able to select a collection approach, justify a survey mode and sampling method, and identify likely survey errors.

Define what the study will measure

- **Target population:** The people or items to which the conclusion should apply; specify place and time.
- **Observational unit:** The person, transaction, branch or other item described by one observation.
- **Operational definition:** The exact rule used to measure a variable.

Illustration: “Service speed” could mean minutes from joining a queue to reaching a teller, or minutes from reaching the teller to completing a transaction. These are different variables.

State the rule and unit before collecting values. A clear variable name alone is insufficient.

Primary and secondary data

Primary data are collected for the current study. A new customer survey can ask exactly what management needs, but requires time, access and resources.

Secondary data already exist for another purpose. Sales records, administrative records and published statistics may be faster to obtain, but their definitions, dates and coverage may not fit the question.

Check who collected the data, why, when, how and for whom. Internal business records can be secondary data for a new analysis; “secondary” does not mean “external.”

Which collection approach fits the question?

Approach	Useful when we need to know…	Main limitation
Survey	What people report, prefer or believe	Reports may differ from actual behavior
Direct observation	What people do or how a process operates	Motives may remain unknown; privacy matters
Business or administrative records	What was recorded during operations	Missing events and changing definitions can mislead
Designed experiment	Whether an assigned intervention changes an outcome	Requires feasible, ethical treatment assignment and control

Random assignment in a well-designed experiment supports causal comparisons. An association in routine observational data alone does not establish causation.

Census, sample and sampling frame

A **census** attempts to collect data from every population unit. A **sample** includes a selected subset. Sampling can reduce cost and time; a census may be feasible for a small workforce.

The **sampling frame** identifies the units available for selection, such as a current employee list.

Distinguish **eligible population → frame → selected sample → respondents**. They need not contain the same people.

Inviting everyone is an attempted census; receiving replies from only some people does not eliminate nonresponse concerns.

Reference: ABS: Census and sample.

Probability sampling: who is selected?

Each population unit has a known, positive selection probability; probabilities need not be equal.

Method	How it works	Useful setting and caution
Simple random	Randomly select units from a complete frame	An up-to-date customer list; small groups may be missed by chance
Systematic	Choose a random start, then every *k*th unit	An ordered transaction list; avoid a repeating pattern aligned with *k*
Stratified	Divide into groups; randomly sample within every group	Comparing departments or regions; unequal sampling fractions need weights for overall estimates
Cluster	Randomly select groups, then study all or a sample of units within them	Dispersed branches or locations; similar units within clusters can reduce precision

Strata contribute sampled units from **every** group; cluster sampling selects **some** groups. Neither design guarantees error-free results.

Nonprobability samples: useful, with limits

- **Convenience:** Ask people who are easy to reach. Useful for testing a questionnaire quickly.
- **Voluntary response:** Invite anyone interested to reply. Useful for gathering suggestions; strongly motivated people may dominate.
- **Purposive:** Select people for relevant expertise. Useful for exploring a specialist problem.

These approaches can support exploration, but selection probabilities are unknown. A large response count alone does not justify population-wide percentages or a conventional probability-sampling margin of error.

An online questionnaire can use a probability sample or an opt-in sample. The communication channel does not determine the sampling design.

Choosing a sampling method: a bank example

Hypothetical planning example; no observed data. A bank wants to describe service satisfaction among all its current adult retail customers.

Method	What the bank would do	Why choose it?
Simple random	Randomly draw customers from the complete customer list	Suitable for an overall estimate when precise comparisons between customer groups are not the main objective. Small groups can receive few selections by chance.
Stratified random	Divide customers into age groups—18–29, 30–59 and 60+—then randomly select customers within **every** group	Suitable when management needs to compare age groups. Their service needs or digital experiences may differ, and a small group could otherwise have too few responses for a useful comparison.
Systematic	Choose a random starting position among the first 20 names, then select every 20th customer	Convenient for working through a long list. Check that the ordering has no repeating pattern aligned with the selection interval.
Cluster	Randomly select some branches, then randomly sample customers assigned to those branches	Suitable when interviews require branch visits: fewer locations reduce travel. Customers within a branch may be similar, so precision can be lower than with a similarly sized simple random sample.
Nonprobability	Ask available branch visitors to test a draft questionnaire, or deliberately interview customers with relevant experience	Suitable for piloting questions or exploring a problem. These people can provide useful feedback, but their satisfaction percentage should not be treated as representative of all customers.

Customer segments, strata and clusters

Customer segments are groups defined by a characteristic relevant to the business question, such as age, region or account type. For stratified sampling, use groups that do not overlap and together cover the target population.

In the bank example, the age groups are **strata**. Sampling within every age group ensures each enters the selected sample; this helps management study each group's experience. It does not mean surveying every customer, taking equal numbers from every group, or guaranteeing that everyone selected responds.

If a small group is deliberately sampled more heavily, use appropriate **weights** when combining results: its sample responses should not count as though that group makes up an equally large share of all customers.

The branches in the cluster example are **clusters**. Assign each customer to one designated branch for selection, so customers using several branches are not counted repeatedly. Visit only the randomly selected branches; the purpose is to reduce collection cost, rather than obtain a separate estimate for every branch.

Remember: stratified sampling selects customers from **every group**; cluster sampling selects **some groups**, then studies customers within them.

Survey modes: online and telephone

Mode	Why and where we might use it	What to watch
Online / email invitation	Fast repeated feedback from customers with reliable digital access; automated routing and visual response options	Digital exclusion, ignored invitations and self-selection if a link is openly posted
Telephone interview	Short interviews when telephone contact details are available; an interviewer can clarify instructions	Unanswered calls, interviewer effects and difficulty presenting long visual lists

For either mode, specify how invitees are selected. An accessible survey is of little use if the contact list omits the population of interest.

Reference: AAPOR: Survey research practices.

Survey modes: face-to-face and paper

Mode	Why and where we might use it	What to watch
Face-to-face interview	Complex questions, physical demonstrations or populations difficult to reach remotely; interviewers can assist	Travel and staffing cost, interviewer influence and privacy during the interview
Paper questionnaire	Respondents with limited internet access, or settings where paper is practical; postal return permits private completion	Slow returns, manual entry errors and reading or language barriers

An interviewer may improve comprehension but also increase socially desirable answers. Private self-completion can help with sensitive questions when respondents can use it comfortably.

References: AAPOR; Pew: Mode effects.

Choosing a mode and combining modes

Choose using **population access, question complexity, sensitivity, cost and available time**.

Illustrative decision: A bank studying all retail customers could invite a selected sample online, then offer telephone or paper completion to people who cannot respond online.

Mixed modes can improve access, but require consistent meaning and response options, controlled follow-up, and checks for duplicate submissions.

Changing mode can change answers even if opinions have not changed. Record the mode and examine comparability when tracking trends.

References: AAPOR; Pew: Mode effects.

Questionnaire design and a small pilot

Problem	Better approach
“How excellent was our fast, friendly service?”	Use neutral wording; ask about speed and friendliness separately
“Do you shop here regularly?”	Specify a recall period: “How many purchases did you make here in the past seven days?”
Overlapping numerical categories	Use non-overlapping, complete ranges; include zero where relevant

Offer “not applicable” or “don’t know” when meaningful. Do not force an opinion that the respondent does not have.

Pretest interpretation with potential respondents. **Pilot** the full process—invitation, routing, completion, recording and export—before launching it widely.

Reference: Pew: Writing survey questions.

Four sources of survey error

Error	Diagnostic question
Coverage	Could the intended population enter the sampling frame correctly?
Nonresponse	Did selected people provide the requested answers?
Sampling	How much might the estimate vary across repeated probability samples?
Measurement	Do recorded answers measure the intended characteristic?

Coverage, nonresponse and measurement errors are **non-sampling errors**. They can also affect a census. Processing mistakes, such as coding errors, are another source of non-sampling error.

Reference: ABS: Types of error.

Coverage error: who cannot enter?

The frame may omit eligible units, include ineligible units or list units more than once.

Illustration: A bank wants all customers’ views, but selects only from its mobile-app user list. Customers who do not use the app cannot be selected.

Improve it: Compare the frame with the target population; update contact information, remove duplicates and add missing groups through suitable sources.

Sending more invitations from the same incomplete frame does not correct the omission.

Nonresponse error: who does not answer?

Unit nonresponse: No questionnaire is returned. **Item nonresponse:** Some answers are missing.

Illustration: Selected dissatisfied customers may ignore a bank's invitation. The resulting satisfaction estimate could be too high if respondents and nonrespondents differ on satisfaction.

Improve it: Use clear invitations, reasonable length, reminders and alternative modes. Compare available characteristics of respondents and nonrespondents.

A response rate alone does not establish the size or direction of bias. Weighting may help with observed differences; it cannot guarantee that all bias is removed.

Sampling error: variation without a mistake

Repeated probability samples can give different estimates even when the same correct procedure is followed.

Larger samples usually reduce sampling variability under comparable designs. They do not automatically remove systematic coverage, nonresponse or measurement problems.

A confidence interval or margin of error describes specified sampling uncertainty under its assumptions; it does not normally include every source of survey error.

Precision concerns variability. **Bias** concerns systematic deviation. An estimate can be precise and biased.

Measurement error: what answer was obtained?

Recorded values may differ from the characteristic we intended to measure because of unclear wording, faulty recall, interviewer influence or a faulty instrument.

Illustration: A customer reports an approximate queue time from memory; the study needs elapsed time measured by timestamps. These measurements may differ systematically.

Improve it: Define units and recall periods, use neutral wording, check translations, train interviewers and pretest questions. Use an appropriate direct measure when available.

Correctly entering an answer does not prove that the underlying measurement is valid.

Before analysis: quality checks and responsible collection

- Explain purpose, voluntary participation and data use; collect only necessary identifiers and protect access.
- Promise anonymity only when responses cannot be linked to individuals; otherwise explain confidentiality accurately.
- Check duplicates, impossible values, units, missing-value codes and routing errors. Preserve raw records and document changes.
- Report the population, frame, selection method, modes, collection dates, achieved responses and known limitations.

Do not replace a missing answer with zero unless zero is its true meaning. Cleaning cannot recover people excluded by the study design.

Reference: AAPOR: Survey research practices.

Discussion: design and diagnose a customer survey

Hypothetical scenario; no observed data: A bank wants to describe service satisfaction among all current retail customers. It posts a questionnaire inside its app. The first question asks, “How satisfied are you with our excellent service?”

1. Define the population, observational unit and satisfaction measure.
2. Is the app a survey mode, a sampling frame, or both in this design? Who might be absent?
3. Propose a selection method and a response mode; explain why each fits.
4. Identify potential coverage, nonresponse and measurement problems. Could a larger response count fix them?
5. Rewrite the opening question and describe one pilot check.

After these design checks, we can organize the collected data using tables, charts and numerical summaries.

References for the added material

- Chapter 1 summary and teaching guide, *Defining and Collecting Data*, supplied in the earlier course. Used as a planning reference; textbook page claims and publisher datasets were not independently verified.
- Australian Bureau of Statistics: Census and sample.
- Australian Bureau of Statistics: Types of error.
- AAPOR: Best practices for survey research.
- Pew Research Center: Writing survey questions.
- Pew Research Center: Phone versus online mode effects.

Added course explanations and hypothetical illustrations, 3 November 2026. No original slide text or case observations were replaced.

Descriptive Statistics Tools

Statistical Tools Used in Descriptive Statistics

- Tables (Data Tables, Frequency Distribution Tables).
- Visualization & Charts.
- Numerical Measures (Measures of Central Location, Dispersion, Shape, Position, Association).



Tables in Descriptive Statistics

- Tables organize and display data in a structured format.
- They are essential for summarizing large data sets efficiently.

Types of Tables in Statistics

- Frequency Distribution Tables: Show how often each value in a set of data occurs.
- Summary Tables: Provide key statistics like mean, median, mode, range, etc., at a glance.
- Contingency Tables/Cross-tabulation: Display the multivariate frequency distribution for categorical data.

In the next section, we present tables examples using students dataset.

Introducing Students Dataset

#	stud.id	name	gender	age	height	weight	country	nc.score	semester	major	minor	score1	score2	av_scor	online.tutorial	graduated	salary
1	833917	Gonzales, (	Female	19	160	64.8	Turkey	1.91	1st	Political Sc	Social Scie	NA	NA	0	0	0	NA
2	898539	Lozano, TH	Female	19	172	73	Other	1.56	2nd	Social Scie	Mathemati	NA	NA	0	0	0	NA
3	379678	Williams, H	Female	22	168	70.6	UK	1.24	3rd	Social Scie	Mathemati	45	46	45.5	0	0	NA
4	807564	Nem, Deniz	Male	19	183	79.7	Other	1.37	2nd	Environme	Mathemati	NA	NA	0	0	0	NA
5	383291	Powell, He	Female	21	175	71.4	Italy	1.46	1st	Environme	Mathemati	NA	NA	0	0	0	NA
6	256074	Perez, Jadi	Male	19	189	85.8	Italy	1.34	2nd	Political Sc	Mathemati	NA	NA	0	0	0	NA
7	754591	Clardy, Ani	Female	21	156	65.9	UK	1.11	2nd	Political Sc	Social Scie	NA	NA	0	0	0	NA
8	146494	Allen, Rebe	Female	21	167	65.7	Other	2.03	3rd	Political Sc	Economics	58	62	60	0	0	NA
9	723584	Tracy, Robi	Male	18	195	94.4	Other	1.29	3rd	Economics	Environme	57	67	62	0	0	NA
10	314281	Nimmons,	Female	18	165	66	Russa	1.19	2nd	Environme	Mathemati	NA	NA	0	0	0	NA
11	200803	Lang, Mack	Female	22	162	66.8	Other	1.04	4th	Economics	Environme	62	61	61.5	1	1	45254.11
12	444907	Rodriguez,	Female	18	172	66.8	Other	3.81	3rd	Environme	Economics	76	82	79	0	0	NA
13	354271	Covar Orer	Male	23	185	84.6	Russa	1	4th	Environme	Mathemati	71	76	73.5	1	1	40552.79
14	317812	Lopez, Mor	Female	20	158	64.4	Italy	2.5	6th	Environme	Social Scie	66	70	68	1	1	27007.03
15	604115	Davis, Shaç	Female	19	157	66.3	Russa	1.92	2nd	Economics	Political Sc	NA	NA	0	0	0	NA
16	889551	Adams, Jos	Male	20	172	73.9	Other	3.61	4th	Mathemati	Political Sc	87	91	89	1	0	NA
17	350040	Hines, Hail	Female	22	156	61.7	Other	2.27	6th	Political Sc	Biology	57	54	55.5	0	1	33969.16
18	240279	Daugherty,	Male	22	182	82.1	Italy	1.42	1st	Economics	Environme	NA	NA	0	0	0	NA
19	865835	Roybal, Ebo	Female	21	162	69.2	Italy	1.32	3rd	Political Sc	Environme	69	46	57.5	1	0	NA
20	137196	Baysinger,	Female	22	168	70.9	UK	2.33	2nd	Environme	Political Sc	NA	NA	0	0	0	NA
21	708242	Phillips, La	Female	20	167	68.5	Other	1.79	4th	Biology	Economics	77	80	78.5	1	0	NA
22	499002	Culbertson	Male	37	175	70.4	UK	1.97	2nd	Political Sc	Environme	NA	NA	0	0	0	NA
23	873149	O Reilly, Jo	Male	19	164	70.3	UK	1.68	2nd	Political Sc	Environme	NA	NA	0	0	0	NA
24	807361	Johnson, S	Female	38	155	67	Italy	2.3	2nd	Environme	Biology	NA	NA	0	0	0	NA

R equivalent (added)

```r
# read.csv() imports the supplied CSV; stringsAsFactors=FALSE keeps category labels as text.
# head() previews six rows; height is in cm, weight in kg, and age in years.
# Find the shared data folder from either the course root or this notebook folder.
data_dir <- if (dir.exists("data")) "data" else file.path("..", "data")
students <- read.csv(file.path(data_dir, "descriptive-statistics/students.csv"), stringsAsFactors =
        FALSE)
head(students)
```

```
  gender age height weight country                major
1 Female  19    160   64.8  Turkey      Political Science
2 Female  19    172   73.0   Other         Social Sciences
3 Female  22    168   70.6      UK         Social Sciences
4   Male  19    183   79.7   Other Environmental Sciences
5 Female  21    175   71.4   Italy Environmental Sciences
6   Male  19    189   85.8   Italy      Political Science
```

Frequency Distribution Tables

Used to show the distribution of data across different categories or intervals.

Frequency Types:

- Absolute Frequency: The actual count of occurrences for each category.
- Relative Frequency: The proportion of the total count represented by each category, often expressed as a percentage.

- Cumulative Frequency: A running total of frequencies through the categories, showing the number of observations up to a certain category.

Row Labels	Number of Students	Relative Frequency	Cumulative Frequency
Italy	2797	33.95%	33.95%
Other	2688	32.63%	66.57%
UK	1839	22.32%	88.89%
Russa	585	7.10%	95.99%
Turkey	330	4.01%	100.00%
Grand Total	**8239**	**100.00%**	

Frequency of Students by Country

R equivalent (added)

```
# table() counts students in each country category; prop.table() divides by all counted students.
# Multiplying by 100 gives percentages; cumsum() is meaningful only when categories have an order.
country_counts <- table(students$country)
country_counts
```

```
Italy  Other  Russa Turkey     UK 
 2797   2688    585    330   1839 
```

```
100 * prop.table(country_counts)
```

```
    Italy     Other     Russa    Turkey        UK 
33.948295 32.625319  7.100376  4.005340 22.320670 
```

Converting Numerical Values into Categorical

Row Labels	Sum of age
<20	23415
20-29	144001
30-39	3785
40-49	5433
50-59	5947
60-70	3139
Grand Total	**185720**

Average Category	Description	Number of Students
Undefined	value undefined	3347
F	< 50	587
D	>= 50 and < 60	767
C	>= 60 and < 70	931
B	>= 70 and < 80	1575
A	>= 80	1032
	Total	8239

Converting average into average groups

Converting age into age groups

R equivalent (added)

```
# cut() converts ages in years into intervals; right=FALSE includes the lower boundary, not the upper.
# table() counts each interval; these explicitly chosen bands are an added R illustration.
age_groups <- cut(students$age, breaks = c(0, 20, 25, 30, Inf), right = FALSE)
table(age_groups)
```

```
age_groups
  [0,20)  [20,25)  [25,30) [30,Inf)
    1246     6167      433      393
```

Cross-tabulation or Contingency Tables

Count of gender	Column Labels		
Row Labels	Female	Male	Grand Total
Biology	959	638	1597
Economics and Finance	461	863	1324
Environmental Sciences	745	881	1626
Mathematics and Statistics	276	949	1225
Political Science	978	477	1455
Social Sciences	691	321	1012
Grand Total	**4110**	**4129**	**8239**

Used to show the **relationship between two categorical variables**.

Major by Gender in our example.


R equivalent (added)

```
# xtabs() counts combinations: the formula ~ major + gender creates a two-way frequency table.
# addmargins() adds row and column totals without changing the underlying counts.
major_gender <- xtabs(~ major + gender, data = students)
addmargins(major_gender)
```

major	gender		
	Female	Male	Sum
Biology	959	638	1597
Economics and Finance	461	863	1324
Environmental Sciences	745	881	1626
Mathematics and Statistics	276	949	1225
Political Science	978	477	1455
Social Sciences	691	321	1012
Sum	4110	4129	8239

Data Visualization


21



Visualizing Qualitative Data

Bar Chart, Pie Chart, Pareto Plots, Side by Side Bars Chart

Bar Charts




- A bar chart is a type of chart that presents data in rectangular bars with lengths proportional to the values they represent.
- Bar charts are commonly used to compare the magnitudes of different categories or groups of data.

Row Labels	Count by country
Italy	2797
Other	2688
Russa	585
Turkey	330
UK	1839
Grand Total	**8239**

We need to construct a frequency table first.


R equivalent (added)

```
# barplot() draws a bar for each category count supplied by table().
# las=2 rotates country labels; ylab states that bar heights are numbers of students.
barplot(table(students$country), las = 2, col = "#356B8C", ylab = "Number of students")
```




Pie Chart: Visualizing Proportions

- A pie chart is a circular chart that is divided into segments to represent numerical proportions or percentages.
- Each segment of the pie chart represents a different category or group, and the size of each segment is proportional to the value or percentage it represents.
- Pie charts are commonly used to display data that can be divided into categories, such as market share, demographic distributions, or survey results.




R equivalent (added)

```
# pie() divides the circle according to the country counts computed by table().
# The denominator is all students with a recorded country category, including Other.
pie(table(students$country))
```




Pareto Charts: Two Graphs in a Single Chart



- A Pareto chart is a type of chart that combines a bar graph with a line graph (Combo Chart).

- It is used to display the relative frequency or size of different categories or groups, and to identify the most important factors affecting a particular outcome or result.


Side by Side Bar Charts: : Visualizing Two Variables



- A side-by-side chart is used to compare two or more variables or categories.
- The chart displays the data side-by-side in separate columns or bars, making it easy to compare the values of each variable or category.
- Side-by-side chart allows viewers to quickly and easily compare the values of different variables and identify patterns or trends in the data.

R equivalent (added)

```r
# table() cross-classifies gender and country; barplot(beside=TRUE) places gender bars side by side.
# legend.text=TRUE labels the groups; counts, rather than percentages, determine bar heights.
barplot(table(students$gender, students$country), beside = TRUE, legend.text = TRUE,
        col = c("#356B8C", "#C47735"), ylab = "Number of students", las = 2)
```




Visualizing Quantitative Data, Frequency Distributions and Histograms

Histograms

- Histograms provide a way to visualize the distribution of a numeric variable.
- Histograms group numeric responses into bins and display the frequency of responses in each.
- The x-axis of a histogram reflects the range of values of a numeric variable, while the y-axis can reflect either the frequencies (counts) or relative frequencies.

R equivalent (added)


```r
# hist() groups ages into bins; breaks specifies yearly boundaries at half-year positions.
# freq=TRUE displays counts rather than probability density; recorded ages are measured in years.
hist(students$age, breaks = seq(min(students$age) - 0.5, max(students$age) + 0.5, by = 1),
     freq = TRUE, col = "#356B8C", main = "", xlab = "Age (years)")
```




Histograms

Histograms tell us the following about our data:


- Shape – what is the shape of the distribution?
- Center – what is an “average” value?
- Spread – how far away from the center do values tend to fall?
- Unique features – are there any outliers?






Number of Modes

- The first distinguishing feature apparent in a histogram is the number of modes, or, in the distribution.
- A unimodal distribution only has one peak in the distribution, a bimodal distribution has two peaks, and a multimodal distribution has three or more peaks.




Skewness

- Histograms can show us if the data is skewed or symmetric.
- Symmetric data should look nearly identical if folded in half at the center point of the distribution.
- Skewed data indicates that there is a large portion of the data collected on one side of the chart and only a small portion on the other side.



Outliers

- Outliers are responses that fall well away from the rest of the values.
- Histograms can be useful in identifying outliers in our data.
- Defining an observation as an outlier is subjective and should lead to an investigation of that value (not an automatic removal from the dataset).




Scotter Plots

Scatter plots can be used to examine of there is a relation between two variables.

Consider for instance, how:


- Incomes vary with education.
- Sales vary with advertising expenditures.
- Stock prices vary with corporate profits.
- Crop yields vary with the use of fertilizer.
- Cholesterol levels vary with dietary intake.

- Price varies with reliability.




R equivalent (added)

```
# plot(x,y) makes a scatterplot, with height on the horizontal axis and weight on the vertical.
# pch=16 uses filled points; no fitted line or causal relationship is assumed.
plot(students$height, students$weight, pch = 16, col = "#356B8C",
     xlab = "Height (cm)", ylab = "Weight (kg)")
```



Relations between Two Variables



Numerical Measures in Descriptive Statistics

Numerical Measures in Descriptive Statistics

- Numerical measures in descriptive statistics are used to summarize and describe the main features of a dataset through numbers.
- These measures provide insights into the distribution, central tendency, dispersion, and shape of the data.
- Understanding these measures helps in interpreting the data more effectively.

Examples

- Rates in a Society (unemployment rate, crime rate, divorce rate).
- Average Income, Maximum age, Minimum score…etc.

Measures of Central Location

Mean, Median and Mode

The Mean

- The **Arithmetic Mean** is the primary measure of central location.
- Generally, we refer to the arithmetic mean as simply the **Mean** or the **Average**.
- In order to calculate the mean of a data set, we simply add up the observations and divide by the number of observations in the population or sample.
- **However, the mean is sensitive to Outliers. Consider the following sample dataset:**

Data = ($1.00, $2.00, $3.00, $3.00, $3.00, $1.00, $100.00)

- **Most of the data values are around $3, yet their average is $16.14**

The Median

MEASURE OF CENTRAL LOCATION: THE MEDIAN

The median is the middle value of a data set. The data are arranged in ascending order (smallest to largest) and the median is calculated as

- The middle value if the number of observations is odd, or
- The average of the two middle values if the number of observations is even.

The median is especially useful when outliers are present.

1, 3, 3, **6**, 7, 8, 9

Median = 6

1, 2, 3, **4**, **5**, 6, 8, 9

Median = (4 + 5) ÷ 2

= 4.5

$$\text{Med}(X) = \begin{cases} X[\frac{n+1}{2}] & \text{if n is odd} \\ \frac{X[\frac{n}{2}]+X[\frac{n}{2}+1]}{2} & \text{if n is even} \end{cases}$$

X = ordered list of values in data set

n = number of values in data set

The Mode

MEASURE OF CENTRAL LOCATION: THE MODE

The mode is the most frequently occurring value in a data set. A data set may have no mode or more than one mode.

Excel Functions for Measuring Central Location

Mean	=AVERAGE(array)
Median	=MEDIAN(array)
Mode	=MODE(array)
Minimum	=MIN(array)
Maximum	=MAX(array)

Practical Exercise

Use the students.xlsx file:

- Compare the Mean, Median for the “age” column for “Italy” and “Russa”.
- Find the Mode “age” column for “UK”.

Note: In Excel you can filter out rows by converting a Range to a Table and using the filtering list in each header of a column.

	A	B	C	D	E	F	G	H
	stud.id	name	gender	age	height	weight	country
1	833917	Gonzales,	Female	19	160	64.8	Turkey
2	898539	Lozano, T	Female	19	172	73	Other
3	379678	Williams,	Female	22	168	70.6	UK
4	807564	Nem, Denz	Male	19	183	79.7	Other
5	383291	Powell, He	Female	21	175	71.4	Italy
6	256074	Perez, Jadı	Male	19	189	85.8	Italy
7	754591	Clardy, Ani	Female	21	156	65.9	UK
8	146494	Allen, Rebe	Female	21	167	65.7	Other
9	723584	Tracy, Robı	Male	18	195	94.4	Other
10	314281	Nimmons,	Female	18	165	66	Russa
11	200803	Lang, Mack	Female	22	162	66.8	Other
12	444907	Rodriguez,	Female	18	172	66.8	Other
13	354271	Covar Orer	Male	23	185	84.6	Russa
14	317812	Lopez, Mor	Female	20	158	64.4	Italy
15	604115	Davis, Shaı	Female	19	157	66.3	Russa
16	889551	Adams, Jos	Male	20	172	73.9	Other
17	350040	Hines, Hail	Female	22	156	61.7	Other
18	240279	Daugherty,	Male	22	182	82.1	Italy
19	865835	Roybal, Ebı	Female	21	162	69.2	Italy
20	137196	Baysinger,	Female	22	168	70.9	UK
21	708242	Phillips, La	Female	20	167	68.5	Other
22	499002	Culbertson	Male	37	175	70.4	UK
23	873149	O Reilly, Jo	Male	19	164	70.3	UK
24	807361	Johnson, S	Female	38	155	67	Italy

R equivalent (added)

```
# mean() and median() summarize ages; table() counts each recorded age to identify every tied mode.
# names() returns the ages whose counts equal max(), the greatest frequency.
mean(students$age)
```

```
[1] 22.54157
```

```
median(students$age)
```

```
[1] 21
```

```
age_counts <- table(students$age)
names(age_counts)[age_counts == max(age_counts)]
```

```
[1] "21"
```

Measures of Dispersion

Range, Mean Absolution Difference, Variance and Standard Deviation

---

Measures of Dispersion

- Average alone is not sufficient to summarize and describe our data.
- Consider the following 5 sample datasets.
- Although the data values are very different, they all have the same average.

	S1	S2	S3	S4	S5
	100	200	0	-600	-2000
	200	200	600	600	2500
	300	200	0	600	100
Average	**200.00**	**200.00**	**200.00**	**200.00**	**200.00**

Measures of Dispersion – The Range

- The **Range** is the simplest measure of dispersion, it is the difference between the maximum value and the minimum value in a data set:

Range = Maximum Value – Minimum Value

- Going back to the previous sample datasets, we can see that they have the same average, but their ranges are very different.

	S1	S2	S3	S4	S5
	100	200	0	-600	-2000
	200	200	600	600	2500
	300	200	0	600	100
Average	200.00	200.00	200.00	200.00	200.00
Range	**200.00**	**0.00**	**600.00**	**1200.00**	**4500.00**

- Therefore, using the **Range** with the **Average** helped us to better describe and summarize our data (Better Understand Our Data).
- **However, the Range is sensitive to Outliers. Consider the following sample dataset:**

Data = ($1.00, $2.00, $3.00, $3.00, $3.00, $1.00, $100.00)

- Most of the data values are around $3, yet their Range is $99.00

Measures of Dispersion - The Mean Absolute Deviation

- A good measure of dispersion should consider differences of all observations from the mean.
- If we simply average all differences from the mean, the positives and the negatives will cancel out, even though they both contribute to dispersion, and the resulting average will equal zero.
- The **mean absolute deviation** (MAD) is an average of the absolute differences between the observations and the mean.


MEASURE OF DISPERSION: THE MEAN ABSOLUTE DEVIATION (MAD)

For sample values, $x_1, x_2, \ldots, x_n$, the sample MAD is computed as


$$\text{Sample MAD} = \frac{\Sigma|x_i - \bar{x}|}{n}.$$

For population values, $x_1, x_2, \ldots, x_N$, the population MAD is computed as


$$\text{Population MAD} = \frac{\Sigma|x_i - \mu|}{N}.$$



Measures of Dispersion - The Variance and the Standard Deviation



- MAD weights large and small differences equally.
- Better measures are the **variance** and the **standard deviation.**
- Instead of calculating the average of the absolute differences from the mean, as in MAD, we calculate the average of the squared differences from the mean.
- The squaring of differences from the mean emphasizes larger differences more than smaller ones.
- The variance has squared units, therefore, to return the values to their original units of measurement, we compute the square root of the variance, which gives us the standard deviation.



Measures of Dispersion - The Variance and the Standard Deviation

Sample

For **sample** values $x_1, x_2, \ldots, x_n$, the sample variance s^2 and the sample standard deviation s are computed as

$$s^2 = \frac{\sum_{i=1}^{n} (x_i - \bar{x})^2}{n-1} \quad \text{and} \quad s = \sqrt{s^2}.$$

Population

For **population** values $x_1, x_2, \ldots, x_N$, the population variance σ^2 (the Greek letter sigma, squared) and the population standard deviation σ are computed as

$$\sigma^2 = \frac{\sum_{i=1}^{N} (x_i - \mu)^2}{N} \quad \text{and} \quad \sigma = \sqrt{\sigma^2}.$$

> **Note:** The sample variance uses $n - 1$ rather than n in the denominator to ensure that the sample variance is an unbiased estimator for the population variance, a topic discussed in Chapter 8.

Excel Functions for Measuring Dispersion

Range	=MAX(array)-MIN(array)
Mean Absolute Deviation	=AVEDEV(array)
Sample Variance	=VAR.S(array)
Sample Standard Deviation	=STDEV.S(array)
Population Variance	=VAR.P(array)
Population Standard Deviation	=STDEV.P(array)

Practical Exercise

Use the students.xlsx file to find the Range, MAD, Variance and Standard Deviation for the age column.

Note: In Excel you can filter out rows by converting a Range to a Table and using the filtering list in each header of a column.

	A	B	C	D	E	F	G	H
	stud.id	name	gender	age	height	weight	country	
1	833917	Gonzales,	Female	19	160	64.8	Turkey	
2	898539	Lozano, T	Female	19	172	73	Other	
3	379678	Williams,	Female	22	168	70.6	UK	
4	807564	Nem, Denz	Male	19	183	79.7	Other	
5	383291	Powell, He	Female	21	175	71.4	Italy	
6	256074	Perez, Jadr	Male	19	189	85.8	Italy	
7	754591	Clardy, Ani	Female	21	156	65.9	UK	
8	146494	Allen, Rebe	Female	21	167	65.7	Other	
9	723584	Tracy, Robi	Male	18	195	94.4	Other	
10	314281	Nimmons,	Female	18	165	66	Russa	
11	200803	Lang, Mack	Female	22	162	66.8	Other	
12	444907	Rodriguez,	Female	18	172	66.8	Other	
13	354271	Covar Orer	Male	23	185	84.6	Russa	
14	317812	Lopez, Mor	Female	20	158	64.4	Italy	
15	604115	Davis, Shaq	Female	19	157	66.3	Russa	
16	889551	Adams, Jos	Male	20	172	73.9	Other	
17	350040	Hines, Hail	Female	22	156	61.7	Other	
18	240279	Daugherty,	Male	22	182	82.1	Italy	
19	865835	Roybal, Ebo	Female	21	162	69.2	Italy	
20	137196	Baysinger,	Female	22	168	70.9	UK	
21	708242	Phillips, La	Female	20	167	68.5	Other	
22	499002	Culbertson	Male	37	175	70.4	UK	
23	873149	O Reilly, Jo	Male	19	164	70.3	UK	
24	807361	Johnson, S	Female	38	155	67	Italy	


R equivalent (added)

```
# diff(range()) gives maximum minus minimum; mean(abs()) calculates mean absolute deviation.
# var() and sd() use the sample denominator n-1; R's mad() instead computes median absolute deviation.
age <- students$age
c(range = diff(range(age)), mean_absolute_deviation = mean(abs(age - mean(age))),
  sample_variance = var(age), sample_standard_deviation = sd(age))
```

```
                    range   mean_absolute_deviation           sample_variance
                46.000000                  2.902923                 36.785568
sample_standard_deviation
                 6.065111
```

Practical Exercise: using Excel's data analysis tools option

Select: Data > Data Analysis > Descriptive Statistics > OK.



The Coefficient of Variation

The Coefficient of Variation

- In some instances, we need to compare the variability of two or more data sets that have different means or different units of measurement.
- Consider the following example:
- We cannot tell which dataset has more variance in its values since both datasets have the same standard deviation.

	DS1	DS2
	$10	$1010
	$20	$1020
	$30	$1030
	$40	$1040
	$50	$1050
Average	**$30**	**$1030**
Standard Deviation	**15.81**	**15.81**

The Coefficient of Variation

- For such cases, we can use the Coefficient of Variation Measure which serves as a relative measure of dispersion.
- It can be calculated by dividing a data set's standard deviation by its mean:

- When comparing two datasets, the dataset that have larger coefficient of variation indicates that its values varies more than the other dataset. Therefore, it is less stable or less uniform.

$$CV = \frac{\sigma}{\mu}$$

The Coefficient of Variation

- Going back to the previous example, we can calculate the coefficient of variation for both sets to help us in determining which dataset has more variability in its values.
- Using the Coefficient of Variation, we can tell that DS1 varies more than DS2, and that DS2 is more stable than DS1.

	DS1	**DS2**
	$10	$1010
	$20	$1020
	$30	$1030
	$40	$1040
	$50	$1050
Average	**$30**	**$1030**
Standard Deviation	**$15.81**	**$15.81**
Coefficient of Variation	**0.430**	**0.013**

R equivalent (added)

```
# c() creates the exact two datasets shown in the original table; sd() uses the sample denominator n-
        1.
# sd()/mean() expresses CV as a ratio, without multiplying by 100; these means are positive.
DS1 <- c(10, 20, 30, 40, 50)
DS2 <- c(1010, 1020, 1030, 1040, 1050)
c(DS1 = sd(DS1) / mean(DS1), DS2 = sd(DS2) / mean(DS2))
```

```
       DS1        DS2
0.52704628 0.01535086
```

Coefficient of Variation Uses in Finance

- In finance, the coefficient of variation is used to measure the risk per unit of return.
- For example, assume that the average monthly return on stock A is 2980 with a standard deviation of 111.06, and the average monthly return on stock B is 9.8 with a standard deviation of 1.3
- We can use the coefficient of variation to determine which stock is more stable:

$$= \frac{111.6}{2980} = 0.037$$


$$= \frac{1.3}{9.8} = 0.134$$

Measures of Association

Covariance and Coefficient of Correlation

Measures of Association

- We previously introduced the idea of a scatter plots to visually assess whether two variables had some type of linear relationship.
- In this section, we present two numerical measures of association that quantify the direction and strength of the linear relationship between two variables, x and y.
- These two numerical values are the Covariance and the Correlation.




Note: Covariance and Correlation works when the relation between the two variables is LINEAR.


Covariance


- Covariance measures the joint variability of two random variables.
- Indicates the direction of the linear relationship between variables.

$$cov_{x,y} = \frac{\sum(x_i - \bar{x})(y_i - \bar{y})}{N - 1}$$

- Positive Covariance: X and Y move in the same direction.
- Negative Covariance: X and Y move in opposite directions.
- Zero Covariance: No linear relationship between X and Y.







Limitations of Covariance

Covariance is sensitive to the units of X and Y.

Changing units changes the covariance value, making direct comparisons difficult.


	Height m	Height cm	Weight KG
	1.6	160	55
	1.7	170	68
	1.8	180	74
Covariance	**.95**	**95**	Which one is bigger?
Correlation	**.98**	**.98**	


The Correlation Coefficient

- Correlation coefficient can be used to determine the Direction and the Strength of a relationship between two numerical variables.
- The correlation coefficient is a dimensionless metric and its value ranges from -1 to +1.
- The closer it is to +1 or -1, the more closely the two variables are related.

$$r = \frac{cov(x, y)}{\sigma_x \cdot \sigma_y}$$

- Correlation Formula





Differences between Covariance and Correlation

Covariance	Correlation
Covariance indicates the direction of the linear relationship between variables.	Correlation measures both the strength and the direction of the linear relationship between two variables.
Covariance value can vary between -∞ and +∞	Correlation value ranges between -1 and +1
Covariance unit is based on the product of the units of the two variables.	Correlation is dimensionless, i.e. It's a unit-free measure of the relationship between variables.

Practical Exercise

Use the students.xlsx file:

- Compute the covariance and correlation between heights and weights of the students.
- Compute the covariance and correlation between heights and ages of the students.

#	stud.id	name	gender	age	height	weight
1	833917	Gonzales, (	Female	19	160	64.8
2	898539	Lozano, TH	Female	19	172	73
3	379678	Williams, H	Female	22	168	70.6
4	807564	Nem, Denz	Male	19	183	79.7
5	383291	Powell, He	Female	21	175	71.4
6	256074	Perez, Jadr	Male	19	189	85.8
7	754591	Clardy, Ani	Female	21	156	65.9
8	146494	Allen, Rebe	Female	21	167	65.7
9	723584	Tracy, Robe	Male	18	195	94.4

=CORREL(Table1[age],Table1[height])

=COVARIANCE.S(Table1[age],Table1[height])

=COVARIANCE.P(Table1[age],Table1[height])

R equivalent (added)

```
# cov() calculates sample covariance and cor() calculates Pearson's linear correlation by default.
# use="complete.obs" omits pairs with missing values; units remain cm, kg, and years.
cov(students$height, students$weight, use = "complete.obs")
```

```
[1] 89.89815
```

```
cor(students$height, students$weight, use = "complete.obs")
```

```
[1] 0.9398042
```

```
cov(students$height, students$age, use = "complete.obs")
```


```
[1] -0.8861261
```

```
cor(students$height, students$age, use = "complete.obs")
```

```
[1] -0.01318906
```

Practical Exercise

We can also use the Data Analysis add-in for this purpose.




#	stud.id	name	gender	age	height	weight
1	833917	Gonzales, (	Female	19	160	64.8
2	898539	Lozano, T'H	Female	19	172	73
3	379678	Williams, H	Female	22	168	70.6
4	807564	Nem, Denz	Male	19	183	79.7
5	383291	Powell, He	Female	21	175	71.4
6	256074	Perez, Jadr	Male	19	189	85.8
7	754591	Clardy, Ani	Female	21	156	65.9
8	146494	Allen, Rebe	Female	21	167	65.7
9	723584	Tracy, Robe	Male	18	195	94.4
10	314281	Nimmons,	Female	18	165	66
11	200803	Lang, Mack	Female	22	162	66.8
12	444907	Rodriguez,	Female	18	172	66.8
13	354271	Covar Orer	Male	23	185	84.6

Measures of Distribution


Symmetry and Skewness

Skewness



- **Definition**: Skewness is a measure of the asymmetry of the probability distribution of a real-valued random variable about its mean.


- **Key Point**: Indicates whether the data are spread out more to one side of the mean.
- **Positive Skewness (Right-skewed): Tail is longer on the right side of the distribution. Mean is greater than the median.**
- **Negative Skewness (Left-skewed**): Tail is longer on the left side of the distribution. Mean is less than the median.
- Symmetrical Distribution: Skewness is around 0, indicating no skew.
- Use histograms or density plots to illustrate examples of positively skewed, negatively skewed, and symmetrical distributions.



Skewness Coefficient

- A positive skewness coefficient implies that extreme values are concentrated in the right tail of the distribution, pulling the mean up, relative to the median and the bulk of values lie to the left of the mean.
- Similarly, a negative skewness coefficient implies that extreme values are concentrated in the left tail of the distribution, pulling the mean down, relative to the median, and the bulk of values lie to the right of the mean.


Formula: $Skewness = \frac{1}{n}\sum_{i=1}^{n}\left(\frac{X_i-\mu}{\sigma}\right)^3$



- **A skewness value greater than 1 or less than -1 indicates a highly skewed distribution.**
- **A value between 0.5 and 1 or -0.5 and -1 is moderately skewed.**
- **A value between -0.5 and 0.5 indicates that the distribution is fairly symmetrical.**
- **A skewness coefficient of zero indicates that the values are evenly distributed on both sides of the mean.**

Practical Example

- Using the students.xlsx file, find the Skewness value for the “age” column.
- You can use Excel’s skew() function to calculate the skewness for a sample, and the skew.p() function to calculate the skewness for a population.




R equivalent (added)

```
# length() supplies n; sum() adds cubed standardized deviations, using sample sd() with n-1.
# n/((n-1)*(n-2)) gives adjusted sample skewness, corresponding to Excel SKEW for n>2.
age <- students$age
n <- length(age)
n / ((n - 1) * (n - 2)) * sum(((age - mean(age)) / sd(age))^3)
```

```
[1] 4.427823
```


Practical Exercise: using Excel’s data analysis tools option

Select: Data > Data Analysis > Descriptive Statistics > OK.



Percentiles, Quartiles and Boxplots


Percentiles



- A **percentile** is a measure that helps us understand the position or relative standing of a particular value within a dataset.
- Percentiles help us divide the data into equal-sized groups and determine the position of a value within those groups.
- For example, if someone's score is at the 80th percentile, it means they performed better than 80% of the people in the dataset.
- Percentiles provide insights into how a specific value compares to others, allowing us to assess performance, rankings, or distributions.
- They are commonly used in various fields, such as education, healthcare, and finance, to understand and communicate data in a meaningful way.

Practical Exercise

- Using the students.xlsx file, find the 90% Percentile value for the “height” column.
- You can use percentile.inc or percentile.exc to compute percentiles in Excel.



	A	B	C	D	E	F	G
1	age	height	weight				
2	19	160	64.8				
3	19	172	73			90% Percentile for Students Height	
4	22	168	70.6		percentile.inc	=PERCENTILE.INC(Table2[height],0.9)	186
5	19	183	79.7		percentile.ex	=PERCENTILE.EXC(Table2[height],0.9)	186
6	21	175	71.4				
7	19	189	85.8				
8	21	156	65.9				
9	21	167	65.7				
10	18	195	94.4				

R equivalent (added)

```
# quantile() finds the requested percentile; probs=0.90 means the 90th percentile of height in cm.
# type=7 matches Excel PERCENTILE.INC; type=6 matches PERCENTILE.EXC for this sample and probability.
quantile(students$height, probs = 0.90, type = 7)
```


```
90%
186
```

```
quantile(students$height, probs = 0.90, type = 6)
```

```
90%
186
```

Quartiles


A Quartile divides a sorted data set into 4 equal parts, so that each part represents ¼ of the data set.



Practical Exercise

- Using the students.xlsx file, find the 1st and the 2nd quartiles for the “height” column.
- You can use quartile.inc or quartile.exc to compute percentiles in Excel.

	A	B	C	D	E
1	height				
2	160		1st Quartile INC	=QUARTILE.INC(A2:A8240,1)	163
3	172		2nd Quartile EXC	=QUARTILE.EXC(A3:A8241,2)	171
4	168				
5	183				



R equivalent (added)

```
# quantile(probs=c(0.25,0.50)) returns the first and second quartiles of height in cm.
# type=7 uses Excel's inclusive convention; type=6 uses its exclusive convention for these
        probabilities.
quantile(students$height, probs = c(0.25, 0.50), type = 7)
```


```
25% 50%
163 171
```

```
quantile(students$height, probs = c(0.25, 0.50), type = 6)
```

```
25% 50%
163 171
```

Boxplots – Box and Whiskers Plots


- A boxplot (box-and-whisker plot) is a standardized way of displaying the distribution of data based on a five number summary.
- The five number summary include the minimum value (Min), quartiles (Q1, Q2, and Q3), and the maximum value (Max).
- Boxplots are useful in showing us the range and the skewness of our data as well as outliers.



Practical Exercise


Use the students.xlsx file:

- Create a box plot that compares the height of male students vs female students.
- Create a box plot that compares the height of Turkish male students vs Russian male students.




R equivalent (added)

```r
# boxplot(height~gender) compares groups; whiskers extend to the last data within 1.5 IQR of the
        hinges.
# subset() selects Turkish/Russian males; R's default boxplot hinges can differ from type-7 quartiles.
boxplot(height ~ gender, data = students, ylab = "Height (cm)", col = "#D4D4D4")
```






```r
# The source spells Russia as "Russa"; match that original label without changing the data.
males <- subset(students, gender == "Male" & country %in% c("Turkey", "Russa"))
boxplot(height ~ country, data = males, ylab = "Height (cm)", col = "#D4D4D4")
```




Outliers

- An outlier is an observation that lies an abnormal distance from other values in a random sample from a population.
- We can identify potential outliers using Histograms, scatter plots, boxplots…etc.







The Interquartile Range (IQR)

- The IQR is like the range, and it shows how the data is spread about the median.
- It is less susceptible than the range to outliers and can, therefore, be more helpful.
- The IQR is calculated according to the following formula:

IQR = Q3 – Q1.

Using the IQR to Find Outliers

Using the IQR to find potential outliers:

- Calculate the IQR for the data.
- Multiply the interquartile range (IQR) by 1.5 (a constant used to discern outliers).
- Add the (1.5 x (IQR)) to the third quartile, any number greater than this is a suspected outlier.

- Subtract the (1.5 x (IQR)) from the first quartile, any number less than this is a suspected outlier.

However, any potential outlier obtained by the interquartile method should be examined in the context of the entire set of data.

We should always follow up our outlier analysis by studying the resulting outliers to see if they make sense.

	A	B	C	D	E	F
1	gender	height				
2	Male	183		Height of Male Students		
3	Male	189		Q1	=QUARTILE.INC(Table3[[#All],[height]],1)	174
4	Male	195		Q3	=QUARTILE.EXC(Table3[[#All],[height]],3)	184
5	Male	185		IQR = Q3 - Q1	=F4-F3	10
6	Male	172		Upper Bound = 1.5 * IQR + Q3	=1.5*F5+F4	199
7	Male	182		Lower Bound = Q1 - 1.5 * IQR	=F3-1.5*F5	159
8	Male	175				
9	Male	164				

R equivalent (added)

```r
# quantile(type=7) supplies inclusive quartiles; the 1.5-IQR fences flag values for investigation.
# sum() counts TRUE flags and does not remove observations; these age fences are measured in years.
q <- quantile(students$age, probs = c(0.25, 0.75), type = 7)
iqr <- q[2] - q[1]
lower <- q[1] - 1.5 * iqr
upper <- q[2] + 1.5 * iqr
c(lower_fence = unname(lower), upper_fence = unname(upper))
```

```
lower_fence upper_fence
       15.5        27.5
```

```r
sum(students$age < lower | students$age > upper)
```

```
[1] 429
```

Outliers

Impacts of Outliers

1. Negative Impacts


- Influence on Measures of Central Tendency: A single extreme outlier can pull the mean in its direction, making it unrepresentative of the majority of the data.
- Impact on Dispersion Measures: The presence of outliers can inflate the measures like the standard deviation and the interquartile range (IQR), making them larger than they would be without outliers.
- Skewing Data Distributions: Positive outliers can result in right-skewed distributions, while negative outliers can result in left-skewed distributions. This can affect the interpretation of the data.

- Misleading Summary Statistics: Outliers can distort the interpretation of summary statistics.
- Impact on Hypothesis Testing: Outliers can affect the results of hypothesis tests. They can lead to incorrect conclusions, such as detecting significant differences when none exist or failing to detect real differences when outliers mask them.

2. Positive Impacts


- Detection of Anomalies: Outliers can signal the presence of anomalies or rare events in a dataset. Identifying these anomalies can be valuable in various fields, including fraud detection, quality control, and outlier detection in scientific experiments.
- Robust Modeling: In some cases, outliers can be genuine observations that are important to model. For example, in financial modeling, extreme stock price movements may contain valuable information for predicting market trends.

Methods to screen for outliers in a dataset



- There are several methods to screen for outliers in a dataset, ranging from graphical techniques to statistical tests. Here are some commonly used methods:
- Box Plots (Box-and-Whisker Plots):
  Box plots provide a visual representation of the distribution of data, including the identification of potential outliers. In a box plot, outliers are typically shown as individual data points beyond the whiskers of the plot.

Methods to screen for outliers in a dataset



Scatterplots:

- Scatterplots are particularly useful for identifying outliers in bivariate or multivariate data.
- Outliers can appear as data points that are far from the main cluster of points in the scatterplot.


Methods to screen for outliers in a dataset

- Z-Scores:
  Z-scores (standard scores) measure how many standard deviations a data point is away from the mean. Data points with high absolute Z-scores (typically greater than 2 or 3) are often considered potential outliers.
- IQR (Interquartile Range) Method:
  The IQR method involves calculating the interquartile range (IQR = Q3 – Q1) and then identifying values that fall below Q1 – 1.5 × IQR or above Q3 + 1.5 × IQR as potential outliers.

Methods to screen for outliers in a dataset

Visual Inspection:

- Sometimes, simple visual inspection of the data through histograms, QQ plots (quantile-quantile plots), or other visualization techniques can reveal the presence of outliers.



Handling Outliers in Datasets

- Handling outliers is an important step in data preprocessing to ensure they do not unduly influence the results of analysis or modeling. The approach depends on the nature of the data, the context of the analysis, and the specific objectives. Here are several methods for handling outliers:
- **Data Truncation or Removal:**
  - Simply remove outliers from the dataset.
  - Should be done cautiously, especially if the outliers represent valid and important observations.
  - Appropriate when outliers are likely due to data entry or measurement errors.
- **Data Transformation:**
  - Transforming the data can mitigate the impact of outliers.
  - Common transformations: logarithmic, square root, or inverse.
  - These compress the range of extreme values.
- **Winsorization:**
  - Involves capping or limiting extreme values by replacing them with a specified percentile value.
  - Example: Replace values above the 95th percentile with the value at the 95th percentile.
- **Imputation:**
  - For missing values (not extreme outliers), impute using methods like:
    - Mean imputation
    - Median imputation
    - Advanced methods (e.g., regression imputation).

Handling Outliers in Datasets

- **Robust Statistics:**
  - Use statistical methods less sensitive to outliers.
  - Examples:
    - Replace mean with median.
    - Use interquartile range (IQR) instead of standard deviation.
- **Model-Based Approaches:**
  - In predictive modeling, use algorithms less sensitive to outliers.
  - Examples: robust regression methods, ensemble methods (e.g., random forests).
- **Domain Knowledge:**
  - Rely on domain expertise to interpret outliers.
  - What looks like an outlier may actually be an important and valid data point.
- **Reporting and Transparency:**
  - Clearly document how outliers were handled.
  - Ensures reproducibility and interpretability of results.

Practice Datasets



Practice Statistics Using the following Datasets

Suggested Datasets:

https://github.com/msfasha/307102-Descriptive-Statistics-for-Business/blob/main/Practical/additional/useful_datasets.md

Suggested Analysis Scenario:

https://github.com/msfasha/307102-Descriptive-Statistics-for-Business/blob/main/Practical/additional/superstore_analysis_scenario.md

---