# Solar Position Algorithm for Solar Radiation Applications

Ibrahim Reda and Afshin Andreas



1617 Cole Boulevard Golden, Colorado 80401-3393

NREL is a U.S. Department of Energy Laboratory Operated by Midwest Research Institute • Battelle • Bechtel

Contract No. DE-AC36-99-GO10337

# Solar Position Algorithm for Solar Radiation Applications

#### Ibrahim Reda and Afshin Andreas

Prepared under Task No. WU1D5600



1617 Cole Boulevard Golden, Colorado 80401-3393

NREL is a U.S. Department of Energy Laboratory Operated by Midwest Research Institute • Battelle • Bechtel

Contract No. DE-AC36-99-GO10337

#### Acknowledgment

We thank Bev Kay for all her support by manually typing all the data tables in the report into text files, which made it easy and timely to transport to the report text and all of our software code. We also thank Daryl Myers for all his technical expertise in solar radiation applications.

#### NOTICE

This report was prepared as an account of work sponsored by an agency of the United States government. Neither the United States government nor any agency thereof, nor any of their employees, makes any warranty, express or implied, or assumes any legal liability or responsibility for the accuracy, completeness, or usefulness of any information, apparatus, product, or process disclosed, or represents that its use would not infringe privately owned rights. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise does not necessarily constitute or imply its endorsement, recommendation, or favoring by the United States government or any agency thereof. The views and opinions of authors expressed herein do not necessarily state or reflect those of the United States government or any agency thereof.

Available electronically at <a href="http://www.osti.gov/bridge">http://www.osti.gov/bridge</a>

Available for a processing fee to U.S. Department of Energy and its contractors, in paper, from:

U.S. Department of Energy Office of Scientific and Technical Information P.O. Box 62 Oak Ridge, TN 37831-0062 phone: 865.576.8401

fax: 865.576.5728

email: reports@adonis.osti.gov

Available for sale to the public, in paper, from:

U.S. Department of Commerce National Technical Information Service 5285 Port Royal Road Springfield, VA 22161

Springfield, VA 22161 phone: 800.553.6847 fax: 703.605.6900

email: orders@ntis.fedworld.gov

online ordering: <a href="http://www.ntis.gov/ordering.htm">http://www.ntis.gov/ordering.htm</a>



### **Table of Contents**

Abstractv
Introduction
Time Scale
Procedure
SPA Evaluation and Conclusion
List of Figures
Figure 1. Uncertainty of cosine the solar zenith angle resulting from 0.01° and 0.0003° uncertainty in the angle calculation
Figure 2. Difference between the Almanac and SPA for the ecliptic longitude & latitude, and the apparent right ascension & declination on the second day of each month at 0-TT for the years 1994, 1995, 1996, and 2004
Figure 3. Difference between the Almanac and SPA for the solar zenith and azimuth angles on the second day of each month at 0-TT for the years 1994, 1995, 1996, and 2004 15
References
Appendix
Equation of Time
Sunrise, Sun Transit, and Sunset
Calculation of Calendar Date from Julian Day
Example A-15
C Code: SPA header file (SPA.h)
C Code: SPA source file (SPA.c)

### List of Appendix Figures

Figure A2.1. Difference between the Almanac and SPA for the Ephemeris Transit on the second day of each month at 0-TT for the years 1994, 1995, 1996, and 2004
List of Appendix Tables
Table A4.1. Examples for Testing any Program to Calculate the Julian Day
Table A4.2. Earth Periodic Terms
Table A4.3. Periodic Terms for the Nutation in Longitude and Obliquity
Table A5.1. Results for Example

#### Abstract

There have been many published articles describing solar position algorithms for solar radiation applications. The best uncertainty achieved in most of these articles is greater than  $\pm 0.01^{\circ}$  in calculating the solar zenith and azimuth angles. For some, the algorithm is valid for a limited number of years varying from 15 years to a hundred years. This report is a step by step procedure for implementing an algorithm to calculate the solar zenith and azimuth angles in the period from the year -2000 to 6000, with uncertainties of  $\pm 0.0003^{\circ}$ . The algorithm is described by Jean Meeus [3]. This report is written in a step by step format to simplify the complicated steps described in the book, with a focus on the sun instead of the planets and stars in general. It also introduces some changes to accommodate for solar radiation applications. The changes include changing the direction of measuring azimuth angles to be measured from north and eastward instead of being measured from south and eastward, and the direction of measuring the observer's geographical longitude to be measured as positive eastward from Greenwich meridian instead of negative. This report also includes the calculation of incidence angle for a surface that is tilted to any horizontal and vertical angle, as described by Iqbal [4].

#### 1. Introduction

With the continuous technological advancements in solar radiation applications, there will always be a demand for smaller uncertainty in calculating the solar position. Many methods to calculate the solar position have been published in the solar radiation literature, nevertheless, their uncertainties have been greater than  $\pm$  0.01° in solar zenith and azimuth angle calculations, and some are only valid for a specific number of years[1]. For example, Michalsky's calculations are limited to the period from 1950 to 2050 with uncertainty of greater than  $\pm$  0.01° [2], and the calculations of Blanco-Muriel et al.'s are limited to the period from 1999 to 2015 with uncertainty greater than  $>\pm$  0.01° [1].

An example emphasizing the importance of reducing the uncertainty of calculating the solar position to lower than  $\pm$  0.01°, is the calibration of pyranometers that measure the global solar irradiance. During the calibration, the responsivity of the pyranometer is calculated at zenith angles from 0° to 90° by dividing its output voltage by the reference global solar irradiance (G), which is a function of the cosine of the zenith angle (cos  $\theta$ ). Figure 1 shows the magnitude of errors that the 0.01° uncertainty in  $\theta$  can contribute to the calculation of cos  $\theta$ , and consequently G that is used to calculate the responsivity. Figure 1 shows that the uncertainty in cos  $\theta$  exponentially increases as  $\theta$  reaches 90° (e.g. at  $\theta$  equal to 87°, the uncertainty in cos  $\theta$  is 0.7%, which can result in an uncertainty of 0.35% in calculating G; because at such large zenith angles the normal incidence irradiance is approximately equal to half the value of G). From this arises the need to use a solar position algorithm with lower uncertainty for users that are interested in measuring the global solar irradiance with smaller uncertainties in the full zenith angle range from 0° to 90°.

In this report we describe a procedure for a Solar Position Algorithm (SPA) to calculate the solar zenith and azimuth angle with uncertainties equal to  $\pm 0.0003^\circ$  in the period from the year -2000 to 6000. Figure 1 shows that the uncertainty of the reference global solar irradiance, resulting from  $\pm 0.0003^\circ$  in calculating the solar zenith angle in the range from  $0^\circ$  to  $90^\circ$  is negligible. The procedure is adopted from *The Astronomical Algorithms* [3], which is based on the Variations Sèculaires des Orbites Planètaires Theory (VSOP87) that was developed by P. Bretagnon in 1982 then modified in 1987 by Bretagnon and Francou [3]. In this report, we summarize the complex algorithm elements scattered throughout the book to calculate the solar position, and introduce some modification to the algorithm to accommodate solar radiation applications. For example, in *The Astronomical Algorithms* [3], the azimuth angle is measured westward from south, but for solar radiation applications, it is measured eastward from north. Also, the observer's geographical longitude is considered positive west, or negative east from Greenwich, while for solar radiation applications, it is considered negative west, or positive east from Greenwich.

We start this report by:

- Describing the time scales because of the importance of using the correct time in the SPA
- Providing a step by step procedure to calculate the solar position and the solar incidence angle for an arbitrary surface orientation using the methods described in *An Introduction*

- to Solar Radiation [4]
- Evaluating the SPA against the *Astronomical Almanac* (AA) data for the years 1994, 1995, 1996, and 2004.

Because of the complexity of the algorithm we included some examples, in the Appendix, to give the users confidence in their step by step calculations. We also included in the Appendix an explanation of how to calculate the equation of time, sun transit (solar noon), sunrise, sunset, and how to change the Julian Day to a Calendar Date. We also included a C source code with header file, for all the calculations in this report (except for the Julian Day to Calendar Date conversion). The users can incorporate this module into their own code by including the header file, declaring the SPA structure, filling in the required input parameters into the structure, and then call the SPA calculation function. This function will calculate all the output values and fill in the SPA structure for the user.

The users should note that this report is used to calculate the solar position for solar radiation applications only, and that it is purely mathematical and not meant to teach astronomy or to describe the Earth rotation. For more description about the astronomical nomenclature that is used through out the report, the user is encouraged to review the definitions in the *Astronomical Almanacs*, or other astronomical reference.

#### 2. Time Scale

The following are the internationally recognized time scales:

- The Universal Time (UT), or Greenwich civil time, is based on the Earth's rotation and counted from 0-hour at midnight; the unit is mean solar day [3]. UT is the time used to calculate the solar position in the described algorithm. It is sometimes referred to as UT1.
- The International Atomic Time (TAI) is the duration of the System International Second (SI-second) and based on a large number of atomic clocks [5].
- The Coordinated Universal Time (UTC) is the bases of most radio time signals and the legal time systems. It is kept to within 0.9 seconds of UT1 (UT) by introducing one second steps to its value (leap second); to date the steps are always positive.
- The Terrestrial Dynamical or Terrestrial Time (TDT or TT) is the time scale of ephemerides for observations from the Earth surface.

The following equations describe the relationship between the above time scales (in seconds):

$$TT = TAI + 32.184$$
 (1)

$$UT = TT - \Delta T \quad , \tag{2}$$

where  $\Delta T$  is the difference between the Earth rotation time and the Terrestrial Time (TT). It is derived from observation only and reported yearly in the *Astronomical Almanac* [5].

$$UT = UT1 = UTC + \Delta UT1 \quad . \tag{3}$$

where  $\Delta$ UT1 is a fraction of a second, positive or negative value, that is added to the UTC to adjust for the Earth irregular rotational rate. It is derived from observation, but predicted values are transmitted in code in some time signals, e.g. weekly by the U.S. Naval Observatory (USNO) [6].

#### 3. Procedure

#### 3.1. Calculate the Julian and Julian Ephemeris Day, Century, and Millennium:

The Julian date starts on January 1, in the year - 4712 at 12:00:00 UT. The Julian Day (JD) is calculated using UT and the Julian Ephemeris Day (JDE) is calculated using TT. In the following steps, note that there is a 10-day gap between the Julian and Gregorian calendar where the Julian calendar ends on October 4, 1582 (JD = 2299160), and after 10-days the Gregorian calendar starts on October 15, 1582.

#### 3.1.1 Calculate the Julian Day (JD),

$$JD = INT(365.25*(Y + 4716)) + INT(30.6001*(M+1)) + D + B - 1524.5 ,$$
(4)

where,

- INT is the Integer of the calculated terms (e.g. 8.7 = 8, 8.2 = 8, and -8.7 = -8..etc.).
- *Y* is the year (e.g. 2001, 2002, ..etc.).
- M is the month of the year (e.g. 1 for January, ..etc.). Note that if M > 2, then Y and M are not changed, but if M = 1 or 2, then Y = Y 1 and M = M + 12.
- D is the day of the month with decimal time (e.g. for the second day of the month at 12:30:30 UT, D = 2.521180556).
- B is equal to 0, for the Julian calendar {i.e. by using B = 0 in Equation 4, JD < 2299160}, and equal to (2 A + INT (A/4)) for the Gregorian calendar {i.e. by using B = 0 in Equation 4, JD > 2299160}, where A = INT(Y/100).

For users who wish to use their local time instead of UT, change the time zone to a fraction of a day (by dividing it by 24), then subtract the result from JD. Note that the fraction is subtracted from JD calculated before the test for  $B \le 2299160$  to maintain the Julian and Gregorian periods.

Table A4.1 shows examples to test any implemented program used to calculate the JD.

3.1.2. Calculate the Julian Ephemeris Day (*JDE*),

$$JDE = JD + \frac{\Delta T}{86400} \quad . \tag{5}$$

3.1.3. Calculate the Julian century (*JC*) and the Julian Ephemeris Century (*JCE*) for the 2000 standard epoch,

$$JC = \frac{JD - 2451545}{36525} \quad , \tag{6}$$

$$JCE = \frac{JDE - 2451545}{36525} \quad . \tag{7}$$

3.1.4. Calculate the Julian Ephemeris Millennium (JME) for the 2000 standard epoch,

$$JME = \frac{JCE}{10} \quad . \tag{8}$$

## 3.2. Calculate the Earth heliocentric longitude, latitude, and radius vector (*L*, *B*, and *R*):

"Heliocentric" means that the Earth position is calculated with respect to the center of the sun.

3.2.1. For each row of Table A4.2, calculate the term *L0*<sub>i</sub> (in radians),

$$L0_i = A_i * \cos(B_i + C_i * JME) , \qquad (9)$$

where,

- i is the i<sup>th</sup> row for the term L0 in Table A4.2.
- $A_i$ ,  $B_i$ , and  $C_i$  are the values in the i<sup>th</sup> row and A, B, and C columns in Table A4.2, for the term L0 (in radians).
- 3.2.2. Calculate the term *L0* (in radians),

$$L0 = \sum_{i=0}^{n} L0_{i} \quad , \tag{10}$$

where n is the number of rows for the term L0 in Table A4.2.

3.2.3. Calculate the terms *L*1, *L*2, *L*3, *L*4, and *L*5 by using Equations 9 and 10 and changing the 0 to 1, 2, 3, 4, and 5, and by using their corresponding values in

columns A, B, and C in Table A4.2 (in radians).

3.2.4. Calculate the Earth heliocentric longitude, *L* (in radians),

$$L = \frac{L0 + L1 * JME + L2 * JME^{2} + L3 * JME^{3} + L4 * JME^{4} + L5 * JME^{5}}{10^{8}} \quad . \tag{11}$$

3.2.5. Calculate *L* in degrees,

$$L (in Degrees) = \frac{L (in Radians)*180}{\pi} , \qquad (12)$$

where  $\pi$  is approximately equal to 3.1415926535898.

- 3.2.6. Limit L to the range from 0° to 360°. That can be accomplished by dividing L by 360 and recording the decimal fraction of the division as F. If L is positive, then the limited L = 360 \* F. If L is negative, then the limited L = 360 360 \* F.
- 3.2.7. Calculate the Earth heliocentric latitude, *B* (in degrees), by using Table A4.2 and steps 3.2.1 through 3.2.5 and by replacing all the *L*s by *B*s in all equations. Note that there are no *B2* through *B5*, consequently, replace them by zero in steps 3.2.3 and 3.2.4.
- 3.2.8. Calculate the Earth radius vector, *R* (in Astronomical Units, AU), by repeating step 3.2.7 and by replacing all *L*s by *R*s in all equations. Note that there is no *R5*, consequently, replace it by zero in steps 3.2.3 and 3.2.4.
- 3.3. Calculate the geocentric longitude and latitude ( $\Theta$  and  $\beta$ ):

"Geocentric" means that the sun position is calculated with respect to the Earth center.

3.3.1. Calculate the geocentric longitude,  $\Theta$  (in degrees),

$$\Theta = L + 180 \quad . \tag{13}$$

- 3.3.2. Limit  $\Theta$  to the range from 0° to 360° as described in step 3.2.6.
- 3.3.3. Calculate the geocentric latitude,  $\beta$  (in degrees),

$$\beta = -B \quad . \tag{14}$$

#### 3.4. Calculate the nutation in longitude and obliquity ( $\Delta \psi$ and $\Delta \varepsilon$ ):

3.4.1. Calculate the mean elongation of the moon from the sun,  $X_0$  (in degrees),

$$X_0 = 297.85036 + 445267.111480 * JCE -$$

$$0.0019142 * JCE^2 + \frac{JCE^3}{189474} .$$
(15)

3.4.2. Calculate the mean anomaly of the sun (Earth),  $X_1$  (in degrees),

$$X_{1} = 357.52772 + 35999.050340 * JCE -$$

$$0.0001603 * JCE^{2} - \frac{JCE^{3}}{300000}$$
 (16)

3.4.3. Calculate the mean anomaly of the moon,  $X_2$  (in degrees),

$$X_{2} = 134.96298 + 477198.867398 * JCE +$$

$$0.0086972 * JCE^{2} + \frac{JCE^{3}}{56250} .$$

$$(17)$$

3.4.4. Calculate the moon's argument of latitude,  $X_3$  (in degrees),

$$X_3 = 93.27191 + 483202.017538 * JCE -$$

$$0.0036825 * JCE^2 + \frac{JCE^3}{327270} .$$
(18)

3.4.5. Calculate the longitude of the ascending node of the moon's mean orbit on the ecliptic, measured from the mean equinox of the date,  $X_4$  (in degrees),

$$X_4 = 125.04452 - 1934.136261 * JCE +$$

$$0.0020708 * JCE^2 + \frac{JCE^3}{450000}$$
(19)

3.4.6. For each row in Table A4.3, calculate the terms  $\Delta \psi_i$  and  $\Delta \varepsilon_i$  (in 0.0001of arc seconds),

$$\Delta \psi_i = (a_i + b_i * JCE) * \sin(\sum_{j=0}^4 X_j * Y_{i,j}) , \qquad (20)$$

$$\Delta \varepsilon_i = (c_i + d_i * JCE) * \cos(\sum_{j=0}^4 X_j * Y_{i,j}) , \qquad (21)$$

where,

- $a_i$ ,  $b_i$ ,  $c_i$ , and  $d_i$  are the values listed in the i<sup>th</sup> row and columns a, b, c, and d in
- $X_j$  is the j<sup>th</sup> X calculated by using Equations 15 through 19.  $Y_{i,j}$  is the value listed in i<sup>th</sup> row and j<sup>th</sup> Y column in Table A4.3.
- 3.4.7. Calculate the nutation in longitude,  $\Delta \psi$  (in degrees),

$$\Delta \psi = \frac{\sum_{i=0}^{n} \Delta \psi_{i}}{36000000} \quad , \tag{22}$$

where n is the number of rows in Table A4.3 (n equals 63 rows in the table).

3.4.8. Calculate the nutation in obliquity,  $\Delta \varepsilon$  (in degrees),

$$\Delta \varepsilon = \frac{\sum_{i=0}^{n} \Delta \varepsilon_i}{36000000} \quad . \tag{23}$$

- 3.5. Calculate the true obliquity of the ecliptic,  $\varepsilon$  (in degrees):
- 3.5.1. Calculate the mean obliquity of the ecliptic,  $\mathcal{E}_0$  (in arc seconds),

$$\varepsilon_0 = 84381.448 - 4680.93U - 1.55U^2 + 1999.25U^3 - 51.38U^4 - 249.67U^5 - 39.05U^6 + 7.12U^7 + 27.87U^8 + 5.79U^9 + 2.45U^{10} ,$$
(24)

where U = JME/10.

3.5.2. Calculate the true obliquity of the ecliptic,  $\varepsilon$  (in degrees),

$$\varepsilon = \frac{\varepsilon_0}{3600} + \Delta \varepsilon \quad . \tag{25}$$

3.6. Calculate the aberration correction,  $\Delta \tau$  (in degrees):

$$\Delta \tau = -\frac{20.4898}{3600 * R} \quad . \tag{26}$$

3.7. Calculate the apparent sun longitude,  $\lambda$  (in degrees):

$$\lambda = \Theta + \Delta \psi + \Delta \tau \quad . \tag{27}$$

- 3.8. Calculate the apparent sidereal time at Greenwich at any given time,  $\nu$  (in degrees):
- 3.8.1. Calculate the mean sidereal time at Greenwich,  $V_0$  (in degrees),

$$\nu_0 = 280.46061837 + 360.98564736629 * (JD - 2451545) + 0.000387933 * JC^2 - \frac{JC^3}{38710000}$$
 (28)

- 3.8.2. Limit  $v_0$  to the range from  $0^{\circ}$  to  $360^{\circ}$  as described in step 3.2.6.
- 3.8.3. Calculate the apparent sidereal time at Greenwich, V (in degrees),

$$v = v_0 + \Delta \psi * \cos(\varepsilon) \quad . \tag{29}$$

- 3.9. Calculate the geocentric sun right ascension,  $\alpha$  (in degrees):
- 3.9.1. Calculate the sun right ascension,  $\alpha$  (in radians),

$$\alpha = Arc \tan 2 \left( \frac{\sin \lambda * \cos \varepsilon - \tan \beta * \sin \varepsilon}{\cos \lambda} \right) , \qquad (30)$$

where Arctan2 is an arctangent function that is applied to the numerator and the denominator (instead of the actual division) to maintain the correct quadrant of the  $\alpha$  where  $\alpha$  is in the rage from  $-\pi$  to  $\pi$ .

- 3.9.2. Calculate  $\alpha$  in degrees using Equation 12, then limit it to the range from 0° to 360° using the technique described in step 3.2.6.
- 3.10. Calculate the geocentric sun declination,  $\delta$  (in degrees):

$$\delta = Arc\sin(\sin\beta^*\cos\varepsilon + \cos\beta^*\sin\varepsilon^*\sin\lambda) \quad , \tag{31}$$

where  $\delta$  is positive or negative if the sun is north or south of the celestial equator, respectively. Then change  $\delta$  to degrees using Equation 12.

#### 3.11. Calculate the observer local hour angle, *H* (in degrees):

$$H = v + \sigma - \alpha \quad , \tag{32}$$

Where  $\sigma$  is the observer geographical longitude, positive or negative for east or west of Greenwich, respectively.

Limit H to the range from  $0^{\circ}$  to  $360^{\circ}$  using step 3.2.6 and note that it is measured westward from south in this algorithm.

#### 3.12. Calculate the topocentric sun right ascension $\alpha'$ (in degrees):

"Topocentric" means that the sun position is calculated with respect to the observer local position at the Earth surface.

#### 3.12.1. Calculate the equatorial horizontal parallax of the sun, $\xi$ (in degrees),

$$\xi = \frac{8.794}{3600 * R} \quad , \tag{33}$$

where R is calculated in step 3.2.8.

#### 3.12.2. Calculate the term u (in radians),

$$u = Arc \tan (0.99664719 * \tan \varphi)$$
 , (34)

where  $\varphi$  is the observer geographical latitude, positive or negative if north or south of the equator, respectively. Note that the 0.99664719 number equals (1 - f), where f is the Earth's flattening.

#### 3.12.3. Calculate the term x,

$$x = \cos u + \frac{E}{6378140} * \cos \varphi \quad , \tag{35}$$

where E is the observer elevation (in meters). Note that x equals  $\rho * \cos \varphi$  where  $\rho$  is the observer's distance to the center of Earth, and  $\varphi$  is the observer's geocentric latitude.

3.12.4. Calculate the term y,

$$y = 0.99664719 * \sin u + \frac{E}{6378140} * \sin \varphi \quad , \tag{36}$$

note that y equals  $\rho * \sin \varphi'$ ,

3.12.5. Calculate the parallax in the sun right ascension,  $\Delta \alpha$  (in degrees),

$$\Delta \alpha = Arc \tan 2 \left( \frac{-x * \sin \xi * \sin H}{\cos \delta - x * \sin \xi * \cos H} \right) \quad . \tag{37}$$

Then change  $\Delta \alpha$  to degrees using Equation 12.

3.12.6. Calculate the topocentric sun right ascension  $\alpha'$  (in degrees),

$$\alpha' = \alpha + \Delta \alpha$$
 (38)

3.12.7. Calculate the topocentric sun declination,  $\delta$  (in degrees),

$$\delta' = Arc \tan 2 \left( \frac{(\sin \delta - y * \sin \xi) * \cos \Delta \alpha}{\cos \delta - x * \sin \xi * \cos H} \right) . \tag{39}$$

3.13. Calculate the topocentric local hour angle, H' (in degrees),

$$H' = H - \Delta \alpha \quad . \tag{40}$$

- 3.14. Calculate the topocentric zenith angle,  $\theta$  (in degrees):
- 3.14.1. Calculate the topocentric elevation angle without atmospheric refraction correction, e<sub>0</sub> (in degrees),

$$e_0 = Arc\sin(\sin\varphi^*\sin\delta' + \cos\varphi^*\cos\delta'^*\cos H') \quad . \tag{41}$$

Then change  $e_0$  to degrees using Equation 12.

3.14.2. Calculate the atmospheric refraction correction,  $\Delta e$  (in degrees),

$$\Delta e = \frac{P}{1010} * \frac{283}{273 + T} * \frac{1.02}{60 * \tan (e_0 + \frac{10.3}{e_0 + 5.11})},$$
(42)

where,

- P is the annual average local pressure (in millibars).
- T is the annual average local temperature (in  $^{\circ}$ C).
- $e_0$  is in degrees. Calculate the tangent argument in degrees, then convert to radians if required by calculator or computer.
- 3.14.3. Calculate the topocentric elevation angle, e (in degrees),

$$e = e_0 + \Delta e \quad . \tag{43}$$

3.14.4. Calculate the topocentric zenith angle,  $\theta$  (in degrees),

$$\theta = 90 - e \quad . \tag{44}$$

- 3.15. Calculate the topocentric azimuth angle,  $\phi$  (in degrees):
- 3.15.1. Calculate the topocentric astronomers azimuth angle,  $\Gamma$  (in degrees),

$$\Gamma = Arc \tan 2 \left( \frac{\sin H'}{\cos H'^* \sin \varphi - \tan \delta'^* \cos \varphi} \right) , \qquad (45)$$

Change  $\Gamma$  to degrees using Equation 12, then limit it to the range from  $0^{\circ}$  to  $360^{\circ}$  using step 3.2.6. Note that  $\Gamma$  is measured <u>westward from south</u>.

3.15.2. Calculate the topocentric azimuth angle,  $\phi$  for navigators and solar radiation users (in degrees),

$$\Phi = \Gamma + 180 \quad , \tag{46}$$

Limit  $\Phi$  to the range from  $0^{\circ}$  to  $360^{\circ}$  using step 3.2.6. Note that  $\Phi$  is measured <u>eastward</u> from north.

## 3.16. Calculate the incidence angle for a surface oriented in any direction, *I* (in degrees):

$$I = Arc\cos(\cos\theta * \cos\omega + \sin\omega * \sin\theta * \cos(\Gamma - \gamma)) , \qquad (47)$$

where,

- $\omega$  is the slope of the surface measured from the horizontal plane.
- $\gamma$  is the surface azimuth rotation angle, measured from south to the projection of the surface normal on the horizontal plane, positive or negative if oriented east or west from south, respectively.

#### 4. SPA Evaluation and Conclusion

Because the solar zenith, azimuth, and incidence angles are not reported in the *Astronomical Almanac* (AA), the following sun parameters are used for the evaluation: The main parameters (ecliptic longitude and latitude for the mean Equinox of date, apparent right ascension, apparent declination), and the correcting parameters (nutation in longitude, nutation in obliquity, obliquity of ecliptic, and true geometric distance). Exact trigonometric functions are used with the AA reported sun parameters to calculate the solar zenith and azimuth angles, therefore it is adequate to evaluate the SPA uncertainty using these parameters. To evaluate the uncertainty of the SPA, we chose the second day of each month, for each of the years 1994, 1995, 1996, and 2004, at 0-hour Terrestrial Time (TT). Figures 2 shows that the maximum difference between the AA and SPA main parameters is -0.00015°. Figure 3 shows that the maximum difference between the AA and SPA for calculating the zenith or azimuth angle is 0.00003° and 0.00008°, respectively. This implies that the SPA is well within the stated uncertainty of ± 0.0003°.




Figure 1. Uncertainty of cosine the solar zenith angle resulting from 0.01° and 0.0003° uncertainty in the angle calculation




Figure 2. Difference between the Almanac and SPA for the ecliptic longitude, ecliptic latitude, apparent right ascension, and apparent declination on the second day of each month at 0-TT for the years 1994, 1995, 1996, and 2004




Figure 3. Difference between the Almanac and SPA for the solar zenith and azimuth angles on the second day of each month at 0-TT for the years 1994, 1995, 1996, 2004

#### References

- 1. Blanco-Muriel, M., et al. "Computing the Solar Vector". *Solar Energy*. Vol. 70, No. 5, 2001; pp. 431-441, 2001, Great Britain.
- 2. Michalsky, J. J. "The Astronomical Almanac's Algorithm for Approximate Solar Position (1950-2050)". *Solar Energy*. Vol. 40, No. 3, 1988; pp. 227-235, USA.
- 3. Meeus, J. "Astronomical Algorithms". Second edition 1998, Willmann-Bell, Inc., Richmond, Virginia, USA.
- 4. Iqbal, M. "An Introduction to Solar Radiation". New York: 1983; pp. 23-25.
- 5. The Astronomical Almanac. Norwich:2004.
- 6. The U.S. Naval Observatory. Washington, DC, <a href="http://www.usno.navy.mil/">http://www.usno.navy.mil/</a>.

#### **Appendix**

Note that some of the symbols used in the appendix are independent from the symbols used in the main report.

#### A.1. Equation of Time

The Equation of Time, E, is the difference between solar apparent and mean time. Use the following equation to calculate E (in degrees),

$$E = M - 0.0057183 - \alpha + \Delta \psi * \cos \varepsilon , \qquad (A1)$$

where,

- M is the sun's mean longitude (in degrees),

$$M = 280.4664567 + 360007.6982779 * JME + 0.03032028 * JME2 + 
\frac{JME3}{49931} - \frac{JME4}{15300} - \frac{JME5}{2000000} ,$$
(A2)

where JME is the Julian Ephemeris Millennium calculated from Equation 8, and M is limited to the range from  $0^{\circ}$  to  $360^{\circ}$  using step 3.2.6.

- $\alpha$  is the geocentric right ascention, from Equation 30 (in degrees).
- $\Delta \psi$  is the nutation in longitude, from Equation 22 (in degrees).
- $\varepsilon$  is the obliquity of the ecliptic, from Equation 25 (in degrees).

Multiply *E* by 4 to change its unit from degrees to minutes of time. Limit E if its absolute value is greater than 20 minutes, by adding or subtracting 1440.

#### A.2. Sunrise, Sun Transit, and Sunset

The value of  $0.5667^{\circ}$  is typically adopted for the atmospheric refraction at sunrise and sunset times. Thus for the sun radius of  $0.26667^{\circ}$ , the value  $-0.8333^{\circ}$  of sun elevation ( $h'_{\theta}$ ) is chosen to calculate the times of sunrise and sunset. On the other hand, the sun transit is the time when the center of the sun reaches the local meridian.

- A.2.1. Calculate the apparent sidereal time at Greenwich at 0 UT,  $\nu$  (in degrees), using Equation 29.
- A.2.2. Calculate the geocentric right ascension and declination at 0 TT, using Equations 30 and 31, for the day before the day of interest  $(D_{-1})$ , the day of interest  $(D_0)$ ,

$$\alpha_{-1}$$
  $\delta_{-1}$ 

then the day after (D\_+1). Denote the values as  $~\alpha_0~\delta_0~$  , in degrees.

$$\alpha_{{}_{\scriptscriptstyle +1}}$$
  $\delta_{{}_{\scriptscriptstyle +}}$ 

A.2.3. Calculate the approximate sun transit time,  $m_0$ , in fraction of day,

$$m_0 = \frac{\alpha_0 - \sigma - \nu}{360} \quad , \tag{A3}$$

where  $\sigma$  is the observer geographical longitude, in degrees, positive east of Greenwich...

A.2.4. Calculate the local hour angle corresponding to the sun elevation equals -  $0.8333^{\circ}$ ,  $H_0$ ,

$$H_0 = Arc\cos\left(\frac{\sin h'_0 - \sin \varphi * \sin \delta_0}{\cos \varphi * \cos \delta_0}\right) , \qquad (A4)$$

where,

 $-h'_{0}$  equals  $-0.8333^{\circ}$ .

-  $\varphi$  is the observer geographical latitude, in degrees, positive north of the equator.

Note that if the argument of the Arccosine is not in the range from -1 to 1, it means that the sun is always above or below the horizon for that day.

Change  $H_0$  to degrees using Equation 12, then limit it to the range from  $0^{\circ}$  to  $180^{\circ}$  using step 3.2.6 and replacing 360 by 180.

A.2.5. Calculate the approximate sunrise time,  $m_1$ , in fraction of day,

$$m_1 = m_0 - \frac{H_0}{360} \quad .$$
(A5)

A.2.6. Calculate the approximate sunset time,  $m_2$ , in fraction of day,

$$m_2 = m_0 + \frac{H_0}{360} \quad . \tag{A6}$$

A.2.7. Limit the values of  $m_0$ ,  $m_1$ , and  $m_2$  to a value between 0 and 1 fraction of day using step 3.2.6 and replacing 360 by 1.

A.2.8. Calculate the sidereal time at Greenwich, in degrees, for the sun transit, sunrise, and sunset,  $V_i$ ,

$$v_i = v + 360.985647 * m_i \quad , \tag{A7}$$

where i equals 0, 1, and 2 for sun transit, sunrise, and sunset, respectively.

A.2.9. Calculate the terms  $n_i$ ,

$$n_i = m_i + \frac{\Delta T}{86400} \quad , \tag{A8}$$

where  $\Delta T = TT-UT$ .

A.2.10. Calculate the values  $\alpha_i$  and  $\delta_i$ , in degrees, where i equals 0, 1, and 2,

where,

$$\alpha'_{i} = \alpha_{0} + \frac{n_{i}(a+b+c*n_{i})}{2}$$
 , (A9)

and,

$$\delta'_{i} = \delta_{0} + \frac{n_{i}(a'+b'+c'*n_{i})}{2}$$
, (A10)

where,

- a and a' equal  $(\alpha_0 - \alpha_{-1})$  and  $(\delta_0 - \delta_{-1})$ , respectively.

- *b* and *b*' equal  $(\alpha_{+1} - \alpha_0)$  and  $(\delta_{+1} - \delta_0)$ , respectively.

- c and c' equal (b - a) and (b' - a'), respectively.

If the absolute value of a, a, b, or b is greater than 2, then limit its value between 0 and 1 as shown in step A.2.7.

A.2.11. Calculate the local hour angle for the sun transit, sunrise, and sunset,  $H'_{i}$  (in degrees),

$$H'_{i} = v_{i} + \sigma - \alpha'_{i} \quad . \tag{A11}$$

 $H'_i$  in this case is measured as positive westward from the meridian, and negative eastward from the meridian. Thus limit  $H'_i$  between -180° and 180°. To preserve the quadrant sign of  $H'_i$  limit it to  $\pm$  360° first, then if  $H'_i$  is less than or equal -180°, then add 360° to force it's value to be between 0° and 180°. And if  $H'_i$  is greater than or equal 180°, then add -360° to force it's value to be between 0° and -180°.

A.2.12. Calculate the sun altitude for the sun transit, sunrise, and sunset,  $h_i$  (in degrees),

$$h_i = Arc\sin\left(\sin\varphi * \sin\delta'_i + \cos\varphi * \cos\delta'_i * \cos H'_i\right) . \tag{A12}$$

A.2.13. Calculate the sun transit, *T* (in fraction of day),

$$T = m_0 - \frac{H'_0}{360} \quad . \tag{A13}$$

A.2.14. Calculate the sunrise, R (in fraction of day),

$$R = m_1 + \frac{h_1 - h'_0}{360 * \cos \delta'_1 * \cos \varphi * \sin H'_1}$$
 (A14)

A.2.15. Calculate the sunset, S (in fraction of day), by using Equation A14 and replacing R by S, and replacing the suffix number 1 by 2.

The fraction of day value is changed to UT by multiplying the value by 24.

To evaluate the uncertainty of the SPA, we chose the second day of each month, for each of the years 1994, 1995, 1996, and 2004, at 0-hour Terrestrial Time (TT). Figure A2.1 shows that the maximum difference between the AA and SPA sun transit time is -0.23 seconds.

Because the sunrise and sunset are recorded in the AA to a one minute resolution, we compared the SPA calculations at only three data points at Greenwich meridian at 0-UT. The comparison result in Table A2.1 shows that the maximum difference between AA and SPA is 15.4 seconds (0.26 minute), which is well within the AA resolution of one minute.

Note that UT can be changed to local time by adding the time zone as a fraction of a day (time zone is divided by 24), and limiting the result to the range from 0 to 1.

Table A2.1. The AA and SPA Results for Sunrise and Sunset at Greenwich Meridian at 0-UT

D 4	Observer	Sunrise		Sun	set
Date	Latitude	AA	SPA	AA	SPA
January 2, 1994	35°	7:08	7:08:12.8	17:00	16:59:55.9
July 5, 1996	-35°	7:08	7:08:15.4	17:00	17:01:04.5
December 4, 2004	-35°	4:39	4:38:57.1	19:02	19:02:2.5




Figure A2.1. Difference between the Almanac and SPA for the Ephemeris Transit on the second day of each month at 0-TT for the years 1994, 1995, 1996, 2004

#### A.3. Calculation of Calendar Date from Julian Day

- A.3.1. Add 0.5 to the Julian Day (*JD*), then record the integer of the result as *Z*, and the fraction decimal as *F*.
- A.3.2. If Z is less than 2299161, then record A equals Z. Else, calculate the term B,

$$B = INT \left( \frac{Z - 1867216.25}{36524.25} \right) \quad , \tag{A15}$$

Then calculate the term A,

$$A = Z + 1 + B - INT\left(\frac{B}{4}\right) \quad . \tag{A16}$$

A.3.3. Calculate the term C,

$$C = A + 1524$$
 . (A17)

A.3.4. Calculate the term D,

$$D = INT \left( \frac{C - 122.1}{365.25} \right) \quad . \tag{A18}$$

A.3.5. Calculate the term G,

$$G = INT (365.25 * D)$$
 (A19)

A.3.6. Calculate the term I,

$$I = INT \left( \frac{C - G}{306001} \right) \quad . \tag{A20}$$

A.3.7. Calculate the day number of the month with decimals, d,

$$d = C - G - INT (30.6001*I) + F . (A21)$$

A.3.8. Calculate the month number, *m*,

$$m = I - 1$$
,  $IF I < 14$  ,  
 $m = I - 13$ ,  $IF I \ge 14$  . (A22)

#### A.3.9. Calculate the year, y,

$$y = D - 4716$$
,  $IF m > 2$ ,  
 $y = D - 4715$ ,  $IF m \le 2$ . (A23)

Note that if local time is used to calculate the JD, then the local time zone is added to the JD in step A.3.1 to calculate the local Calendar Date.

#### A.4. Tables

Table A4.1. Examples for Testing any Program to Calculate the Julian Day

Date	UT	JD	Date	UT	JD
January 1, 2000	12:00:00	2451545.0	December 31, 1600	00:00:00	2305812.5
January 1, 1999	00:00:00	2451179.5	April 10, 837	07:12:00	2026871.8
January 27, 1987	00:00:00	2446822.5	December 31, -123	00:00:00	1676496.5
June 19, 1987	12:00:00	2446966.0	January 1, -122	00:00:00	1676497.5
January 27, 1988	00:00:00	2447187.5	July 12, -1000	12:00:00	1356001.0
June 19, 1988	12:00:00	2447332.0	February 29, -1000	00:00:00	1355866.5
January 1, 1900	00:00:00	2415020.5	August 17, -1001	21:36:00	1355671.4
January 1, 1600	00:00:00	2305447.5	January 1, -4712	12:00:00	0.0

**Table A4.2. Earth Periodic Terms** 

Term	Row Number	A	В	C
L0	0	175347046	0	0
	1	3341656	4.6692568	6283.07585
	2	34894	4.6261	12566.1517
	3	3497	2.7441	5753.3849
	4	3418	2.8289	3.5231
	5	3136	3.6277	77713.7715
	6	2676	4.4181	7860.4194
	7	2343	6.1352	3930.2097
	8	1324	0.7425	11506.7698
	9	1273	2.0371	529.691
	10	1199	1.1096	1577.3435

11	990	5.233	5884.927
12	902	2.045	26.298
13	857	3.508	398.149
14	780	1.179	5223.694
15	753	2.533	5507.553
16	505	4.583	18849.228
17	492	4.205	775.523
18	357	2.92	0.067
19	317	5.849	11790.629
20	284	1.899	796.298
21	271	0.315	10977.079
22	243	0.345	5486.778
23	206	4.806	2544.314
24	205	1.869	5573.143
25	202	2.458	6069.777
26	156	0.833	213.299
27	132	3.411	2942.463
28	126	1.083	20.775
29	115	0.645	0.98
30	103	0.636	4694.003
31	102	0.976	15720.839
32	102	4.267	7.114
33	99	6.21	2146.17
34	98	0.68	155.42
35	86	5.98	161000.69
36	85	1.3	6275.96
37	85	3.67	71430.7
38	80	1.81	17260.15
39	79	3.04	12036.46
40	75	1.76	5088.63
41	74	3.5	3154.69
42	74	4.68	801.82
43	70	0.83	9437.76
44	62	3.98	8827.39

	45	61	1.82	7084.9
	46	57	2.78	6286.6
	47	56	4.39	14143.5
	48	56	3.47	6279.55
	49	52	0.19	12139.55
	50	52	1.33	1748.02
	51	51	0.28	5856.48
	52	49	0.49	1194.45
	53	41	5.37	8429.24
	54	41	2.4	19651.05
	55	39	6.17	10447.39
	56	37	6.04	10213.29
	57	37	2.57	1059.38
	58	36	1.71	2352.87
	59	36	1.78	6812.77
	60	33	0.59	17789.85
	61	30	0.44	83996.85
	62	30	2.74	1349.87
	63	25	3.16	4690.48
L1	0	628331966747	0	0
	1	206059	2.678235	6283.07585
	2	4303	2.6351	12566.1517
	3	425	1.59	3.523
	4	119	5.796	26.298
	5	109	2.966	1577.344
	6	93	2.59	18849.23
	7	72	1.14	529.69
	8	68	1.87	398.15
	9	67	4.41	5507.55
	10	59	2.89	5223.69
	11	56	2.17	155.42
	12	45	0.4	796.3
	13	36	0.47	775.52
	14	29	2.65	7.11

	15	21	5.34	0.98
	16	19	1.85	5486.78
	17	19	4.97	213.3
	18	17	2.99	6275.96
	19	16	0.03	2544.31
	20	16	1.43	2146.17
	21	15	1.21	10977.08
	22	12	2.83	1748.02
	23	12	3.26	5088.63
	24	12	5.27	1194.45
	25	12	2.08	4694
	26	11	0.77	553.57
	27	10	1.3	6286.6
	28	10	4.24	1349.87
	29	9	2.7	242.73
	30	9	5.64	951.72
	31	8	5.3	2352.87
	32	6	2.65	9437.76
	33	6	4.67	4690.48
L2	0	52919	0	0
	1	8720	1.0721	6283.0758
	2	309	0.867	12566.152
	3	27	0.05	3.52
	4	16	5.19	26.3
	5	16	3.68	155.42
	6	10	0.76	18849.23
	7	9	2.06	77713.77
	8	7	0.83	775.52
	9	5	4.66	1577.34
	10	4	1.03	7.11
	11	4	3.44	5573.14
	12	3	5.14	796.3
	13	3	6.05	5507.55
	14	3	1.19	242.73

	15	3	6.12	529.69
	16	3	0.31	398.15
	17	3	2.28	553.57
	18	2	4.38	5223.69
	19	2	3.75	0.98
L3	0	289	5.844	6283.076
	1	35	0	0
	2	17	5.49	12566.15
	3	3	5.2	155.42
	4	1	4.72	3.52
	5	1	5.3	18849.23
	6	1	5.97	242.73
L4	0	114	3.142	0
	1	8	4.13	6283.08
	2	1	3.84	12566.15
L5	0	1	3.14	0
B0	0	280	3.199	84334.662
	1	102	5.422	5507.553
	2	80	3.88	5223.69
	3	44	3.7	2352.87
	4	32	4	1577.34
B1	0	9	3.9	5507.55
	1	6	1.73	5223.69
R0	0	100013989	0	0
	1	1670700	3.0984635	6283.07585
	2	13956	3.05525	12566.1517
	3	3084	5.1985	77713.7715
	4	1628	1.1739	5753.3849
	5	1576	2.8469	7860.4194
	6	925	5.453	11506.77
	7	542	4.564	3930.21
	8	472	3.661	5884.927
	9	346	0.964	5507.553
	10	329	5.9	5223.694

	4	31	2.84	5507.55
	3	32	1.02	18849.23
	2	702	3.142	0
	1	1721	1.0644	12566.1517
R1	0	103019	1.10749	6283.07585
	39	26	4.59	10447.39
	38	28	1.9	6279.55
	37	28	1.21	6286.6
	36	32	1.78	398.15
	35	32	0.18	5088.63
	34	33	0.24	7084.9
	33	35	1.84	2942.46
	32	36	1.67	12036.46
	31	37	4.9	12139.55
	30	37	0.83	19651.05
	29	38	2.39	8827.39
	28	39	5.36	4694
	27	43	6.01	6275.96
	26	45	5.54	9437.76
	25	47	2.58	775.52
	24	49	3.25	2544.31
	23	56	5.24	71430.7
_	22	57	2.01	83996.85
	21	63	0.92	529.69
	20	65	0.27	17260.15
	19	86	1.27	161000.69
	18	86	5.69	15720.84
	17	98	0.89	6069.78
	16	110	5.055	5486.778
	15	175	3.012	18849.228
	14	186	5.022	10977.079
	13	212	5.847	1577.344
	12	243	4.273	11790.629

	5	25	1.32	5223.69
	6	18	1.42	1577.34
	7	10	5.91	10977.08
	8	9	1.42	6275.96
	9	9	0.27	5486.78
R2	0	4359	5.7846	6283.0758
	1	124	5.579	12566.152
	2	12	3.14	0
	3	9	3.63	77713.77
	4	6	1.87	5573.14
	5	3	5.47	18849.23
R3	0	145	4.273	6283.076
	1	7	3.92	12566.15
R4	0	4	2.56	6283.08

Table A4.3. Periodic Terms for the Nutation in Longitude and Obliquity

Coefficients for Sin terms					Coefficien	ts for $\Delta\psi$	Coefficier	nts for $arDelta\epsilon$
Y0	Y1	Y2	<b>Y3</b>	Y4	a	b	c	d
0	0	0	0	1	-171996	-174.2	92025	8.9
-2	0	0	2	2	-13187	-1.6	5736	-3.1
0	0	0	2	2	-2274	-0.2	977	-0.5
0	0	0	0	2	2062	0.2	-895	0.5
0	1	0	0	0	1426	-3.4	54	-0.1
0	0	1	0	0	712	0.1	-7	
-2	1	0	2	2	-517	1.2	224	-0.6
0	0	0	2	1	-386	-0.4	200	
0	0	1	2	2	-301		129	-0.1
-2	-1	0	2	2	217	-0.5	-95	0.3
-2	0	1	0	0	-158			
-2	0	0	2	1	129	0.1	-70	
0	0	-1	2	2	123		-53	
2	0	0	0	0	63			
0	0	1	0	1	63	0.1	-33	
2	0	-1	2	2	-59		26	_

0	0	-1	0	1	-58	-0.1	32	
0	0	1	2	1	-51		27	
-2	0	2	0	0	48			
0	0	-2	2	1	46		-24	
2	0	0	2	2	-38		16	
0	0	2	2	2	-31		13	
0	0	2	0	0	29			
-2	0	1	2	2	29		-12	
0	0	0	2	0	26			
-2	0	0	2	0	-22			
0	0	-1	2	1	21		-10	
0	2	0	0	0	17	-0.1		
2	0	-1	0	1	16		-8	
-2	2	0	2	2	-16	0.1	7	
0	1	0	0	1	-15		9	
-2	0	1	0	1	-13		7	
0	-1	0	0	1	-12		6	
0	0	2	-2	0	11			
2	0	-1	2	1	-10		5	
2	0	1	2	2	-8		3	
0	1	0	2	2	7		-3	
-2	1	1	0	0	-7			
0	-1	0	2	2	-7		3	
2	0	0	2	1	-7		3	
2	0	1	0	0	6			
-2	0	2	2	2	6		-3	
-2	0	1	2	1	6		-3	
2	0	-2	0	1	-6		3	
2	0	0	0	1	-6		3	
0	-1	1	0	0	5			
-2	-1	0	2	1	-5		3	
-2	0	0	0	1	-5		3	
0	0	2	2	1	-5		3	
-2	0	2	0	1	4			

-2	1	0	2	1	4		
0	0	1	-2	0	4		
-1	0	1	0	0	-4		
-2	1	0	0	0	-4		
1	0	0	0	0	-4		
0	0	1	2	0	3		
0	0	-2	2	2	-3		
-1	-1	1	0	0	-3		
0	1	1	0	0	-3		
0	-1	1	2	2	-3		
2	-1	-1	2	2	-3		
0	0	3	2	2	-3		
2	-1	0	2	2	-3		

## A.5. Example

The results for the following site parameters are listed in Table A5.1:

- Date = October 17, 2003. Time = 12:30:30 Local Standard Time (LST).
- Time zone(TZ) = -7 hours. Longitude =  $-105.1786^{\circ}$ .
- Latitude = 39.742476°. Pressure = 820 mbar.
- Elevation = 1830.14 m. Temperature = 11°C.
- Surface slope =  $30^{\circ}$ . Surface azimuth rotation =  $-10^{\circ}$ .
- $\Delta T = 67$  Seconds.

LST must be changed to UT by subtracting TZ from LST, and changing the date if necessary.

**Table A5.1. Results for Example** 

JD	2452930.312847				
LO	172067561.526586	L1	628332010650.051147		
L2	61368.682493	L3	-26.902819		
L4	-121.279536	L5	-0.999999		
L	24.0182616917°				
<i>B0</i>	-176.502688	B1	3.067582		
В	-0.0001011219°				
RO	99653849.037796	R1	100378.567146		

R2	-1140.953507	R3	-141.115419			
R4	1.232361					
R	(	0.9965422974 AU				
Θ	204.0182616917°	β	0.0001011219°			
Δψ	-0.00399840°	$\Delta arepsilon$	0.00166657°			
$oldsymbol{arepsilon}$	23.440465°	λ	204.0085519281°			
α	202.22741°	δ	-9.31434°			
Н	11.105900°	H'	11.10629°			
α'	202.22704°	δ'	-9.316179°			
$\theta$	50.11162°	φ	194.34024°			
I	25.18700°	M	205.8971722516°			
E	14.641503 minutes	Transit	18:46:04.97 UT			
Sunrise	13:12:43.46 UT	Sunset	00:20:19.19 UT			

```
A.6. C Code: SPA header file (SPA.h)
            HEADER FILE for SPA.C
       Solar Position Algorithm (SPA)
                    for
         Solar Radiation Application
                May 12, 2003
    Filename: SPA.H
,,
||
||
||
    Afshin Michael Andreas
    afshin andreas@nrel.gov (303)384-6383
//
    Measurement & Instrumentation Team
     Solar Radiation Research Laboratory
11
    National Renewable Energy Laboratory
// 1617 Cole Blvd, Golden, CO 80401 //
// Usage:
     1) In calling program, include this header file,
      by adding this line to the top of file:
//
             #include "spa.h"
//
//
//
//
//
//
//
//
//
     2) In calling program, declare the SPA structure:
            spa data spa;
     3) Enter the required input values into SPA structure
       (input values listed in comments below)
     4) Call the SPA calculate function and pass the SPA structure
       (prototype is declared at the end of this header file):
            spa calculate(&spa);
    Selected output values (listed in comments below) will be
     computed and returned in the passed SPA structure. Output
     will based on function code selected from enumeration below.
    Note: A non-zero return code from spa_calculate() indicates that //
           one of the input values did not pass simple bounds tests.
//
           The valid input ranges and return error codes are also
           listed below.
#ifndef __solar_position_algorithm header
#define __solar_position_algorithm header
//enumeration for function codes to select desired final outputs from SPA
enum {
    SPA ZA,
                      //calculate zenith and azimuth
                     //calculate zenith, azimuth, and incidence
//calculate zenith, azimuth, and sun rise/transit/set values
    SPA ZA INC,
   SPA ZA RTS,
   SPA ALL,
                      //calculate all SPA output values
typedef struct
    //----INPUT VALUES-----
                        // 4-digit year,
                                             valid range: -2000 to 6000, error code: 1
   int year;
                        // 2-digit month,
// 2-digit day,
                                            valid range: 1 to 12, error code: 2
   int month;
                                                   valid range: 1 to 31, error code: 3
    int day;
                        // Observer local hour, valid range: 0 to 24, error code: 4
// Observer local minute, valid range: 0 to 59, error code: 5
// Observer local second, valid range: 0 to 59, error code: 6
   int hour;
    int minute;
    int second;
   double delta t;
                        // Difference between earth rotation time and terrestrial time
                         // It is derived from observation only and is reported in this
                         // bulletin: http://maia.usno.navy.mil/ser7/ser7.dat,
                         // where delta_t = 32.184 + (TAI-UTC) + DUT1
                         // valid range: -8000 to 8000 seconds, error code: 7
                         // Observer time zone (negative west of Greenwich)
   double timezone;
```

```
// valid range: -12 to 12 hours, error code: 8
   double longitude;
                         // Observer longitude (negative west of Greenwich)
                         // valid range: -180 to 180 degrees, error code: 9
                        // Observer latitude (negative south of equator)
    double latitude;
                        // valid range: -90 to 90 degrees, error code: 10
                        // Observer elevation [meters]
    double elevation;
                        // valid range: -6500000 or higher meters,
                                                                     error code: 11
   double pressure;
                        // Annual average local pressure [millibars]
                        // valid range: 0 to 5000 millibars,
                                                                      error code: 12
   double temperature; // Annual average local temperature [degrees Celsius]
                         // valid range: -273 to 6000 degrees Celsius, error code; 13
                         // Surface slope (measured from the horizontal plane)
    double slope;
                         // valid range: -360 to 360 degrees, error code: 14
   double azm rotation; // Surface azimuth rotation (measured from south to projection of
                            surface normal on horizontal plane, negative west)
                         // valid range: -360 to 360 degrees, error code: 15 \,
   double atmos refract; // Atmospheric refraction at sunrise and sunset (0.5667 deg is typical)
                         // valid range: -5 to 5 degrees, error code: 16
                        // Switch to choose functions for desired output (from enumeration)
    int function;
    //-----Intermediate OUTPUT VALUES-----
   double jd;
                       //Julian day
   double jc;
double jde;
                       //Julian century
                       //Julian ephemeris day
   double jce;
                      //Julian ephemeris century
   double jme;
                       //Julian ephemeris millennium
   double 1;
                      //earth heliocentric longitude [degrees]
   double b;
                       //earth heliocentric latitude [degrees]
                       //earth radius vector [Astronomical Units, AU]
   double r;
   double theta;
                      //geocentric longitude [degrees]
   double beta;
                       //geocentric latitude [degrees]
   double x0;
                       //mean elongation (moon-sun) [degrees]
   double x1;
                       //mean anomaly (sun) [degrees]
                       //mean anomaly (moon) [degrees]
//argument latitude (moon) [degrees]
   double x2;
   double x3;
   double x4;
                       //ascending longitude (moon) [degrees]
                       //nutation longitude [degrees]
   double del psi;
   double del epsilon; //nutation obliquity [degrees]
   double epsilon0; //ecliptic mean obliquity [arc seconds]
                       //ecliptic true obliquity [degrees]
   double ensilon:
   double del tau;
                       //aberration correction [degrees]
   double lamda;
                       //apparent sun longitude [degrees]
   double nu0;
                       //Greenwich mean sidereal time [degrees]
   double nu;
                       //Greenwich sidereal time [degrees]
   double alpha;
                       //geocentric sun right ascension [degrees]
   double delta;
                      //geocentric sun declination [degrees]
   double h;
                       //observer hour angle [degrees]
   double xi;
                       //sun equatorial horizontal parallax [degrees]
   double del alpha; //sun right ascension parallax [degrees]
   double delta prime; //topocentric sun declination [degrees]
   double alpha prime; //topocentric sun right ascension [degrees]
   double h prime;
                       //topocentric local hour angle [degrees]
   double e0;
                       //topocentric elevation angle (uncorrected) [degrees]
   double del_e;
                      //atmospheric refraction correction [degrees]
   double e;
                       //topocentric elevation angle (corrected) [degrees]
                       //equation of time [minutes]
   double eot;
   double srha;
                       //sunrise hour angle [degrees]
   double ssha;
                       //sunset hour angle [degrees]
   double sta;
                       //sun transit altitude [degrees]
   //----Final OUTPUT VALUES-----
   double zenith;
                       //topocentric zenith angle [degrees]
   double azimuth180; //topocentric azimuth angle (westward from south) [-180 to 180 degrees]
   double azimuth;
                       //topocentric azimuth angle (eastward from north) [ 0 to 360 degrees]
                       //surface incidence angle [degrees]
   double incidence;
   double suntransit; //local sun transit time (or solar noon) [fractional hour]
   double sunrise;  //local sunrise time (+/- 30 seconds) [fractional hour]
double sunset;  //local sunset time (+/- 30 seconds) [fractional hour]
   double sunset;
} spa data;
//Calculate SPA output values (in structure) based on input values passed in structure
int spa calculate(spa data *spa);
#endif
```

```
A.7. C Code: SPA source file (SPA.c)
       Solar Position Algorithm (SPA)
                    for
         Solar Radiation Application
11
                May 12, 2003
    Filename: SPA.C
//
//
    Afshin Michael Andreas
    afshin andreas@nrel.gov (303)384-6383
//
    Measurement & Instrumentation Team
    Solar Radiation Research Laboratory
//
    National Renewable Energy Laboratory
    1617 Cole Blvd, Golden, CO 80401
See the SPA.H header file for usage
    This code is based on the NREL
    technical report "Solar Position
    Algorithm for Solar Radiation
// Application" by I. Reda & A. Andreas //
//
    NOTICE
//This solar position algorithm for solar radiation applications (the "data") was produced by
//the National Renewable Energy Laboratory ("NREL"), which is operated by the Midwest Research
//Institute ("MRI") under Contract No. DE-AC36-99-G010337 with the U.S. Department of Energy
//(the "Government").
//Reference herein, directly or indirectly to any specific commercial product, process, or
//service by trade name, trademark, manufacturer, or otherwise, does not constitute or imply
//its endorsement, recommendation, or favoring by the Government, MRI or NREL.
//THESE DATA ARE PROVIDED "AS IS" AND NEITHER THE GOVERNMENT, MRI, NREL NOR ANY OF THEIR
//EMPLOYEES, MAKES ANY WARRANTY, EXPRESS OR IMPLIED, INCLUDING THE WARRANTIES OF MERCHANTABILITY
//AND FITNESS FOR A PARTICULAR PURPOSE, OR ASSUMES ANY LEGAL LIABILITY OR RESPONSIBILITY FOR THE
//ACCURACY, COMPLETENESS, OR USEFULNESS OF ANY SUCH INFORMATION DISCLOSED IN THE ALGORITHM, OR
//OF ANY APPARATUS, PRODUCT, OR PROCESS DISCLOSED, OR REPRESENTS THAT ITS USE WOULD NOT INFRINGE
//PRIVATELY OWNED RIGHTS.
// Revised 27-FEB-2004 Andreas
          Added bounds check on inputs and return value for spa calculate().
// Revised 10-MAY-2004 Andreas
          Changed temperature bound check minimum from -273.15 to -273 degrees C.
// Revised 17-JUN-2004 Andreas
          Corrected a problem that caused a bogus sunrise/set/transit on the equinox.
// Revised 18-JUN-2004 Andreas
          Added a "function" input variable that allows the selecting of desired outputs.
// Revised 21-JUN-2004 Andreas
          Added 3 new intermediate output values to SPA structure (srha, ssha, & sta).
  Revised 23-JUN-2004 Andreas
          Enumerations for "function" were renamed and 2 were added.
//
          Prevented bound checks on inputs that are not used (based on function).
// Revised 01-SEP-2004 Andreas
          Changed a local variable from integer to double.
// Revised 12-JUL-2005 Andreas
          Put a limit on the EOT calculation, so that the result is between -20 and 20.
// Revised 26-OCT-2005 Andreas
          Set the atmos. refraction correction to zero, when sun is below horizon.
          Made atmos refract input a requirement for all "functions".
          Changed atmos refract bound check from +/- 10 to +/- 5 degrees.
// Revised 07-NOV-2006 Andreas
          Corrected 3 earth periodic terms in the L TERMS array.
          Corrected 2 earth periodic terms in the R TERMS array.
```

```
// Revised 10-NOV-2006 Andreas
           Corrected a constant used to calculate topocentric sun declination.
           Put a limit on observer hour angle, so result is between 0 and 360.
// Revised 13-NOV-2006 Andreas
           Corrected calculation of topocentric sun declination.
//
           Converted all floating point inputs in spa structure to doubles.
// Revised 27-FEB-2007 Andreas
// Minor correction made as to when atmos. refraction correction is set to zero.
#include <math.h>
#include "spa.h"
                  3.1415926535897932384626433832795028841971
#define PI
#define SUN RADIUS 0.26667
#define L COUNT 6
#define B COUNT 2
#define R COUNT 5
#define Y COUNT 63
#define L MAX SUBCOUNT 64
#define B MAX SUBCOUNT 5
#define R MAX SUBCOUNT 40
enum {TERM A, TERM B, TERM C, TERM COUNT};
enum {TERM X0, TERM X1, TERM X2, TERM X3, TERM X4, TERM X COUNT};
enum {TERM_PSI_A, TERM_PSI_B, TERM_EPS_C, TERM_EPS_D, TERM_PE_COUNT};
enum {JD_MINUS, JD_ZERO, JD_PLUS, JD_COUNT};
enum {SUN TRANSIT, SUN RISE, SUN SET, SUN COUNT};
#define TERM Y COUNT TERM X COUNT
const int 1 subcount[L COUNT] = {64,34,20,7,3,1};
const int b subcount[B COUNT] = {5,2};
const int r subcount [R COUNT] = \{40, 10, 6, 2, 1\};
/// Earth Periodic Terms
const double L TERMS[L COUNT][L MAX SUBCOUNT][TERM COUNT]=
        {175347046.0,0,0},
        {3341656.0, 4.6692568, 6283.07585},
        {34894.0, 4.6261, 12566.1517},
        {3497.0,2.7441,5753.3849},
        {3418.0,2.8289,3.5231},
        {3136.0,3.6277,77713.7715},
        {2676.0,4.4181,7860.4194},
        {2343.0,6.1352,3930.2097}
        {1324.0,0.7425,11506.7698},
        {1273.0,2.0371,529.691},
        {1199.0,1.1096,1577.3435},
        {990,5.233,5884.927},
        {902,2.045,26.298},
        {857,3.508,398.149},
        {780,1.179,5223.694},
         [753,2.533,5507.553],
        {505,4.583,18849.228},
        {492,4.205,775.523},
        {357,2.92,0.067},
        {317,5.849,11790.629},
        {284,1.899,796.298},
        {271,0.315,10977.079},
        {243,0.345,5486.778},
        {206,4.806,2544.314},
        {205,1.869,5573.143},
        {202,2.458,6069.777},
        {156,0.833,213.299},
        {132,3.411,2942.463},
        {126,1.083,20.775},
        {115,0.645,0.98},
        {103,0.636,4694.003}
        {102,0.976,15720.839},
        {102,4.267,7.114},
```

```
{99,6.21,2146.17},
{98,0.68,155.42},
{86,5.98,161000.69},
{85,1.3,6275.96},
{85,3.67,71430.7},
{80,1.81,17260.15},
{79,3.04,12036.46},
{75,1.76,5088.63},
{74,3.5,3154.69},
{74,4.68,801.82},
[70,0.83,9437.76],
[62,3.98,8827.39],
{61,1.82,7084.9},
{57,2.78,6286.6},
{56,4.39,14143.5},
{56,3.47,6279.55},
{52,0.19,12139.55},
{52,1.33,1748.02},
{51,0.28,5856.48},
{49,0.49,1194.45},
{41,5.37,8429.24},
{41,2.4,19651.05},
{39,6.17,10447.39},
{37,6.04,10213.29},
{37,2.57,1059.38},
{36,1.71,2352.87},
[36,1.78,6812.77],
{33,0.59,17789.85},
{30,0.44,83996.85},
{30,2.74,1349.87},
{25,3.16,4690.48}
{628331966747.0,0,0},
{206059.0,2.678235,6283.07585},
{4303.0,2.6351,12566.1517},
{425.0,1.59,3.523},
{119.0,5.796,26.298}
{109.0,2.966,1577.344},
{93,2.59,18849.23},
{72,1.14,529.69},
(68,1.87,398.15),
{67,4.41,5507.55},
{59,2.89,5223.69},
{56,2.17,155.42},
{45,0.4,796.3},
{36,0.47,775.52},
{29,2.65,7.11},
{21,5.34,0.98},
{19,1.85,5486.78},
{19,4.97,213.3},
{17,2.99,6275.96},
{16,0.03,2544.31},
{16,1.43,2146.17},
{15,1.21,10977.08},
{12,2.83,1748.02},
{12,3.26,5088.63},
{12,5.27,1194.45},
{12,2.08,4694},
{11,0.77,553.57},
{10,1.3,6286.6}
{10,4.24,1349.87},
{9,2.7,242.73},
{9,5.64,951.72},
{8,5.3,2352.87},
{6,2.65,9437.76},
{6,4.67,4690.48}
{52919.0,0,0},
{8720.0,1.0721,6283.0758},
{309.0,0.867,12566.152},
{27,0.05,3.52},
{16,5.19,26.3},
{16,3.68,155.42},
{10,0.76,18849.23},
```

```
{9,2.06,77713.77},
         {7,0.83,775.52},
{5,4.66,1577.34},
         {4,1.03,7.11},
         {4,3.44,5573.14},
         {3,5.14,796.3},
         {3,6.05,5507.55},
         {3,1.19,242.73},
         {3,6.12,529.69},
         {3,0.31,398.15},
         {3,2.28,553.57},
         {2,4.38,5223.69},
         {2,3.75,0.98}
         {289.0,5.844,6283.076},
         {35,0,0},
         {17,5.49,12566.15},
         {3,5.2,155.42},
         {1,4.72,3.52},
{1,5.3,18849.23},
         {1,5.97,242.73}
         {114.0,3.142,0},
         {8,4.13,6283.08},
         {1,3.84,12566.15}
         {1,3.14,0}
};
const double B TERMS[B COUNT][B MAX SUBCOUNT][TERM COUNT]=
         {280.0,3.199,84334.662},
{102.0,5.422,5507.553},
         {80,3.88,5223.69},
         {44,3.7,2352.87},
         {32,4,1577.34}
         {9,3.9,5507.55},
{6,1.73,5223.69}
};
const double R_TERMS[R_COUNT][R_MAX_SUBCOUNT][TERM_COUNT]=
         {100013989.0,0,0},
{1670700.0,3.0984635,6283.07585},
         {13956.0,3.05525,12566.1517},
         {3084.0,5.1985,77713.7715},
         {1628.0,1.1739,5753.3849},
{1576.0,2.8469,7860.4194},
         {925.0,5.453,11506.77},
         {542.0,4.564,3930.21},
         {472.0,3.661,5884.927},
         {346.0,0.964,5507.553},
         {329.0,5.9,5223.694},
         {307.0,0.299,5573.143},
         {243.0,4.273,11790.629},
         {212.0,5.847,1577.344},
         {186.0,5.022,10977.079},
         {175.0,3.012,18849.228},
         {110.0,5.055,5486.778},
         {98,0.89,6069.78},
         {86,5.69,15720.84},
         {86,1.27,161000.69},
         {65,0.27,17260.15},
         {63,0.92,529.69},
         {57,2.01,83996.85},
         {56,5.24,71430.7},
         {49,3.25,2544.31},
```

```
{47,2.58,775.52},
        {45,5.54,9437.76},
        {43,6.01,6275.96},
       {39,5.36,4694},
        {38,2.39,8827.39},
        {37,0.83,19651.05},
        {37,4.9,12139.55},
        {36,1.67,12036.46},
       {35,1.84,2942.46},
        {33,0.24,7084.9},
        {32,0.18,5088.63},
       {32,1.78,398.15},
       {28,1.21,6286.6},
       {28,1.9,6279.55},
       {26, 4.59, 10447.39}
       {103019.0,1.10749,6283.07585},
        {1721.0,1.0644,12566.1517},
       {702.0,3.142,0},
        {32,1.02,18849.23},
       {31,2.84,5507.55},
       {25,1.32,5223.69},
       {18,1.42,1577.34},
       {10,5.91,10977.08},
        {9,1.42,6275.96},
       {9,0.27,5486.78}
       {4359.0,5.7846,6283.0758},
       {124.0,5.579,12566.152},
       {12,3.14,0},
       {9,3.63,77713.77},
       {6,1.87,5573.14},
       {3,5.47,18849.23}
       {145.0,4.273,6283.076},
       {7,3.92,12566.15}
       {4,2.56,6283.08}
};
/// Periodic Terms for the nutation in longitude and obliquity
const int Y TERMS[Y COUNT][TERM Y COUNT]=
   {0,0,0,0,1},
{-2,0,0,2,2},
    {0,0,0,2,2},
    {0,0,0,0,2},
    {0,1,0,0,0},
   {0,0,1,0,0},
    {-2,1,0,2,2},
    {0,0,0,2,1},
    {0,0,1,2,2},
   {-2,-1,0,2,2},
{-2,0,1,0,0},
    {-2,0,0,2,1},
    \{0,0,-1,2,2\},
    {2,0,0,0,0},
    {0,0,1,0,1},
    {2,0,-1,2,2},
    \{0,0,-1,0,1\},
    {0,0,1,2,1},
{-2,0,2,0,0},
    \{0,0,-2,2,1\},
    {2,0,0,2,2},
    {0,0,2,2,2},
    {0,0,2,0,0},
    {-2,0,1,2,2},
    {0,0,0,2,0},
```

```
{-2,0,0,2,0},
     {0,0,-1,2,1},
{0,2,0,0,0},
     {2,0,-1,0,1},
{-2,2,0,2,2},
     {0,1,0,0,1},
     \{-2,0,1,0,1\},
     {0,-1,0,0,1},
     {0,0,2,-2,0},
     {2,0,-1,2,1},
     {2,0,1,2,2},
     {0,1,0,2,2},
     \{-2,1,1,0,0\},
     {0,-1,0,2,2},
     {2,0,0,2,1},
     {2,0,1,0,0},
     {-2,0,2,2,2},
     {-2,0,1,2,1},
     {2,0,-2,0,1},
     {2,0,0,0,1},
     {0,-1,1,0,0},
{-2,-1,0,2,1},
     {-2,0,0,0,1},
     {0,0,2,2,1},
     {-2,0,2,0,1},
     {-2,1,0,2,1},
     {0,0,1,-2,0},
     {-1,0,1,0,0},
{-2,1,0,0,0},
{1,0,0,0,0},
     {0,0,1,2,0},
     \{0,0,-2,2,2\},
     {-1,-1,1,0,0},
     {0,1,1,0,0},
     \{0, -1, 1, 2, 2\},\
     {2,-1,-1,2,2},
{0,0,3,2,2},
     \{2,-1,0,2,2\},
};
const double PE_TERMS[Y_COUNT][TERM_PE_COUNT]={
     {-171996,-174.2,92025,8.9},
     \{-13187, -1.6, 5736, -3.1\},\
     \{-2274, -0.2, 977, -0.5\},
     {2062,0.2,-895,0.5},
{1426,-3.4,54,-0.1},
     {712,0.1,-7,0},
{-517,1.2,224,-0.6},
     {-386,-0.4,200,0},
     {-301,0,129,-0.1},
     {217,-0.5,-95,0.3},
     {-158,0,0,0},
{129,0.1,-70,0},
{123,0,-53,0},
     {63,0,0,0},
     {63,0.1,-33,0},
     {-59,0,26,0},
     {-58,-0.1,32,0},
     {-51,0,27,0},
     {48,0,0,0},
     {46,0,-24,0},
{-38,0,16,0},
     {-31,0,13,0},
     {29,0,0,0},
     {29,0,-12,0},
     {26,0,0,0},
     {-22,0,0,0},
     {21,0,-10,0},
     {17,-0.1,0,0},
{16,0,-8,0},
     \{-16, 0.1, 7, 0\},\
     {-15,0,9,0},
     {-13,0,7,0},
     \{-12,0,6,0\},
     {11,0,0,0},
     \{-10,0,5,0\},
```

```
{-8,0,3,0},
    {7,0,-3,0},
{-7,0,0,0},
    {-7,0,3,0},
{-7,0,3,0},
    {6,0,0,0},
    \{6,0,-3,0\},\
    {6,0,-3,0},
    {-6,0,3,0},
    {-6,0,3,0},
    {5,0,0,0},
    \{-5,0,3,0\},
    {-5,0,3,0},
    \{-5,0,3,0\},
    {4,0,0,0},
    {4,0,0,0},
    {4,0,0,0},
    {-4,0,0,0},
    {-4,0,0,0},
    {-4,0,0,0},
    {3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
    {-3,0,0,0},
};
double rad2deg(double radians)
    return (180.0/PI)*radians;
double deg2rad(double degrees)
    return (PI/180.0)*degrees;
double limit degrees (double degrees)
    double limited;
    degrees /= 360.0;
    limited = 360.0*(degrees-floor(degrees));
    if (limited < 0) limited += 360.0;</pre>
    return limited;
double limit_degrees180pm(double degrees)
    double limited;
    degrees /= 360.0;
    limited = 360.0*(degrees-floor(degrees));
    if (limited < -180.0) limited += 360.0;
else if (limited > 180.0) limited -= 360.0;
    return limited;
double limit degrees180 (double degrees)
    double limited;
    degrees /= 180.0;
    limited = 180.0*(degrees-floor(degrees));
    if (limited < 0) limited += 180.0;</pre>
    return limited;
```

```
double limit zero2one(double value)
    double limited;
   limited = value - floor(value);
if (limited < 0) limited += 1.0;</pre>
   return limited;
double limit minutes(double minutes)
   double limited=minutes;
   if (limited < -20.0) limited += 1440.0;
else if (limited > 20.0) limited -= 1440.0;
   return limited;
double dayfrac to local hr (double dayfrac, double timezone)
    return 24.0*limit zero2one(dayfrac + timezone/24.0);
double third order polynomial (double a, double b, double c, double d, double x)
   return ((a*x + b)*x + c)*x + d;
int validate inputs(spa_data *spa)
                         < -2000) || (spa->year
                                                        > 6000)) return 1;
    if ((spa->year
                                                        > 12 )) return 2;
> 31 )) return 3;
    if ((spa->month
                         < 1 ) || (spa->month
    if ((spa->day
                         < 1
                                 ) || (spa->day
                                                      > 24 )) return 4;
> 59 )) return 5;
> 59 )) return 6;
> 5000)) return 12;
                         < 0    ) || (spa->hour
< 0    ) || (spa->minute)
    if ((spa->hour
    if ((spa->minute
                        < 0     ) || (spa->second
< 0     ) || (spa->pressure
    if ((spa->second
    if ((spa->pressure
    if ((spa->temperature <= -273) || (spa->temperature > 6000)) return 13;
                     == 24 ) && (spa->minute > 0 )) return 5;
== 24 ) && (spa->second > 0 )) return 6;
    if ((spa->hour
    if ((spa->hour
                                 > 8000
    if (fabs(spa->delta t)
                                           ) return 7;
    if (fabs(spa->timezone) > 12
if (fabs(spa->longitude) > 180
                                           ) return 8;
                                           ) return 9;
                                          ) return 10;
                                 > 90
    if (fabs(spa->latitude)
    if (fabs(spa->atmos_refract) > 5
                                           ) return 16;
                                 < -6500000) return 11;
            spa->elevation
    if ((spa->function == SPA ZA INC) || (spa->function == SPA ALL))
                                    > 360) return 14;
        if (fabs(spa->slope)
        if (fabs(spa->azm rotation) > 360) return 15;
    return 0;
double julian_day (int year, int month, int day, int hour, int minute, int second, double tz)
    double day decimal, julian day, a;
    day decimal = day + (hour - tz + (minute + second/60.0)/60.0)/24.0;
    if (month < 3) {
       month += 12;
        year--;
    julian day = floor(365.25*(year+4716.0)) + floor(30.6001*(month+1)) + day decimal - 1524.5;
    if (julian day > 2299160.0) {
        a = floor(year/100);
        julian day += (2 - a + floor(a/4));
```

```
return julian day;
double julian century(double jd)
    return (jd-2451545.0)/36525.0;
double julian ephemeris day (double jd, double delta t)
    return jd+delta t/86400.0;
double julian ephemeris century(double jde)
    return (jde - 2451545.0)/36525.0;
double julian ephemeris millennium (double jce)
   return (jce/10.0);
double earth periodic term summation(const double terms[][TERM COUNT], int count, double jme)
   double sum=0;
    for (i = 0; i < count; i++)
       sum += terms[i][TERM A]*cos(terms[i][TERM B]+terms[i][TERM C]*jme);
   return sum;
double earth values (double term sum[], int count, double jme)
    int i;
   double sum=0;
   for (i = 0; i < count; i++)</pre>
        sum += term sum[i]*pow(jme, i);
   sum /= 1.0e8;
   return sum;
double earth heliocentric longitude (double jme)
   double sum[L COUNT];
   int i;
    for (i = 0; i < L_COUNT; i++)</pre>
        sum[i] = earth periodic term summation(L TERMS[i], l subcount[i], jme);
    return limit degrees(rad2deg(earth values(sum, L COUNT, jme)));
double earth heliocentric latitude (double jme)
    double sum[B COUNT];
   int i;
    for (i = 0; i < B COUNT; i++)</pre>
        sum[i] = earth periodic term summation(B TERMS[i], b subcount[i], jme);
    return rad2deg(earth_values(sum, B_COUNT, jme));
double earth_radius_vector(double jme)
    double sum[R COUNT];
```

```
int i;
    for (i = 0; i < R COUNT; i++)
        sum[i] = earth periodic term summation(R TERMS[i], r subcount[i], jme);
    return earth values(sum, R COUNT, jme);
double geocentric longitude (double 1)
    double theta = 1 + 180.0;
    if (theta >= 360.0) theta -= 360.0;
   return theta;
double geocentric latitude (double b)
   return -b;
double mean elongation moon sun(double jce)
    return third order polynomial (1.0/189474.0, -0.0019142, 445267.11148, 297.85036, jce);
double mean anomaly sun(double jce)
    return third order polynomial (-1.0/300000.0, -0.0001603, 35999.05034, 357.52772, jce);
double mean anomaly moon (double jce)
   return third order polynomial (1.0/56250.0, 0.0086972, 477198.867398, 134.96298, jce);
double argument latitude moon (double jce)
    return third order polynomial(1.0/327270.0, -0.0036825, 483202.017538, 93.27191, jce);
double ascending longitude moon(double jce)
    return third_order_polynomial(1.0/450000.0, 0.0020708, -1934.136261, 125.04452, jce);
double xy_term_summation(int i, double x[TERM X COUNT])
    int j;
    double sum=0;
    for (j = 0; j < TERM_Y_COUNT; j++)</pre>
       sum += x[j]*Y TERMS[i][j];
   return sum;
void nutation longitude and obliquity (double jce, double x[TERM X COUNT], double *del psi,
                                                                           double *del epsilon)
    int i;
   double xy_term_sum, sum psi=0, sum epsilon=0;
    for (i = 0; i < Y COUNT; i++) {</pre>
        xy term_sum = deg2rad(xy term summation(i, x));
        sum psi += (PE TERMS[i][TERM PSI A] + jce*PE TERMS[i][TERM PSI B])*sin(xy term sum);
        sum epsilon += (PE TERMS[i][TERM EPS C] + jce*PE TERMS[i][TERM EPS D])*cos(xy term sum);
    *del psi
                = sum psi
                              / 36000000.0;
    *del_epsilon = sum_epsilon / 36000000.0;
double ecliptic mean obliquity (double jme)
```

```
double u = jme/10.0;
   return 84381.448 + u*(-4680.96 + u*(-1.55 + u*(1999.25 + u*(-51.38 + u*(-249.67 +
                       u^*(-39.05 + u^*(7.12 + u^*(27.87 + u^*(5.79 + u^*2.45)))))));
double ecliptic true obliquity (double delta epsilon, double epsilon0)
   return delta epsilon + epsilon0/3600.0;
double aberration correction (double r)
   return -20.4898 / (3600.0*r);
double apparent sun longitude (double theta, double delta psi, double delta tau)
   return theta + delta psi + delta tau;
double greenwich mean sidereal time (double jd, double jc)
   return limit degrees (280.46061837 + 360.98564736629 * (jd - 2451545.0) +
                                       jc*jc*(0.000387933 - jc/38710000.0));
double greenwich sidereal time (double nu0, double delta psi, double epsilon)
   return nu0 + delta psi*cos(deg2rad(epsilon));
double geocentric sun right ascension (double lamda, double epsilon, double beta)
   double lamda rad
                      = deg2rad(lamda);
   double epsilon rad = deg2rad(epsilon);
   return limit degrees (rad2deg (atan2 (sin (lamda rad) *cos (epsilon rad) -
                                       tan(deg2rad(beta))*sin(epsilon rad), cos(lamda rad))));
double geocentric sun declination (double beta, double epsilon, double lamda)
   double beta_rad
                      = deg2rad(beta);
   double epsilon rad = deg2rad(epsilon);
   return rad2deg(asin(sin(beta rad)*cos(epsilon rad) +
                        cos(beta_rad)*sin(epsilon_rad)*sin(deg2rad(lamda))));
double observer hour angle (double nu, double longitude, double alpha deg)
    return limit degrees (nu + longitude - alpha deg);
double sun equatorial horizontal parallax (double r)
   return 8.794 / (3600.0 * r);
void sun right ascension parallax and topocentric dec (double latitude, double elevation,
               double xi, double h, double delta, double *delta alpha, double *delta prime)
   double delta_alpha_rad;
   double lat_rad = deg2rad(latitude);
                   = deg2rad(xi);
= deg2rad(h);
   double xi rad
   double h rad
   double delta rad = deg2rad(delta);
   double u = atan(0.99664719 * tan(lat rad));
   double y = 0.99664719 * \sin(u) + elevation*sin(lat rad)/6378140.0;
   double x =
                           cos(u) + elevation*cos(lat rad)/6378140.0;
   delta alpha rad =
                           atan2(
                                                  - x*sin(xi_rad) *sin(h_rad),
                                  cos(delta rad) - x*sin(xi rad) *cos(h rad));
```

```
*delta prime = rad2deg(atan2((sin(delta rad) - y*sin(xi rad))*cos(delta alpha rad),
                                cos(delta rad) - x*sin(xi rad) *cos(h rad)));
   *delta alpha = rad2deg(delta alpha rad);
double topocentric sun right ascension (double alpha deg, double delta alpha)
   return alpha deg + delta alpha;
double topocentric local hour angle (double h, double delta alpha)
   return h - delta alpha;
double topocentric elevation angle (double latitude, double delta prime, double h prime)
   double lat rad
                          = deg2rad(latitude);
   double delta prime rad = deg2rad(delta prime);
   return rad2deg(asin(sin(lat_rad)*sin(delta_prime_rad) +
                       cos(lat rad) *cos(delta prime rad) * cos(deg2rad(h prime))));
double atmospheric refraction correction (double pressure, double temperature,
                                       double atmos refract, double e0)
   double del e = 0;
   if (e0 >= -1* (SUN RADIUS + atmos refract))
       del e = (pressure / 1010.0) * (283.0 / (273.0 + temperature)) *
                1.02 / (60.0 * tan(deg2rad(e0 + 10.3/(e0 + 5.11))));
   return del e;
double topocentric elevation angle corrected (double e0, double delta e)
   return e0 + delta e;
double topocentric zenith angle (double e)
   return 90.0 - e;
double topocentric azimuth angle neg180 180 (double h prime, double latitude, double delta prime)
   double h_prime_rad = deg2rad(h_prime);
   double lat rad
                    = deg2rad(latitude);
   return rad2deg(atan2(sin(h_prime_rad),
                        cos(h_prime_rad)*sin(lat_rad) -
                        tan(deg2rad(delta prime))*cos(lat rad)));
double topocentric azimuth angle zero 360 (double azimuth180)
   return azimuth180 + 180.0;
double surface incidence angle (double zenith, double azimuth180, double azm rotation,
                                                               double slope)
   double zenith rad = deg2rad(zenith);
   double slope rad = deg2rad(slope);
   azm rotation)));
double sun mean longitude (double jme)
   return limit degrees (280.4664567 + jme*(360007.6982779 + jme*(0.03032028 +
```

```
jme*(1/49931.0 + jme*(-1/15300.0)
                                                       + jme*(-1/2000000.0)))));
double eot (double m, double alpha, double del psi, double epsilon)
    return limit minutes(4.0*(m - 0.0057183 - alpha + del psi*cos(deg2rad(epsilon))));
double approx_sun_transit_time(double alpha zero, double longitude, double nu)
   return (alpha zero - longitude - nu) / 360.0;
double sun hour angle at rise set (double latitude, double delta zero, double h0 prime)
                         = -999999:
   double latitude rad = deg2rad(latitude);
   double delta zero_rad = deg2rad(delta zero);
   double argument
                       = (sin(deg2rad(h0 prime)) - sin(latitude rad)*sin(delta zero rad)) /
                                                   (cos(latitude rad)*cos(delta zero rad));
   if (fabs(argument) <= 1) h0 = limit degrees180(rad2deg(acos(argument)));</pre>
   return h0;
void approx sun rise and set(double *m rts, double h0)
   double h0 dfrac = h0/360.0;
   m_rts[SUN_RISE] = limit_zero2one(m_rts[SUN_TRANSIT] - h0_dfrac);
   m_rts[SUN_SET] = limit_zero2one(m_rts[SUN_TRANSIT] + h0_dfrac);
m_rts[SUN_TRANSIT] = limit_zero2one(m_rts[SUN_TRANSIT]);
double rts alpha delta prime (double *ad, double n)
    double a = ad[JD ZERO] - ad[JD MINUS];
   double b = ad[JD PLUS] - ad[JD ZERO];
   if (fabs(a) >= 2.0) a = limit zero2one(a);
   if (fabs(b) >= 2.0) b = limit zero2one(b);
   return ad[JD ZERO] + n * (a + b + (b-a)*n)/2.0;
double rts sun altitude (double latitude, double delta prime, double h prime)
   double latitude rad
                        = deg2rad(latitude);
   double delta prime rad = deg2rad(delta prime);
   return rad2deg(asin(sin(latitude rad)*sin(delta prime rad) +
                       cos(latitude rad)*cos(delta prime rad)*cos(deg2rad(h prime))));
return m rts[sun] + (h rts[sun] - h0 prime) /
          (360.0*cos(deg2rad(delta prime[sun]))*cos(deg2rad(latitude))
          *sin(deg2rad(h prime[sun])));
// Calculate required SPA parameters to get the right ascension (alpha) and declination (delta)
// Note: JD must be already calculated and in structure
void calculate geocentric sun right ascension and declination (spa data *spa)
   double x [TERM X COUNT];
   spa->jc = julian century(spa->jd);
   spa->jde = julian ephemeris day(spa->jd, spa->delta t);
   spa->jce = julian ephemeris century(spa->jde);
```

```
spa->jme = julian ephemeris millennium(spa->jce);
   spa->l = earth heliocentric longitude(spa->jme);
   spa->b = earth heliocentric latitude (spa->jme);
   spa->r = earth radius vector (spa->jme);
   spa->theta = geocentric longitude(spa->l);
   spa->beta = geocentric latitude(spa->b);
   x[TERM X0] = spa->x0 = mean elongation moon sun(spa->jce);
   x[TERM X1] = spa->x1 = mean anomaly sun(spa->jce);
   x[TERM X2] = spa->x2 = mean anomaly moon(spa->jce);
   x[TERM_X3] = spa->x3 = argument_latitude_moon(spa->jce);
   x[TERM X4] = spa->x4 = ascending longitude moon(spa->jce);
   nutation longitude and obliquity (spa->jce, x, & (spa->del psi), & (spa->del epsilon));
   spa->epsilon0 = ecliptic mean obliquity(spa->jme);
   spa->epsilon = ecliptic true obliquity(spa->del epsilon, spa->epsilon0);
   spa->del tau
                 = aberration correction(spa->r);
                 = apparent sun longitude (spa->theta, spa->del psi, spa->del tau);
   spa->lamda
   spa->nu0
                 = greenwich mean sidereal time (spa->jd, spa->jc);
   spa->nu
                  = greenwich sidereal time (spa->nu0, spa->del psi, spa->epsilon);
   spa->alpha = geocentric sun right ascension(spa->lamda, spa->epsilon, spa->beta);
   spa->delta = geocentric sun declination(spa->beta, spa->epsilon, spa->lamda);
// Calculate Equation of Time (EOT) and Sun Rise, Transit, & Set (RTS) \,
////////////////-
void calculate eot and sun rise transit set (spa data *spa)
   spa data sun rts = *spa;
   double nu, m, h0, n;
   double alpha[JD COUNT], delta[JD COUNT];
   double m rts[SUN COUNT], nu rts[SUN COUNT], h rts[SUN COUNT];
   double alpha_prime[SUN_COUNT], delta_prime[SUN_COUNT], h_prime[SUN_COUNT];
   double h0 prime = -1*(SUN RADIUS + spa->atmos refract);
   int i;
           = sun mean longitude(spa->jme);
   spa->eot = eot(m, spa->alpha, spa->del psi, spa->epsilon);
   sun rts.hour = sun rts.minute = sun rts.second = sun rts.timezone = 0;
   sun_rts.jd = julian_day (sun_rts.year, sun_rts.month, sun_rts.day,
                           sun rts.hour, sun rts.minute, sun rts.second, sun rts.timezone);
   calculate geocentric sun right ascension and declination (&sun rts);
   nu = sun rts.nu;
   sun rts.delta t = 0;
   sun rts.id--;
    for (i = 0; i < JD COUNT; i++) {
       calculate geocentric sun right ascension and declination (&sun rts);
       alpha[i] = sun rts.alpha;
       delta[i] = sun_rts.delta;
       sun rts.jd++;
   m_rts[SUN_TRANSIT] = approx_sun_transit_time(alpha[JD ZERO], spa->longitude, nu);
   ho = sun_hour_angle_at_rise_set(spa->latitude, delta[JD_ZERO], ho_prime);
   if (h0 >= 0) {
       approx sun rise and set (m rts, h0);
       for (i = 0; i < SUN COUNT; i++) {
           nu rts[i]
                        = nu + 360.985647*m rts[i];
                         = m rts[i] + spa->delta t/86400.0;
```

```
alpha prime[i] = rts alpha delta prime(alpha, n);
          delta prime[i] = rts alpha delta prime(delta, n);
          h prime[i]
                        = limit degrees180pm(nu rts[i] + spa->longitude - alpha prime[i]);
                        = rts sun altitude(spa->latitude, delta prime[i], h prime[i]);
          h rts[i]
       spa->srha = h prime[SUN RISE];
       spa->ssha = h prime[SUN SET];
       spa->sta = h rts[SUN TRANSIT];
       spa->suntransit = dayfrac_to_local_hr(m_rts[SUN_TRANSIT] - h prime[SUN TRANSIT] / 360.0,
                                          spa->timezone);
       spa->sunrise = dayfrac to local hr(sun rise and set(m rts, h rts, delta prime,
                       spa->latitude, h prime, h0 prime, SUN RISE), spa->timezone);
       spa->sunset = dayfrac to local hr(sun rise and set(m rts, h rts, delta prime,
                       spa->latitude, h prime, h0 prime, SUN SET), spa->timezone);
   } else spa->srha= spa->ssha= spa->sta= spa->suntransit= spa->sunrise= spa->sunset= -99999;
// Calculate all SPA parameters and put into structure
// Note: All inputs values (listed in header file) must already be in structure
int spa calculate(spa data *spa)
   int result;
   result = validate inputs(spa);
   if (result == 0)
       spa->jd = julian_day (spa->year, spa->month, spa->day,
                           spa->hour, spa->minute, spa->second, spa->timezone);
       calculate geocentric sun right ascension and declination (spa);
       spa->h = observer hour angle(spa->nu, spa->longitude, spa->alpha);
       spa->xi = sun equatorial horizontal parallax(spa->r);
       sun right ascension parallax and topocentric dec (spa->latitude, spa->elevation, spa->xi,
                                 spa->h, spa->delta, &(spa->del alpha), &(spa->delta prime));
       spa->alpha_prime = topocentric_sun_right_ascension(spa->alpha, spa->del_alpha);
       spa->h prime
                      = topocentric local hour angle (spa->h, spa->del alpha);
                  = topocentric elevation angle (spa->latitude, spa->delta prime,
       spa->h prime);
       spa->del e = atmospheric refraction correction(spa->pressure, spa->temperature,
                                                    spa->atmos refract, spa->e0);
       spa->e
                   = topocentric elevation angle corrected (spa->\overline{0}, spa->del e);
       spa->zenith
                     = topocentric zenith angle(spa->e);
       spa->azimuth180 = topocentric azimuth angle neg180 180 (spa->h prime, spa->latitude,
                                                                      spa->delta prime);
                     = topocentric azimuth angle zero 360 (spa->azimuth180);
       spa->azimuth
       if ((spa->function == SPA ZA INC) || (spa->function == SPA ALL))
           spa->incidence = surface incidence angle(spa->zenith, spa->azimuth180,
                                                 spa->azm rotation, spa->slope);
       if ((spa->function == SPA ZA RTS) || (spa->function == SPA ALL))
          calculate eot and sun rise transit set (spa);
   return result;
.
.
```

REPORT DOCUMEN	Form Approved OMB NO. 0704-0188				
Public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302, and to the Office of Management and Budget, Paperwork Reduction Project (0704-0188), Washington, DC 20503.					
AGENCY USE ONLY (Leave blank)	ERED				
TITLE AND SUBTITLE     Solar Position Algorithm for S	5. FUNDING NUMBERS WU1D5600				
6. AUTHOR(S)					
Ibrahim Reda & Afshin Andreas					
7. PERFORMING ORGANIZATION NAM National Renewable Energy L	PERFORMING ORGANIZATION     REPORT NUMBER				
1617 Cole Blvd. Golden, CO 80401-3393	NREL/TP-560-34302				
9. SPONSORING/MONITORING AGENC	10. SPONSORING/MONITORING AGENCY REPORT NUMBER				
11. SUPPLEMENTARY NOTES					
NREL Technical Monitor: Ibra	ahim Reda				
12a. DISTRIBUTION/AVAILABILITY STA National Technical Informa U.S. Department of Commo 5285 Port Royal Road Springfield, VA 22161	12b. DISTRIBUTION CODE				
13. ABSTRACT (Maximum 200 words)					
1. This report is a step-by-step procedure for implementing an algorithm to calculate the solar zenith and azimuth angles in the period from the year -2000 to 6000, with uncertainties of ±0.0003/. It is written in a step-by-step format to simplify otherwise complicated steps, with a focus on the sun instead of the planets and stars in general. The algorithm is written in such a way to accommodate solar radiation applications.					
14. SUBJECT TERMS	15. NUMBER OF PAGES				
algorithm; solar position algo zenith angles; solar azimuth	16. PRICE CODE				
17. SECURITY CLASSIFICATION OF REPORT Unclassified	18. SECURITY CLASSIFICATION OF THIS PAGE Unclassified	19. SECURITY CLASSIFICATION OF ABSTRACT Unclassified	20. LIMITATION OF ABSTRACT  UL		