



Exercise During Pregnancy

Safe and Beneficial for Most

Raul Artal, Carl Sherman & Nicholas A. DiNubile

To cite this article: Raul Artal, Carl Sherman & Nicholas A. DiNubile (1999) Exercise During Pregnancy, The Physician and Sportsmedicine, 27:8, 51-75, DOI: [10.3810/psm.1999.08.947](https://doi.org/10.3810/psm.1999.08.947)

To link to this article: <https://doi.org/10.3810/psm.1999.08.947>



Published online: 19 Jun 2015.



Submit your article to this journal [↗](#)



Article views: 532



View related articles [↗](#)



Citing articles: 1 View citing articles [↗](#)



Exercise During Pregnancy

Safe and Beneficial for Most

Raul Artal, MD, with Carl Sherman

Series Editor: Nicholas A. DiNubile, MD

Photo: © 1999. Tim Defrisco/ALLSPORT

In brief

Regular, moderate exercise does not significantly affect length of gestation or birth weight, and it may ease pregnancy and labor. Active pregnant women should maintain adequate hydration and nutrition, avoid hyperthermia, and take precautions if obstetric or medical complications are present. Supine exercise, contact sports, high-altitude exertion, and scuba diving should be avoided, and some women, including those who lift weights or do high-impact exercise, may need to modify their routines. Most women can begin or continue to exercise during pregnancy.

As attitudes toward exercise have changed in recent decades, the shift with regard to exercise during pregnancy has been particularly marked.


In the 1950s, prevailing standards of care allowed a pregnant woman 1 mile of walking daily, preferably broken up into several sessions.¹ In

continued



For CME credit, see page 77

Dr Artal is professor and chairman of the Department of Obstetrics and Gynecology at St Louis University in St Louis. He is a fellow of the American College of Obstetricians and Gynecologists and the American College of Sports Medicine. Mr Sherman is a freelance writer in New York City. Dr DiNubile is an orthopedic surgeon in private practice in Havertown, Pennsylvania, specializing in sports medicine and arthroscopy. He is director of Sports Medicine and Wellness at the Crozer-Keystone Healthplex in Springfield, Pennsylvania, a clinical assistant professor in the department of orthopedic surgery at the University of Pennsylvania in Philadelphia, and the orthopedic consultant to the Philadelphia 76ers basketball team and the Pennsylvania Ballet. Drs Artal and DiNubile are members of the editorial board of THE PHYSICIAN AND SPORTSMEDICINE.



Women who have been active before pregnancy can often continue an exercise routine with modifications. Though pregnancy is not a time to begin a new sport or intensive training, it can offer an opportunity for sedentary women to initiate positive changes.

1985, the American College of Obstetrics and Gynecology (ACOG) endorsed the safety of most aerobic exercise but placed definite limits on duration (15 minutes) and heart rate (140 beats per minute).² A decade later the organization issued recommendations, based on physiologic evidence, that women with uncomplicated pregnancies can exercise safely with virtually the same safeguards as nonpregnant women.³ Many women of childbearing age, 15% by one estimate, now exercise regularly, and strongly desire to continue doing so.⁴ These changes reflect cultural developments such as the “fitness boom” of the 1970s and the evolving role of women as much as advances in scientific understanding.⁵

Data are inconclusive, but it appears that birth weight is not affected in exercising women who have proper calorie intake.

In fact, our knowledge in this area is far from complete. Concerns about the physiologic interactions between the changes of pregnancy and the demands of exercise have not been entirely laid to rest.⁶ As with exercise recommendations for patients who have

any medical condition, those for pregnant women must balance the risks and benefits—to both mother and fetus.

Physiologic Concerns and Adaptations

Cardiovascular. Relative to prepregnancy levels, increases of up to 50% in cardiac output, 15 beats per minute in resting heart rate, and 45% in blood volume can take place by the third trimester. These increases are offset to some extent by an increased capacity of the veins, resulting in stable or reduced blood pressure, and greater vasodilation at the skin, resulting in greater heat loss and a reduced tendency for hyperthermia.⁷

The cardiovascular impact of the dual stresses of exercise and pregnancy has been the focus of much investigation. The demands of exercising muscle divert blood flow from the uteroplacental unit, raising the theoretical risk of fetal hypoxia. In actuality, compensatory changes with exercise, such as increased maternal he-

matocrit and oxygen extraction, appear to prevent impairment of fetal oxygenation.⁵ Furthermore, repeated studies have shown, at most, short-term, limited fetal heart rate increases of 5 to 15 beats per minute, which do not pose a threat to the fetus. Episodes of fetal bradycardia are relatively rare in the absence of obstetric or medical complications and do not appear to affect pregnancy outcome.⁸

The supine position is associated with decreased maternal cardiac output after the first trimester; this has led to the caveat against exercise in this position.

In addition, studies of women living at high altitudes during their pregnancies⁹ showed reduced uterine blood flow. These results suggest that women should use caution, avoiding exertion at altitudes over 8,250 ft (2,500 m), at least until after 4 to 5 days of exposure.^{10,11}

Metabolic. Hyperthermia is a concern, given that both exercise and pregnancy increase metabolic rate and that maternal core temperatures in excess of 102.6°F (39.2°C) are potentially teratogenic in the first trimester.¹² Here too, adaptive changes associated with pregnancy, such as increased minute ventilation and skin blood flow, augment heat dissipation and offset somewhat the potential hyperthermic effects of exercise.¹³ Still, sensible precautions are indicated, such as maintaining adequate hydration and avoid exercising in very hot, humid environments.³

Probably because of the combined effect of limited expansion of the diaphragm (owing to an enlarged uterus) and increased fetal oxygen demand,¹⁴ $\dot{V}O_2$ max is significantly decreased in pregnant women during exercise as compared with nonpregnant women.¹ As a result, carbohydrates are used preferentially (as indicated by indirect calorimetry),¹⁵ resulting in significantly reduced plasma glucose concentrations during continuous prolonged (longer than 45 minutes) or strenuous bouts.¹⁶ Therefore, to reduce the risk of hypoglycemia, consuming adequate calories and limiting workout sessions to less than 45 minutes should be recommended.

Musculoskeletal. Marked musculoskeletal alterations in the pregnant woman's body could potentially increase the risk of injury during ex-

continued

Pregnant Athletes Test the Limits

Although a moderate exercise schedule is generally recommended during pregnancy, some highly conditioned women appear able to train safely at considerably more demanding levels.

One recent study¹ found that a rigorous protocol continued nearly to term had no adverse effect on fetal growth. It involved 42 well-trained women, some competitive at national or international levels (three world-class in bicycle, marathon, and biathlon competitions), who followed exercise protocols that included strength training, interval training, and endurance training 6 days a week. A "high-intensity" group's regimen consisted of two 2½-hour endurance training bouts weekly; "moderate" exercisers' workouts lasted 1½ hours.

In both groups, most women continued to exercise to within 4 days of labor. There was no significant difference in mean birth weights between the moderate- and high-intensity exercise groups (for both groups, mean birth weights were about 7 lb, 15 oz or 3,600 g), and these weights were within or higher than birth-

weight ranges among healthy women recorded elsewhere. Neither the rate of complications nor any other indices of pregnancy outcome were adversely affected.

A case report² described a 33-year-old elite marathon runner (personal best time: 2 hours, 34 minutes for the 42.2 km race) who continued to train an average of 66.5 miles (107 km) weekly up to 3 days before the birth of twins by elective cesarean section. "These data show that an endurance athlete can continue to train intensively during a twin pregnancy with no apparent effect on maternal and fetal health," the authors wrote.

Competitive athletes, however, will likely experience a decrease in their performance during pregnancy.³

References

1. Kardel KR, Kase T: Training in pregnant women: effects on fetal development and birth. *Am J Obstet Gynecol* 1998;178(2):280-286
2. Bailey DM, Davies B, Budgett R, et al: Endurance training during a twin pregnancy in a marathon runner. *Lancet* 1998;351(9110):1182
3. Hale RW, Milne L: The elite athlete and exercise in pregnancy. *Semin Perinatol* 1996;20(4):277-284

ercise. Growth of the breasts, uterus, and fetus increases lumbar lordosis and shifts the center of gravity forward, putting strain on the lower back, and hormonal changes lead to increased joint laxity and mobility.¹⁷ Though increases in athletic injury rates for pregnant women have not been documented,¹² caution and moderation are important as pregnancy progresses. Prudence dictates that activities such as contact sports be avoided, and that weight-bearing exercise be limited in women affected by joint-laxity changes.¹²

Effects on the Fetus

Most research has focused on the effects of exercise on fetal growth and length of gestation.

Birth weight. With regard to fetal weight in women who exercise, findings have been inconsistent. One widely cited study¹⁸ found mean

birth weight to be substantially lower when women exercised at or above 50% of preconception levels, compared with nonexercisers, while another study¹⁹ found no difference between birth weights of offspring of vigorous exercisers and those of sedentary women. A prospective study²⁰ of more than 800 pregnant women found that the babies of those who expended a mean of 2,000 kcal/wk in leisure-time physical activity (a level that does not necessarily reflect *intense* activity) were significantly heavier at birth than those of nonexercisers. It appears, though, that birth weight is not affected in women who have a proper calorie intake.

One review²¹ concluded that "current evidence appears to indicate participation in moderate to vigorous activity throughout pregnancy may enhance birth weight," with a caution that more severe regimens could result in lighter off-

continued

spring. This reviewer also suggested that careful quantification of calorie balance—the key to interpretation and missing from most studies—is needed for more definitive conclusions.

Gestation length. A similar picture emerges with regard to preterm delivery. The concern is that increased norepinephrine and prostaglandin output during exercise could stimulate uterine activity and premature labor,¹⁵ and, in fact, a number of epidemiologic studies²² have associated risk with various kinds of occupational activity. Other investigations²³ suggest that prolonged standing at work—rather than active movement—is the critical factor. Generally, studies of occupational activity are complicated, if not confounded, by such variables as socioeconomic status, stress, and fatigue.⁵

The few investigations that focused on recreational activity per se have tended to associate exercise with positive effects on term length. In one poorly controlled series²⁴ of 876 consecutive prenatal patients from Pennsylvania and New York, long-term heavy exercisers who expended more than 1,000 kcal/wk doing aerobic activity were more likely to deliver close to term than those who exercised less or not at all; they showed a marked reduction in risk of early delivery, and late deliveries in this group were significantly more likely to be spontaneous. “The observed reduction in risk of preterm delivery in a general obstetrical population is evidence of the safety, as well as the potential benefits, of exercise during pregnancy,” the authors concluded.

Effects on the Mother

Course of pregnancy. Whether exercise has a positive effect on the course of pregnancy—beyond the general benefit of maintaining fitness—is open to question, but at least some data suggest it does. Attenuation of such symptoms as nausea, fatigue, and back pain has been reported,²⁵ as well as improvements in self-esteem and other psychological parameters.²⁶ One observational study¹⁹ of 398 women found an inverse association between exercise and symptoms, with an increase in symptoms after exercise was reduced.

Labor and delivery. The effect of exercise on

labor is uncertain. Many women report feeling that being fit made labor easier,²⁷ and a number of studies have confirmed this. One poorly controlled, nonrandomized study²⁸ found shorter active labor (223 vs 302 minutes) among women who had exercised throughout pregnancy; in another,²⁹ a significant inverse correlation between fitness and labor duration emerged for multiparas but not primiparas. Some other investigators,⁵ however, have found no correlation between fitness, training level, and labor.

Recommendations

As a general rule, I recommend exercise to all my pregnant patients. Those who have been active will want to stay fit, and, though pregnancy is not a time to begin training intensively for a new sport, for sedentary women it often represents a window of opportunity for positive lifestyle changes.

Beginners. For women who have previously been sedentary, non-weight-bearing activities (eg, swimming, stationary cycling) pose the least risk of injury, and moderate walking is also safe for most. A schedule of one 20- to 30-minute exercise session daily at a comfortable pace is a reasonable goal. Because of the theoretical increase in risk of injury to joints, a jogging program should not be initiated at this time.

Previously active women. In the absence of obstetric or medical complications, most active women can continue to exercise in accustomed ways at close to prepregnancy levels. Participation in a full range of activities—sports such as volleyball and tennis, as well as aerobic exercise—are generally safe, but contact sports and scuba diving should be avoided, as well as any activity with the potential for causing abdominal trauma. Exercise during the first few days of exposure to high altitude is also not recommended because altitude, as well as the requirements of pregnancy, reduces oxygen availability.

Running. While most habitual runners can

Many women report feeling that being fit made labor easier. A number of studies have confirmed this, but some have found no correlation.

Exercise and Gestational Diabetes

Gestational diabetes mellitus (GDM) is the most common complication of pregnancy, affecting 3% to 6% of pregnant Americans.¹ It is the consequence of hormone-related insulin resistance. As in type 2 diabetes, exercise has a role in treatment and, perhaps, prevention.

The American Diabetes Association (ADA) has, in fact, endorsed exercise as "a helpful adjunctive therapy" for GDM when euglycemia is not effected by diet alone.² One study³ cited in an ADA report compared GDM patients who did aerobic arm exercise with patients who used diet alone, and found that glycemic levels began to diverge by the fourth week, primarily because of the low-intensity exercise regimen. By week 6, hemoglobin A_{1c}, fasting plasma glucose, and response to glucose challenge had normalized in the exercising group, while no improvement had occurred among the controls.

Another randomized study,⁴ which used a higher-intensity exercise regimen, resulted in normoglycemia within 1 week. The protocol used in this study is currently implemented in clinic settings.

Women who have GDM must take particu-

lar precautions with exercise, including monitoring blood glucose, regulating meal times, scheduling periods of rest, and carefully tracking fetal activity and uterine contractions.

Whether exercise can prevent GDM in populations at risk has been the subject of limited investigation. One study⁵ showed a significant reduction in risk among exercising women who were morbidly obese before pregnancy (body mass index above 33), but no difference among those who weighed less.

References

1. Artal R: Exercise: an alternative therapy for gestational diabetes. *Phys Sportsmed* 1996;24(3):54-66
2. Jovanovic L: American Diabetes Association's Fourth International Workshop-Conference on gestational diabetes mellitus: summary and discussion: therapeutic interventions. *Diabetes Care* 1998;21(suppl 2):B131-B137
3. Jovanovic Peterson L, Durak EP, Peterson CM: Randomized trial of diet versus diet plus cardiovascular conditioning on glucose levels in gestational diabetes. *Am J Obstet Gynecol* 1989;161(2):415-419
4. Bung P, Artal R, Khodiguian N, et al: Exercise in gestational diabetes: an optional therapeutic approach? *Diabetes* 1991;40(suppl 2):182-185
5. Dye TD, Knox KL, Artal R, et al: Physical activity, obesity, and diabetes in pregnancy. *Am J Epidemiol* 1997;146(11):961-965

continue to participate until late in pregnancy, those who are simply trying to maintain fitness (as opposed to competing) are well advised to reduce their running time to under 45 minutes.

Weight training. Until recently, weight training during pregnancy was unheard of. Now, many women want to continue their established routine and should be instructed in how to do so safely. Relatively light weights and moderate repetitions will maintain flexibility and muscle tone while minimizing the risk of ligament and joint injury. Lifting of heavy weights should be avoided, except under proper prescription and supervision.

General precautions. All pregnant women, regardless of their activity or fitness level, should take precautions against dehydration and hyper-

thermia. They should avoid exertion in hot, humid environments and should drink adequate fluids before, during, and after exercise. Appropriate clothing and other strategies (eg, using a small fan while using an exercise cycle) will facilitate heat dissipation. This is especially important in the first trimester. Prolonged use of hot tubs and whirlpools may lead to episodes of hypotension, which could be detrimental to the fetus.

To avoid compromising fetal growth, caloric intake must be adequate to offset the combined demands of pregnancy (an additional 300 kcal/day) and exercise.

Except in the case of competitive athletes who are determined to maintain high performance for as long as possible (see "Pregnant Athletes Test the Limits," page 54), I counsel pa-

continued

tients to avoid strenuous activity in favor of moderation; in other words, they should be able to converse while working out. During pregnancy, there are no known added benefits, and more potential complications, with vigorous exertion.

Oxygen cost during pregnancy is higher, limiting the ability to perform high-intensity tasks. Regular exercisers may therefore need to be advised to heed signs of fatigue and curtail workouts short of exhaustion to reduce the potential risk of hypoglycemia. Sessions for recreational athletes should generally be limited to 45 minutes.

Warning signs. Pregnant women should be alerted to signs that should prompt them to stop exercising and seek medical attention. In addition to excessive fatigue, these include pain (particularly in the back or pubic area), dizziness, shortness of breath, palpitations, decreased fetal movement, persistent contractions,¹³ rupture of membranes, and vaginal bleeding.

Contraindications. Some women should not exercise during pregnancy. According to the ACOG guidelines,³ careful evaluation should be done to determine if exercise is appropriate for pregnant women who have cardiac disease or constrictive lung disease. Second- or third-trimester bleeding, pregnancy-induced hypertension, preterm labor with present or previous pregnancies, intrauterine growth retardation, incompetent cervix, placenta previa, and premature rupture of membranes are absolute contraindications to exercise.³

Patients with multiple gestation are at signifi-

cant risk for premature labor, and, if they have additional risk factors, the aggregate risk is even higher. Exercise programs for women with multiple gestation should therefore be individualized and medically supervised.

Women who are excessively obese or significantly underweight should exercise under supervision and with extreme caution, if at all, as should those who have type 1 diabetes, pregnancy-induced hypertension, seizure disorder, or anemia. Women who have gestational diabetes mellitus are often helped by exercise, and such programs have been endorsed by the American Diabetes Association and the Second and Third International Workshop-Conference on gestational diabetes.³⁰ (See "Exercise and Gestational Diabetes," page 58.)

Monitor and Encourage

Although risks are minimal with moderation, even healthy, active women should be examined periodically to assess the effect of their exercise programs on the developing fetus, and their regimen should be adjusted or discontinued if necessary. Nonetheless, the consensus seems clear: In the absence of pregnancy complications, exercise has minimal risks and demonstrated benefits. **FSM**

Address correspondence to Raul Artal, MD, Professor and Chairman, St Louis University School of Medicine, Dept of Ob/Gyn, 1031 Bellevue Ave, Suite 290, St Louis, MO 63117.

References

1. Artal R, Gardin K: Historical perspectives, in Artal R, Wiswell RA (eds): *Exercise in Pregnancy*. Baltimore, Williams & Wilkins, 1986, pp 1-6
2. ACOG Home Exercise Programs: *Exercise During Pregnancy and the Postnatal Period*. Washington DC, American College of Obstetricians and Gynecologists, 1985
3. Exercise during pregnancy and the postpartum period. ACOG Technical Bulletin Number 189—February 1994. *Int J Gynaecol Obstet* 1994;45(1):65-70
4. Clapp JF III: Exercise in pregnancy: good, bad or indifferent? in *Current Obstetric Medicine*. St Louis, Mosby Year Book, 1991, pp 25-49, vol 2
5. Sternfeld B: Physical activity and pregnancy outcome: review and recommendations. *Sports Med* 1997;23(1):33-47
6. Stevenson L: Exercise in pregnancy: part 1: update on pathophysiology. *Can Fam Physician* 1997;43(1):97-104
7. Araujo D: Expecting questions about exercise and pregnancy? *Phys Sportsmed* 1997;25(4):85-93
8. Artal R, Rutherford S, Romem Y, et al: Fetal heart rate responses to maternal exercise. *Am J Obstet Gynecol* 1986;155(4):729-733
9. Zamudio S, Palmer SK, Droma T, et al: Effect of altitude on uterine artery blood flow during normal pregnancy. *J Appl Physiol* 1995;79(1):7-14
10. Huch R: Physical activity at altitude in pregnancy. *Semin Perinatol* 1996;20(4):303-314
11. Artal R, Fortunato V, Welton A, et al: A comparison of cardiopulmonary adaptations to exercise in preg-

continued on page 75

MRI is also helpful in diagnosing ligamentous injury (such as to the ACL), which, in one study,¹⁰ occurred in 38% of patients (6 of 16) who had femoral physeal fractures. MRI provides excellent visualization of the knee ligaments and may help confirm the extent of these associated injuries when physical exam findings are equivocal.

Treatment. Treatment for a nondisplaced distal femoral physeal fracture should include immobilizing the knee in a long leg cast, prescribing non-weight-bearing use of crutches, and following up with weekly radiographs to assess for fracture displacement. If displacement is suspected or diagnosed, the patient should be referred to an orthopedic specialist. **FSM**

Address correspondence to Kenneth R. Veenema, MD, University Sports Medicine, 2180 South Clinton Avenue, Rochester, New York 14618.

References

1. Mann DC, Rajmaira S: Distribution of physeal and nonphyseal fractures in 2,650 long-bone fractures in children aged 0-16 years. *J Pediatr Orthop* 1990;10(6):713-716
2. Pritchett JW: Longitudinal growth and growth-plate activity in the lower extremity. *Clin Orthop* 1992; Feb(275):274-279
3. Lombardo SJ, Harvey JP Jr: Fractures of the distal femoral epiphyses. Factors influencing prognosis: a review of 34 cases. *J Bone Joint Surg (Am)* 1977;59(6):742-751
4. Beaty JH, Kumar A: Fractures about the knee in children. *J Bone Joint Surg (Am)* 1994;76(12):1870-1880
5. Thomson JD, Stricker SJ, Williams MM: Fractures of the distal femoral epiphyseal plate. *J Pediatr Orthop* 1995;15(4):474-478
6. Simpson WC Jr, Fardon DF: Obscure distal femoral epiphyseal injury. *South Med J* 1976;69(10):1338-1340
7. Torg JS, Pavlov H, Morris VB: Salter-Harris type-III fracture of the medial femoral condyle occurring in the adolescent athlete. *J Bone Joint Surg (Am)* 1981; 63(4):586-591
8. Smith BG, Rand F, Jaramillo D, et al: Early MR imaging of lower-extremity physeal fracture-separations: a preliminary report. *J Pediatr Orthop* 1994;14(4): 526-533
9. Jaramillo D, Hoffer FA, Shapiro F, et al: MR imaging of fractures of the growth plate. *AJR Am J Roentgenol* 1990;155(6):1261-1265
10. Bertin KC, Goble EM: Ligament injuries associated with physeal fractures about the knee. *Clin Orthop* 1983;Jul(177):188-195

- nancy at sea level and altitude. *Am J Obstet Gynecol* 1995;172(4 pt 1):1170-1180
12. Karzel RP Jr, Friedman MJ: Orthopedic injuries in pregnancy, in Artal R, Wiswell RA, Drinkwater BL (eds): *Exercise in pregnancy*, ed 2. Baltimore, Williams & Wilkins, 1991, pp 123-132
13. Stevenson L: Exercise in pregnancy: part 2: recommendations for individuals. *Can Fam Physician* 1997;43(1):107-111
14. Wang TW, Apgar BS: Exercise during pregnancy. *Am Fam Physician* 1998;57(8):1846-1852
15. Artal R: Hormonal responses to exercise in pregnancy, in Artal R, Wiswell RA, Drinkwater BL (eds): *Exercise in pregnancy*, ed 2. Baltimore, Williams & Wilkins, 1991, pp 175-184
16. Soultanakis HN, Artal R, Wiswell RA: Prolonged exercise in pregnancy: glucose homeostasis, ventilatory and cardiovascular responses. *Semin Perinatol* 1996;20(4):315-327
17. Calguneri M, Bird HA, Wright V: Changes in joint laxity occurring during pregnancy. *Ann Rheum Dis* 1982;41(2):126-128
18. Clapp JF III, Capeless EL: Neonatal morphometrics after endurance exercise during pregnancy. *Am J Obstet Gynecol* 1990;163(6 pt 1):1805-1811
19. Sternfeld B, Quesenberry CP Jr, Eskenazi B, et al: Exercise during pregnancy and pregnancy outcome. *Med Sci Sports Exerc* 1995;27(5):634-640
20. Hatch MC, Stein ZA: Work and exercise during pregnancy: epidemiological studies, in Artal R, Wiswell RA, Drinkwater BL (eds): *Exercise in pregnancy*, ed 2. Baltimore, Williams & Wilkins, 1991, pp 279-286
21. Pivarnik JM: Potential effects of maternal physical activity on birth weight: brief review. *Med Sci Sports Exerc* 1998;30(3):400-406
22. Misra DP, Strobino DM, Stashinko EE, et al: Effects of physical activity on preterm birth. *Am J Epidemiol* 1998;147(7):628-635
23. Teitelman AM, Welch LS, Hellenbrand KG, et al: Effect of maternal work activity on preterm birth and low birth weight. *Am J Epidemiol* 1990;131(1):104-113
24. Hatch M, Levin B, Shu XO, et al: Maternal leisure-time exercise and timely delivery. *Am J Public Health* 1998;88(10):1528-1533
25. Wallace AM, Boyer DB, Dan A, et al: Aerobic exercise, maternal self-esteem, and physical discomforts during pregnancy. *J Nurse Midwifery* 1986;31(6): 255-262
26. Clapp JF III, Rokey R, Treadway JL, et al: Exercise in pregnancy. *Med Sci Sports Exerc* 1992;24(6 suppl): S294-S300
27. Wolfe LA, Hall P, Webb KA, et al: Prescription of aerobic exercise during pregnancy. *Sports Med* 1989;8(5):273-301
28. Clapp JF III: The course of labor after endurance exercise during pregnancy. *Am J Obstet Gynecol* 1990; 163(6 pt 1):1799-1805
29. Pomerance JJ, Gluck L, Lynch VA: Physical fitness in pregnancy: its effect on pregnancy outcome. *Am J Obstet Gynecol* 1974;119(7):867-876
30. Artal R, Bellman D, Deckert T, et al: Summary and recommendations of the Second International Workshop-Conference on gestational diabetes mellitus: therapeutic strategies. *Diabetes* 1985;34(2):125