

Statistical Properties of Electron Curtain Precipitation Derived with AeroCube-6

M. Shumko¹, A.T. Johnson¹, T.P. O'Brien², D.L. Turner³, J.G. Sample¹, J.B.
Blake², L.W. Blum⁴, A.J. Halford⁴

¹Department of Physics, Montana State University, Bozeman, Montana, USA

²Space Science Applications Laboratory, The Aerospace Corporation, El Segundo, California USA

³Johns Hopkins Applied Physics Laboratory, Laurel, Maryland, USA

⁴NASA's Goddard Space Flight Center, Greenbelt, Maryland, USA

Key Points:

- We used the dual AeroCube-6 CubeSats to identify stationary, narrow, and persistent > 30 keV precipitation in low Earth orbit
- 90% of curtains observed are narrower than 21 kilometers in latitude
- A few curtains persistently scattered into the atmosphere for at least six seconds

Abstract

Curtains are stationary, persistent, and narrow in latitude electron precipitation phenomena recently discovered in low Earth orbit over sequential passes of the dual AeroCube-6 CubeSats. Curtains with electron energies > 30 keV were stationary over a variety of spacecraft separations, and observed by the follower spacecraft for up to 65 seconds after the leader. This study expands the recent curtain discovery and quantifies statistical properties of 1634 curtains observed over three years. We found that in low Earth orbit, many curtains are narrower than 10 kilometers in latitude and 90% are less than 21 kilometers wide. We also found that curtains are an outer radiation belt phenomena that are observed in the late morning and midnight magnetic local time, with a higher occurrence rate at midnight. Furthermore curtains are observed more often at lower geomagnetic activity than microbursts. We compare every statistical result to microbursts to test the hypothesis that curtains are drifting remnants of microbursts. Lastly, we found a few curtains in the bounce loss cone region in the north Atlantic Ocean where particle drift motion is impossible. In one such example, a curtain was continuously scattered for at least six seconds so curtains can be a significant source of > 30 keV electrons into the atmosphere.

1 Plain Language Summary

2 Introduction

Curtains are a stationary electron precipitation phenomena observed in low Earth orbit (LEO). They are narrow in latitude and appear stationary for up to a minute between subsequent satellite passes. Blake and O'Brien (2016) recently discovered curtains with the > 30 keV electron dosimeters onboard the dual AeroCube-6 (AC6) CubeSats that operated together between 2014 and 2017. This discovery was possible due to AC6's actively maintained in-track separation between a few hundred meters and a few hundred kilometers. Besides the Blake and O'Brien (2016) discovery study not much is known about curtains including what they are, how are they generated, their statistical properties, and their impact on the atmosphere. Answering these questions is an essential next step towards a more complete understanding of how curtains, and particle precipitation in general, affect the magnetosphere and Earth.

In low Earth orbit (LEO) curtains are narrower than a few tens of kilometers in latitude so a LEO satellite, such as AC6, will pass through them in a second or less. Curtains appear spiky in the electron data. AC6 also observes similar-looking transient precipitation called electron microbursts. Both microbursts and curtains appear appear spiky in the AC6 data for different reasons: microbursts primarily for being temporally short, and curtains primarily for being narrow in latitude. Hence AC6, and other recently developed multi-spacecraft missions, are necessary to identify and distinguish between curtains and microbursts.

Microbursts have been observed since mid 1960s by high altitude balloons and satellites and are also a spiky increase of electrons shorter than a second but are not spatially stationary (e.g. Anderson & Milton, 1964; Lorentzen, Blake, et al., 2001; O'Brien et al., 2003; Douma et al., 2017). The companion study to this work by Shumko et al. (2019) calculated the spatial size of microbursts using simultaneous observations of microbursts observe by AC6. The impact of microbursts on the environment is substantial. Lorentzen, Looper, and Blake (2001), Thorne et al. (2005), Breneman et al. (2017), and Douma et al. (2019)—among others—estimated that microbursts can deplete the outer radiation belt electrons in about a day. Furthermore, Seppälä et al. (2018) modeled a 6 hour microburst storm and concluded that microbursts depleted mesospheric ozone by roughly 10%. Thus, it is important to understand the connection, if any, between microbursts and curtains. Curtains and microbursts can be easily misidentified from a single spacecraft so we need to reevaluate single-satellite microburst studies. If curtains are numer-

ous then the estimated microburst occurrence rates are overestimated. Furthermore, the microburst impact on the atmosphere and the outer radiation belt is overestimated.

Blake and O’Brien (2016) proposed the following hypothesis that explains the curtain-microburst relationship. If a microburst is not completely lost in the atmosphere after the initial scatter, the remaining microburst electrons will spread out (bounce phase disperse) along the entire magnetic field line over a few bounce periods. Concurrently these electrons drift to the east, with higher energy electrons drifting at a faster rate. Therefore, if this hypothesis is true, the initially localized microburst is spread out in longitude into the shape of a curtain. The idea of curtains is not entirely new, and Lehtinen et al. (2000) presented a different hypothesis that curtains can be created by energetic runaway beams driven by lightning.

The AC6 dosimeters lack the necessary pitch angle resolution to differentiate between drifting and precipitating electrons to test the Blake and O’Brien (2016) hypothesis. Instead we use AC6’s position and Earth’s asymmetric magnetic field to differentiate particles that are either nearly-trapped in the drift loss cone, and particles immediately lost in the bounce loss cone (BLC). These concepts are described below.

This study expands the Blake and O’Brien (2016) study by estimating statistical properties of curtains. We use 1634 confirmed curtain observations to study the distributions of: the curtain width in latitude, the geomagnetic conditions favorable to curtains, and curtain distribution in L and magnetic local time (MLT). Lastly we will address the hypothesis that curtains are drifting remnants of microbursts by showing examples of curtains observed in the BLC region.

3 Instrumentation

The AC6 mission was a pair of 0.5U (10x10x5 cm) CubeSats built by The Aerospace Corporation designed to measure the electron and proton environment in low Earth orbit (O’Brien et al., 2016). AC6 was launched on 19 June 2014 into a 620x700 km, 98° inclination orbit. The AC6 orbit over the three year mission lifetime was roughly dawn-dusk, and precessed only a few hours in MLT; 8-12 MLT in dawn and 20-24 MLT in dusk. The two AC6 spacecraft, designated as AC6-A and AC6-B, separated after launch and were in proximity for the duration of the three year mission—maintained by an active attitude control system. The attitude control system allowed them to precisely control the amount of atmospheric drag experienced by each AC6 unit using the surface area of their solar panel “wings”. By changing their orientation, AC6 was able to maintain a separation between 2-800 km, confirmed with the Global Positioning System. The two AC6 units were in a string of pearls configuration so one unit, typically unit A, was leading the other by an in-track lag—the time it would take the following spacecraft to catch up to the position of the leading spacecraft. To convert between the AC6 in-track separation and in-track lag, we use a typical 7.5 km/s LEO velocity. The in-track lag was readily available with the Global Positioning System which makes it easy to study precipitation phenomena observed at the same time, and at the same position by shifting one time series by the in-track lag.

Each AC6 unit contains three Aerospace microdosimeters (licensed to Teledyne Microelectronics, Inc) that measure the electron and proton dose in orbit (O’Brien et al., 2016). The dosimeter used for this study is dos1 with a 30 keV electron threshold. dos1 is used for this study because the other dosimeters either responded primarily to protons or were not identical between unit A and B. All dosimeters sample at 1 Hz in survey mode, and 10 Hz in burst mode. 10 Hz data was readily available from both AC6 units from June 2014 to May 2017 while their in-track lag was less than 65 seconds, and at times was a fraction of a second. **Show a distribution of the in-track lag when they had 10 Hz data?** The variety of AC6 separations and data availability over the three-

year mission makes it possible to study transient electron microburst precipitation (Shumko et al., 2019) and now stationary electron curtain precipitation.

4 Methodology

4.1 Curtain Identification

The 10 Hz data was used to identify curtains with the following two criteria: a high spatial correlation, and spiky. The first criterion quantifies the similarity of the feature between both AC6 units, and the second criterion checks that highly correlated times were spiky. Before we applied the identification criteria, the AC6-B time series was shifted by the in-track lag to spatially align it with the AC6-A time series.

The first identification criterion is a 1-second rolling Pearson correlation applied to both time series. Spatial features with a correlation greater than 0.8 are considered highly correlated.

The second identification criterion, applied to the highly correlated locations identified by the first criterion, identifies locations where both AC6 units observed spiky precipitation. Similar to how precipitation bands were identified by Blum et al. (2015) and microbursts by Greeley et al. (2019), we find spiky precipitation by quantifying the number of Poisson standard deviations, σ , that a *dos1* count rate is above a 10-second centered running average, B_{10} . Locations where *dos1* is at least two σ above B_{10} , in other words $\text{dos1} > 2\sqrt{B_{10}} + B_{10}$, are spiky.

The locations where the two criteria are met are curtain candidates and the time of the peak count rates are saved. We tuned the detection parameters to have as many candidate curtains as possible while being feasible to inspect every detection. To check the quality of the data set, one author visually checked every candidate curtain and 1634 curtains were confirmed. Four curtain examples are shown in Fig. 1. In each example, the unmodified time series is shown in the top row and the spatially-aligned time series the bottom row. The in-track lag used to shift the bottom row is annotated by dt , corresponding to an AC6 in-track separation annotated by s . The top row is uncorrelated so these events were not microbursts. The bottom row is correlated at the same location after 3 to 26 seconds. The correlated curtains in the bottom row are peculiar—their fine width is on a 10-kilometer scale and was observed to persist for at least 26 seconds.

4.2 Differentiating Between Drifting and Precipitating Curtains

As mentioned in the introduction, AC6 lacks pitch angle resolution that would allow it to easily differentiate between drifting and precipitating curtain electrons. Fortunately we can use the South Atlantic Anomaly (SAA) and AC6's location in LEO to differentiate between the two possibilities.

Earth's magnetic field is spatially shifted towards Singapore which creates a region of weaker magnetic field in the South Atlantic Ocean called the South Atlantic Anomaly. Most particles observed in LEO are barely-trapped: they bounce and drift around the Earth until they reach the SAA. In the SAA the weaker magnetic field strength lowers the electron's mirror point into the atmosphere where collisions with the atmosphere are more numerous and the particle is lost. Particles that drift and are lost in the SAA have pitch angles in the drift loss cone. Particles with small enough pitch angles to be lost in the atmosphere within one bounce are in the BLC. Traditionally, we define a particle with a mirror point at or below and altitude of 100 km to be in the BLC. In this study we will use this definition of the BLC as well as a more strict definition of a mirror point below sea level.




Figure 1. Four examples showing the AC6 > 30 keV electron data taken by AC6 at the same time in the top row and at the same position in the bottom row. AC6-A, whose data is shown with red curves, was s kilometers ahead of AC6-B. To show the data at the same position the time series data from one spacecraft was shifted by the in-track lag and annotated by dt . These examples show that curtain precipitation was highly correlated for up to 26 seconds.

Without pitch angle resolution AC6 can not easily differentiate between trapped, drift loss cone, and bounce loss cone electrons. In the SAA, AC6 observes a combination of trapped, drift loss cone, and BLC electrons. In the region conjugate to the SAA in the North Atlantic, AC6 only observes particles in the BLC. A particle is in the BLC if its conjugate mirror point is below 100 km. **Too much hand holding or appropriate?** If an electron makes it to AC6's altitude, it can be in the local loss cone and precipitate in the local hemisphere. Alternatively, the electron will mirror at or below AC6 and gyrate into the SAA where the equivalent mirror point magnetic field strength is deep in the atmosphere or below sea level. Therefore, any precipitation observed in the BLC region in the North Atlantic must rapidly precipitate. Lastly, in all other regions in LEO AC6 will observe electrons in both the drift and bounce loss cones.

5 Results

In the spirit of brevity, we limited the scope of this statistical study to answer three questions:

1. How narrow are curtains?
2. When and where are curtains observed?
3. Are curtains drifting or locally precipitating?

For each of these questions we will compare the curtain distribution with the > 30 keV microburst distribution from Shumko et al. (2019). Lastly, we will show evidence that suggests that some curtains are continuously scattered and not drifting.

5.1 Curtain Width

We quantified curtain width in time as the width at half of the curtain's topographic prominence: the height of the peak above the lowest contour that encircles the peak but contains no higher peak. The spatial width of a curtain is the product of the temporal width and the 7.5 km/s orbital velocity mostly in latitude. The distribution of curtain widths is shown in Fig. 2 by the thick black curve. Curtains are very narrow in latitude. Many curtains are less than 10 km wide, and 90% are narrower than 21 km.

We compared the curtain width distribution to the microburst size distribution estimated in Shumko et al. (2019). Shumko et al. (2019) estimated the microburst size distribution by finding microbursts that were observed simultaneously by both AC6 units so the microburst size much be larger than the AC6 separation. The red curve in Fig. 2 shows the microburst distribution estimated from the ratio of the number of concurrent and nonconcurrent microbursts observed in each separation bin.

The curtain and microburst size distributions are very similar with one notable difference. Curtain widths are observed directly and measure the entire region of instability hence their distribution is peaked, while the microburst distribution is estimated from concurrent observations that sample a sub-region of a microburst so their distribution is peaked at 0 km AC6 separation.

5.2 When and Where Are Curtains Observed

The distribution of curtains in L and MLT is shown in Fig. 3. Figure 3a shows the distribution of the 1634 observed curtains and Fig. 3b shows the normalized distribution. The white bins, mostly in the early morning and evening MLT regions, had no observed curtains because the AC6 orbit did not sample there. Figure 3c confirms this by showing the distribution of the number of quality (data quality flag = 0) 10 Hz samples.

After scaling by the uneven sampling in the late morning and midnight MLT regions, the normalized distribution in Fig. 3b shows enhanced curtain occurrence in the outer radiation belt in late morning and midnight MLT regions.

Now we quantify the geomagnetic conditions favorable for curtains. Figure 4 shows the distribution of the Auroral Electroject (AE) index between 2014 and 2017, and the distribution of the AE index when curtains and microbursts were observed. The distribution of the AE index when > 30 keV microbursts were observed is shown with a dashed green curve. Curtains are observed typically when AE is enhanced. Microbursts are observed when AE is enhanced more than for curtains.

5.3 Local Atmospheric Precipitation

The evidence presented so far hint at, but not directly confirm that, curtains are connected to microbursts. But a few curtains observed in the bounce loss cone put the hypothesis into question. We estimated the BLC region in the North Atlantic Ocean using the IRBEM-Lib magnetic field library (Boscher et al., 2012). We defined a latitude-longitude grid spanning the North Atlantic, at 700 kilometer altitude that is typical for AC6, and estimated the local magnetic field strength. For each latitude-longitude point we traced the magnetic field line to the SAA and found the position along that field line with an field strength equivalent to the local field strength. Thus for each grid location a locally-mirroring particle will gyrate to the southern conjugate mirror point. If this mirror point is at or below 100 km, the latitude-longitude grid point is in the bounce loss cone. A particle could have a different pitch angle, but because it was observed by AC6, its pitch angle must be small enough and mirror at or below 100 km in the SAA. It is reasonable that the electron will have an even smaller pitch angle and mirror deeper in the atmosphere. Lastly, we saved the grid locations where the conjugate mirror point altitude is below sea level.

Since we are considering particles mirroring at the grid points in North Atlantic, this estimate is the upper bound altitude that the particle can mirror. Particles mirroring above the grid will not be observed, and the particles that are observed will mirror at or below the estimated altitude in the SAA. If the corresponding field strength position is at 100 kilometers or lower in the SAA we considered that grid point in the North Atlantic in the BLC region. For the more rigorous BLC criteria, we saved the grid locations where the equivalent field strength altitude in the SAA is below sea level.




Figure 2. Distribution of curtain width in latitude shown in black and sizes of microbursts when they were simultaneously observed by AC6 is shown in red. Microburst distribution adopted from Shumko et al. (2019). The error bars represent the Poisson standard error.




Figure 3. Distribution of observed curtains as a function of L and MLT in panel a and normalized number of curtains in panel b. Panel c shows the number of quality 10 Hz samples taken at the same time that was used to normalize panel b. The white bins in panels a have 0 curtain observations. In panel b the white shaded bins had either 0 detections or very little 10 Hz samples (less than 10,000).



Figure 4. The distribution of the Auroral Electrojet (AE) index from 2014 to 2017. The black curve shows the distribution for the entire 2014-2017 AE data set, the dotted blue curve shows the AE distribution for curtains, and the dashed green curve shows AE the distribution for microbursts studied in Shumko et al. (2019).

Figure 5a shows a map of the BLC region in the North Atlantic. The solid blue line is the north bound of the region where a particle observed at 700 kilometers at that location will mirror at 100 kilometers in the SAA. Immediately south of the solid blue line the SAA mirror altitude rapidly decreases towards, and below, sea level. The dashed blue line is the set of latitude-longitude points where the SAA mirror point altitude is at sea level. AC6 takes about 30 seconds to move between the solid and dashed blue curves. We superposed two dotted black lines on Fig. 5a that represent the boundary of the outer radiation belt defined between L shells of 4 and 8. The BLC region here closely matches the region shown in Comess et al. (2013, Figure 1) and Dietrich et al. (2010, Figure 3). We used the Olson-Pfitzer magnetic field model (Olson & Pfitzer, 1982) to estimate the BLC boundary. The same analysis using the Tsyganenko 1989 model (Tsyganenko, 1989) yielded similar boundaries.

We found 4 good curtains that were observed inside the BLC regions and included the shifted time series plots in Fig. 5b-e, with the AC6 in-track lag, L and MLT of the observations annotated. The AC6 locations where these curtains were observed are shown in 5a with red stars and the corresponding panel labels. The curtains shown in Fig. 5c and e were observed near the sea level SAA mirror altitude curve thus they were not drifting and were precipitating as much as 6 seconds as shown in Fig. 5e. For reference, the precipitation persisted for longer than the typical ≈ 1.5 second bounce period of 30 keV electrons in this region.

These examples were observed near the sea level BLC curve, so at a minimum, those electrons would have gyrated down to sea level in the South Atlantic Anomaly. This is a very conservative estimate because the observed electrons likely mirrored below AC6 and thus below sea level (if Earth was not in the way). Figure 3 casts doubt on the microburst-curtain hypothesis but more work will be necessary to fully conclude if microbursts and curtains are related. These examples were observed near the sea level BLC curve, so at a minimum, those electrons would have gyrated down to sea level in the South Atlantic Anomaly. This is a very conservative estimate because the observed electrons likely mirrored below AC6 and thus below sea level (if Earth was not in the way). Figure 3 casts doubt on the microburst-curtain hypothesis but more work will be necessary to fully conclude if microbursts and curtains are related.

6 Discussion

6.1 Curtain Width In Latitude

Curtains are very narrow in latitude. Figure 2 shows the width in latitude of many curtains are on the order of 10 kilometers and 90% are narrower than 21 km. Scaled to the magnetic equator, where we presume curtains are generated, these latitudinal widths correspond to a source with a radial scale size of a few hundred kilometers. As shown in Fig. 1f and 1h; it is remarkable that some curtains maintain a fine structure after multiple seconds with little observable difference. Sometimes curtains appear to be slightly and systematically shifted in latitude, while maintaining their shape (not shown).

If curtains are remnants of microbursts then the distribution of curtain widths in latitude correspond to the microburst size distribution. Figure 2 shows a good correspondence between the distribution of curtain widths in latitude and the microburst size distribution from Shumko et al. (2019). Therefore, it is reasonable to believe that curtains and microbursts are related, but this result needs to be closely inspected for sources of bias.

The microburst scale size distribution, as described in Shumko et al. (2019), is the fraction of microbursts observed simultaneously to all microbursts observed either simultaneously or by only one AC6 unit. A microburst observed simultaneously must be larger than the spacecraft separation so the microburst distribution represents a lower bound.




Figure 5. Curtains observed inside the bounce loss cone region. Panel a shows a map of the North Atlantic region with the outer radiation belt, defined by an L shell range between 4 and 8, shown with the dotted black curves. The solid blue curve shows the northern boundary of the bounce loss cone region. Along this curve, electrons observed at a 700 kilometer altitude in the North Atlantic will mirror at 100 kilometers in the SAA. A more strict bounce loss cone criteria is the dashed blue curve that represents a mirror point altitude at sea level in the SAA. The 4 red stars with labels show the locations of the curtain examples shown in panels b-e. The panels b-e show the 4 example curtains with the AC6-A shown by the red line, and AC6-B with the blue line. AC6-A was ahead in all examples except panel d.

Shumko et al. (2019) attempted to account for this bias but it is difficult. This bias that shifts the microburst distribution to smaller sizes so in the microburst size distribution shown in Fig. 2 is underestimated.

Furthermore the detection algorithm described in section 4.1 has a width bias. For wide curtains with a similar width to the detection algorithm’s 10-second baseline, the 10-second baseline is then calculated using the enhanced curtain counts instead of the background and is increased above the true background and thus the curtain peak is less pronounced relative to the baseline. As a result, the curtain detection algorithm is less sensitive to wider curtains. The result of this bias is similar to the bias inherent in the microburst distribution—the curtain widths are underestimated.

Both of these biases underestimate the true size of curtains and microbursts, so this evidence agrees with the Blake and O’Brien (2016) microburst-curtain hypothesis.

6.2 When and Where Are Curtains Observed

Figure 3 shows that curtain phenomena originates in the outer radiation belt, and observed relatively more in the evening than morning regions. Unfortunately, the limited AC6 coverage prevents a complete curtain distribution in MLT. From the MLT information we have in Fig. 3b, curtains are most often observed near midnight MLT with relatively less observed in the late morning MLT. This distribution, though limited, appears to be similar to the L-MLT distribution of microbursts from prior studies (e.g. O’Brien et al., 2003; Douma et al., 2017).

Curtains also exhibit a preference to disturbed geomagnetic conditions as shown in Fig. 4, but not as disturbed as microbursts from Shumko et al. (2019). A possible explanation: during quiet conditions the remnant microburst electrons are more likely to drift undisturbed and AC6 is more likely to observe the fine, highly-correlated curtain structure. In contrast, during active conditions curtain electrons are still drifting, but the dynamics of an actively-changing magnetosphere can easily perturb curtain electrons until AC6 no longer observes a highly correlated structure at the same location.

6.3 Curtains Observed in The Bounce Loss Cone

Lastly we address curtains observed in the bounce loss cone. What can cause continuous > 30 keV electron precipitation lasting for multiple seconds? This mechanism must be radially localized near the magnetic equator, on a scale of a few hundred kilometers. A candidate mechanism is a direct current electric field that is parallel to the background magnetic field that lowers the electron mirror point to AC6 altitudes. To find the minimum potential we assume the electron is barely trapped and has a mirror point at 100 kilometers in the SAA (and will likewise mirror above AC6’s altitude in the conjugate point). This condition implies that the particle’s mirror point in the bounce loss cone is above AC6.

To find the parallel potential we use the kinetic energy, W , of a 30 keV particle at its initial mirror point at a magnetic field strength of B_i . The kinetic energy at the initial mirror point can be written as $W_i = \mu B_i$ where μ is the first adiabatic invariant. Now when a parallel potential acts on the electron of charge q and does $q\Phi$ amount of work, the electron will mirror closer to Earth’s surface and mirror at a field strength B_f where its final energy is $W_f = \mu B_f$. Now we relate the initial and final kinetic energy of the electron,

$$\mu B_f = \mu B_i + q\Phi. \quad (1)$$

Then we substitute μ and the above expression simplifies to

$$q\Phi = W \frac{(B_f - B_i)}{B_i}. \quad (2)$$

Also, comment that needed Φ depends proportionally on initial energy, and that AC6 dos1 response increases sharply with energy up to a peak response around 100 keV.

In the aurora papers, do we need to worry about the *sign* of Φ ? In the FAST papers, is it pointing the right way to pull electrons down?

We again use IRBEM-Lib to estimate $q\Phi$. For each example curtain in Fig. 5, we first estimated the local magnetic field, B_f for this derivation. Then we traced the field line from AC6 into the SAA. We found B_i at 100 kilometers altitude in the SAA for barely trapped electrons. With the initial and final B , along with $W = 30$ keV, the minimum potential must be $q\Phi = 1 - 4$ kV. Recalculate the potentials for the four cases.. This range of potentials is typical for the aurora. Partamies et al. (2008) used the observations made by the Fast Auroral SnapshoT (FAST) mission and reported that the inverted-V auroral structures, observed up to a few tens of keV, were accelerated by 2-4 kV potentials. While the aurora and curtains are likely different, the aurora is observed at higher L and near midnight MLT region, they do share a number of similarities. If AC6 had differential energy channels below 30 keV, then it would be possible to test a possible relationship between curtains and inverted-V structures.

Outside of the BLC, the lack of pitch angle information makes the AC6 electron data ambiguous, but the curtains observed in the BLC suggest that some curtains continuously precipitate for multiple seconds. Curtains could be a significant source of energetic particles into the atmosphere. Energetic electron precipitation generates odd Nitrogen (HO_x) that is currently underestimated by atmospheric models such as the widely-used Whole Atmosphere Community Climate Model using Specified Dynamics (SD-WACCM) (e.g. Randall et al., 2015). A comprehensive study of the curtain impact on the atmosphere should be done with an AC6-like mission with pitch angle and energy resolution.

7 Conclusions

The 1634 confirmed curtains allowed us to make the following statistical inferences:

1. Curtains are very narrow—90% are less than 21 kilometers wide in latitude.
2. Curtains are observed in the outer radiation belt, predominately in the midnight and the late morning MLT regions, during periods of moderate geomagnetic activity.
3. Some curtains continuously precipitate into the atmosphere for multiple seconds.

Curtains are remarkably narrow with fine structure that persist for multiple seconds. Either the scattering mechanism that contentiously generates curtains is physically static for multiple seconds, or the curtain electron drift is often undisturbed.

The curtain-microburst relationship hypothesized in Blake and O'Brien (2016) is not clear. The two results in support of the hypothesis are: curtain width and microburst size distributions are very similar, and the limited AC6 sampling in MLT shows that both occur in similar locations in the magnetosphere. But the bounce loss cone result complicates this interpretation. Some curtains continuously precipitate for at least a few seconds and can be a significant source of energetic electron precipitation into the atmosphere. Furthermore, the continuous scatter of curtain electrons can be explained by a parallel direct current electric field and could be related to the aurora.

Acknowledgments

This work was made possible with the help from the many engineers and scientists at The Aerospace Corporation who designed, built, and operated AC6. M. Shumko was supported by NASA Headquarters under the NASA Earth and Space Science Fellowship Program - Grant 80NSSC18K1204. D.L. Turner is thankful for support from the Van Allen Probes mission and a NASA grant (Prime award number: 80NSSC19K0280). The work at The Aerospace Corporation was supported in part by RBSP-ECT funding provided by JHU/APL contract 967399 under NASA's Prime contract NAS501072. The AC6 data is available at <http://rbspgway.jhuapl.edu/ac6> and the IRBEM-Lib version used for this analysis can be downloaded from <https://sourceforge.net/p/irbem/code/616/tree/>.

8 Homeless Words

Title: Statistical Properties of Curtains–Latitudinally-Narrow and Persistent Electron Precipitation Phenomena

This study leverages AC6, a multi-spacecraft mission, to interpret and understand particle precipitation in a way that is impossible with a single spacecraft.

This study leverages the asymmetry in Earth's magnetic field. The asymmetric magnetic field results in the SAA and the BLC, two very related and unique regions

Particles that impact the atmosphere are lost during that bounce motion. We found curtains in the bounce loss cone, a region in the North Atlantic near and above Iceland.

The bounce loss cone is magnetically connected to the SAA, where Earth's magnetic field is weakest near Earth's surface. A particle observed in the blc in the northern hemisphere will descend below 100 km altitude. At sub-100 km altitudes the particle has a high chance of encountering and scattering with the atmosphere and be lost.

We found curtain electrons that, when given the chance to execute their cyclical bounce motion, will descend below Earth's surface in the SAA. An electrons can not survive that trip.

Write the paper and ask the question: "What is this paper really about?" Not just curtains, but uncovering something unexpected that has been observed and overlooked for decades.

Are curtains related to aurora? This is a good question—one that is not pertinent here (idea from *The Elements of Style* p.68).

Here are two parting questions that are not considered here. Why were some curtains shifted slightly? Perhaps it was due to the movement of the magnetic field lines. Also do curtains have a corresponding visual signature on the ground? The answer to this question will show if curtains are related to the aurora.

References

- Anderson, K. A., & Milton, D. W. (1964). Balloon observations of X rays in the auroral zone: 3. High time resolution studies. *Journal of Geophysical Research*, 69(21), 4457–4479. Retrieved from <http://dx.doi.org/10.1029/JZ069i021p04457> doi: 10.1029/JZ069i021p04457
- Blake, J. B., & O'Brien, T. P. (2016). Observations of small-scale latitudinal structure in energetic electron precipitation. *Journal of Geophysical Research: Space Physics*, 121(4), 3031–3035. Retrieved from <http://dx.doi.org/10.1002/2015JA021815> (2015JA021815) doi: 10.1002/2015JA021815
- Blum, L., Li, X., & Denton, M. (2015). Rapid MeV electron precipitation as ob-

- served by SAMPEX/HILT during high-speed stream-driven storms. *Journal of Geophysical Research: Space Physics*, 120(5), 3783–3794. Retrieved from <http://dx.doi.org/10.1002/2014JA020633> (2014JA020633) doi: 10.1002/2014JA020633
- Boscher, D., Bourdarie, S., O'Brien, P., Guild, T., & Shumko, M. (2012). *Irbem-lib library*.
- Breneman, A., Crew, A., Sample, J., Klumpar, D., Johnson, A., Agapitov, O., ... others (2017). Observations directly linking relativistic electron microbursts to whistler mode chorus: Van allen probes and FIREBIRD II. *Geophysical Research Letters*.
- Comess, M., Smith, D., Selesnick, R., Millan, R., & Sample, J. (2013). Duskside relativistic electron precipitation as measured by sampex: A statistical survey. *Journal of Geophysical Research: Space Physics*, 118(8), 5050-5058. Retrieved from <https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/jgra.50481> doi: 10.1002/jgra.50481
- Dietrich, S., Rodger, C. J., Clilverd, M. A., Bortnik, J., & Raita, T. (2010). Relativistic microburst storm characteristics: Combined satellite and ground-based observations. *Journal of Geophysical Research: Space Physics*, 115(A12).
- Douma, E., Rodger, C., Blum, L., O'Brien, T., Clilverd, M., & Blake, J. (2019). Characteristics of relativistic microburst intensity from sampex observations. *Journal of Geophysical Research: Space Physics*.
- Douma, E., Rodger, C. J., Blum, L. W., & Clilverd, M. A. (2017). Occurrence characteristics of relativistic electron microbursts from SAMPEX observations. *Journal of Geophysical Research: Space Physics*, 122(8), 8096–8107. Retrieved from <http://dx.doi.org/10.1002/2017JA024067> (2017JA024067) doi: 10.1002/2017JA024067
- Greeley, A., Kanekal, S., Baker, D., Klecker, B., & Schiller, Q. (2019). Quantifying the contribution of microbursts to global electron loss in the radiation belts. *Journal of Geophysical Research: Space Physics*.
- Lehtinen, N. G., Inan, U. S., & Bell, T. F. (2000). Trapped energetic electron curtains produced by thunderstorm driven relativistic runaway electrons. *Geophysical research letters*, 27(8), 1095–1098.
- Lorentzen, K. R., Blake, J. B., Inan, U. S., & Bortnik, J. (2001). Observations of relativistic electron microbursts in association with VLF chorus. *Journal of Geophysical Research: Space Physics*, 106(A4), 6017–6027. Retrieved from <http://dx.doi.org/10.1029/2000JA003018> doi: 10.1029/2000JA003018
- Lorentzen, K. R., Looper, M. D., & Blake, J. B. (2001). Relativistic electron microbursts during the GEM storms. *Geophysical Research Letters*, 28(13), 2573–2576. Retrieved from <http://dx.doi.org/10.1029/2001GL012926> doi: 10.1029/2001GL012926
- O'Brien, T. P., Blake, J. B., & W., G. J. (2016, May). *Aerocube-6 dosimeter data readme* (Tech. Rep. No. TOR-2016-01155). The Aerospace Corporation.
- O'Brien, T. P., Lorentzen, K. R., Mann, I. R., Meredith, N. P., Blake, J. B., Fennell, J. F., ... Anderson, R. R. (2003). Energization of relativistic electrons in the presence of ULF power and MeV microbursts: Evidence for dual ULF and VLF acceleration. *Journal of Geophysical Research: Space Physics*, 108(A8). Retrieved from <http://dx.doi.org/10.1029/2002JA009784> doi: 10.1029/2002JA009784
- Olson, W. P., & Pfitzer, K. A. (1982). A dynamic model of the magnetospheric magnetic and electric fields for july 29, 1977. *Journal of Geophysical Research: Space Physics*, 87(A8), 5943–5948. Retrieved from <http://dx.doi.org/10.1029/JA087iA08p05943> doi: 10.1029/JA087iA08p05943
- Partamies, N., Donovan, E., & Knudsen, D. (2008). Statistical study of inverted-v structures in fast data. In *Annales geophysicae* (Vol. 26, pp. 1439–1449).
- Randall, C. E., Harvey, V. L., Holt, L. A., Marsh, D. R., Kinnison, D., Funke, B.,

- 474 & Bernath, P. F. (2015). Simulation of energetic particle precipitation effects
 475 during the 2003–2004 arctic winter. *Journal of Geophysical Research: Space*
 476 *Physics*, 120(6), 5035–5048.
- 477 Seppälä, A., Douma, E., Rodger, C., Verronen, P., Clilverd, M. A., & Bortnik, J.
 478 (2018). Relativistic electron microburst events: Modeling the atmospheric
 479 impact. *Geophysical Research Letters*, 45(2), 1141–1147.
- 480 Shumko, M., Johnson, A., Sample, J., Griffith, B. A., Turner, D. L., O’Brien, T. P.,
 481 ... Claudepierre, S. G. (2019). Electron microburst size distribution de-
 482 rived with aerocube-6. *Journal of Geophysical Research: Space Physics*,
 483 e2019JA027651.
- 484 Thorne, R. M., O’Brien, T. P., Shprits, Y. Y., Summers, D., & Horne, R. B. (2005).
 485 Timescale for MeV electron microburst loss during geomagnetic storms. *Jour-*
 486 *nal of Geophysical Research: Space Physics*, 110(A9). Retrieved from [http://](http://dx.doi.org/10.1029/2004JA010882)
 487 dx.doi.org/10.1029/2004JA010882 (A09202) doi: 10.1029/2004JA010882
- 488 Tsyganenko, N. (1989). A solution of the chapman-ferraro problem for an el-
 489 lipsoidal magnetopause. *Planetary and Space Science*, 37(9), 1037 - 1046.
 490 Retrieved from [http://www.sciencedirect.com/science/article/pii/](http://www.sciencedirect.com/science/article/pii/0032063389900767)
 491 [0032063389900767](http://www.sciencedirect.com/science/article/pii/0032063389900767) doi: [http://dx.doi.org/10.1016/0032-0633\(89\)90076-7](http://dx.doi.org/10.1016/0032-0633(89)90076-7)