## Microburst Scale Size Distribution Derived with AeroCube-6

M. Shumko<sup>1</sup>, T.P. O'Brien<sup>2</sup>, J. Sample<sup>1</sup>, A. Johnson<sup>1</sup>, D.L. Turner<sup>2</sup>, J.B. Blake<sup>2</sup>, B.A. Griffith<sup>1</sup>, S. Claudepierre<sup>2</sup>, O. Agapitov<sup>3</sup>

 $^1\mathrm{Department}$ of Physics, Montana State University, Bozeman, Montana, USA  $^2\mathrm{Space}$  Science Applications Laboratory, The Aerospace Corportation, El Segundo, California, USA  $^3\mathrm{Space}$  Sciences Laboratory, University of California berkeley, Berkeley, California, USA

## **Key Points:**

11

- Microburst scale size in low Earth orbit and the magnetic equator was estimated.
- Majority of microbursts in low Earth orbit have a scale size on the order of 10 km.
- The majority of microbursts correspond to the correlation scale of high amplitude? whistler-mode chorus waves at the magnetic equator.

Corresponding author: M. Shumko, msshumko@gmail.com

- 13 Abstract
- enter abstract here
- 1 Plain Language Summary
- https://sharingscience.agu.org/creating-plain-language-summary/
- 2 = enter section title =
- 18 Acknowledgments
- Enter acknowledgments, including your data availability statement, here.