

```

\documentclass[GRL,proof,RL]{wb_agu_v2}

%==Extracted from Meta-XML
\hypersetup{
pdftitle={Microburst scale size derived from multiple bounces of a
microburst simultaneously observed with the FIREBIRD-II CubeSats},
pdfauthor={Mykhaylo Shumko, John Sample, Arlo Johnson, J.~Bernard
Blake, Alexander,~B. Crew, Harlan,~E. Spence, David Klumpar, Oleksiy
Agapitov and Matthew,~Lee Handley}, pdfkeywords={10.1029/2018GL078925
and }, pdfsubject={},
}

\artid{57707}
\jid{57707}
\jyear{2018}
\doi{10.1029/2018GL078925}
\jvol{45}
\jissue{0}
\jmonth{7}
\copyeditor{RMAP}
\articletype{Research Letter}

\suppinfo{
\item Supporting Information S1

\papernum{2018GL078925}
\subsetone{Research Letters}
\subettwo{Space Sciences (SSC)}

%\pointsto{This is a companion to [10.1029/2018gl078925].}
%==End Meta-XML

\startpage{1}

\wordcount{1000}
%\draftfalse

%\journalname{Geophysical Research Letters}

%%Note to Pre-editors: Check whether the below supporting info. query
need to be present or not
\AQ{Author: Supporting Information has been supplied with this paper
but no citation is provided in the main text of the article. Please
provide citation.}

%\usepackage{amsrefs}
\begin{document}
\citation{\cname{\author{Shumko, M.}, \author{Sample, J.},
\author{Johnson, A.}, \author{Blake, J.B.}, \author{Crew, A.B.},
\author{Spence, H.E.}, \& \author{others}} (\cyear{2018}).
\ctitle{Microburst scale size derived from multiple bounces of a
microburst simultaneously observed with the FIREBIRD-II CubeSats},
\cjournal{Geophysical Research Letters}, \cvol{45}.)

\ttitle{Microburst Scale Size Derived From Multiple Bounces of a
Microburst Simultaneously Observed With the FIREBIRD-II CubeSats}

```

Deleted: Scale Size Derived

Deleted: Multiple Bounces

Deleted: Microburst Simultaneously Observed

Deleted: {}

Deleted: 2018GL078925-sup-0001-Figure_SI-S01.pdf

Deleted: Scale Size Derived

Deleted: Multiple Bounces

Deleted: Microburst Simultaneously Observed

Deleted: from

Deleted: with

\authormark{SHUMKO ET AL.}
\titlemark{}

\author*[1]{Mykhaylo Shumko}\orcid{0000-0002-0437-7521}\email{msshumko@gmail.com}
\corres{M. Shumko}{}
\author[1]{John Sample}\orcid{0000-0002-9516-9292}
\author[1]{Arlo Johnson}\orcid{0000-0002-7220-1288}
\author[2]{Bern Blake}\orcid{0000-0001-7439-4362}
\author[3]{Alex Crew}\orcid{0000-0002-7493-2682}
\author[4]{Harlan Spence}\orcid{0000-0002-2526-2205}
\author[1]{David Klumpar}
\author[5]{Oleksiy Agapitov}\orcid{0000-0001-6427-1596}
\author[1]{Matthew Handley}\orcid{0000-0001-5341-5072}

\address[1]{\orgdiv{Department of Physics}, \orgname{Montana State University}, \orgaddress{\city{Bozeman}, \state{MT}, \country{USA}}}
\address[2]{\orgdiv{Space Science Applications Laboratory}, \orgname{The Aerospace Corporation}, \orgaddress{\city{Los Angeles}, \state{CA}, \country{USA}}}
\address[3]{\orgname{The Johns Hopkins University Applied Physics Laboratory LLC}, \orgaddress{\city{Laurel}, \state{MD}, \country{USA}}}
\address[4]{\orgdiv{Institute for the Study of Earth, Oceans, and Space}, \orgname{University of New Hampshire}, \orgaddress{\city{Durham}, \state{N.H}, \country{USA}}}
\address[5]{\orgdiv{Space Sciences Laboratory}, \orgaddress{\orgname{UC Berkeley}, \city{Berkeley}, \state{CA}, \country{USA}}}

\abstract{We present the observation of a spatially large microburst with multiple bounces made simultaneously by the [Focused Investigation of Relativistic Electron Bursts: Intensity, Range, and Dynamics II](#) (FIREBIRD-II) CubeSats on 2 February 2015. This is the first observation of a microburst with a subsequent decay made by two [coorbiting](#) but spatially separated spacecraft. From these unique measurements, we place estimates on the lower bounds of the spatial scales as well as quantify the electron bounce periods. The microburst's lower bound latitudinal scale size was 29 ± 1 km and the longitudinal scale size was 51 ± 1 km in low [Earth](#) orbit. We mapped these scale sizes to the magnetic equator and found that the radial and azimuthal scale sizes were at least 500 ± 10 km and 530 ± 10 km, respectively. These lower bound equatorial scale sizes are similar to whistler mode chorus wave source scale sizes, which supports the hypothesis that microbursts are a product of electron scattering by chorus waves. Lastly, we estimated the bounce periods for 200- to 800-keV electrons and found good agreement with four common magnetic field models.

\abstract[Plain Language Summary]{Microbursts are a subsecond impulsive increase of electron precipitation from the outer Van Allen radiation belt into the atmosphere, believed to be an important loss process of radiation belt electrons. Here we present an observation of

Deleted: Montana

Deleted: California

Deleted: Maryland

Deleted: New Hampshire

Deleted: California

Deleted: 2nd,

Deleted: co-orbiting

Deleted: \$

Deleted: \pm 1\$

Deleted: earth

Deleted: \$

Deleted: \$

Deleted: -

Deleted:

a microburst observed simultaneously by the twin [Focused Investigation of Relativistic Electron Bursts: Intensity, Range, and Dynamics](#) II CubeSats. This unique observation allowed us to calculate the microburst's spatial scale size and electron bounce periods. The spatial scale size in low [Earth](#) orbit was found to be a few tens of kilometers in size, one of the largest reported in literature. We then magnetically mapped this scale size to the region near the microburst's generation region and found it to be around 500~km, similar to the spatial scale size of the waves that are believed to be responsible for microburst generation. This observation shows an example of how large microbursts can be, and it sheds light on its scattering mechanism.}

```
\keypoints{
\item Multiple bounces from a microburst were observed by the two
FIREBIRD-II Cube-Sats at LEO,
\item The lower bounds on the microburst scale size at LEO were 29
{plus minus} 1~km (latitudinal)and 51 {plus minus} 11~km
(longitudinal)
\item Deduced lower bound equatorial scale size was similar to the
whistler~mode chorussource scale, }
```

```
\received{25 May 2018}
\revised{28 June 2018}
\accepted{29 June 2018}
```

```
\maketitle
```

```
\section{Introduction}\label{Intro}
The dynamics of radiation belt electrons are complex, and are driven by
competition between source and loss processes. A few possible loss
processes are radial diffusion \cite{Shprits2004}, magnetopause
shadowing \cite{Ukhorskiy2006}, and pitch angle and energy diffusion
due to scattering of electrons by plasma waves \citelist{e.g.,
\cite{Abel1998_1}; \cite{Horne2003}; \cite{Meredith2002};
\cite{Mozer2018}; \cite{Selesnick2003}; \cite{Summers1998};
\cite{Thorne2005}}. There are a variety of waves that cause pitch
angle scattering, including electromagnetic ion cyclotron waves,
plasmaspheric hiss, and chorus \cite{Millan2007, Thorne2010}. Chorus
predominantly occurs in the dawn sector (6--12 magnetic local times,
MLT; Li et al., \citeyear{Li2009}) where it accelerates electrons with
large equatorial pitch angles and scatters electrons with small
equatorial pitch angles \cite{Horne2003}. Some of these electrons may
be impulsively scattered into the loss cone, where they result in
short-duration ( $\sim 100$ ~ms) enhancements in precipitating flux
called microbursts.
```

```
\ocite{Anderson1964} coined the term microburst to describe high-
altitude balloon observations of  $\sim 100$ ~ms duration enhancements of
bremsstrahlung X-rays emitted from scattered microburst electrons
impacting the atmosphere. Since then, nonrelativistic (less than a few
hundred keV) microbursts have been routinely observed with other
balloon missions \citelist{e.g., \cite{Anderson2017};
\cite{Parks1967}; \cite{Woodger2015}}. A review of the literature
shows no reports of microbursts above a few hundreds of
kiloelectronvolts observed by balloons \cite{Millan2002, Woodger2015}.
```

Deleted: FIREBIRD-

Deleted: earth

Deleted:

Deleted: .

Deleted:

Deleted:

Deleted:) .

Deleted: -

Deleted: .

Deleted: ,

Deleted: .

Deleted: -

Deleted: (

Deleted:)) \cite

Deleted: }

Deleted: \$

Deleted:

Deleted:

Deleted: \$

Deleted: non-relativistic

Deleted: .

Deleted: hundred keV

This lack of observation may be explained by relatively weaker pitch angle scattering of relativistic electrons by chorus \cite{Lee2012}.

In addition to the X-ray signature for bursts of electron precipitation, the precipitating relativistic and nonrelativistic electrons have been measured in situ by spacecraft orbiting in low Earth orbit (LEO). Hereinafter, we refer to these electron signatures observed by LEO spacecraft also as microbursts. Microbursts have been observed with, for example, the Solar Anomalous and Magnetospheric Particle Explorer's (SAMPEX) ~ 150 -keV and > 1 -MeV channels \citelist{\cite{Blake1996}; \cite{Blum2015}; Lorentzen, Blake, et al., \citeyear{Lorentzen2001a}; Lorentzen, Looper, \& Blake, \citeyear{Lorentzen2001b}; \cite{Nakamura1995}, \citeyear{Nakamura2000}; \cite{O'Brien2003}, \citeyear{O'Brien2004}} and Focused Investigation of Relativistic Electron Bursts: Intensity, Range, and Dynamics (FIREBIRD-II) with its > 200 -keV energy channels \cite{Anderson2017, Breneman2017, Crew2016}.

Understanding microburst precipitation and its scattering mechanism is important to radiation belt dynamics. The scattering mechanism has been observationally studied by, for example, Lorentzen, Looper, and Blake (\citeyear{Lorentzen2001b}) who found that microbursts and chorus waves predominantly occur in the dawn sector and \ocite{Breneman2017} made a direct observational link between individual microbursts and chorus elements. Microbursts have been modeled and empirically estimated to be capable of depleting the relativistic electron population in the outer radiation belt on the order of a day \cite{Breneman2017, O'Brien2004, Shprits2007, Thorne2005}. An important parameter in this estimation of instantaneous radiation belt electron losses due to microbursts is their scale size. \ocite{Parks1967} used balloon measurements of bremsstrahlung X-rays to estimate the high-altitude scale size of predominantly low-energy microbursts to be 40 ± 14 km. In \ocite{Blake1996} a microburst with multiple bounces was observed by SAMPEX, and the microburst's latitudinal scale size in LEO was estimated to have been at least a few tens of kilometers. \ocite{Blake1996} concluded that typically microbursts are less than a few tens of electron gyroradii in size (at $L = 5$ at LEO, the gyroradii of 1-MeV electrons is on the order of 100 m). \ocite{Dietrich2010} used SAMPEX along with ground-based very low frequency stations to conclude that during one SAMPEX pass, the observed microbursts had scale sizes less than 4 km.

Since 1 February, 2015, microbursts have been observed by FIREBIRD-II, a pair of CubeSats in LEO. Soon after launch, when the two FIREBIRD-II spacecraft were at close range, a microburst with a scale size greater than 11 km was observed \cite{Crew2016}. On the same day, FIREBIRD-II simultaneously observed a microburst with multiple bounces. The microburst decay was observed over a period of a few seconds, while the spacecraft were traveling predominantly in latitude. Here we present the analysis and results of the latitude and longitude scale sizes and bounce periods of the first microburst with multiple bounces observed with the two FIREBIRD-II spacecraft.

\section{Spacecraft and Observation} \label{obs}

The FIREBIRD missions are composed of a pair of identically instrumented 1.5U CubeSats ($15 \times 10 \times 10$ cm) that are

Deleted: non-relativistic

Deleted: earth

Deleted: e.g.

Deleted:) >

Deleted:

Deleted: >

Deleted:

Deleted: Inan, \& Bortnik,

Deleted:); \cite

Deleted:); \cite

Deleted: >

Deleted:

Deleted: e.g. \ocite

Deleted:)

Deleted:

Deleted:

Deleted: \$

Deleted: ``

Deleted: " .

Deleted:

Deleted:

Deleted:

Deleted: 1st,

Deleted:

Deleted: comprised

Deleted: -

Deleted: \$\

Deleted: \$

Deleted: x

Deleted:

designed to measure electron precipitation in LEO \cite{Klumpar2015, Spence2012}. The second mission, termed FIREBIRD-II, was launched on 31 January 2015. The two FIREBIRD-II CubeSats, identified as Flight Unit 3 (FU3) and Flight Unit 4 (FU4), were placed in a 632-km apogee, 433-km perigee, and 99° inclination orbit \cite{Crew2016}. FU3 and FU4 are orbiting in a string of pearls configuration with FU4 ahead, to resolve the space-time ambiguity of microbursts. Each FIREBIRD-II unit has two solid state detectors: one is mounted essentially at the spacecraft surface, covered only by a thin foil acting as a sun shade, with a field of view of 90° (surface detector), and the other is beneath a collimator that restricts the field of view to 54° (collimated detector). Only FU3 has a functioning surface detector, so this analysis utilizes the collimated detectors. FU3's surface and collimated detectors, as well as FU4's collimated detector, observe electron fluxes in six energy channels from ~ 230 keV to > 51 MeV. FIREBIRD-II's High-Resolution (HiRes) electron flux data are gathered with an adjustable sampling period of 18.75 ms by default and can be as fast as 12.5 ms.

On 2 February 2015 at 06:12 UT, both FIREBIRD-II spacecraft simultaneously observed an initial microburst, followed by subsequent periodic electron enhancements of diminishing amplitude shown in Figure~\ref{hires_plot}. This is thought to be the signature of a single burst of electrons, some of which precipitate, but the rest mirror near the spacecraft then bounce to the conjugate hemisphere where they mirror again and the subsequent bounces produce a train of decaying peaks \cite{Blake1996, Thorne2005}. This bounce signature occurred during the transition between the main and recovery phases of a storm with a minimum Dst of -544 nT ($K_p = 4$, and $AE \sim \approx 400$ nT). At this time, the HiRes data were sampled at 18.75 ms. Five peaks were observed by both spacecraft. The fifth peak observed by FU4 was comparable to the Poisson noise and was not used in this analysis. This microburst was observed from the first energy channel (≈ 200 – 300 keV), to the fourth energy channel (≈ 500 – 700 keV), and FU3's surface detector observed the microburst up to the fifth energy channel (683– 950 keV).

```
\begin{figure}
\SPFIG{\framebox[10cm]{Figure 1}}{\caption{High-resolution data of
the microburst observed at 2 February 2015 at 06:12:53 UT, smoothed
with a 150-ms rolling average. The subsequent bounces showed some
energy dispersion. As discussed in the supporting information, a time
correction of  $-2.28$  s was applied to FU3. While the flux from five
energy channels is shown, only channels with reasonable counting
statistics were used for the spatial scale analysis. Vertical colored
bars show the  $\sqrt{N}$  error every 10th data point and vertical
black bars are lined up with the peaks in the 220- to 283-keV energy
channel to help identify dispersion.\label{hires_plot}}}
\end{figure}
```

The HiRes data in Figure~\ref{hires_plot} show signs of energy dispersion, characterized by higher-energy electrons arriving earlier than the lower energies. This time of flight energy dispersion tends to smear out the initial sharp burst upon each subsequent bounce. The first peak does not appear to be dispersed, and subsequent peaks show a dispersion trend consistent across energy channels. The black vertical bars have been added to Figure~\ref{hires_plot} to highlight this energy dispersion. This dispersion signature and amplitude decay

Deleted: 31st

Deleted:

Deleted:

Deleted: which

Deleted:

Deleted: \$

Deleted: \$>

Deleted: \$

Deleted:

Deleted: is

Deleted:

Deleted:

Deleted: 2nd,

Deleted: -

Deleted:

Deleted: (

Deleted: ,

Deleted: \${\

Deleted:)\$

Deleted: was

Deleted:

Deleted:

Deleted: -

Deleted: \$

Deleted:

Deleted: -

Deleted: \$

Deleted: -

Deleted:

Deleted: HiRes

Deleted: 2nd,

Deleted:

Deleted: -

Deleted:

Deleted:

Deleted: shows

Deleted:

implies that the first peak was observed soon after the electrons were scattered, followed by decaying bounces.

At this time, in magnetic coordinates, FIREBIRD-II was at McIlwain $L = 4.7$ and $\text{MLT} = 8.3$, calculated with the Tsyganenko 1989 (T89) magnetic field model \cite{Tsyganenko1989} using IRBEM-Lib \cite{irbem}. Geographically, they were above Sweden, $\text{latitude} = 63^\circ\text{N}$, $\text{longitude} = 15^\circ\text{E}$, $\text{altitude} = 650$ -km. This geographic location is magnetically conjugate to the east of the so-called South Atlantic Anomaly (SAA). The SAA is the location where the mirror points of electrons tend to occur at locations deeper in the atmosphere owing to the offset of the dipole magnetic field from the Earth's center. Electrons with pitch angles within the drift loss cone will encounter the SAA and be removed from their eastward longitudinal drift paths \cite{Comess2013, Dietrich2010}. FU3 and FU4 are therefore both in regions where the particles in the drift loss cone have recently precipitated, leaving only particles that were recently scattered. At the spacecraft location, locally mirroring electrons would have mirrored at 95-km in the opposite hemisphere, with more field-aligned electrons mirroring at even lower altitudes. From the analysis done by \cite{Fang2010}, the peak in the total ionization rate in the atmosphere for 100-keV electrons is around 80-km altitude, while the total ionization rate from 1-MeV electrons peaks around 60-km altitude. It is, therefore, expected that a fraction of the microburst electrons will survive each encounter with the atmosphere. By plotting the peak flux as a function of bounce (not shown), it was found that 40--60\% of the microburst electrons were lost on the first bounce, similar to the 33\% loss per bounce observed for a bouncing microburst observed by SAMPEX \cite{Thorne2005}.

\section{Analysis} \label{analysis}

At the beginning of the FIREBIRD-II mission, two issues prevented the proper analysis of the microburst's spatial scale size: the spacecraft clocks were not synchronized, and their relative positions were not accurately known. We addressed these issues with a cross-correlation time lag analysis described in detail in the supporting information (SI) \cite{sgp4}. From this analysis, the time correction was 2.28 ± 0.12 s (applied to Figure~\ref{hires_plot}) and the separation was 19.9 ± 0.9 km at the time of the microburst observation.

\subsection{Electron Bounce Period} \label{t_b}

We used this unique observation of bouncing electrons to calculate the bounce period, t_b as a function of energy and compare it to the energy-dependent t_b curves derived from four magnetic field models, the results of which are shown in Figure~\ref{tb_plot}. The observed t_b and uncertainties were calculated by fitting the baseline-subtracted HiRes flux. The baseline flux used in this analysis is given in \cite{O'Brien2004} as the flux at the 10th percentile over a specified time interval, which in this analysis was taken to be 0.5 s. The flux was fitted with a superposition of Gaussians for each energy channel, and the uncertainty in flux was calculated using the Poisson error from the microburst and baseline fluxes summed in quadrature. Using the fit parameters, the mean t_b for the lowest four energy channels is shown in Figure~\ref{tb_plot}. The trend of decreasing t_b as a function of energy is evident in Figure~\ref{tb_plot}, which further supports the

Deleted: ,

Deleted: = \$

Deleted: = \$

Deleted:

Deleted: (DLC)

Deleted: DLC

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted:

Deleted: -

Deleted:

Deleted: \$

Deleted: \$

Deleted: seconds

assumption that the subsequent peaks are bounces, and not a train of microbursts scattered by bouncing chorus.

```
\begin{figure}
\SPFIG{\framebox[10cm]{Figure 2}}{\caption{Observed and theoretical
 $\phi$  for electrons of energies from 200 to 770 keV. The solid black
line is  $\phi$  in a dipole magnetic field, derived in
\cite{Schulz1974}. The red dotted and cyan dashed lines are the
 $\phi$  derived using the T89, and T04 magnetic field models with
IRBEM-Lib. Lastly, the blue dot-dash curve is the  $\phi$  derived
using the Olson and Pfitzer Quiet model. The green and purple
rectangles represent the observed  $\phi$  for Flight Unit 3 (FU3) and
Flight Unit 4 (FU4) using a Gaussian fit, respectively. The blue
rectangles represent the observed  $\phi$  calculated with the minima
between the bounces. The width of the boxes represent the width of
those energy channels, and the height represents the uncertainty from
the fit.} \label{tb_plot}}
\end{figure}
```

The decaying peaks in the 231- to 408-keV electron flux observed by FU3's lowest two energy channels (see Figure~\ref{hires_plot}) were right-skewed. One explanation is that there was in-channel energy dispersion within those channels. Since ϕ of higher-energy electrons is shorter, a right-skewed peak implies that higher-energy electrons were more abundant within that channel, for example, in FU3's 231- to 300-keV channel, the 300-keV electrons will arrive sooner than the 231-keV electrons, but will they will be binned in the same channel. A Gaussian fit cannot account for this in-channel dispersion, and as a first-order correction, minima between peaks were used to calculate ϕ and are shown in Figure~\ref{tb_plot}. The observed energy-dependent dispersion shown in Figure~\ref{tb_plot} is consistent with higher-energy peaks returning sooner. This dispersion consistency further supports the assumption that the subsequent peaks are bounces, and not a train of microbursts scattered by bouncing chorus.

To compare the observed and modeled ϕ , we superposed ϕ curves for various models including an analytical solution in a dipole \cite{Schulz1974}, and numerical models: T89, Tsyganenko 2004 (T04; \citeauthor{Tsyganenko2005}, \citeyear{Tsyganenko2005}), and Olson and Pfitzer Quiet \cite{Olson1982} in Figure~\ref{tb_plot}. The numerical ϕ curves were calculated using a wrapper for IRBEM-Lib. This code traces the magnetic field line between mirror points and calculates ϕ assuming conservation of energy and the first adiabatic invariant for electrons mirroring at FIREBIRD-II. With the empirical ϕ , the models agree within FIREBIRD-II's uncertainties, but the T04 model has the largest discrepancy compared to the other models.

\subsection{Microburst Energy Spectra}

Next, we investigated the energy spectra of this microburst. The energy spectra were modeled with an exponential that was fit to the peak flux derived from the Gaussian fit parameters in section~\ref{t_b} to all but the highest energy channel. We found that the ϕ -folding energy, $E_0 \sim 100$ keV. These spectra are similar to spectra showed by \cite{Lee2005} from STSAT-1 and \cite{Datta1997} from sounding rocket measurements. The energy spectra are soft for a typical microburst observed with FIREBIRD-II,

Deleted:

Deleted: \&

Deleted:

Deleted: -

Deleted:

Deleted:

Deleted: e.g.

Deleted:

Deleted:

Deleted:

Deleted:

Deleted: was

Deleted: },

Deleted: is

Deleted:

Deleted:) \cite

Deleted: },

Deleted: \&

Deleted: ,

Deleted: was

Deleted: E-

Deleted: \$

Deleted: This

Deleted: is

Deleted: show

Deleted: is

and there was no statistically significant change in $E_{\{0\}}$ for subsequent bounces.

`\subsection{Microburst Scale Sizes} \label{scale_size}`
 Lastly, after we applied the time and separation corrections detailed in the SI, we mapped the locations of FU3 and FU4 in Figure~\ref{map_plot}. The locations where FU3 saw peaks 1--5 and where FU4 saw peaks 1--4 are shown as P1--5 and P1--4, respectively. The lower bound on the latitudinal extent of the microburst was the difference in latitude between P1 on FU3 and P4 on FU4 and was found to be 29 ± 1 km. The uncertainty was estimated from the spacecraft separation uncertainty described in the SI. This scale size is the largest reported by FIREBIRD-II.

`\begin{figure}`
`\SPIFIG{\framebox[10cm]{Figure 3}}{\caption{The topology of the`
`FIREBIRD-II orbit and the multiple bounces of the microburst projected`
`onto latitude and longitude with axis scaled to equal distance.`
`Attributes relating to FU3 shown in red dashed lines, and FU4 with`
`blue solid lines. The spacecraft path is shown with the diagonal`
`lines, starting at the upper right corner. The labels P1--4 for FU4`
`and P1--5 for FU3 indicate where the spacecraft were when the Nth`
`peak was seen in the lowest energy channel in the High-resolution`
`data. The stars with the accompanying energy labels represent the`
`locations of the electrons with that energy that started at time of P1`
`and were seen at the last peak on each spacecraft. The rectangles`
`represent the lower bound of the microburst scale size, assuming that`
`the majority of the electrons were in the upper boundary of energy`
`channel 4.\label{map_plot}}}`
`\end{figure}`

In section~\ref{t_b}, we showed that the observed decaying peaks were likely due to bouncing, so we assume that the observed electrons in subsequent bounces were the drifted electrons from the initial microburst. Under this assumption, the scattered electrons observed in the last bounce by FIREBIRD-II, must have drifted east from their initial scattering longitude, allowing us to calculate the minimum longitudinal scale size. Following geometrical arguments, the distance that electrons drift east in a single bounce is a product of the circumference of the drift shell foot print, and the fraction of the total drift orbit traversed in a single bounce and is given by

`\begin{equation}`
 `$d_{\{az\}} = 2 \pi (R_{\{E\}} + A) \cos(\lambda) \frac{t_{\{b\}}}{\angle T_{\{d\}}}$`
`\angle`
`\label{bounce_drift}`
`\end{equation}`

where $R_{\{E\}}$ is the Earth's radius, A is the spacecraft altitude, λ is the magnetic latitude, $t_{\{b\}}$ is the electron bounce period, and $\angle T_{\{d\}}$ is the electron drift period. \ocite{Parks2003} derived $\angle T_{\{d\}}$ to be,

`\begin{equation}`
 `$\angle T_{\{d\}} \approx$`
`\left{\right.`
`\begin{array}{ll}`
`43.8 / (L \cdot E) & \text{if } \alpha_{\{0\}} = 90^\circ \\`
`62.7 / (L \cdot E) & \text{if } \alpha_{\{0\}} = 0^\circ`
`\end{array}\right.`
`\label{drift}`

Deleted: -

Deleted: -

Deleted: -

Deleted: -

Deleted: \$

Deleted: -

Deleted: -

Deleted: HiRes

Deleted: ,

Deleted: ,

Deleted: ,

\end{equation}

where E is the electron energy in MeV, L is the L shell, and α_0 is the equatorial pitch angle. Electrons mirroring at FIREBIRD-II have $\alpha_0 \approx 3.7^\circ$ and so the $\alpha_0 = 0^\circ$ limit was used.

The microburst's longitudinal scale size is defined as the distance the highest energy electrons drifted in the time between the observations of the first and last peaks. This scale size is given by $D_{az} = n \lambda_{az}$ where n is the number of bounces observed. The stars in Figure~\ref{map_plot} (with labels corresponding to energy channel boundaries) represent the locations when the microburst was observed at P1, such that an electron of that energy would drift eastward to be seen at P5 for FU3 and P4 for FU4. Since FU3 observed more peaks it observed the larger longitudinal scale size which is shown with the red dashed box in Figure~\ref{map_plot}. FU3's fourth energy channel's bounds are 555 and 771 keV, which correspond to longitudinal distances of 39 ± 1 km and 51 ± 1 , respectively. The uncertainty was estimated by propagating the uncertainty in the bounce time [equation~\eqref{bounce_drift}](#). While the observed minimum longitudinal scale size is dependent on FIREBIRD-II's energy channels, the true scale size may not be.

To investigate how the microburst scale size compares to the scale sizes of chorus waves near the magnetic equator, the microburst's longitudinal and latitudinal scale sizes and their uncertainties in LEO were mapped to the magnetic equator with T89. The radial scale size (latitudinal scale mapped from LEO) was greater than 500 ± 10 km. The azimuthal scale size (longitudinal scale mapped from LEO) of 555-keV electrons was greater than 450 ± 10 km, and for the 771-keV electrons it was greater than 530 ± 10 km. The lower bound microburst scale size is similar to the chorus scale sizes derived by \citeauthor{Agapitov2011b} (\citeyear{Agapitov2011b}), \citeauthor{Agapitov2017a} and is discussed below.

\section{Discussion and Conclusions} \label{discussion}

We presented the first observation of a large microburst with multiple bounces made possible by the twin FIREBIRD-II CubeSats. The microburst's lower bound LEO latitudinal and longitudinal scale sizes of 29 ± 1 km and 51 ± 1 km make it one of the largest observed. The microburst's LEO scale size was larger than the latitudinal scale sizes of typical >1 -MeV microbursts reported in \cite{Blake1996}, approximately 10 times larger than reported in \cite{Dietrich2010}, and approximately 2.6 times larger than other simultaneous microbursts observed by FIREBIRD-II \cite{Crew2016}. Lastly, the scale sizes derived here were similar to the scale sizes of >15 -keV microbursts observed with a high-altitude balloon \cite{Parks1967}. No energy dependence on the minimum latitudinal scale size was observed, while the observed energy dependence of the minimum longitudinal scale size is an artifact of the technique we used to estimate their drift motion.

The microburst scale size obtained in [section~\ref{scale_size}](#) and scaled to the geomagnetic equator can be compared with the scales of chorus waves presumably responsible for the rapid burst electron precipitation. Early direct estimates of the chorus source scales were made by the coordinated measurement by ISEE-1, 2. The wave power correlation scale was estimated to be about several hundred kilometers

Deleted: keV

Deleted:

Deleted: \$

Deleted: Equation

Deleted: \$

Deleted:

Deleted: \$

Deleted:

Deleted: \$

Deleted:) ,

Deleted: \$

Deleted: \$

Deleted: >

Deleted:

Deleted: >

Deleted:

Deleted:

Deleted: Section

across the background magnetic field \cite{Gurnett1979}. Furthermore, \ocite{Santolik2003} determined the correlation lengths of chorus-type whistler waves to be around 100~km based on multipoint CLUSTER Wide Band Data measurements near the chorus source region at $L \approx 4$, during the magnetic storm of 18 April 2002. \citeauthor{Agapitov2010} (\citeyear{Agapitov2017a}, \citeyear{Agapitov2011b}, \citeyear{Agapitov2010}) recently showed that the spatial extent of chorus source region can be larger, ranging from 600~km in the outer radiation belt to more than 1,000~km in the outer magnetosphere. The lower bound azimuthal and latitudinal scales obtained in [section~\ref{scale_size}](#) and scaled to the magnetic equator are similar to the whistler mode chorus source scale sizes reported in \citeauthor{Agapitov2011b} (\citeyear{Agapitov2017a}, \citeyear{Agapitov2011b}).

No wave measurements from nearby spacecraft were available at this time. Nevertheless, during the hours before and after this observation, the Van Allen Probes' \cite{Mauk2013} Electric and Magnetic Field Instrument and Integrated Science \cite{Kletzing2013} observed strong wave power in the lower band chorus frequency range, inside the outer radiation belt between 22 and 2 MLT. Furthermore, AE ~ 400 nT at this time, and relatively strong chorus waves were statistically more likely to be present at FIREBIRD-II's MLT \cite{Li2009}.

The empirically estimated and modeled σ_{b} in this study agree within FIREBIRD-II's uncertainties, confirming that the energy-dependent dispersion was due to bouncing. The σ_{b} curves are a proxy for field line length, and this agreement implies that they are comparable. This is expected since the magnetosphere is not drastically compressed at 8 MLT, but we expect a larger discrepancy near midnight, where the magnetosphere is more stretched and difficult to accurately model. In future studies, this analysis can be used as a diagnostic tool to validate field line lengths, and improve magnetic field models.

The similarity of the microburst and chorus source region scale sizes, as well as magnetospheric location and conditions, further [supports](#) the causal relationship between microbursts and chorus.

\backmatter

\begin{bm}[Acknowledgments]

This work was made possible with help from the FIREBIRD team, and the members of the Space Sciences and Engineering Laboratory at Montana State University for their hard work to make this mission a success. In addition, M. Shumko acknowledges Drew Turner for his suggestions regarding the bounce period calculations, and Dana Longcope for his proofreading feedback. The FIREBIRD-II data are available at <http://solar.physics.montana.edu/FIREBIRD\II/>. This analysis is supported by the \fundingAgency{National Science Foundation} under [grants](#) \fundingNumber{0838034} and \fundingNumber{1339414}. Furthermore, the work of O. Agapitov was supported by the \fundingAgency{NASA} grant \fundingNumber{NNX16AF85G}.

\end{bm}

\begin{thebibliography}{49}

Deleted:

Deleted: },

Deleted: Agapitov2010

Deleted: Agapitov2017a

Deleted: ~

Deleted:

Deleted: 1000

Deleted: Section

Deleted: ,

Deleted: -

Deleted: }, \citeyear{Agapitov2017a}

Deleted: \$

Deleted: ,

Deleted: support

Deleted: Grant Numbers

```
%=Ref1
\bib{Abell1998_1}{article}{
author={Abel, B.},
author={Thorne, R. M.},
date={1998},
title={Electron scattering loss in Earth's inner magnetosphere: 1.
Dominant physical processes},
journal={Journal of Geophysical Research: Space Physics},
volume={103},
number={A2},
pages={2385--2396},
}
```

Deleted: earth's

Deleted: dominant

```
%=Ref2
\bib{Agapitov2017a}{article}{
author={Agapitov, O.},
author={Blum, L. W.},
author={Mozer, F. S.},
author={Bonnell, J. W.},
author={Wygant, J.}
title={Chorus whistler wave source scales as determined from
multipoint Van Allen Probe measurements},
date={2017},
volume={44}
pages={2634-2642}
publisher={Geophysical Research Letters},
note={\url{https://doi.org/10.1002/2017GL072701}},
}
```

Deleted: van allen probe

```
%=Ref3
\bib{Agapitov2011b}{article}{
author={Agapitov, O.},
author={Krasnoselskikh, V.}
author={Dudok de Wit, T.}
author={Khotyaintsev, Y.}
author={Pickett, J. S.}
author={Santolik, O.}
author={Rolland, G.}
date={2011},
title={Multispacecraft observations of chorus emissions as a tool for
the plasma density fluctuations' remote sensing},
journal={Journal of Geophysical Research: Space Physics},
volume={116}
pages={A09222}
note={\url{https://doi.org/10.1029/2011JA016540}}
}
```

```
%=Ref4
\bib{Agapitov2010}{article}{
author={Agapitov, O.},
author={Krasnoselskikh, V.},
author={Zaliznyak, Y.},
author={Angelopoulos, V.},
author={Le Contel, O.},
author={Rolland, G.},
date={2010},
```

title={Chorus source region localization in the [Earth's](#) outer magnetosphere using [THEMIS](#) measurements},
journal={Annales Geophysicae},
volume={28},
number={6},
pages={1377--1386},
note={\url{https://doi.org/10.5194/angeo-28-1377-2010}}

Deleted: earth's

Deleted: themis

%=Ref5
\bib{Anderson1964}{article}{
author={Anderson, K. A.},
author={Milton, D. W.},
date={1964},
title={Balloon observations of [X](#) rays in the auroral zone: 3. [High](#) time resolution studies},
journal={Journal of Geophysical Research: Space Physics},
volume={69},
number={21},
pages={4457--4479},
note={\url{https://doi.org/10.1029/JZ069i021p04457}},
}

Deleted: x

Deleted: high

%=Ref6
\bib{Anderson2017}{article}{
author={Anderson, B.},
author={Shekhar, S.},
author={Millan, R.},
author={Crew, A.},
author={Spence, H.},
author={Klumpar, D.},
author={others},
title={Spatial scale and duration of one microburst region on 13 [August](#) 2015},
date={2017},
volume={122}
journal={Journal of Geophysical Research: Space Physics},
pages={5949--5964}
}

Deleted: august

%=Ref7
\bib{Blake1996}{article}{
author={Blake, J.},
author={Looper, M.},
author={Baker, D.},
author={Nakamura, R.},
author={Klecker, B.},
author={Hovestadt, D.},
date={1996},
title={New high temporal and spatial resolution measurements by [SAMPEX](#) of the precipitation of relativistic electrons},
journal={Advances in Space Research},
volume={18}

Deleted: sampex

```
number={8},
pages={171--186},
note={\url{https://doi.org/10.1016/0273-1177(95)00969-8}},
}
```

```
%=Ref8
\bib{Blum2015}{article}{
author={Blum, L.},
author={Li, X.},
author={Denton, M.},
date={2015},
title={Rapid MeV electron precipitation as observed by SAMPEX/HILT
during high-speed stream-driven storms},
journal={Journal of Geophysical Research: Space Physics},
volume={120},
pages={3783--3794},
note={\url{https://doi.org/10.1002/2014JA020633}},
}
```

Deleted: mev

Deleted: sampex/hilt

```
%=Ref9
\bib{irbem}{misc}{
author={Boscher, D.},
author={Bourdarie, S.},
author={O'Brien, P.},
author={Guild, T.},
author={Shumko, M.},
date={2012},
note={\a href="#">IRBEM-LIB library},
}
```

Deleted: Irbem-lib

```
%=Ref10
\bib{Breneman2017}{article}{
author={Breneman, A.},
author={Crew, A.},
author={Sample, J.},
author={Klumpar, D.},
author={Johnson, A.},
author={Agapitov, O.},
author={others},
date={2017},
title={Observations directly linking relativistic electron microbursts
to whistler mode chorus: Van Allen Probes and FIREBIRD II},
journal={Geophysical Research Letters},
volume={44}
pages={11,265--11272}
}
```

Deleted: allen probes

Deleted: firebird ii

```
%=Ref11
\bib{Comess2013}{article}{
author={Comess, M.},
author={Smith, D.},
author={Selesnick, R.},
author={Millan, R.},
author={Sample, J.},
date={2013},
title={Duskside relativistic electron precipitation as measured by
SAMPEX: A statistical survey},
journal={Journal of Geophysical Research: Space Physics},
```

Deleted: sampex

```
volume={118},
pages={5050--5058},
note={\url{https://doi.org/10.1002/jgra.50481}},
}
```

```
%=Ref12
\bib{Crew2016}{article}{
author={Crew, A. B.},
author={Spence, H. E.},
author={Blake, J. B.},
author={Klumpar, D. M.},
author={Larsen, B. A.},
author={O'Brien, T. P.},
author={others},
date={2016},
title={First multipoint in situ observations of electron microbursts:
{Initial} results from the NSF FIREBIRD II mission},
journal={Journal of Geophysical Research: Space Physics},
volume={121},
pages={5272--5283},
note={\url{https://doi.org/10.1002/2016JA022485}},
}
```

Deleted: nsf firebird ii

```
%=Ref13
\bib{Datta1997}{article}{
author={Datta, S.},
author={Skoug, R.}
author={McCarthy, M.}
author={Parks, G.}
date={1997},
title={Modeling of microburst electron precipitation using pitch angle
diffusion theory},
journal={Journal of Geophysical Research: Space Physics},
volume={102},
number={A8},
pages={17,325--17,333}
}
```

```
%=Ref14
\bib{Dietrich2010}{article}{
author={Dietrich, S.},
author={Rodger, C. J.},
author={Clilverd, M. A.},
author={Bortnik, J.},
author={Raita, T.},
date={2010},
title={Relativistic microburst storm characteristics: Combined
satellite and ground-based observations},
journal={Journal of Geophysical Research: Space Physics},
volume={115},
pages={A12240},
}
```

```
%=Ref15
\bib{Fang2010}{article}{
author={Fang, X.},
author={Randall, C. E.},
author={Lummerzheim, D.},
```

```
author={Wang, W.},
author={Lu, G.},
author={Solomon, S. C.},
author={Frahm, R. A.},
date={2010},
title={Parameterization of monoenergetic electron impact ionization},
journal={Geophysical Research Letters},
volume={37},
pages={L22106}
}
```

```
%=Ref16
\bib{Gurnett1979}{article}{
author={Gurnett, D.},
author={Anderson, R.},
author={Scarf, F.},
author={Fredricks, R.},
author={Smith, E.},
date={1979},
title={Initial results from the ISEE-1 and-2 plasma wave
investigation},
journal={Space Science Reviews},
volume={23},
number={1},
pages={103--122},
}
```

Deleted: isee

```
%=Ref17
\bib{sgp4}{report}{
author={Hoots, F. R.},
author={Roehrich, R. L.}
date={1980},
title={Models for \AQ{AUTHOR: Please provide the city location of
publisher for Reference Hoots and Roehrich (1980).}propagation of
norad element sets},
number={Tech. Rep. 3},
organization={Spacetrack}
}
```

Deleted: /CED

```
%=Ref18
\bib{Horne2003}{article}{
author={Horne, R. B.},
author={Thorne, R. M.}
date={2003},
title={Relativistic electron acceleration and precipitation during
resonant interactions with whistler-mode chorus},
journal={Geophysical Research Letters},
volume={30}
number={10}
pages={1527}
note={\url{https://doi.org/10.1029/2003GL016973}}
}
```

```
%=Ref19
\bib{Kletzing2013}{article}{
author={Kletzing, C.},
author={Kurth, W.},
```

```

author={Acuna, M.},
author={MacDowall, R.},
author={Torbert, R.},
author={Averkamp, T.},
author={others},
date={2013},
title={The electric and magnetic field instrument suite and integrated
science (EMFISIS) on Rbsp},
journal={Space Science Reviews},
volume={179},
number={1-4},
pages={127--181},
}

```

Deleted: emfisis

Deleted: rbsp

```

%=Ref20
\bib{Klumpar2015}{proceedings}{
author={Klumpar, D.},
author={Springer, L.},
author={Mosleh, E.},
author={Mashburn, K.},
author={Berardinelli, S.},
author={Gunderson, A.},
author={others},
date={2015},
title={Flight system \AQ{AUTHOR: Please check that Reference Klumpar
et al. (2015) is presented correctly.}technologies enabling the twin-
CubeSat FIREBIRD II scientific mission},
conference={Proceedings of the 29th Annual AIAA/USU Conference on
Small Satellites}
address={Logan, Utah, USA}
}

```

Deleted: /CED

Deleted: if

Deleted: if captured

Deleted: cubesat firebird-ii

```

%=Ref21
\bib{Lee2012}{article}{
author={Lee, J. J.},
author={Parks, G. K.},
author={Lee, E.},
author={Tsurutani, B. T.},
author={Hwang, J.},
author={Cho, K. S.},
author={others},
date={2012},
title={Anisotropic pitch angle distribution of 100 kev microburst
electrons in the loss cone: Measurements from STSAT-1},
journal={Annales Geophysicae},
volume={30},
number={11},
pages={1567--1573},
note={\url{https://doi.org/10.5194/angeo-30-1567-2012}},
}

```

Deleted: stsat

```

%=Ref22
\bib{Lee2005}{article}{
author={Lee, J.-J.},
author={Parks, G. K.}
author={Min, K. W.}
author={Kim, H. J.}
}

```


author={Park, J.}
author={Hwang, J.}
author={others}
date={2005},
title={Energy spectra of 170-360 kev electron microbursts measured by the [Korean STSAT-1](#)},
journal={Geophysical Research Letters},
volume={32}
pages={113106}
note={\url{https://doi.org/10.1029/2005GL022996}}
}

Deleted: korean stsat

%=Ref23
\bib{Li2009}{article}{
author={Li, W.},
author={Thorne, R. M.}
author={Angelopoulos, V.}
author={Bortnik, J.}
author={Cully, C. M.}
author={Ni, B.}
author={others}
date={2009},
title={Global distribution of whistler-mode chorus waves observed on the themis spacecraft},
journal={Geophysical Research Letters},
volume={36}
note={\url{https://doi.org/10.1029/2009GL037595}}
pages={109104}
}

%=Ref24
\bib{Lorentzen2001a}{article}{
author={Lorentzen, K. R.},
author={Blake, J. B.},
author={Inan, U. S.},
author={Bortnik, J.},
date={2001},
title={Observations of relativistic electron microbursts in association with [VLF](#) chorus},
journal={Journal of Geophysical Research: Space Physics},
volume={106},
number={A4},
pages={6017--6027},
doi={10.1029/2000JA003018},
}

Deleted: 2001a

Deleted: vlf

%=Ref25
\bib{Lorentzen2001b}{article}{
author={Lorentzen, K. R.},
author={Looper, M. D.},
author={Blake, J. B.},
date={2001},
title={Relativistic electron microbursts during the [GEM](#) storms},
journal={Geophysical Research Letters},
volume={28},
number={13},
pages={2573--2576},
note={\url{https://doi.org/10.1029/2001GL012926}},

Deleted: 2001b

Deleted: gem

}

%=Ref26

```
\bib{Mauk2013}{article}{
author={Mauk, B.},
author={Fox, N. J.},
author={Kaneval, S.},
author={Kessel, R.},
author={Sibeck, D.},
author={Ukhorskiy, A.},
date={2013},
title={Science objectives and rationale for the Radiation Belt Storm Probes mission},
journal={Space Science Reviews},
volume={179},
number={1-4},
pages={3--27},
}
```

Deleted: radiation belt storm probes

%=Ref27

```
\bib{Meredith2002}{article}{
author={Meredith, N.},
author={Horne, R.},
author={Summers, D.},
author={Thorne, R.},
author={Iles, R.},
author={Heynderickx, D.},
author={Anderson, R.},
date={2002},
title={Evidence for acceleration of outer zone electrons to relativistic energies by whistler mode chorus},
journal={Annales Geophysicae},
volume={20},
pages={967--979},
}
```

%=Ref28

```
\bib{Millan2002}{article}{
author={Millan, R. M.},
author={Lin, R.},
author={Smith, D.},
author={Lorentzen, K.},
author={McCarthy, M.},
date={2002},
title={X-ray observations of MeV electron precipitation with a balloon-borne germanium spectrometer},
journal={Geophysical research letters},
volume={29},
number={24},
pages={47-1-47-4}
}
```

Deleted: mev

%=Ref29

```
\bib{Millan2007}{article}{
author={Millan, R.},
author={Thorne, R.}
date={2007},
title={Review of radiation belt relativistic electron losses},
```

```

journal={Journal of Atmospheric and Solar-Terrestrial Physics},
volume={69}
number={3},
pages={362--377}
note={\url{https://doi.org/10.1016/j.jastp.2006.06.019}, Global
Aspects of Magnetosphere-Ionosphere CouplingGlobal Aspects of
Magnetosphere-Ionosphere Coupling}
}

```

Deleted: global

```

%=Ref30
\bib{Mozer2018}{article}{
author={Mozer, F. S.},
author={Agapitov, O. V.},
author={Blake, J. B.},
author={Vasko, I. Y.},
date={2018},
title={Simultaneous observations of \AQ{AUTHOR: Please provide article
volume and page range for Reference Mozer, et al. (2018).}lower band
chorus emissions at the equator and microburst precipitating electrons
in the ionosphere},
journal={Geophysical Research Letters},
note={\url{https://doi.org/10.1002/2017GL076120}},
}

```

```

%=Ref31
\bib{Nakamura1995}{article}{
author={Nakamura, R.},
author={Baker, D. N.},
author={Blake, J. B.},
author={Kanekeal, S.},
author={Klecker, B.},
author={Hovestadt, D.},
date={1995},
title={Relativistic electron precipitation enhancements near the outer
edge of the radiation belt},
journal={Geophysical Research Letters},
volume={22},
number={9},
pages={1129--1132},
note={\url{https://doi.org/10.1029/95GL00378}},
}

```

```

%=Ref32
\bib{Nakamura2000}{article}{
author={Nakamura, R.},
author={Isowa, M.}
author={Kamide, Y.}
author={Baker, D.}
author={Blake, J.}
author={Looper, M.}
date={2000},
title={Observations of relativistic electron microbursts in
association with VLF chorus},
journal={Journal of Geophysical Research},
volume={105},
pages={15,875--15,885}
}

```

Deleted: vlf

%=Ref33
\bib{O'Brien2004}{article}{
author={O'Brien, T. P.},
author={Looper, M. D.}
author={Blake, J. B.}
date={2004},
title={Quantification of relativistic electron microburst losses
during the [GEM](#) storms},
journal={Geophysical Research Letters},
volume={31}
note={\url{https://doi.org/10.1029/2003GL018621}}
pages={104802}
}

Deleted: gem

%=Ref34
\bib{O'Brien2003}{article}{
author={O'Brien, T. P.},
author={Lorentzen, K. R.}
author={Mann, I. R.}
author={Meredith, N. P.}
author={Blake, J. B.}
author={Fennell, J. F.}
author={others}
date={2003},
title={Energization of relativistic electrons in the presence of [ULF](#)
power and [MeV](#) microbursts: Evidence for dual [ULF](#) and [VLF](#)
acceleration},
journal={Journal of Geophysical Research: Space Physics},
volume={108}
number={A8}
note={\url{https://doi.org/10.1029/2002JA009784}}
pages={1329}
}

Deleted: ulf

Deleted: mev

Deleted: ulf

Deleted: vlf

%=Ref35
\bib{Olson1982}{article}{
author={Olson, W. P.},
author={Pfitzer, K. A.},
date={1982},
title={A dynamic model of the magnetospheric magnetic and electric
fields for [July](#) 29, 1977},
journal={Journal of Geophysical Research: Space Physics},
volume={87},
number={A8},
pages={5943--5948},
note={\url{https://doi.org/10.1029/JA087iA08p05943}},
}

Deleted: july

%=Ref36
\bib{Parks1967}{article}{
author={Parks, G. K.},
date={1967},
title={Spatial characteristics of auroral-zone [X](#)-ray microbursts},

Deleted: x

```
journal={Journal of Geophysical Research: Space Physics},
volume={72},
number={1},
pages={215--226},
}
```

```
%=Ref37
\bib{Parks2003}{book}{
author={Parks, G.},
date={2003},
title={Physics of space plasmas: An introduction},
edition={Second Edition},
publisher={Westview Press},
address={Boulder, Colorado}
}
```

Deleted: Of Space Plasmas

Deleted: Introduction

```
%=Ref38
\bib{Santolik2003}{article}{
author={Santolik, O.},
author={Gurnett, D.},
author={Pickett, J.},
author={Parrot, M.},
author={Cornilleau-Wehrlin, N.},
date={2003},
title={Spatio-temporal structure of storm-time chorus},
journal={Journal of Geophysical Research: Space Physics},
volume={108},
number={A7},
pages={1278}
}
```

```
%=Ref39
\bib{Schulz1974}{book}{
author={Schulz, M.},
author={Lanzerotti, L. J.}
title={Particle diffusion in the radiation belts},
date={1974},
address={Berlin},
publisher={Springer},
}
```

Deleted: Diffusion

Deleted: Radiation Belts

```
%=Ref40
\bib{Selesnick2003}{article}{
author={Selesnick, R. S.},
author={Blake, J. B.}
author={Mewaldt, R. A.}
date={2003},
title={Atmospheric losses of radiation belt electrons},
journal={Journal of Geophysical Research: Space Physics},
volume={108}
number={A12}
pages={1468}
note={\url{https://doi.org/10.1029/2003JA010160}}
}
```

```
%=Ref41
\bib{Shprits2007}{article}{
```

```
author={Shprits, Y. Y.},
author={Meredith, N. P.}
author={Thorne, R. M.}
date={2007},
title={Parameterization of radiation belt electron loss timescales due
to interactions with chorus waves},
journal={Geophysical Research Letters},
volume={34}
note={\url{https://doi.org/10.1029/2006GL029050}}
pages={111110}
}
```

```
%=Ref42
\bib{Shprits2004}{article}{
author={Shprits, Y. Y.},
author={Thorne, R. M.}
date={2004},
title={Time dependent radial diffusion modeling of relativistic
electrons with realistic loss rates},
journal={Geophysical Research Letters},
volume={31}
note={\url{https://doi.org/10.1029/2004GL019591}}
pages={108805}
}
```

```
%=Ref43
\bib{Spence2012}{article}{
author={Spence, H. E.},
author={Blake, J. B.},
author={Crew, A. B.},
author={Driscoll, S.},
author={Klumpar, D. M.},
author={Larsen, B. A.},
author={others},
date={2012},
title={Focusing on size and energy dependence of electron microbursts
from the Van Allen radiation belts},
journal={Space Weather},
volume={10},
pages={S11004}
note={\url{https://doi.org/10.1029/2012SW000869}},
}
```

```
%=Ref44
\bib{Summers1998}{article}{
author={Summers, D.},
author={Thorne, R. M.}
author={Xiao, F.}
date={1998},
title={Relativistic theory of wave-particle resonant diffusion with
application to electron acceleration in the magnetosphere},
journal={Journal of Geophysical Research: Space Physics},
volume={103}
number={A9},
pages={20,487--20,500}
}
```

Deleted: van allen

```

%=Ref45
\bib{Thorne2010}{article}{
author={Thorne, R. M.}
date={2010},
title={Radiation belt dynamics: The importance of wave-particle
interactions}, journal={Geophysical Research Letters},
volume={37}
pages={122107}
note={\url{https://doi.org/10.1029/2010GL044990}},
}

%=Ref46
\bib{Thorne2005}{article}{
author={Thorne, R. M.},
author={O'Brien, T. P.}
author={Shprits, Y. Y.}
author={Summers, D.}
author={Horne, R. B.}
date={2005},
title={Timescale for MeV electron microburst loss during geomagnetic
storms}, journal={Journal of Geophysical Research: Space Physics},
volume={110}
pages={a09202}
note={\url{https://doi.org/10.1029/2004JA010882}}
}

%=Ref47
\bib{Tsyganenko1989}{article}{
author={Tsyganenko, N.},
date={1989},
title={A solution of the Chapman-Ferraro problem for an ellipsoidal
magnetopause},
journal={Planetary and Space Science},
volume={37}
number={9},
pages={1037--1046},
note={\url{https://doi.org/10.1016/0032-0633(89)90076-7}},
}

%=Ref48
\bib{Tsyganenko2005}{article}{
author={Tsyganenko, N. A.},
author={Sitnov, M. I.}
date={2005},
title={Modeling the dynamics of the inner magnetosphere during strong
geomagnetic storms},
journal={Journal of Geophysical Research: Space Physics},
volume={110}
pages={a03208}
note={\url{https://doi.org/10.1029/2004JA010798}}
}

%=Ref49
\bib{Ukhorskiy2006}{article}{
author={Ukhorskiy, A. Y.},
author={Anderson, B. J.}
author={Brandt, P. C.}
author={Tsyganenko, N. A.}

```

Deleted: mev

Deleted: chapman-ferraro

```
date={2006},
title={Storm time evolution of the outer radiation belt: Transport and
losses}, journal={Journal of Geophysical Research: Space Physics},
volume={111}
pages={a11S03}
note={\url{https://doi.org/10.1029/2006JA011690}}
}
```

```
%=Ref50
```

```
\bib{Woodger2015}{article}{
author={Woodger, L.},
author={Halford, A.},
author={Millan, R.},
author={McCarthy, M.},
author={Smith, D.},
author={Bowers, G.},
author={others},
date={2015},
title={A summary of the barrel campaigns: Technique for studying
electron precipitation},
journal={Journal of Geophysical Research: Space Physics},
volume={120},
number={6},
pages={4922--4935},
}
\end{thebibliography}
```

```
\printquery
\end{document}
```