

CONCLUSIONS AND FUTURE WORK

1. Microburst scattering mechanism and relation to prior work
2. Microburst sizes and the elliptical shape, mention the curtain-microburst ambiguity.

In this dissertation we have explored the microburst scattering mechanism directly in Chapter ?? and indirectly in Chapters ?? and ??. In Chapter ?? we used numerous particle and wave instruments on the Van Allen Probes and found signatures of microbursts with the Magnetic Electron Ion Spectrometer. To these observations we applied the relativistic theory of wave-particle resonant diffusion and found that the motion of the microburst electrons was not along single-wave particle characteristics in momentum phase space, given the spacecraft position and orientation and most probable wave and plasma parameters. This result at first appears to contradict the belief that many members of the community hold, that microburst precipitation is due to a diffusive process. In reality both are probably valid, just on different time scales. Individual microbursts are probably not scattered diffusively, but the combined contribution of an ensemble of microbursts will have properties that are well modeled as a diffusion process.

The microburst sizes estimated in prior literature as well as Chapters ?? and ?? show that there is a large variability in microburst sizes. The AC6 study in Chapter ?? showed that in LEO, most microbursts were observed while the AC6 separation was less than a few tens of km while a minority of microbursts were observed up to ≈ 100 km separation. These conclusions agree with prior literature from high altitude balloons and LEO spacecraft. One unanswered question is what shape is a

microburst? A circular microburst is easy to interpret and model due to its symmetry,
but surely the real shape is

Future Work

28

- In future work mention how curtains should be studied and outline that project.
Mention the SAA and amplitude idea.
- Bouncing packet idea
- Mention inverse microburst analysis and compare to Saito.
- Mention REAL and how it plans to look at the microburst flux in PA space.
- Mention BOOMS and how it plans to image microbursts to determine their
shape without ambiguity.

An extension of the case study in Chapter ?? will be a statistical study using
the Van Allen Probes. Other microburst-like events have already been identified
by eye. These other events are also simultaneously observed with enhanced wave
activity, hence they may be related and a further investigation is warranted. A
microburst detection scheme similar to the one used in Chapter ?? can be easily
implemented to automatically identify other microbursts for further study. A few
compelling questions that can be addressed with this study are: what is the typical
pitch angle extent of microbursts? Do these microbursts have a similar MLT extent
to microbursts observed in LEO? What fraction of microbursts were observed during
enhanced wave activity? What wave modes and properties are observed during these
events? And lastly, what fraction of microbursts can be modeled with a diffusive
process?

48 With these results, it is always important to keep our methods and their biases
49 in mind. Werner Heisenberg once wrote “What we observe is not nature itself, but
50 nature exposed to our method of questioning.” Heisenberg (1958)

- 52 Heisenberg, W. (1958). Physics and philosophy.