## MeV Dusk Microbursts


Mykhaylo Shumko

16 March 2023

#### Methodology


- I wrote a program to generate summary plots of each microburst in the dusk MLTs that were poorly and well fit
- The program is in a "202303 dusk" development branch of the "sampex microburst widths" package
- Filtered the pre-existing microburst catalog (microburst catalog 04.csv) by
  - 16 < MLT < 22
  - "Good" fits which are adj\_R^2>0.9
  - "Bad" fits which are adj\_R^2<0.5</li>
- Stats:
  - 16083 microbursts in mlt\_range=[16, 22]
  - 4515 microbursts in mlt\_range=[16, 22] and with adj\_R^2>0.9
  - 2047 microbursts in mlt\_range=[16, 22] and with adj\_R^2<0.5
- I produced the ~6500 plots and I upload them to <u>Google Drive</u>. Take a look at your leisure!

#### Compare good vs. bad fits & dawn vs. dusk



#### Where are the bad fits?


#### SAMPEX >1 MeV microbursts | Distribution of Good vs. Bad Fits





MLT


MLT


#### Interesting poor fits













Each x-axis tick is 100 ms!



SAMPEX-HILT | >1 MeV Microburst Validation

... I recall finding these dusk microbursts using the FIRBEIRD-II data too

#### More short microburst examples...



# Did I rediscover precipitation bands, or are these still microbursts?

