

Lecture Notes: Relativity

Based on Youtube series by eigenchris

Contents

1	Introduction to Relativity	2
1.1	Galilean Relativity	2
1.2	Special Relativity	3

1 Introduction to Relativity

Before we even start by exploring relativity, either it being Galilean or Einstein one, we should ask ourselves: what is relativity?

Relativity is how the laws of motion work in different reference frames.

1.1 Galilean Relativity

Now, the Galilean principle of relativity, states that **the laws of motion are the same in all inertial frames of reference.**

And an inertial reference frame, is a reference frame that's not accelerating.

If we think about a very simple experiment from a scientist dropping a ball on the floor in his office room, and a car passing by this office and observing the experiment:

- The scientist will just see the ball falling down straight to the ground
- A potential car passing (at a constant velocity) by the office and looking at it, will see the ball moving along a curved trajectory towards the floor

At the end, they would disagree on the velocity, on the momentum, kinetic energy as well.

So like, what are they actually agreeing on? Well, they would agree on:

- The time passed
- The size of the ball
- The mass of the ball

Eventually, this leads to them agreeing on the laws of motion, the Newton's three laws of motion and the gravitational law:

$$\vec{F} = m\vec{a} \tag{1}$$

$$\vec{F} = G \frac{m_1 m_2}{r^2} (-\hat{r}) \tag{2}$$

And if we copy the experiment of dropping the ball inside the car passing by at constant velocity, the experiment will be exactly the same as the scientist run in his own office. So the laws of motion are exactly the same in all inertial reference frames.

There's no experiment you can do to determine if your reference frame is stationary, or moving at constant speed. And there's no special reference frame truly stationary

And what would instead be an example of non-inertial reference frame? Well an elevator accelerating, or a car accelerating forward, a car turning a corner.

And in all these non-inertial situations, the passenger (man in the elevator, or driver in the car) always feels a mysterious fictitious force pulling them in the opposite direction of the acceleration. **This means that a passenger in an accelerating reference frame, will have different set of laws of motion!** Everytime they use Newton's second law, they need to add an extra term for the fictitious force in order to predict the motion of objects correctly.

1.2 Special Relativity

This type of relativity is the one we experience when moving to speed close to the speed of light.

The story starts in the 19th century with Maxwell's equations of electro-magnetism:

$$\begin{aligned}\nabla \cdot \vec{E} &= \rho \\ \nabla \times \vec{E} &= -\frac{\partial \vec{B}}{\partial t} \\ \nabla \cdot \vec{B} &= 0 \\ \nabla \times \vec{B} &= \mu_0 \vec{J} + \mu_0 \epsilon_0 \frac{\partial \vec{E}}{\partial t}\end{aligned}$$

There's a problem with these equations, when we combine them together, we obtain equations for electromagnetic waves, or in other words..light.

$$\nabla^2 \vec{E} = \frac{1}{c^2} \left(\frac{\partial^2 \vec{E}}{\partial t^2} \right) \qquad \nabla^2 \vec{B} = \frac{1}{c^2} \left(\frac{\partial^2 \vec{B}}{\partial t^2} \right)$$

These equations say that light, in a vacuum, travels at the speed of $c = \frac{1}{\sqrt{\mu_0 \epsilon_0}}$, and this speed, if you plug in the two constants for the magnetic permeability $\mu_0 = 4\pi \times 10^{-7} H/m$, and electric permittivity $\epsilon_0 = 8.854 \times 10^{-12} F/m$, is equal to $c \approx 3 \times 10^8 m/s$.

If you think about this, and try to correlate to other speeds, like the speed of sound, you know that that's measured relative to air, or the speed of ocean's waves, that's relative to water, etc., but then what's the speed of light relative to, if it travels in the vacuum? Remember that the Galilean relativity states that there's no special reference frame! And all motion is always relative to something else.

So what are we measuring the speed of light relative to in the vacuum, if the vacuum is just empty space?? To get around this problem, scientists predicted the existence of the "aether", a special stationary reference frame that light travels in. But experiments such as

the "Michelson-Morley" one showed that aether does not exist.

And this is where Einstein comes in, proposing that the speed of light c is the same in all reference frames, basically treating the speed of light as a law of physics. So in an attempt to build a new theory of relativity, Einstein began by agreeing with Galileo and said that the laws of motion are the same in all inertial reference frames, but he expanded this principle by including the Maxwell's equations in the laws of motion that the reference frames agree upon, and made the speed of light as a constant that everyone agree upon.

Finally, the two Einstein postulates of Special Relativity are:

- Laws of physics are the same in all inertial reference frames;
- Speed of light in a vacuum c is equal for all observers in inertial reference frames (a.k.a speed of light in a vacuum is included in laws of physics).

Unfortunately, these postulates are not compatible with the Galilean relativity! To see why, we can simply think about a similar experiment we thought about above, a scientist on the ground, and a car passing by at constant velocity v to the right, but this time with a giant flash light attached on its roof.

In the car's frame, the driver sees the scientist pass by with a speed of v to the left, and the light coming out from the flash light traveling to the right at speed c . But in the frame of the scientist, since the car is traveling right with speed v , the light coming out of the flash light, would appear to have the speed of light c , plus the car speed v , and this violates the postulate that the speed of light is equal to all inertial frames observers.

The only way to fix this, is to take time and space, which Galilean relativity stated as point of agreement among observers in different inertial reference frames, and change them from things that everyone agrees on, to things that everyone can disagree on. In the theory of special relativity, time and space become relative and depend on the observer's reference frame, just like the velocity, momentum and kinetic energy.


To understand this, let's consider the example of a train passing by a station at constant speed v , this speed is measured relative to the platform on the ground where a scientist is standing. And now let's see how a beam of light behave on the train.

Imagine inside the train, a beam of light sent from one of the wagon's floors, towards the roof perpendicular to the train travel direction, and on the roof there's a mirror reflecting the light back to the floor. The distance between the floor and roof of the wagon is d .

We can use the time it takes to travel this distance as a kind of clock, so instead of ticking once per second, this clock will tick once every time the beam crosses the train wagon in one direction. This would be given by:

$$\Delta t = \frac{d}{c} \tag{3}$$

Now consider this from the point of reference of the scientist on the platform ground. Since the train is passing by at speed v on the right, he sees the beam of light traveling in a triangle shape because of the train motion.



So in the frame of the platform, the light takes $\Delta\tilde{t}$ time to cross the train:

$$\begin{aligned} d^2 + (v\Delta\tilde{t})^2 &= (c\Delta\tilde{t})^2 \\ \frac{d^2}{c^2} + \frac{v^2}{c^2} (\Delta\tilde{t})^2 &= (\Delta\tilde{t})^2 \\ \frac{d^2}{c^2} &= (\Delta\tilde{t})^2 \left(1 - \frac{v^2}{c^2}\right) \end{aligned}$$

which means that:

$$\begin{aligned} (\Delta t)^2 &= (\Delta\tilde{t})^2 \left(1 - \frac{v^2}{c^2}\right) \\ \Delta\tilde{t} &= \Delta t \left(\frac{1}{\sqrt{1 - \frac{v^2}{c^2}}} \right) = \Delta t \gamma_v \end{aligned}$$

This factor that you see, it's called the Lorentz factor, and we often write it as γ_v .

This tells us that, the time calculated by the scientist on the ground platform $\Delta\tilde{t}$ is actually greater than the time calculated by the moving train Δt .

In other words, the travel time for the light beam in each reference frame is not equal. When we measure the time between clock ticks for a moving clock, and measure the time between clock ticks for a stationary clock, we find that the time between clock ticks for the moving clock is larger.

This effect is called time dilation.

There is a similar effect called length contraction, where rods that are moving appear to have a shorter length than the same rods when they are standing still. However length contraction only happens along the direction of motion, in the direction perpendicular to motion, no length contraction happens.

$$\Delta\tilde{l} = \Delta l \frac{1}{\gamma_v} \tag{4}$$

References

- [1] eigenchris, *Relativity*, YouTube video series,
<https://www.youtube.com/playlist?list=PLJHszsWbB6hqlw73QjgZcFh4DrkQLSCQa>