

Checking for $w \in H^1(\Omega); v \in H^3(\Omega); v_0 \in H^1$

Contents

1	The H^1 regularity of C using the H^3 regularity of the horizontal velocity v	2
2	Using the upgraded, from L^2 to H^1 regularity for the initial data v_0 to have strong convergence of u_3^ξ.	2
3	$w \in H^1(\Omega)$ — Can we have strong convergence for w not assuming anything more than the original L^2 regularity for u_0?	3
3.1	$v = u_H \in H^2$	3
3.2	Estimate ∇w by Δv	3
3.3	Obtaining the weak convergence of Cu_3 by $u_3 \rightarrow u_3$ strongly in $L_t^2 L_x^2$	4

1 The H^1 regularity of C using the H^3 regularity of the horizontal velocity v

The strong convergence of the concentration C can be obtained through an H^3 horizontal velocity v following the idea on p34 in the non matching horizontal vertical viscosity and diffusivity article. In the Strong Justification article there is a theorem on p15 that states L^2H^3 regularity for a velocity function — but starting from an H^2 initial velocity v_0 for the **full, anisotropic Navier-Stokes equations**, and just checking the boundary conditions / periodicity constraint what we arrive to is not there, as that is in chapter 3, and that regularity result is for the **primitive equations**. If we check what results are stated, specifically for the full Navier-Stokes, they are stated in Theorem 1.1, 1.2 and Remark 1.1 / iii), and they don't state obtaining a higher order H^k regularity for v than the regularity to begin with for v_0 . For the result on p15 (L^2H^3) they use again these ideas we had seen before for the case of primitive equations: the pressure is a function of only the horizontal variables, can be expressed in special ways and brought of integrals. This suggests that if we want to use an H^3 -regular v , we need $v_0 \in H^3$ to begin with.

2 Using the upgraded, from L^2 to H^1 regularity for the initial data v_0 to have strong convergence of u_3^ϵ .

Theorem 1.1 in the Strong Justification article and Remark 1.1 / iii) state a result, for periodic functions that we could use. This states the strong convergence of the vertical velocity for the case of periodic velocities. The higher order boundary conditions for the velocity needs to be checked but this looks maybe the most realistic / probable path, as this result is stated for the anisotropic scaled Navier-Stokes equations.

3 $w \in H^1(\Omega)$ — Can we have strong convergence for w not assuming anything more than the original L^2 regularity for u_0 ?

3.1 $v = u_H \in H^2$

We have obtained $L_t^2 H_x^2$ regularity for the horizontal velocity v . Using the Gronwall inequality we arrive to:

$$\int_0^t \|\Delta v\|_2^2, \int_0^t \|\partial_{33} v\|_2^2 \leq K_v(t).$$

3.2 Estimate ∇w by Δv

We need w to be in $H^1(\Omega)$ in space (with that we have the complete $\mathbf{u}$ in H_x^1), so we need the **full** gradient of the vertical velocity to be in L_x^2 .

We use the

$$w(x, y, z, t) = - \int_0^z \nabla_H \cdot v(x, y, \xi, t) d\xi$$

equation, or, in the case of $\partial_z w$, we can use directly the divergence-free condition.

i) $\operatorname{div}(\mathbf{u}) = \nabla_H \cdot v + \partial_z w = 0$, thus we have $\partial_z w = -\nabla_H \cdot v$. As v is in $L^2 H^2$, we have that $\partial_z w = -\nabla_H \cdot v \in L^2 H^1$.

ii) For $i = 1, 2$

$$\partial_i w(x, y, z, t) = -\partial_i \int_0^z \nabla_H \cdot v(x, y, \xi, t) d\xi = - \int_0^z \partial_1 \partial_i v_1 + \partial_2 \partial_i v_2$$

$$\int_{\Omega} |\partial_i w(x, y, z, t)| \leq \int_{\Omega} \int_0^h |\partial_1 \partial_i v_1| + |\partial_2 \partial_i v_2| = \int_M \int_0^h \int_0^h |\partial_1 \partial_i v_1| + |\partial_2 \partial_i v_2|$$

$$\int_{\Omega} |\partial_i w(x, y, z, t)|^2 \leq C \cdot h \int_{\Omega} |\partial_1 \partial_i v_1|^2 + |\partial_2 \partial_i v_2|^2 \leq C \|\Delta_H v\|_2^2$$

As a consequence we have a bound for the horizontal derivatives of w as well:

$$\int_0^t \|\nabla_H w\|_2^2 \leq C \int_0^t \|\Delta_H v\|_2^2.$$

With $\partial_z w$ in $L^2 H^1$ and $\nabla_H w$ in $L^2 L^2$ we obtain that w is in $L_t^2 H_x^1$.

3.3 Obtaining the weak convergence of Cu_3 by $u_3 \rightarrow u_3$ strongly in $L_t^2 L_x^2$

We have the weak convergence of C^ϵ to C in $L_t^2 L_x^2$. We need $u_3 \rightarrow u_3$ strongly in $L_t^2 L_x^2$ in order to obtain the weak convergence of the product.

How to obtain the strong convergence of u_3^ϵ using its H^1 space regularity?

- Is it [through a similar way as in the original, Azerad-Guillen article?](#) The perturbation theorem or something similar to the Aubin-Lions lemma I think / it seems / does not work, because we need $\|\tau u_3^\epsilon - u_3^\epsilon\|_{L^2(0, T-h, (H^2)^*)}$, but the equation for $\partial_t u_3^\epsilon$ contains an ϵ^2 , with which we would have to divide.

$$\epsilon^2 \int_0^{T-h} |\tau_h u_3 - u_3| \leq \int_0^{T-h} \epsilon^2 (\|\tau_h u_3\|_2 + \|u_3\|_2) \|v_3\|_2 \leq \epsilon^2 \int_0^{T-h} C \|v_3\|_2,$$

which gives an ϵ -independent

$$\int_0^{T-h} |\tau_h u_3 - u_3| \leq \int_0^{T-h} C \|v_3\|_2,$$

however it should go to zero as $\epsilon \rightarrow 0$.

Similarly, the Aubin-Lions lemma requires the time derivative $\partial_t u_3$ to be in $L^2 H^{2*}$, but we have a division by ϵ^2 .

$$\left(\frac{\partial u_3}{\partial t}, \tilde{u}_3 \right) = \dots + \frac{1}{\epsilon} \left(\beta u_1, \tilde{u}_3 \right) - \frac{1}{\epsilon^2} \left(\partial_3 p, \tilde{u}_3 \right)$$

- Is it [like from around the proof that is on page 25 \(where the \$\|\nabla w\|\$ idea was used\)](#)? That is for strong solutions and H^2 initial data.
- Is it [as for the case of \$H^1\$ initial velocity](#)? This also uses $v_0 \in H^1$.

At this point, the strong convergence of the vertical velocity could either be obtained

- i) through assuming the original L^2 regularity of v_0 and using the additionally obtained L^2H^2 regularity for v , or,
- ii) if that (and w being in H_x^1) does not result to be sufficient, use a minimal increase, to H^1 from L^2 in the regularity of the initial data v_0 , combined with some additional Neumann boundary conditions, to be able to apply the analogous version of the first theorem in the Strong Convergence Justification article, which is still for weak solutions (the second one is for strong solutions). This option still seems to be less demanding compared to proving H^1 regularity for C , which requires an H^3 regularity for the horizontal velocity.