

A generalised weak convergence result for the mathematical equations of the polluted atmosphere expressed in downwind matching coordinates

Donatella Donatelli and Nóra Juhász *

Department of Information Engineering, Computer Science and Mathematics
University of L'Aquila
67100 L'Aquila, Italy.
donatella.donatelli@univaq.it, norjuh@univaq.it

Abstract

A widely used approach to mathematically describe the atmosphere is to consider it as a geophysical fluid in a shallow domain — to model it using classical fluid dynamical equations combined with the explicit inclusion of an ϵ parameter representing the small aspect ratio of the physical domain. In our previous paper we proved a weak convergence theorem for the polluted atmosphere described by the Navier-Stokes equations extended by an advection-diffusion equation. We obtained a justification of the generalised hydrostatic limit model including the pollution effect described in the case of a classical, east-north oriented Cartesian coordinate system. Here we give an improvement of this statement: we consider a meteorologically more meaningful coordinate system, and verify that the weak convergence still holds for the governing equations altered because of the consequences of the coordinate change.

Keywords: downwind-matching coordinate system, Navier-Stokes equations, shallow domains, pollution evolution equation, hydrostatic approximation, compactness, weak solutions.

*The research of DD and NJ leading to these results has received funding from the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie Grant Agreement No 642768 (Project Name: ModCompShock).

Contents

1	Introduction	3
1.1	Notations	3
2	The polluted atmosphere model	4
3	Weak formulation and Main Result	11
4	Proof of the main theorem	13
4.1	Energy inequality	13
4.2	A priori estimates and weak convergence	15
4.3	Passing to the limit	16
5	Concluding remarks	18
A	The existence of weak solution of the hydrostatic system of the polluted atmosphere	22

1 Introduction

A local domain on the surface of the rotating Earth is often described by a Cartesian coordinate system where the x, y, z axes represent the directions towards east, north, and upwards, respectively, as in this specific system the Coriolis force takes a simpler form. However, for modelling wind and pollution effects in the atmosphere there is another naturally distinctive viewpoint for fixing the orientation of the coordinate system: the introduction of downwind and crosswind, and to match the horizontal axes according to these meteorological circumstances. In this paper we take this idea as a foundation, and we migrate the result of our previously stated weak convergence theorem to this more considerably chosen coordinate system.

As in this scenario the advection is dominant compared to diffusion effects along the x axis, this approach naturally brings an additional ϵ term in the Laplacian of the concentration equation, the simplification of the limit model, and brings the challenge of proving a weak convergence result for the product of the vertical velocity and the concentration, as in the energy inequality the $\partial_1 C$ term is replaced by $\epsilon \partial_1 C$. This issue is overcome by the construction of new estimates obtained by multiplying the governing equations by higher order terms.

Additional ϵ ., [9] idea.

Modified boundary conditions. Explain the necessity of changing the sediment functions todo.

The intuitive meaning of the concept of downwind and crosswind, and the velocity for this choice of coordinates.

This paper is organised in the following way.

We end up this section with some notations we will use through the paper.

1.1 Notations

- a) ϵ = the aspect ratio,
- b) ν = viscosity,
- c) Ω_ϵ, Ω = the original and the rescaled (ϵ -independent) domain, respectively,
- d) ∇ stands for the gradient vector; specifically, $(\partial_x, \partial_y, \partial_z)$ before, and $(\partial_1, \partial_2, \partial_3)$ after rescaling,

- e) $\nabla_\nu = (\nu_x^{1/2}\partial_x, \nu_y^{1/2}\partial_y, \nu_z^{1/2}\partial_z)$ on Ω_ϵ , $\nabla_\nu = (\nu_1^{1/2}\partial_1, \nu_2^{1/2}\partial_2, \nu_3^{1/2}\partial_3)$ on the rescaled domain Ω ,
- f) Δ_ν stands for the anisotropic Laplacian operators $\nu_x\partial_{xx}^2 + \nu_y\partial_{yy}^2 + \nu_z\partial_{zz}^2$ and $\nu_1\partial_{11}^2 + \nu_2\partial_{22}^2 + \nu_3\partial_{33}^2$ on the original and rescaled domains, respectively,
- g) $\mathbf{g}$ represents the force due to gravity, which also includes the centrifugal effect,
- h) φ = the gravity potential term, i.e. $\mathbf{g} = \nabla\varphi$,
- i) $\mathbf{w}$ represents the earth rotation angular speed, $l(y)$ = latitude,
- j) $\alpha = 2f \sin(l(x_2))$, $\beta = 2f \cos(l(x_2))$,
- k) $\mathbf{K}$ represents the diffusion matrix,
- l) $(\cdot, \cdot) =$ the scalar product in $L^2(\Omega)^d$ or the duality $L^p(\Omega)$, $L^{p'}(\Omega)$,
- m) $\langle \cdot, \cdot \rangle_\Gamma =$ the scalar product or duality on the boundary curve Γ ($H^{-1/2}(\Gamma)$, $H^{1/2}(\Gamma)$),
- n) $\mathbf{u}_H =$ the horizontal velocity components (u_1, u_2) ,
- o) $b(\mathbf{u}_H) = \alpha(-u_2, u_1)$,
- p) $L_t^p L_x^q$ is the abbreviated notation for $L^p(0, T; L^q(\Omega))$.

2 The polluted atmosphere model

We begin by choosing a coordinate system, set the domain we start with and an appropriate set of equations describing both atmosphere motions and the effect of pollution. We ignore large scale effects such as the curvature of the Earth. For describing the effect of pollution we use a continuity equation written in Cartesian coordinates for the concentration.

We fix a local Cartesian coordinate system where the axes are independent from the wind (x, y and z are oriented towards east, north, and upwards, respectively). Other possibilities and considerations regarding choosing the coordinate system are described briefly in the concluding remarks.

The domain we consider is a local slice of the atmosphere filled with homogeneous air that we assume to be incompressible, and a pollutant of concentration C , therefore it is defined by the set

$$\Omega_\epsilon = \{(x, y, z) \in \mathbb{R}^3; (x, y) \in \omega, 0 < z < \epsilon h(x, y)\}, \quad (1)$$

where ω is a Lipschitz-domain in $\mathbb{R}^2$, h is a nonnegative Lipschitzian application, and ϵ incorporates the shallowness of the domain.

The equations governing the air velocity $\mathbf{v}$ and pressure q in the atmosphere are the incompressible Navier-Stokes equations, in which we use different viscosities according to vertical or horizontal directions, and the density is taken to be identically equal to one:

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} - \Delta_\nu \mathbf{v} + \nabla q + 2\mathbf{w} \times \mathbf{v} = \mathbf{g} \text{ in } \Omega_\epsilon \times (0, T) \quad (2)$$

$$\nabla \cdot \mathbf{v} = 0 \text{ in } \Omega_\epsilon \times (0, T) \quad (3)$$

Contaminants emitted into the atmosphere are transported by the wind, mixed and dispersed by turbulent behaviour, and deposited onto the ground by gravity. The dispersion of pollutants is commonly modelled by a diffusive equation of the form

$$\partial_t P + \mathbf{v} \cdot \nabla P = \nabla \cdot (\mathbf{K} \nabla P) + Q, \quad (4)$$

where P is the pollution concentration and $\mathbf{K}$ is the 3×3 diffusion matrix — the diagonal terms represent the strength of the diffusion in the x, y, z directions, while the non-diagonal terms reflect the correlation of random motions between each pair of principal directions.

Here Q represents the sources of the pollutant in the atmosphere i.e. its emission, removal from the atmosphere by dry and wet deposition, and chemical reactions.

Now, we perform a vertical scaling to make the domain independent of ϵ .

$$x = x_1, \quad y = x_2, \quad z = \epsilon x_3.$$

For simplicity of notation we will use $\partial_i = \partial_{x_i}$ for $i = 1, 2, 3$.

We have the following new, ϵ - independent domain

$$\Omega = \{(x_1, x_2, x_3) \in \mathbb{R}^3; (x_1, x_2) \in \omega, 0 < x_3 < h(x_1, x_2)\}.$$

The corresponding scaling regarding the kinematic, pressure and viscosity terms are

$$v_1 = u_1^\epsilon, \quad v_2 = u_2^\epsilon, \quad v_3 = \epsilon u_3^\epsilon, \quad p = p^\epsilon,$$

$$\nu_x = \nu_1, \quad \nu_y = \nu_2, \quad \nu_z = \epsilon^2 \nu_3,$$

where the later embodies the different eddy viscosity hypothesis that we mentioned earlier, and p is the pressure term including the gravity potential φ , i.e., $p = q - \varphi$. Basically the scaling of the viscosity parameters can be heuristically deduced from the viscosity dimensions: typical length²/typical time, more details on the subject can be found in [7], [6].

Note that from this point we have $\nabla = (\partial_1, \partial_2, \partial_3)$ and $\Delta_\nu = \nu_1 \partial_{11}^2 + \nu_2 \partial_{22}^2 + \nu_3 \partial_{33}^2$.

We use an analogous rescaling methodology for the diffusivity constants in (4),

$$\begin{bmatrix} K_{xx} & K_{xy} & K_{xz} \\ K_{yx} & K_{yy} & K_{yz} \\ K_{zx} & K_{zy} & K_{zz} \end{bmatrix} = \begin{bmatrix} M_{11} & M_{12} & \epsilon M_{13} \\ M_{21} & M_{22} & \epsilon M_{23} \\ \epsilon M_{31} & \epsilon M_{32} & \epsilon^2 M_{33} \end{bmatrix} \quad (5)$$

Here we emphasise that the diffusivity matrix $\mathbf{M}$ is assumed to have the coercivity property

$$\lambda \|x\|^2 \leq (\mathbf{M}x, x) \quad (6)$$

for all $x \in \mathbb{R}^3$ with the coercivity constant $\lambda > 0$. In other words we require that $\lambda = \min \lambda_i > 0$ where λ_i are the eigenvalues of $\mathbf{M}$.

Finally we obtain the rescaling equations for the pollution concentration and the source term by considering their physical dimensions, namely: typical molar density/typical volume, from which we deduce

$$P = \frac{1}{\epsilon} C^\epsilon, \quad Q = \frac{1}{\epsilon} S^\epsilon.$$

In this model we will use localised point sources as suggested in [22] and we use what is called a parametrised nascent delta function (a Gaussian impulse, specifically) to describe the source term S^ϵ .

Let us consider a gas plume with intensity I originating at emission coordinates $\mathbf{x}_s$ beginning at time t_s . At the beginning we assume zero concentration. An abrupt gas emission is modelled by a source term represented as a product of an approximated delta function $\delta_\kappa(\mathbf{x} - \mathbf{x}_s)$ (centered at $\mathbf{x}_s$ with a constant parameter κ) and a Heaviside function s (switching from 0 to I at time t_s .) In fact, this source term models the continuous emission of the gas at constant rate and fixed location. Basically $s(t; t_s)$ is a function which activates the source, namely $s(t; t_s) := IH(t - t_s)$ where $H(\cdot)$ is the

Heaviside step function and $0 \leq t_s \leq T$. We arrive to the complete source term model of the form

$$S_\kappa(t, t_s, x, x_s) = s(t; t_s) \delta_\kappa(\mathbf{x} - \mathbf{x}_s), \quad (7)$$

while in the limit model the source term S naturally stands for the δ distribution.

Eventually, as we mentioned before, we use a Gaussian approximated delta function to represent the source, so it is convenient to reformulate (7) in terms of ϵ , which is the standard deviation parameter of the distribution describing the source. Thus, we choose κ to be ϵ and the following function will give shape to the source term:

$$\delta_\epsilon(\mathbf{x} - \mathbf{x}_s) = \gamma \epsilon^{-3} e^{-\frac{(\mathbf{x} - \mathbf{x}_s)^2}{\epsilon^2}};$$

and finally

$$S^\epsilon = IH(t - t_s) \gamma \epsilon^{-3} e^{-\frac{(\mathbf{x} - \mathbf{x}_s)^2}{\epsilon^2}}, \quad (8)$$

where γ is a normalising constant. Later we will take the limit as ϵ goes to zero, and the above setting will result in having the δ distribution representing the hydrostatic source term, which is a physically natural choice.

Putting this condition and the previously described rescaling terms together to the original equations (2)-(4), we arrive to the merged anisotropic equations of the polluted atmosphere above inland surface:

$$\partial_t u_1^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_1^\epsilon - \Delta_\nu u_1^\epsilon - \alpha u_2^\epsilon + \epsilon \beta u_3^\epsilon + \partial_1 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (9)$$

$$\partial_t u_2^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_2^\epsilon - \Delta_\nu u_2^\epsilon + \alpha u_1^\epsilon + \partial_2 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (10)$$

$$\epsilon^2 \{ \partial_t u_3^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_3^\epsilon - \Delta_\nu u_3^\epsilon \} - \epsilon \beta u_1^\epsilon + \partial_3 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (11)$$


$$\nabla \cdot \mathbf{u}^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (12)$$

$$\partial_t C^\epsilon + \mathbf{u}^\epsilon \cdot \nabla C^\epsilon = \nabla \cdot (\mathbf{M} \nabla C^\epsilon) + S^\epsilon \text{ in } \Omega \times (0, T) \quad (13)$$

The initial condition for the velocity and the pollution concentration are

$$\mathbf{u}^\epsilon(\cdot, t = 0) = \mathbf{u}_0, \quad C^\epsilon(\cdot, t = 0) = C_0 \quad \text{in } \Omega. \quad (14)$$

To make the above set of equations complete, we need to assign the boundary conditions.



Concerning the pollution concentration it is vital to choose boundary conditions that are physically as accurate as possible and at the same time not unnecessarily complicated. As we can see in many related papers, in the case of LAMs (limited area models) finding the right boundary conditions can be a nontrivial task.

In order to describe the boundary condition in a precise way we need to divide $\Gamma = \partial\Omega$ as follows. The upper boundary of the Ω domain is denoted by Γ_U , the lateral boundary is Γ_L , the lower boundary of the domain is Γ_G , and for the collective $\Gamma_L \cup \Gamma_U$ section we use the notion Γ_A (above the ground). As we mentioned before, the case we are investigating is that of inland domains Ω , i.e. the case of landscapes and cities with no significant nearby water surface. This is very important since the boundary conditions for both wind velocity and pollution concentration in the air are different depending whether we are above solid ground or water surface. The boundary conditions we choose to use in the inland scenario are the following:

- The upper boundary Γ_U of the domain we are considering is chosen to be at the isobar $z = h$ representing the planetary boundary layer (see [5]), therefore we assume

$$\mathbf{u}_H^\epsilon = 0, \quad u_3^\epsilon = 0, \quad C^\epsilon = 0 \quad \text{on } \Gamma_U \times (0, T) \quad (15)$$

- On Γ_L we use the same boundary conditions as on the upper level, i.e. we have

$$\begin{array}{ccc}
& \mathbf{u}^\epsilon = 0, C^\epsilon = 0 & \\
\mathbf{u}^\epsilon = 0, C^\epsilon = 0 & \square \quad \Omega & \mathbf{u}^\epsilon = 0, C^\epsilon = 0 \\
& \mathbf{u}^\epsilon = 0, \partial_1 C^\epsilon \text{ and } \partial_2 C^\epsilon \text{ are bounded, } \partial_3 C^\epsilon = 0 &
\end{array}$$

$$\mathbf{u}_H^\epsilon = 0, \quad u_3^\epsilon = 0, \quad C^\epsilon = 0 \quad \text{on } \Gamma_L \times (0, T). \quad (16)$$

Remark 1. *If we wanted to use $C^\epsilon(x, y, z) = c$ with some c constant value, we could make a change of variables distracting that background flow and arrive back to the same mathematical problem with C being zero.*

- In this model we neglect topological variations, and the $z = 0$ plane represents the lower boundary. On the ground level we have

$$\mathbf{u}_H^\epsilon = 0, \quad u_3^\epsilon = 0 \quad \text{on } \Gamma_G \times (0, T). \quad (17)$$

This is the mathematical way of expressing no-slip boundary conditions for the horizontal wind speed and the impervious nature of the ground with respect to the wind. As for the concentration, the way we can choose ground level boundary conditions is the following. Firstly, we assume that the pollution is not absorbed by the ground, so we use

$$\partial_3 C^\epsilon = 0 \text{ on } \Gamma_G$$

(see also for example [21]). As we will see later in the case of the coastal domain, this condition can be relaxed without significantly affecting the proof. For the horizontal partial derivatives of the pollution, we require finiteness in the boundary norm, i.e. we have

$$\partial_1 C^\epsilon, \partial_2 C^\epsilon \text{ are finite in } L^2(0, T; \Gamma_G).$$

Specifically, for the pollution concentration reaching the ground level we use a similar approach as the one classically used for the wind traction reaching the top of the ocean (see for example [6], [19]). We assume that the functions $\partial_1 C^\epsilon, \partial_2 C^\epsilon, \partial_3 C^\epsilon$ are equal to some sufficiently regular functions defined on the respective boundary. More accurately, we assume that we have $\nabla C^\epsilon = \boldsymbol{\sigma}$ on the lower boundary, where $\boldsymbol{\sigma} \in L^2(0, T; L^2(\Gamma_G))$, and the σ_i terms denote the components of $\boldsymbol{\sigma}$. In particular in this inland case σ_3 is equal to zero on the Γ_G .

Finally, the boundary conditions to describe the anisotropic system can be summarised as follows:

$$\begin{aligned} \mathbf{u}^\epsilon &= 0 \text{ on } \Gamma \times (0, T), \\ C^\epsilon &= 0 \text{ on } (\Gamma_U \cup \Gamma_L) \times (0, T), \\ \partial_1 C^\epsilon &= \sigma_1, \partial_2 C^\epsilon = \sigma_2, \partial_3 C^\epsilon = 0 \text{ on } \Gamma_G \times (0, T), \\ \boldsymbol{\sigma} &= (\sigma_1, \sigma_2, 0) \in L^2(0, T; \Gamma_G) \end{aligned} \tag{18}$$

Note that we have a domain where the lower and upper boundaries represent a physically existent layer (i.e. the ground and the planetary boundary layer), but outside the lateral boundary the physical world continues without influence and without any particular external condition. On boundaries of such nature we have to apply OBCs (open boundary conditions), which is not a trivial task, see for example [20]. We need to avoid any spurious reflection or constraint which is unnatural, but at the same time we want the model to remain mathematically manageable.

If we assume $\mathbf{u}^\epsilon = O(1)$, then neglecting the ϵ^2 and ϵ in the anisotropic equations, we formally arrive to the hydrostatic Navier-Stokes equations (i.e., primitive equations) combined with pollution.

$$\partial_t u_1 + \mathbf{u} \cdot \nabla u_1 - \Delta_\nu u_1 - \alpha u_2 + \partial_1 p = 0 \text{ in } \Omega \times (0, T) \tag{19}$$

$$\partial_t u_2 + \mathbf{u} \cdot \nabla u_2 - \Delta_\nu u_2 - \alpha u_1 + \partial_2 p = 0 \text{ in } \Omega \times (0, T) \tag{20}$$

$$\partial_3 p = 0 \text{ in } \Omega \times (0, T) \tag{21}$$

$$\nabla \cdot \mathbf{u} = 0 \text{ in } \Omega \times (0, T) \tag{22}$$

$$\partial_t C + \mathbf{u} \cdot \nabla C = \nabla \cdot (\mathbf{M} \nabla C) + S \text{ in } \Omega \times (0, T) \tag{23}$$

$$u_1 = u_2 = u_3 n_3 = 0 \text{ on } \Gamma \times (0, T) \quad (24)$$

$$u_i(\cdot, t = 0) = u_{0i}, C(\cdot, t = 0) = C_0 \text{ in } \Omega, i = 1, 2 \quad (25)$$

$$\begin{aligned} C &= 0 \text{ on } (\Gamma_U \cup \Gamma_L) \times (0, T), \\ \partial_1 C &= \sigma_1, \partial_2 C = \sigma_2, \partial_3 C = 0, \boldsymbol{\sigma} = (\sigma_1, \sigma_2, 0) \in L^2(0, T; \Gamma_G) \end{aligned} \quad (26)$$

Remark 2. Here S stands for the δ limit distribution. We keep the notation S for generality, since as we will see later, instead of the approximated delta functions and the delta distribution, it is possible to use any sort of "sufficiently smooth" source term (which can be independent from ϵ in the anisotropic case as well) that is bounded in $L_t^2 L_x^2$.

3 Weak formulation and Main Result

In this section we describe the weak formulation of the merged equations in both the anisotropic and hydrostatic case we are going to use in the paper, and we state our main result.

We need to define the following spaces:

$$C_{\Gamma_A}^\infty(\Omega) = \{\varphi \in C^\infty(\bar{\Omega}); \varphi = 0 \text{ on some neighbourhood of } \Gamma_A\}$$

$$H_{\Gamma_A}^k(\Omega) = \overline{C_{\Gamma_A}^\infty(\Omega)}^{H^k(\Omega)} = \{v \in H^k(\Omega); v = 0 \text{ on } \Gamma_A\}$$

$$\mathbf{V} = \{\mathbf{v} \in H_{\Gamma_A}^1(\Omega) \times H_{\Gamma_A}^1(\Omega) \times H_0^1(\Omega); \nabla \cdot \mathbf{v} = 0 \text{ in } \Omega\}$$

$$H(\partial_3, \Omega) = \{v \in L^2(\Omega); \partial_3 v \in L^2(\Omega)\}$$

$$H_0(\partial_3, \Omega) = \overline{C_0^\infty(\Omega)}^{H(\partial_3, \Omega)} = \{v \in H(\partial_3, \Omega); v n_3 = 0 \text{ on } \Gamma\}$$

$$\mathbf{W} = \{\mathbf{u} \in H_{\Gamma_A}^1(\Omega) \times H_{\Gamma_A}^1(\Omega) \times H_0(\partial_3, \Omega); \nabla \cdot \mathbf{u} = 0 \text{ in } \Omega\}$$

Then the weak formulation of the hydrostatic system (19) - (26) takes the following form.

Definition 1. Find $(\mathbf{u}, C)$ such that $\mathbf{u} = (\mathbf{u}_H, u_3) \in L^2(0, T; \mathbf{W})$ with $\mathbf{u}_H \in L^\infty(0, T; L^2(\Omega)^2)$, and $C \in L^\infty(0, T; L^2(\Omega))$ with $C \in L^2(0, T; H^1(\Omega))$, such that

$$\begin{aligned} & \int_0^T \left[-(\mathbf{u}_H, \partial_t \tilde{\mathbf{u}}_H) + (\nabla_\nu \mathbf{u}_H, \nabla_\nu \tilde{\mathbf{u}}_H) - (\mathbf{u}_H, (\mathbf{u} \cdot \nabla) \tilde{\mathbf{u}}_H) + (b(\mathbf{u}_H), \tilde{\mathbf{u}}_H) \right] dt \\ & = (\mathbf{u}_{0H}, \tilde{\mathbf{u}}_H(0)) \end{aligned} \quad (27)$$

and

$$\begin{aligned} & \int_0^T \left[-(C, \partial_t \tilde{C}) - (\mathbf{u}C, \nabla \tilde{C}) + (\mathbf{M} \nabla C, \nabla \tilde{C}) - \int_{\Gamma_G} \boldsymbol{\sigma} \mathbf{M} \tilde{C} \tilde{\mathbf{n}}_{\Gamma_G} \right] dt \\ & = (C(0), \tilde{C}(0)) + \int_0^T (S, \tilde{C}) dt \end{aligned} \quad (28)$$

for all $(\tilde{\mathbf{u}}, \tilde{C})$ with $\tilde{\mathbf{u}} = (\tilde{\mathbf{u}}_H, \tilde{u}_3) \in H^1(0, T, \mathbf{W})$, $\tilde{\mathbf{u}}_H(T) = 0$ and $\partial_3 \tilde{\mathbf{u}}_H \in L^\infty(0, T; L^3(\Omega)^2)$ and $\tilde{C} \in L^2(0, T, H_{\Gamma_A}^2(\Omega)) \cap H^1(0, T, L^2)$, $\tilde{C}(T) = 0$.

Note that the term $\int_0^T (S, \tilde{C}) dt$ makes sense since in this case S denotes the delta distribution and we have

$$(S, \tilde{C}) = \int_\Omega \delta(\mathbf{x} - \mathbf{x}_s) \tilde{C}(\mathbf{x}) d\mathbf{x} = \tilde{C}(\mathbf{x}_s).$$

The existence of weak solution for the hydrostatic system (19) - (26) can be proved in the spirit of [19]. In appendix A we will outline the main steps of the proof and the differences compared to the ocean model.

The weak form of the anisotropic system is as follows.

Definition 2. Find $\mathbf{u}^\epsilon = (\mathbf{u}_H^\epsilon, u_3^\epsilon) \in L^2(0, T; \mathbf{V}) \cap L^\infty(0, T; L^2(\Omega)^3)$ and $C^\epsilon \in L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$, such that

$$\begin{aligned} & \int_0^T \left[-(\mathbf{u}_H^\epsilon, \partial_t \tilde{\mathbf{u}}_H) + (\nabla_\nu \mathbf{u}_H^\epsilon, \nabla_\nu \tilde{\mathbf{u}}_H) - (\mathbf{u}_H^\epsilon, (\mathbf{u}^\epsilon \cdot \nabla) \tilde{\mathbf{u}}_H) + (b(\mathbf{u}_H^\epsilon), \tilde{\mathbf{u}}_H) \right] dt \\ & + \epsilon \beta \int_0^T [(u_3^\epsilon, \tilde{u}_1) - (u_1^\epsilon, \tilde{u}_3)] dt \\ & + \epsilon^2 \int_0^T [-(u_3^\epsilon, \partial_t \tilde{u}_3) + (\mathbf{u}^\epsilon \cdot \nabla u_3^\epsilon, \tilde{u}_3) + (\nabla_\nu u_3^\epsilon, \nabla_\nu \tilde{u}_3)] dt \\ & = (\mathbf{u}_{0H}^\epsilon, \tilde{\mathbf{u}}_H(0)) + \epsilon^2 (u_{03}^\epsilon, \tilde{u}_3(0)) \end{aligned} \quad (29)$$

and

$$\begin{aligned} & \int_0^T \left[- (C^\epsilon, \partial_t \tilde{C}) - (\mathbf{u}^\epsilon C^\epsilon, \nabla \tilde{C}) + (\mathbf{M} \nabla C^\epsilon, \nabla \tilde{C}) - \int_{\Gamma_G} \boldsymbol{\sigma} \mathbf{M} \tilde{C} \vec{\mathbf{n}}_{\Gamma_G} \right] dt \\ &= (C^\epsilon(0), \tilde{C}(0)) + \int_0^T (S^\epsilon, \tilde{C}) dt \end{aligned} \quad (30)$$

for all $\tilde{\mathbf{u}} = (\tilde{\mathbf{u}}_H, \tilde{u}_3) \in H^1(0, T; \mathbf{V})$ with $\tilde{\mathbf{u}}(T) = 0$ and $\tilde{C}$ such that $\tilde{C} \in H^1(0, T, H_{\Gamma_A}^2(\Omega))$ and $\tilde{C}(T) = 0$.

Note that since in our case the source term S^ϵ is given by (7), the last term in (30) is well defined. If we use a source function of a different and more general form, it needs to satisfy the $L_t^2 L_x^2$ boundedness criteria for the weak formulation to make sense.

We will omit the proof of the existence of weak solutions (see Definition 2.) for the anisotropic system (9) - (18) since it follows by standard finite-dimensional Galerkin approximation.

Now we are ready to state our main result.

Theorem 1. *Let $\mathbf{u}_0 \in L^2(\Omega)^3$, with $\nabla \cdot \mathbf{u}_0 = 0$, $\mathbf{u}_0 \cdot \mathbf{n} = 0$ on $\partial\Omega$; let $C_0 \in L^2(\Omega)$, $C_0 = 0$ on $\partial\Omega$, and assume that the boundary conditions (18) hold; then as the aspect ratio ϵ tends to zero, any weak solution $(\mathbf{u}^\epsilon, C^\epsilon)$ of the anisotropic equations (9) - (13) converge to a weak solution $(\mathbf{u}, C)$ of the hydrostatic equations of the polluted atmosphere (19) - (23).*

4 Proof of the main theorem

This section is devoted to the proof of Theorem 1. In the extended case with the pollution equation the proof relies on a priori estimates gained from the energy inequality and, as we will see this is sufficient to pass to the limit in the linear terms. Concerning the nonlinear terms we will prove a uniform in ϵ space time estimate for $\mathbf{u}^\epsilon$ and C^ϵ that will allow us to apply a compactness criterion proved in [6].

4.1 Energy inequality

As we mentioned before, one of the most fundamental tools in this proof is the energy inequality. If we calculate it for the anisotropic system, we obtain that for a.e. $t \in [0, T]$ the following holds:

$$\begin{aligned}
& \frac{1}{2} (\|\mathbf{u}_H^\epsilon(t)\|_{L^2}^2 + \epsilon^2 \|u_3^\epsilon(t)\|_{L^2}^2 + \|C^\epsilon(t)\|_{L^2}^2) + \int_0^t [\|\nabla_\nu \mathbf{u}_H^\epsilon\|_{L^2}^2 + \epsilon^2 \|\nabla_\nu u_3^\epsilon\|_{L^2}^2] d\tau \\
& + \int_0^t (\mathbf{M} \nabla C^\epsilon, \nabla C^\epsilon) d\tau \\
& \leq \int_0^t (S^\epsilon, C^\epsilon) d\tau + \frac{1}{2} (\|\mathbf{u}_H^\epsilon(0)\|_{L^2}^2 + \epsilon^2 \|u_3^\epsilon(0)\|_{L^2}^2 + \|C_0^\epsilon\|_{L^2}^2) \\
& + \int_0^t \int_{\Gamma_G} ((\nabla C^\epsilon) \mathbf{M}) C^\epsilon \vec{\mathbf{n}}_{\Gamma_G} d\tau \\
& = I_1 + I_2 + I_3.
\end{aligned} \tag{31}$$

From the hypothesis on the initial data and since the S^ϵ source term is in $L_x^\infty(\Omega)$, I_1 and I_2 are bounded. The only terms we need to work on further are the expressions in I_3 .

By using the Hölder inequality and the generalised Young inequality for any $\zeta > 0$, we can estimate this term in the following way.

$$\begin{aligned}
I_3 &= \int_0^t \int_{\Gamma_G} \sum_{i=1}^3 \left(\left(\sum_{j=1}^3 M_{ij} \partial_j C^\epsilon \right) C^\epsilon \mathbf{n}_{\Gamma_G, i} \right) d\tau \\
&\leq \sum_{i,j=1}^3 M_{ij} \int_0^t \|\partial_i C^\epsilon\|_{L^2(\Gamma_G)} \|C^\epsilon\|_{L^2(\Gamma_G)} d\tau \\
&\leq \sum_{i,j=1}^3 \frac{1}{\zeta} M_{ij} \frac{1}{2} \int_0^t \|\partial_i C^\epsilon\|_{L^2(\Gamma_G)}^2 d\tau + \zeta M_{ij} \int_0^t \frac{1}{2} \|C^\epsilon\|_{L^2(\Gamma_G)}^2 d\tau \\
&\leq \sum_{i,j=1}^3 \frac{1}{\zeta} M_{ij} \frac{1}{2} \int_0^t \|\partial_i C^\epsilon\|_{L^2(\Gamma_G)}^2 d\tau + \zeta \frac{1}{2} M_{ij} c_T \int_0^t \|C^\epsilon\|_{W^{1,2}(\Omega)}^2 d\tau \\
&\leq \sum_{i,j=1}^3 \left(\frac{1}{\zeta} M_{ij} \frac{1}{2} \int_0^t \|\partial_i C^\epsilon\|_{L^2(\Gamma_G)}^2 d\tau + \zeta \frac{1}{2} M_{ij} c_T \int_0^t \|\nabla C^\epsilon\|_{L^2(\Omega)}^2 d\tau \right. \\
&\quad \left. + \zeta \frac{1}{2} M_{ij} c_T \int_0^t \|C^\epsilon\|_{L^2(\Omega)}^2 d\tau \right),
\end{aligned} \tag{32}$$

where the c_T constant is provided by the trace theorem, and the boundary term is bounded because of the a priori assumptions made on the boundary conditions.

After applying the coercivity property (6) of the diffusion matrix, (31) now takes the form of the following energy inequality,

$$\begin{aligned}
& \frac{1}{2} (\|\mathbf{u}_H^\epsilon(t)\|_{L^2}^2 + \epsilon^2 \|u_3^\epsilon(t)\|_{L^2}^2 + \|C^\epsilon(t)\|_{L^2}^2) + \int_0^t [\|\nabla_\nu \mathbf{u}_H^\epsilon\|_{L^2}^2 + \epsilon^2 \|\nabla_\nu u_3^\epsilon\|_{L^2}^2] d\tau \\
& + \int_0^t \left[\left(\lambda - \sum_{i,j=1}^3 \zeta \frac{1}{2} M_{ij} c_T \right) \|\nabla C^\epsilon\|_{L^2}^2 \right] d\tau \\
& \leq \sum_{i,j=1}^3 \frac{1}{\zeta} M_{ij} \frac{1}{2} \int_0^t \|\sigma_i\|_{L^2(\Gamma_G)}^2 d\tau + \frac{1}{2} (\|\mathbf{u}_H^\epsilon(0)\|_{L^2}^2 + \epsilon^2 \|u_3^\epsilon(0)\|_{L^2}^2 + \|C^\epsilon(0)\|_{L^2}^2) \\
& + \int_0^t (S^\epsilon, C^\epsilon) d\tau + \sum_{i,j=1}^3 \zeta \frac{1}{2} M_{ij} c_T \int_0^t \|C^\epsilon\|_{L^2(\Omega)}^2 d\tau.
\end{aligned} \tag{33}$$

It is now straightforward to see that the coefficients of the pollution terms are positive, since the $\zeta = \zeta(\lambda)$ constant of the generalised Young inequality can be freely chosen in a way such that the requirement $(\lambda - \sum \zeta \frac{1}{2} M_{ij} c_T) > \lambda/2 > 0$ is satisfied.

4.2 A priori estimates and weak convergence

From the final form of the energy inequality we obtain uniform a priori estimates that we summarise in the following proposition.

Proposition 1. *Let $u_1^\epsilon, u_2^\epsilon, u_3^\epsilon$ and C^ϵ be the weak solutions of the system (9) - (13) in the sense of Definition 2. Then it holds,*

$$u_1^\epsilon, u_2^\epsilon, \epsilon u_3^\epsilon, C^\epsilon \quad \text{are bounded in } L^\infty(0, T; L^2(\Omega)), \tag{34}$$

$$u_3^\epsilon \quad \text{is bounded in } L^2(0, T; L^2(\Omega)), \tag{35}$$

$$u_1^\epsilon, u_2^\epsilon, \epsilon u_3^\epsilon \quad \text{are bounded in } L^2(0, T; H^1(\Omega)), \tag{36}$$

$$C^\epsilon \quad \text{is bounded in } L^2(0, T; H^1(\Omega)). \tag{37}$$

Proof. By applying the classical Gronwall inequality, (34), (36) and (37) follow directly by the energy inequality (33). On the other hand, (35) can be verified by using the divergence free condition and that as a consequence, we have $\partial_3 u_3 = -\partial_1 u_1 - \partial_2 u_2$ bounded in $L_t^2 L_x^2$. The bound holds for u_3 itself as well, owing to the Poincaré inequality in the vertical direction, and using that u_3 vanishes on the boundary. $\square$

As a consequence, up to subsequences still denoted by the same way, we have the following weak convergence results.

$$\mathbf{u}_H^\epsilon \rightharpoonup \mathbf{u}_H \quad \text{weakly in } L_t^\infty L_x^2 \cap L_t^2 H_x^1, \quad (38)$$

$$\epsilon^2 u_3^\epsilon \rightarrow 0 \quad \text{strongly in } L_t^\infty L_x^2 \cap L_t^2 H_x^1, \quad (39)$$

$$u_3^\epsilon \rightharpoonup u_3 \quad \text{weakly in } L_t^2 L_x^2, \quad (40)$$

$$C^\epsilon \rightharpoonup C \quad \text{weakly in } L_t^\infty L_x^2 \cap L_t^2 H_x^1 \quad (41)$$

4.3 Passing to the limit

In order to conclude the proof of Theorem 1 we pass to the limit in the weak formulation (29)-(30). To take the limit for the linear velocity terms that do not include the concentration function C^ϵ we can use (38), (39) and (40), and the bounds (35), (36). As for the nonlinear terms, the convergence can be shown using a time compactness theorem that is based on a result by Simon [17]. The key point is to apply a sort of generalisation of the classical translation criterium of Riesz-Frechet-Kolmogorov which enables us to get strong convergence for the horizontal velocities. This is achieved by establishing a bound for the perturbation of $\mathbf{u}_H$ of the form $\|\tau_h \mathbf{u}_H - \mathbf{u}_H\|_{L^p(0, T-h, \mathbf{Y})} \leq \varphi(h) + \psi(\epsilon)$, where $\mathbf{Y}$ is a Banach space, φ and ψ are appropriate functions with their limits vanishing at zero, and $\tau_h \mathbf{u}_H = \mathbf{u}_H(t+h)$. After obtaining strong convergence for $\mathbf{u}_H$, the proof of the convergence for these nonlinear terms can be closed by applying basic interpolation techniques and the generalised Holder inequality. The details are omitted since they are analogous to those described in [6].

For the linear terms including the concentration C^ϵ we can directly apply the weak convergence result of (41).

We get

$$\begin{aligned} \int_0^T (C^\epsilon, \partial_t \tilde{C}) dt &\rightarrow \int_0^T (C, \partial_t \tilde{C}) dt, \\ \int_0^T (\mathbf{M} \nabla C^\epsilon, \nabla \tilde{C}) dt &\rightarrow \int_0^T (\mathbf{M} \nabla C, \nabla \tilde{C}) dt. \end{aligned}$$

For the source term, by applying standard results for distribution functions, we get $(S^\epsilon, \tilde{C}) \rightarrow (S, \tilde{C})$.

Finally we have to deal with the nonlinear term $(\mathbf{u}^\epsilon C^\epsilon, \nabla \tilde{C})$.

By analysing this nonlinear term in a more precise way we see that $u_i^\epsilon C^\epsilon$ for $i = 1, 2$ is weakly convergent since as we mentioned before, we

have the strong convergence of the horizontal velocities, while in the case of $i = 3$ this property does not hold. We overcome this difficulty by proving a compactness property for the pollution concentration C^ϵ . The key tool will be the following compactness criterion (for the proof see Theorem 5.1 in [6]).

Theorem 2. *Let $T > 0$, and let the Banach spaces $\mathbf{X} \hookrightarrow \mathbf{B} \hookrightarrow \mathbf{Y}$, where $\mathbf{X}$ is compactly embedded in $\mathbf{B}$, and $\mathbf{B}$ is continuously embedded in $\mathbf{Y}$. Let $(f_\epsilon)_{\epsilon>0}$ be a family of functions of $L^p(0, T; \mathbf{X})$, $1 \leq p \leq \infty$, with the extra condition $(f_\epsilon)_{\epsilon>0} \subset C(0, T; \mathbf{Y})$ if $p = \infty$, such that*

- a) $(f_\epsilon)_{\epsilon>0}$ is bounded in $L^p(0, T; \mathbf{X})$,
- b) $\|\tau_h f_\epsilon - f_\epsilon\|_{L^p(0, T-h; \mathbf{Y})} \leq \varphi(h) + \psi(\epsilon)$ with

$$\begin{cases} \lim_{h \rightarrow 0} \varphi(h) = 0, \\ \lim_{\epsilon \rightarrow 0} \psi(\epsilon) = 0. \end{cases}$$

Then the family $(f_\epsilon)_{\epsilon>0}$ possesses a cluster point in $L^p(0, T; \mathbf{B})$ and also in $C(0, T; \mathbf{B})$ if $p = \infty$ as $\epsilon \rightarrow 0$.

We will be able to apply the statement of this theorem for $p = 2$ and the spaces $H^1 \hookrightarrow L^2 \hookrightarrow H^{2*}$, where H^{2*} is the dual of H^2 , obtaining that

$$C^\epsilon \rightarrow C \text{ strongly in } L_t^2 L_x^2,$$

and thus we have the weak convergence result $u_3^\epsilon C^\epsilon \rightharpoonup u_3 C$.

In the following we will verify that the theorem's conditions hold for our choice of spaces and p . Since (37) is already given, we only have to show the bound $\|\tau_h C^\epsilon - C^\epsilon\|_{L^2(0, T-h, H^{2*})} \leq \varphi(h) + \psi(\epsilon)$ in order to close the argument regarding the weak convergence of $u_3^\epsilon C^\epsilon$.

Proposition 2. *The estimate $\|\tau_h C^\epsilon - C^\epsilon\|_{L^2(0, T-h, H^{2*})} \leq c \cdot h^{1/4}$ holds.*

Proof. The spatial weak form of the anisotropic system (9) - (13) is

$$\left(\frac{\partial C^\epsilon}{\partial t}, \tilde{C} \right) - (\mathbf{u}^\epsilon C^\epsilon, \nabla \tilde{C}) + (\mathbf{M} \nabla C^\epsilon, \nabla \tilde{C}) = (S^\epsilon, \tilde{C}), \quad (42)$$

where in this case the equality has to hold for all $\tilde{C} \in H_x^2$.

Now we take a test function $\tilde{C} \in H_x^2$ in (42) and integrate over $(t, t+h)$,

i. e. we have

$$(\tau_h C^\epsilon(t) - C^\epsilon(t), \tilde{C}) = \int_t^{t+h} \iota^\epsilon(s) \, ds,$$

where

$$\iota^\epsilon(s) = (\mathbf{u}^\epsilon C^\epsilon, \nabla \tilde{C}) - (\mathbf{M} \nabla C^\epsilon, \nabla \tilde{C}) + (S^\epsilon, \tilde{C}).$$

In the next step we prove the bound

$$\|\iota^\epsilon\|_{L^{4/3}(0,T)} \leq c \|\tilde{C}\|_{H^2} \quad (43)$$

for $\iota^\epsilon(s)$, where c stands for a constant. In order to do this, we estimate each term of ι^ϵ .

$$(S^\epsilon, \tilde{C}) \quad \text{is bounded in } L^2(0, T), \quad (44)$$

$$(\mathbf{M} \nabla C^\epsilon, \nabla \tilde{C}) \leq c \|\tilde{C}\|_{H^1} \|C^\epsilon\|_{H^1} \quad \text{is bounded in } L^2(0, T), \quad (45)$$

$$(\mathbf{u}^\epsilon C^\epsilon, \nabla \tilde{C}) \leq \|\mathbf{u}^\epsilon\|_{L^2} \|C^\epsilon\|_{L^3} \|\tilde{C}\|_{H^2} \quad \text{is bounded in } L^{4/3}(0, T). \quad (46)$$

The first bound regarding the source term holds because of the $L_t^2 L_x^2$ regularity of the source (see (8) and Remark 2.); the bound of the linear term (45) is a direct consequence of the Cauchy-Schwartz inequality; while the ultimate bound can be verified using (34) and (35), moreover considering the interpolation between $L^\infty(0, T; L^2)$ and $L^2(0, T; L^6)$, which yields that C^ϵ is in $L^4(0, T; L^3)$.

Finally, as we have now showed (43), we apply the Holder inequality combined with this latter and arrive to


$$\int_t^{t+h} |\iota^\epsilon(s)| \, ds \leq c \|\tilde{C}\|_{H^2} h^{1/4},$$

which gives the proof of the lemma. □

Eventually, we obtain the strong convergence result $C^\epsilon \rightarrow C$ in $L^2(0, T; L^2)$, and the weak convergence $\mathbf{u}^\epsilon C^\epsilon \rightharpoonup \mathbf{u} C$ of the nonlinear term. This, combined with the previously described linear terms, concludes the proof of the main theorem.

5 Concluding remarks

We are finishing this paper briefly mentioning a few ideas regarding the physical aspects of the model and some additional choices concerning the source term specifically.



Remark 3. As we have mentioned previously, the generic definition of the σ boundary function and the structure of the proof makes it easy to consider a generalised model adapted for seaside conditions where the lower boundary consists of $\Gamma_G = \Gamma_E \cup \Gamma_I$ — here Γ_I is the interface between air and water where rigid and no-slip boundary conditions are switched for ones expressing interactions. The main difference regarding the concentration function is that the σ_3 sediment is not zero in this case. Since we never exploited σ_3 being zero in the inland scenario, and we only used the bounded nature of this function, the generalisation does not bring any additional challenge in this front. For the velocity function the wind traction is represented by the new boundary condition $\nu_3 \partial_3 \mathbf{u}_H^\epsilon = \theta_H, u_3 = 0$ with assuming a finite wind energy, i.e. the function θ being in $L^2(0, T; H^{-1/2}(\Gamma_I))$. This part of the work has already been done by [6].

Remark 4. In the paper [19] it is suggested that the domain we work on has the form of $p_0 < p < p_1$, because for the thin portion of air between the earth and the isobar $p = p_1$, another specific simplified model would be necessary. This means that a way to make the model more precise is to cut the domain at an isobar a little bit off the physical ground. This would of course radically change the boundary conditions we use for the functions (in fact it is not trivial how to change them for this case), because it would take away the advantage of having a wall-like, natural and physical boundary condition for the domain we work on. If we are positively above the ground, $\partial_3 C = 0$ would represent a strange reflective layer.

Remark 5. Note that the aspect ratio going to zero is not only legitimate on a global scale. When we switch from a global model to a local one in order to be able to use the beta plane approximation, we still have hundreds of thousands of square kilometres versus 1—5 kilometres.

Remark 6. We chose the Gaussian nascent delta function to give shape to the source term, but it can be defined in several different ways using the many different pulses that approximate the delta function (choosing only those however that are physically meaningful in our case, for example its values are nonnegative), e.g.

- a) the unit impulse: $\delta_\epsilon(\mathbf{x} - \mathbf{x}_s) = \epsilon/2, |\mathbf{x} - \mathbf{x}_s| < 1/\epsilon,$
- b) the Lorentzian pulse: $\delta_\epsilon(\mathbf{x} - \mathbf{x}_s) = \epsilon/(1 + \pi^2\epsilon^2(\mathbf{x} - \mathbf{x}_s)^2).$

The Gaussian and Lorentzian version of the source term is continuous and bounded on Ω , it is actually in L^∞ in space (and in L^2 as a consequence), while it is easy to see that the piecewise constant unit pulse is in L^2 as well. Since the time dependence of this source is described by the Heaviside function, the time norm is also finite.

Remark 7. However the results of this article were achieved in a classical, fixed Cartesian coordinate system, there are other possibilities to build up the model in. We will briefly consider two of these.

- a) One option is to fix the coordinate system in such a way that the x axis matches the downwind direction ([9]). In other words, z is the vertical direction, while x is the main wind, and y is the crosswind direction. It is a legitimate assumption if we are considering a time interval which is not too long in the sense that it is reasonable to assume that we have a permanent, relatively stable downwind direction. For this scenario we consider a diagonal diffusion matrix, i.e. diffusion coefficients K_x, K_y, K_z , and more strict boundary conditions, namely we require $\partial_2 C$ and $\partial_3 C$ to vanish on the lower boundary. This is necessary because of the changes in the energy inequality caused by the additional ϵ term that we will introduce in the following. The adjusted coordinate system makes it possible to add another equation in the scaling process, which is different in nature from the previous ones in the sense that it is not derived from the shallowness of the domain but rather suggested by the fact that in the downwind direction it is natural to assume that the diffusion is negligible compared to downwind transport, i.e.

$$|u_1 \partial_x C| \gg |K_x \partial_{xx}^2 C| \quad (47)$$

which leads us to using

$$K_x = \epsilon M_1$$

in this convection dominated scenario.

Adding this equation on the one hand makes the concentration equation in the hydrostatic limit model one term shorter, since the first diffusion term drops out. On the other hand it leads to a new mathematical challenge handling the weak convergence of the nonlinear term $\mathbf{u}^\epsilon C^\epsilon$ in a situation where we do not have the H_x^1 regularity of C^ϵ — as we have L_x^2 boundedness only for $\sqrt{\epsilon} \partial_1 C^\epsilon$, and not for $\partial_1 C^\epsilon$ itself. We plan to investigate this scenario in an upcoming article.

- b) Another possibility is to adjust the coordinate system in a way that the x axis matches not with the downwind, but the wind velocity vector function $\mathbf{u}^\epsilon$ itself. This option brings along a $\mathbf{u}^\epsilon$ -dependent coordinate transform matrix in every equation, however it also makes the velocity vector one dimensional in the new coordinate system, which can be a potential benefit in a computational application.

Remark 8. Note that, — with some slight modifications, — the steps made leading to the final form of the energy inequality remain true even if we use a diffusion matrix function with nonconstant coefficients of the form $M(x)$ and such that the uniform ellipticity condition (6) holds in the form

$$\sum_{i,j=1}^3 M_{ij}(x) \xi_i \xi_j \geq \lambda \|\xi\|^2 \quad (48)$$

for almost any $x \in \Omega$ and all $\xi \in \mathbb{R}^3$. Moreover we require $M_{ij}(x) \in L^\infty(\Gamma)$ for any $i, j = 1, 2, 3$. The term I_3 in (32) now can be bounded as

$$\begin{aligned} I_3 &\leq \sum_{i,j=1}^3 \int_0^t \|M_{ij}(x) \partial_i C^\epsilon\|_{L^2(\Gamma_G)} \|C^\epsilon\|_{L^2(\Gamma_G)} d\tau \\ &\leq \sum_{i,j=1}^3 \|M_{ij}\|_{L^\infty(\Gamma_G)}^2 \int_0^t \|\sigma_i\|_{L^2(\Gamma_G)} \|C^\epsilon\|_{L^2(\Gamma_G)} d\tau. \end{aligned} \quad (49)$$

On the other hand, the term that took form in the original version as $\zeta M_{ij} \int_0^t \frac{1}{2} \|C^\epsilon\|_{L^2(\Gamma_G)}^2 d\tau$, now is present without the M_{ij} coefficient, since that has been incorporated into the first term of the product in (32) — and as a consequence, M_{ij} vanishes from the lefthand side of the final form (33) of the inequality as well.

A The existence of weak solution of the hydrostatic system of the polluted atmosphere

In this section for completeness we sketch the proof of the existence of weak solutions of the system (19) - (26).

The idea is that the polluted atmosphere can be viewed as the air-analogous version of the salty sea water in terms of the equations that represent it. In both cases we have a fluid which can be seen approximately as a homogenous incompressible fluid carrying some other particles in a significant concentration, the basis fluid motion is well described by the incompressible Navier-Stokes equations, and the additional concentration is described by an advection-diffusion equation.

The paper of [19] provides an existence result for an arbitrary time interval $[0, t_1]$ for the system describing the salty ocean water with temperature. The structure of the equations themselves describing the system is essentially the same, if we drop the equation describing the temperature, we basically arrive to the same system that describes the polluted atmosphere.

The main difference consists in the boundary conditions: on every boundary they use a zero flux for the salinity. This coincides with our lower boundary conditions, but on the lateral boundary we use a fix constant.

We will follow step by step the structure of the proof in [19], redefining the necessary quantities to be valid for our case and verify that the proof holds in the case of the polluted atmosphere as well (this is necessary not only because of using altered boundary conditions but also because the regularities of the functions are different in some cases).

Step 1. Firstly we redefine the norms and the functional setting in order to make them match our system.

Throughout this section we will use the $V = V_1 \times V_2, H = H_1 \times H_2, \mathcal{V} = \mathcal{V}_1 \times \mathcal{V}_2$, and $V_{(2)}$ function spaces, where V and H essentially represent H^1 and L^2 regularities, while $\mathcal{V}$ and $V_{(2)}$ stand for a subset of the C^∞ function space and its closure in H^2 , respectively. In more details, we introduce

$$V_1 = \{\mathbf{u}_H \in H_{\Gamma_A}^1(\Omega) \times H_{\Gamma_A}^1(\Omega), \nabla \cdot \int_h^0 \mathbf{u}_H \, dz = 0\},$$

$$V_2 = H^1(\Omega),$$

$$H_1 = \{\mathbf{u}_H \in H_{\Gamma_A}^0(\Omega) \times H_{\Gamma_A}^0(\Omega), \nabla \cdot \int_h^0 \mathbf{u}_H \, dz = 0\},$$

and

$$H_2 = L^2(\Omega)$$

Let $\mathcal{V}_1$ be the space of C^∞ vector functions $\mathbf{u}_H$ which vanish in a neighborhood of Γ and such that

$$\nabla \cdot \int_h^0 \mathbf{u}_H \, dz = 0,$$

and let $\mathcal{V}_2$ denote the C^∞ functions on $\bar{\Omega}$.

Finally we introduce the space $V_{(2)}$:

$$V_{(2)} \text{ is the closure of } \mathcal{V} \text{ in } (H^2(\Omega))^3.$$

We denote by U the vector $(\mathbf{u}_H, C)$. The scalar products and norms in the case of the polluted atmosphere are defined as follows.

$$((U, \tilde{U})) = ((\mathbf{u}_H, \tilde{\mathbf{u}}_H))_1 + K_C((C, \tilde{C}))_2$$

$$((\mathbf{u}_H, \tilde{\mathbf{u}}_H))_1 = \int_{\Omega} \nabla_{\nu} \mathbf{u}_H \nabla_{\nu} \tilde{\mathbf{u}}_H \, d\omega$$

$$((C, \tilde{C}))_2 = \int_{\Omega} (\nabla C)^T \mathbf{M} \nabla \tilde{C} \, d\omega$$

$$\|U\| = ((U, U))^{1/2}$$

$$(U, \tilde{U})_H = \int_{\Omega} [-\mathbf{u}_H \tilde{\mathbf{u}}_H - K_C C \tilde{C}] \, d\omega$$

$$|U|_H = |(U, U)_H|^{1/2}$$

Step 2. Now the hydrostatic weak formulation of the polluted atmosphere can be written as

$$\left(\frac{d}{dt} U, \tilde{U} \right)_H + a(U, \tilde{U}) + b(U, U, \tilde{U}) + e(U, \tilde{U}) = l(\tilde{U}) \quad (50)$$

with

$$a(U, \tilde{U}) = a_1(U, \tilde{U}) + K_C a_2(U, \tilde{U})$$

$$a_1(U, \tilde{U}) = \int_{\Omega} \nabla_{\nu} \mathbf{u}_H \nabla_{\nu} \tilde{\mathbf{u}}_H \, d\omega$$

$$a_2(U, \tilde{U}) = \int_{\Omega} (\nabla C)^T \mathbf{M} \nabla \tilde{C} \, d\omega$$

$$b = b_1 + K_C b_2$$

$$b_1(U, \tilde{U}, U^{\#}) = - \int_{\Omega} (\mathbf{u}_H \tilde{\mathbf{u}}_H + \mathbf{u}_H \tilde{u}_3) \nabla \mathbf{u}_H^{\#}$$

$$b_2(U, \tilde{U}, U^{\#}) = - \int_{\Omega} (\mathbf{u}_H \tilde{C} + u_3 \tilde{C}) \nabla C^{\#}$$

$$e(U, \tilde{U}) = \int_{\Omega} b(\mathbf{u}_H) \tilde{\mathbf{u}}_H \, d\omega$$

$$l(\tilde{U}) = - \int_{\Omega} S \tilde{C} \, d\omega + [((\nabla C)^T \mathbf{M}) \tilde{C} \vec{\mathbf{1}}]_{\Gamma_G}.$$

Here the term $b(\mathbf{u}_H) = \alpha(-u_2, u_1)$ represents the cross product, and we use the fact that the vertical velocity w can be expressed as $u_3 = u_3(\mathbf{u}_H)$ because the divergence of the three dimensional velocity is zero.

Step 3. It is straightforward to see that a is trilinear continuous, the coercivity of a is also trivial, moreover e is bilinear continuous and $e(U, U) = 0$.

What changes slightly is the verification of the results available on the form b , since the space derivatives in our case are applied on the test function. In the first two points below we discuss the difference in the proof that guarantees the trilinear continuity of b , that is,

$$|b(U, \tilde{U}, U^{\#})| \leq \begin{cases} c \|U\| \|\tilde{U}\| \|U^{\#}\|_{V_{(2)}}, & \forall U, \tilde{U} \in V, U^{\#} \in V_{(2)}, \\ c \|U\| \|\tilde{U}\|_{V_{(2)}} \|U^{\#}\|, & \forall U, U^{\#} \in V, \tilde{U} \in V_{(2)}; \end{cases} \quad (51)$$

finally we also adjust the original steps that provide the antisymmetric property $b(U, \tilde{U}, U^{\#}) = -b(U, U^{\#}, \tilde{U})$ and an improvement of (51) in this new environment.

- To prove the first bound for the case of $U, \tilde{U} \in V, U^\# \in V_{(2)}$ in (51), let us consider the typical term

$$\int_{\Omega} u_3 \tilde{C} \nabla C^\# d\omega.$$

We observe that we need to change the original choice of constants ($p_1 = p_2 = 2, p_3 = \infty$) for applying the Holder inequality. We can not use the L_∞ bound on the third term, since we would eventually arrive to $\|C^\#\|_{H^3}$, which would prevent us from closing the bound. Instead, we can use

$$\begin{aligned} \int_{\Omega} u_3 \tilde{C} \nabla C^\# d\omega &\leq c \|u_3\|_{L^2} \|\tilde{C}\|_{L^4} \|\nabla C^\#\|_{L^4} \leq \\ &\leq c \|U\| \|\tilde{U}\| \|\nabla C^\#\|_{H^1} \leq c \|U\| \|\tilde{U}\| \|U^\#\|_{V_{(2)}}, \end{aligned}$$

which yields the needed result.

- Similarly, for the case of $U, U^\# \in V, \tilde{U} \in V_{(2)}$ in the second inequality of (51), we need to adjust the space choices according to the present scenario. With the updated L^p constants we arrive to the

$$\int_{\Omega} u_3 \tilde{C} \nabla C^\# d\omega \leq c \|\tilde{C}\|_{L^\infty} \|u_3\|_{L^2} \|\nabla C^\#\|_{L^2} \leq c \|U\| \|\tilde{U}\|_{V_{(2)}} \|U^\#\|$$

new inequality that provides the continuity property.

- The proof of

$$b(U, \tilde{U}, U^\#) = -b(U, U^\#, \tilde{U}) \text{ for } U, \tilde{U}, U^\# \in V \text{ and } \tilde{U} \text{ or } U^\# \text{ in } V_{(2)}$$

can be derived by integration by parts as we have

$$\int_0^T (\mathbf{u} \nabla C, \tilde{C}) dt = - \int_0^T (\mathbf{u} C, \nabla \tilde{C}) dt,$$

observing that the boundary terms disappear.

- To establish the improvement

$$|b(U, \tilde{U}, U^\#)| \leq c \|U\| |\tilde{U}|_H^{1/2} \|\tilde{U}\|^{1/2} \|U^\#\|_{V_{(2)}}$$

of the first bound for $U, \tilde{U} \in V, U^\# \in V_{(2)}$, we consider the most typical term and we bound it by

$$\int_{\Omega} u_3 \tilde{C} \nabla C^\# \, d\omega \leq \|u_3\|_{L^2} \|\tilde{C}\|_{L^3} \|\nabla C^\#\|_{L^6}.$$

Note that since our definition of the form b is different from [19], thus we need to skip the switch of function arguments caused by the application of $|b(U, \tilde{U}, U^\#)| = |b(U, U^\#, \tilde{U})|$, otherwise the bound can not be closed.

Now, using the $\|\varphi\|_{L^3} \leq c\|\varphi\|_{L^2}^{1/2}\|\varphi\|_{H^1}^{1/2}$ Sobolev-Ladyzhenskaya inequality in space dimension 3, we can bound this term by

$$c\|\mathbf{u}_H\| \|U^\#\|_{V_{(2)}} |\tilde{U}|^{1/2} \|\tilde{U}\|^{1/2},$$

which gives the inequality.

This means that all the necessary properties of the forms a, b and e are equally valid for our set of equations (19) - (26), and we can use them to prove a stationary-case existence theorem identically as the proof is constructed in [19].

The weak formulation is as follows.

Definition 3. *Given a source term $S = \delta$ or $S \in L^2(\Omega)$, find $U = (\mathbf{v}, C)$, that satisfies the hypothesis of Definition 1 in the space dimensions, moreover*

$$a(U, \tilde{U}) + b(U, U, \tilde{U}) + e(U, \tilde{U}) = l(\tilde{U}) \quad (52)$$

for every $\tilde{U} \in V_{(2)}$.

Note that the definition of the weak formulation in our case is a bit more general, since we allow not only L^2 functions to represent the source term, but the δ distribution as well. As we have seen before, the weak formulation is still valid in this case because of the basic properties of δ , so this generalisation is mathematically valid.

We have the following theorem establishing the existence of solutions of the stationary equations, which can be proved by Galerkin method using the fundamental results we verified for the functions a, b and e .

Theorem 3. *We are given $S = \delta$ or S in $L^2(\Omega)$, then the problem (52) possesses at least one solution $U \in V$ such that*

$$\|U\| \leq \frac{1}{c_1} \|l\|_{V'},$$

where c_1 is the coercivity constant.

Step 4. In this step we provide the operator form of the equation (50) in the Hilbert space $V'_{(2)}$.

The operators $\frac{dU}{dt}$, A , B and E can be set analogously as in the original version; A for example is defined to be a linear continuous form from V into V' , given by the formula

$$\langle AU, \tilde{U} \rangle = a(U, \tilde{U}), \quad \forall U, \tilde{U} \in V.$$

The operator form of equation (50) becomes

$$\frac{dU}{dt} + AU + B(U, U) + EU = l.$$

Step 5. Finally we discuss the details of the time-dependent case, where we establish the existence, for all time, of the solutions for the system (19)-(23).

The weak formulation in the time-dependent case takes the following form.

Definition 4. Given T , $U_0 \in H$, and a source term $S = \delta$ or $S \in L^2(\Omega)$, find U that satisfies the hypothesis of Definition 1, moreover

$$\begin{aligned} \left(\frac{d}{dt} U, \tilde{U} \right) + a(U, \tilde{U}) + b(U, U, \tilde{U}) + e(U, \tilde{U}) &= l(\tilde{U}) \quad \forall \tilde{U} \in V_{(2)} \\ U(0) &= U_0 \end{aligned} \quad (53)$$

We proceed by using finite differences in time, taking an arbitrary integer N , setting a time step $k = \Delta t = T/N$, and recursively for $n = 1, \dots, N$ considering

$$\begin{aligned} U^0 &= U_0, \\ \frac{1}{\Delta t} (U^n - U^{n-1}, \tilde{U})_H + a(U^n, \tilde{U}) + b(U^n, U^n, \tilde{U}) \\ &\quad + e(U^n, \tilde{U}) = l^n(\tilde{U}), \quad \text{for any } \tilde{U} \in V_{(2)} \end{aligned} \quad (54)$$

Following the proof of the existence of $U^n \in V$, proving some a priori estimates, introducing approximate functions on $(0, T)$, and using the Aubin

compactness theorem in an identical way as described in [19], we pass to the limit and arrive to

$$\begin{aligned} - \int_0^T (U, \tilde{U})_H \psi' dt + \int_0^T [a(U, \tilde{U}) + b(U, U, \tilde{U}) + e(U, \tilde{U})] \psi dt = \\ = (U_0, \tilde{U}) \psi(0) + \int_0^T l(\tilde{U}) \psi dt, \end{aligned}$$

where ψ is a function defined on $[0, T]$, used to create the time-dependent version $\tilde{U}\psi$ of the test function. Renaming this product simply to be $\tilde{U}(t)$, the time-dependent test function, we arrive exactly to

$$\begin{aligned} - \int_0^T (U, \partial_t \tilde{U})_H dt + \int_0^T [a(U, \tilde{U}) + b(U, U, \tilde{U}) + e(U, \tilde{U})] dt = \\ = (U_0, \tilde{U}(0)) + \int_0^T l(\tilde{U}) dt, \end{aligned}$$

which is the same as (53), and which thus finally leads us to the main existence result of this section.

Theorem 4. *Given $T > 0$, $U_0 \in H$, and a source term $S = \delta$ or $S \in L^2(\Omega)$, then there exists $(\mathbf{u}, C)$ with $u_3 = - \int_0^T \nabla \cdot \mathbf{u}_H$, $\mathbf{u} = (\mathbf{u}_H, u_3) \in L^2(0, T; \mathbf{W})$ with $\mathbf{u}_H \in L^\infty(0, T; L^2(\Omega)^2)$, and $C \in L^\infty(0, T; L^2(\Omega)) \cap L^2(0, T; H^1(\Omega))$, which is solution to (27)-(28).*

References

- [1] R. A. Adams and J. J. Fournier. *Sobolev spaces*, volume 140. Academic press, 2003.
- [2] I. Ali, S. Kalla, and H. Khajah. A partial differential equation related to a problem in atmospheric pollution. *Mathematical and computer modelling*, 28(12):1–6, 1998.
- [3] J. D. Anderson and J. Wendt. *Computational fluid dynamics*, volume 206. Springer, 1995.
- [4] G. Andria, A. Lay-Ekuakille, and M. Notarnicola. Mathematical models for atmospheric and industrial pollutant prediction. In *XVI IMEKO World Congress, Vienna, Austria*, 2000.

- [5] P. S. Arya. *Introduction to micrometeorology*, volume 79. Elsevier, 2001.
- [6] P. Azérad and F. Guillén. Mathematical justification of the hydrostatic approximation in the primitive equations of geophysical fluid dynamics. *SIAM Journal on Mathematical Analysis*, 33(4):847–859, 2001.
- [7] O. Besson and M. Laydi. Some estimates for the anisotropic navier-stokes equations and for the hydrostatic approximation. *ESAIM: Mathematical Modelling and Numerical Analysis*, 26(7):855–865, 1992.
- [8] H. E. de Swart. The governing equations and the dominant balances of flow in the atmosphere and ocean. *USPC Summerschool, Utrecht*, 2016.
- [9] P. Goyal and A. Kumar. Mathematical modeling of air pollutants: An application to indian urban city. In *Air Quality-Models and Applications*. InTech, 2011.
- [10] B. Hosseini. Dispersion of pollutants in the atmosphere: a numerical study. Master’s thesis, Simom Fraser University, 2013.
- [11] P. Kathirgamanathan, R. McKibbin, and R. McLachlan. Source term estimation of pollution from an instantaneous point source. *Research Letters in Information Mathematic Science*, Vol. 3, pp. 59-67., 2002.
- [12] P.-L. Lions. *Mathematical Topics in Fluid Mechanics: Volume 2: Compressible Models*, volume 2. Oxford University Press on Demand, 1996.
- [13] P. Marsaleix, F. Auclair, and C. Estournel. Considerations on open boundary conditions for regional and coastal ocean models. *Journal of Atmospheric and Oceanic Technology*, 23(11):1604–1613, 2006.
- [14] A. Monin. On the boundary condition on the earth surface for diffusing pollution. In *Advances in Geophysics*, volume 6, pages 435–436. Elsevier, 1959.
- [15] J. Pedlosky. *Geophysical fluid dynamics*. Springer Science & Business Media, 2013.
- [16] M. Prodanova, J. L. Perez, D. Syrakov, R. San Jose, K. Ganev, N. Miloshev, and S. Roglev. Application of mathematical models to simulate an extreme air pollution episode in the bulgarian city of Stara Zagora. *Applied Mathematical Modelling*, 32(8):1607–1619, 2008.

- [17] J. Simon. Compact sets in the space $L^p(0, t; B)$. *Annali di Matematica pura ed applicata*, 146(1):65–96, 1986.
- [18] R. Temam. *Navier-Stokes equations: Theory and numerical analysis*, volume 343. American Mathematical Soc., 2001.
- [19] R. Temam and M. Ziane. Some mathematical problems in geophysical fluid dynamics. *Handbook of mathematical fluid dynamics*, 3:535–658, 2005.
- [20] Q. Wang, W. Zhou, D. Wang, and D. Dong. Ocean model open boundary conditions with volume, heat and salinity conservation constraints. *Advances in Atmospheric Sciences*, 31(1):188–196, 2014.
- [21] G.-T. Yeh and C.-H. Huang. Three-dimensional air pollutant modeling in the lower atmosphere. *Boundary-Layer Meteorology*, 9(4):381–390, 1975.
- [22] S. Zhuk, T. T. Tchakian, S. Moore, R. Ordóñez-Hurtado, and R. Shorten. On source-term parameter estimation for linear advection-diffusion equations with uncertain coefficients. *SIAM Journal on Scientific Computing*, 38(4):A2334–A2356, 2016.