

A generalised weak convergence result for the mathematical equations of the polluted atmosphere expressed in downwind matching coordinates

Donatella Donatelli and Nóra Juhász *

Department of Information Engineering, Computer Science and Mathematics
University of L'Aquila
67100 L'Aquila, Italy.
donatella.donatelli@univaq.it, norjuh@univaq.it

Abstract

A widely used approach to mathematically describe the atmosphere is to consider it as a geophysical fluid in a shallow domain — and thus to model it using classical fluid dynamical equations combined with the explicit inclusion of an ϵ parameter representing the small aspect ratio of the physical domain. In our previous paper we proved a weak convergence theorem for the polluted atmosphere described by the Navier-Stokes equations extended by an advection-diffusion equation. We obtained a justification of the generalised hydrostatic limit model including the pollution effect described in the case of a classical, east-north-upwards oriented local Cartesian coordinates. Here we give an improvement of this statement: we consider a meteorologically more meaningful coordinate system, incorporate the analytical consequences of this coordinate change into the governing equations, and verify that the weak convergence still holds for the altered velocity – concentration system.

Keywords: downwind-matching coordinate system, Navier-Stokes equations, shallow domains, pollution evolution equation, hydrostatic approximation, compactness, weak solutions.

*The research of DD and NJ leading to these results has received funding from the European Union's Horizon 2020 Research and Innovation Programme under the Marie Skłodowska-Curie Grant Agreement No 642768 (Project Name: ModCompShock).

Contents

1	Introduction	3
1.1	Notations	3
2	The polluted atmosphere model	4
3	Weak formulation and Main Result	8
4	Proof of the main theorem	8
4.1	Estimate for $\Delta \mathbf{u}_h^\epsilon$	8
4.1.1	The p terms fall out from the summed integrals	9
4.1.2	Using the Gronwall inequality to obtain the $\Delta \mathbf{u}_h^\epsilon$ estimate	11
4.2	H_1 estimate for C	12
5	Concluding remarks	12

1 Introduction

A local domain on the surface of the rotating Earth is often described by a Cartesian coordinate system where the x, y, z axes represent the directions towards east, north, and upwards, respectively, as in this specific system the rotation vector, and as a consequence, the Coriolis force takes a simpler form. However, for modelling wind and pollution effects in the atmosphere there is another naturally distinctive viewpoint for fixing the orientation of the coordinate system: the introduction of downwind and crosswind, and to match the horizontal axes according to these meteorological parameters ([10]). In this paper we take this idea as a foundation, and we migrate the result of our previously stated weak convergence theorem to this more considerably chosen coordinate system.

Here talk a little bit about the original convergence theorem to make the new one more self-contained and for details refer to the first article.

As in this scenario the advection is dominant compared to diffusion effects along the x axis, this approach naturally brings an additional ϵ term in the Laplacian of the concentration equation, the simplification of the limit model, and brings the challenge of proving a weak convergence result for the product of the vertical velocity and the concentration, as in the energy inequality the $\|\partial_1 C^\epsilon\|$ term is replaced by $\|\sqrt{\epsilon}\partial_1 C^\epsilon\|$. This issue is overcome by the construction of new estimates obtained by multiplying the governing equations by higher order terms.

Modified boundary conditions. Explain the necessity of changing the sediment functions.

We end up this section with some notations we will use through the paper.

1.1 Notations

- a) ϵ = the aspect ratio,
- b) ν = viscosity,
- c) Ω_ϵ, Ω = the original and the rescaled (ϵ -independent) domain, respectively,
- d) ∇ stands for the gradient vector; specifically, $(\partial_x, \partial_y, \partial_z)$ before, and $(\partial_1, \partial_2, \partial_3)$ after rescaling,
- e) $\nabla_\nu = (\nu_x^{1/2}\partial_x, \nu_y^{1/2}\partial_y, \nu_z^{1/2}\partial_z)$ on Ω_ϵ , $\nabla_\nu = (\nu_1^{1/2}\partial_1, \nu_2^{1/2}\partial_2, \nu_3^{1/2}\partial_3)$ on the rescaled domain Ω ,

- f) Δ_ν stands for the anisotropic Laplacian operators $\nu_x \partial_{xx}^2 + \nu_y \partial_{yy}^2 + \nu_z \partial_{zz}^2$ and $\nu_1 \partial_{11}^2 + \nu_2 \partial_{22}^2 + \nu_3 \partial_{33}^2$ on the original and rescaled domains, respectively,
- g) $\mathbf{g}$ represents the force due to gravity, which also includes the centrifugal effect,
- h) φ = the gravity potential term, i.e. $\mathbf{g} = \nabla \varphi$,
- i) ω represents the Earth rotation angular speed,
- j) $\phi = l(y)$ = latitude, f = the module of the Earth rotation vector
- k) $\alpha = -2f \sin(\theta) \cos(\phi)$, $\beta = 2f \cos(\theta) \cos(\phi)$, $\gamma = 2f \sin(\phi)$,
- l) $\mathbf{c}(\mathbf{u}) = (\mathbf{c}_h(\mathbf{u}), c_3(\mathbf{u})) = (-\gamma u_2^\epsilon + \epsilon \beta u_3^\epsilon, \alpha \epsilon u_3^\epsilon - \gamma u_1^\epsilon, \alpha u_2^\epsilon - \beta u_1^\epsilon)$, the Coriolis force terms
- m) $\mathbf{K}$ represents the diffusion matrix,
- n) $(\cdot, \cdot)$ = the scalar product in $L^2(\Omega)^d$ or the duality $L^p(\Omega)$, $L^{p'}(\Omega)$,
- o) $\langle \cdot, \cdot \rangle_\Gamma$ = the scalar product or duality on the boundary curve Γ ($H^{-1/2}(\Gamma)$, $H^{1/2}(\Gamma)$),
- p) $\mathbf{u}_h$ = the horizontal velocity components (u_1, u_2) ,
- q) $b(\mathbf{u}_h) = -\gamma(u_2, u_1)$,
- r) $L_t^p L_x^q$ is the abbreviated notation for $L^p(0, T; L^q(\Omega))$.

2 The polluted atmosphere model

As we highlighted before, the key point of this new model is to integrate some of the main ideas of a Gaussian plume dispersion model into the system describing the polluted atmosphere. Therefore we begin by fixing the local Cartesian coordinate system to be adjusted to the downwind direction, i.e. z is oriented upwards, while x is the downwind, and y is the crosswind direction. Another assumption the Gaussian dispersion models typically use as well is that advection is more dominant than diffusion along the downwind direction, which leads to the appearance of an additional ϵ term in the concentration equation's Laplacian. Namely, we have

$$|u_1 \partial_1 C| \gg |K_x \partial_{11}^2 C| \quad (1)$$

which leads us to using this additional scaling equation

$$K_x = \epsilon K_1. \quad (2)$$

Remark 1. The inclusion of these two features in a pollution model are legitimate when the following circumstances hold:

- We are considering a time interval which is not too long in the sense that it is reasonable to assume that we have a permanent, relatively stable main wind direction, i.e. fixing the horizontal coordinates is reasonable in the first place,
- There is an adequate, relatively high wind speed — low wind speeds in general lead to a situation where three dimensional diffusion dominates, leading to an error caused by the dominant advection assumption. Fast changing wind direction and low windspeed resulting in fast pollutant settling can make it extremely challenging to obtain reliable results from Gaussian models.

Choosing the coordinate system in this particular way results in changes in three main points of the system. One of these is the Coriolis term, since now the x axis is not necessarily perpendicular to the rotation vector. We need to revisit the boundary conditions as well, and as discussed above, a modification in the pollution concentration equation also has to be taken into account. We will describe these changes and their consequences in detail — the main challenge in proving this paper's convergence result is in fact to handle the effect of this latter change on the energy inequality.

In order to obtain the set of equations that describe the phenomena we start with the original set of equations

$$\partial_t \mathbf{v} + (\mathbf{v} \cdot \nabla) \mathbf{v} - \Delta_\nu \mathbf{v} + \nabla q + 2\omega \times \mathbf{v} = \mathbf{g} \text{ in } \Omega_\epsilon \times (0, T) \quad (3)$$

$$\nabla \cdot \mathbf{v} = 0 \text{ in } \Omega_\epsilon \times (0, T) \quad (4)$$

$$\partial_t P + \mathbf{v} \cdot \nabla P = \nabla \cdot (\mathbf{K} \nabla P) + Q, \quad (5)$$

As for the rescaling process, here we perform the same steps, with the addition of (2).

To expand the $2\omega \times \mathbf{v}$ Coriolis term firstly it is necessary to understand the structure of the new rotation vector. Note that the downwind-matching coordinate system and the original east-north-upwards oriented system are

different only in a rotation around the z axis. Let θ denote the angle between the respective x axes, ϕ denotes the latitude, then the rotation vector expressed in the downwind matching coordinate system becomes

$$\begin{bmatrix} \cos(\theta) & -\sin(\theta) & 0 \\ \sin(\theta) & \cos(\theta) & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{bmatrix} 0 \\ f \cos(\phi) \\ f \sin(\phi) \end{bmatrix} = f \begin{bmatrix} -\sin(\theta) \cos(\phi) \\ \cos(\theta) \cos(\phi) \\ \sin(\phi) \end{bmatrix}. \quad (6)$$

Now, let

$$\alpha = -2f \sin(\theta) \cos(\phi), \quad \beta = 2f \cos(\theta) \cos(\phi), \quad \gamma = 2f \sin(\phi),$$

and using this, the Coriolis force takes the form

$$\begin{bmatrix} \alpha \\ \beta \\ \gamma \end{bmatrix} \times \begin{bmatrix} u_1 \\ u_2 \\ \epsilon u_3 \end{bmatrix} = \begin{bmatrix} \beta \epsilon u_3 - \gamma u_2 \\ \alpha \epsilon u_3 - \gamma u_1 \\ \alpha u_2 - \beta u_1 \end{bmatrix}. \quad (7)$$

Thus, these are the terms that appear in the three velocity equations; and finally, note that the third component is present in the rescaled version of the vertical velocity equation multiplied by ϵ , which is obtained from the ϵ term in the denominator of $\partial_z p$.

Remark 2. The intuitive meaning of what we understand by downwind and the velocity in this new coordinate system. The downwind direction is an approximately steady-in-time direction, it can be seen as the direction marked by the wind direction trackers on an airport or meteorological point. When we consider the downwind, we neglect the small perturbations in its orientation, it can be viewed as a time-averaged direction as long as the fluctuation is relatively small. On the other hand, the velocity is a three dimensional, time dependent vector that changes second after second.

We arrive to the merged anisotropic equations of the polluted atmosphere above inland surface:

$$\partial_t u_1^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_1^\epsilon - \Delta_\nu u_1^\epsilon - \gamma u_2^\epsilon + \epsilon \beta u_3^\epsilon + \partial_1 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (8)$$

$$\partial_t u_2^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_2^\epsilon - \Delta_\nu u_2^\epsilon + \alpha \epsilon u_3^\epsilon - \gamma u_1^\epsilon + \partial_2 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (9)$$

$$\epsilon^2 \{ \partial_t u_3^\epsilon + \mathbf{u}^\epsilon \cdot \nabla u_3^\epsilon - \Delta_\nu u_3^\epsilon \} + \epsilon \alpha u_2^\epsilon - \epsilon \beta u_1^\epsilon + \partial_3 p^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (10)$$

$$\nabla \cdot \mathbf{u}^\epsilon = 0 \text{ in } \Omega \times (0, T) \quad (11)$$

$$\partial_t C^\epsilon + \mathbf{u}^\epsilon \cdot \nabla C^\epsilon = \epsilon K_1 \partial_{11}^2 C^\epsilon + K_2 \partial_{22}^2 C^\epsilon + K_3 \partial_{33}^2 C^\epsilon + S_\epsilon \text{ in } \Omega \times (0, T) \quad (12)$$

The initial condition for the velocity and the pollution concentration are

$$\mathbf{u}^\epsilon(\cdot, t=0) = \mathbf{u}_0, \quad C^\epsilon(\cdot, t=0) = C_0 \quad \text{in } \Omega. \quad (13)$$

Finally, the boundary conditions to describe the anisotropic system can be summarised as follows:

$$\begin{aligned} \mathbf{u}^\epsilon &= 0 \text{ on } \Gamma \times (0, T), \\ C^\epsilon &= 0 \text{ on } (\Gamma_U \cup \Gamma_L) \times (0, T), \\ \partial_2 C^\epsilon &= \partial_3 C^\epsilon = 0 \text{ on } \Gamma_G \times (0, T), \quad \partial_1 C^\epsilon \in L^2 L^2(0, T; \Gamma_G), \end{aligned} \quad (14)$$

combined with the additional

$$J(\mathbf{u}_h^\epsilon) \cdot \vec{\mathbf{n}}_\Gamma = \mathbf{0} \text{ on } \Gamma \times (0, T), \quad (15)$$

$$J(\mathbf{u}_h^\epsilon)^T \cdot \vec{\mathbf{n}}_\Gamma = \mathbf{0} \text{ on } \Gamma \times (0, T), \quad (16)$$

$$\nabla u_3 \cdot \vec{\mathbf{n}}_\Gamma = 0 \text{ on } \Gamma \times (0, T) \quad (17)$$

Neumann boundary conditions, where $J(\mathbf{u}_h^\epsilon)$ is the Jacobian.

We highlight the difference between the boundary conditions of the original, east-north oriented, and the new, downwind-matching model.

The first difference is the kind of boundary conditions that we can "afford" to have on the lower boundary for the concentration comes from the energy inequality. In the east-north-upwards oriented coordinate system we do not have an ϵ in the diffusion matrix, and thus it is not an issue to have any of the $\partial_i C$ terms to be different from 0, not even for an ocean case where $\partial_3 C$ becomes nonzero as well (they are required however to be bounded of course). On the other hand, the downwind-matching coordinate system brings in an ϵ term in front of the K_1 diffusion, which makes the model more sensitive to assigning boundary conditions for C . This means

that we can not have the relatively relaxed, boundedness requirement for any $\partial_i C$ except for $i = 1$. All other $\partial_2 C, \partial_3 C$ derivatives must be zero on the lower boundary, otherwise they would result in an $(\epsilon - \text{const.})$ negative term.

The second difference is the Neumann boundary conditions (15)-(17) for the velocity. We use these additional terms when calculating the estimate for $\mathbf{u}_h$.

If we assume $\mathbf{u}^\epsilon = O(1)$, then neglecting the ϵ^2 and ϵ in the anisotropic equations, we formally arrive to the hydrostatic Navier-Stokes equations (i.e., primitive equations) combined with pollution.

$$\partial_t u_1 + \mathbf{u} \cdot \nabla u_1 - \Delta_\nu u_1 - \gamma u_2 + \partial_1 p = 0 \text{ in } \Omega \times (0, T) \quad (18)$$

$$\partial_t u_2 + \mathbf{u} \cdot \nabla u_2 - \Delta_\nu u_2 - \gamma u_1 + \partial_2 p = 0 \text{ in } \Omega \times (0, T) \quad (19)$$

$$\partial_3 p = 0 \text{ in } \Omega \times (0, T) \quad (20)$$

$$\nabla \cdot \mathbf{u} = 0 \text{ in } \Omega \times (0, T) \quad (21)$$

$$\partial_t C + \mathbf{u} \cdot \nabla C = K_2 \partial_{22}^2 C + K_3 \partial_{33}^2 C + S \text{ in } \Omega \times (0, T) \quad (22)$$

$$u_1 = u_2 = u_3 n_3 = 0 \text{ on } \Gamma \times (0, T) \quad (23)$$

$$u_i(\cdot, t = 0) = u_{0i}, C(\cdot, t = 0) = C_0 \text{ in } \Omega, i = 1, 2 \quad (24)$$

$$\begin{aligned} C &= 0 \text{ on } (\Gamma_U \cup \Gamma_L) \times (0, T), \\ \partial_2 C &= \partial_3 C = 0 \text{ on } \Gamma_G \times (0, T), \partial_1 C \in L^2 L^2(0, T; \Gamma_G) \end{aligned} \quad (25)$$

3 Weak formulation and Main Result

4 Proof of the main theorem

4.1 Estimate for $\Delta \mathbf{u}_h^\epsilon$

In this part of the proof we prove an estimate for $\Delta \mathbf{u}_h^\epsilon$ which we will later need in order to obtain H_1 regularity for C . We use the $\mathbf{u}^\epsilon = (\mathbf{u}_h^\epsilon, u_3^\epsilon)$

partition, and in this section of the proof for better readability Δ stands for the horizontal Laplacian Δ_h , while ∇ represents ∇_h . Firstly, we separate the equations into horizontal and vertical directions and multiply them by the minus Laplacian of $\mathbf{u}_h$ and u_3 , respectively:

the $\mathbf{v}$ notation is already taken in the original velocity equation

$$\begin{aligned} \partial_t \mathbf{u}_h + \mathbf{u}_h \nabla \mathbf{u}_h + u_3 \partial_3 \mathbf{u}_h - \Delta \mathbf{u}_h - \partial_{33} \mathbf{u}_h + \mathbf{c}_h(\mathbf{u}) + \nabla p &= 0 \quad / \cdot -\Delta \mathbf{u}_h \\ \epsilon^2 (\partial_t u_3 + \mathbf{u}_h \nabla u_3 + u_3 \partial_3 u_3 - \Delta u_3 - \partial_{33} u_3) + \epsilon c_3(\mathbf{u}) + \partial_z p &= 0 \quad / \cdot -\Delta u_3 \\ \nabla \cdot \mathbf{u}_h + \partial_3 u_3 &= 0 \end{aligned} \quad (26)$$

We now show that after taking the integral of the horizontal and vertical velocity equations and summing the results, the parts including the p pressure term cancel out.

4.1.1 The p terms fall out from the summed integrals

We have to show that

$$\int_{\Omega} \Delta \mathbf{u}_h \nabla p + \int_{\Omega} \Delta u_3 \nabla p = 0.$$

In order to see this we expand the first term as

$$\begin{aligned} \int_{\Omega} \Delta \mathbf{u}_h \nabla p &= \sum_{i=1}^2 \int_{\Omega} \Delta u_i \partial_i p = \int_{\Omega} \sum_{i,j=1}^2 \partial_{jj} u_i \partial_i p = \\ \int_{\Omega} \left(\sum_{i,j=1}^2 \partial_j (\partial_j u_i \cdot \partial_i p) - \sum_{i,j=1}^2 \partial_j u_i \cdot \partial_j \partial_i p \right) &:= T_1 + T_2 \end{aligned}$$

Now,

$$\begin{aligned} T_1 &= \int_{\Omega} \sum_{i,j=1}^2 \partial_j (\partial_j u_i \cdot \partial_i p) = \int_{\Omega} \sum_{j=1}^2 \partial_j (\partial_j \mathbf{u}_h \cdot \nabla p) \\ &= \int_{\Omega} \operatorname{div}(\partial_1 \mathbf{u}_h \nabla p, \partial_2 \mathbf{u}_h \nabla p) = \int_{\partial\Omega} (\partial_1 \mathbf{u}_h \nabla p, \partial_2 \mathbf{u}_h \nabla p) \cdot \vec{\mathbf{n}}_{\Gamma}, \\ &= \int_{\partial\Omega} \nabla p \cdot (J(\mathbf{u}_h) \vec{\mathbf{n}}_{\Gamma}) = 0, \end{aligned} \quad (27)$$

using the Neumann boundary condition (15) regarding the Jacobian of $\mathbf{u}_h$.

$$T_2 = - \int_{\Omega} \sum_{i,j=1}^2 \partial_j u_i \cdot \partial_j \partial_i p = - \sum_{i,j=1}^2 \int_{\Omega} \partial_i (\partial_j u_i \partial_j p) + \sum_{i,j=1}^2 \int_{\Omega} \partial_j \partial_i u_i \partial_j p.$$

We show that the first term in this expanded version of T_2 disappears, and thus what remains is $\sum_{i,j=1}^2 \int_{\Omega} \partial_j \partial_i u_i \partial_j p$, i.e. $\sum_{j=1}^2 \int_{\Omega} \partial_j (\operatorname{div} \mathbf{u}_h) \partial_j p$.

$$\begin{aligned} & - \sum_{i,j=1}^2 \int_{\Omega} \partial_i (\partial_j u_i \partial_j p) = - \int_{\Omega} \sum_{i=1}^2 \partial_i (\nabla u_i \cdot \nabla p) \\ & = - \int_{\Omega} \operatorname{div}(\nabla u_1 \cdot \nabla p, \nabla u_2 \cdot \nabla p) = - \int_{\partial\Omega} (\nabla u_1 \cdot \nabla p, \nabla u_2 \cdot \nabla p) \vec{\mathbf{n}}_{\Gamma} \quad (28) \\ & = - \int_{\partial\Omega} \nabla p J(\mathbf{u}_h)^T \vec{\mathbf{n}}_{\Gamma} = 0, \end{aligned}$$

using the Neumann boundary condition (16).

From the second equation we have

$$\begin{aligned} \int_{\Omega} \Delta u_3 \partial_3 p &= \int_{\Omega} \sum_{i=1}^2 \partial_i (\partial_i u_3 \partial_3 p) - \int_{\Omega} \sum_{i=1}^2 \partial_i u_3 \partial_3 \partial_i p \\ &= \int_{\Omega} \operatorname{div}(\partial_1 u_3 \partial_3 p, \partial_2 u_3 \partial_3 p) - \int_{\Omega} \sum_{i=1}^2 \partial_3 (\partial_i u_3 \partial_i p) + \sum_{i=1}^2 \int_{\Omega} \partial_i \partial_3 u_3 \cdot \partial_i p \\ &= \int_{\partial\Omega} \partial_3 p \nabla u_3 \vec{\mathbf{n}}_{\Gamma} - \int_{\Omega} \partial_3 (\nabla u_3 \nabla p) - \sum_{i=1}^2 \int_{\Omega} \partial_i \operatorname{div}(\mathbf{u}_h) \cdot \partial_i p \\ &= 0 - \int_{\partial\Omega} \nabla u_3 \nabla p \cdot \mathbf{n}_{3\Gamma} - \sum_{i=1}^2 \int_{\Omega} \partial_i \operatorname{div}(\mathbf{u}_h) \cdot \partial_i p = - \sum_{i=1}^2 \int_{\Omega} \partial_i \operatorname{div}(\mathbf{u}_h) \cdot \partial_i p. \quad (29) \end{aligned}$$

using (17), the fact that $\nabla u_3 \cdot \mathbf{n}_{3\Gamma} = 0$, and (26).

Finally, summing up the remaining terms we have

$$\sum_{j=1}^2 \int_{\Omega} \partial_j (\operatorname{div} \mathbf{u}_h) \cdot \partial_j p - \sum_{i=1}^2 \int_{\Omega} \partial_i \operatorname{div}(\mathbf{u}_h) \cdot \partial_i p = 0,$$

thus the terms including p cancel each other.

4.1.2 Using the Gronwall inequality to obtain the $\Delta \mathbf{u}_h^\epsilon$ estimate

As we have shown previously, the pressure terms add to zero, so we don't need to consider them in the following part of the calculations. Integrating the equations on Ω we have

$$\begin{aligned}
& \frac{1}{2} \partial_t (\|\nabla \mathbf{u}_h\|_2^2 + \|\epsilon \nabla u_3\|_2^2) + \|\Delta \mathbf{u}_h\|_2^2 + \|\nabla \partial_z \mathbf{u}_h\|_2^2 + \|\epsilon \Delta u_3\|_2^2 + \|\epsilon \nabla \partial_z u_3\|_2^2 \\
&= \int_{\Omega} \Delta \mathbf{u}_h \cdot \mathbf{u}_h \nabla \mathbf{u}_h + \Delta \mathbf{u}_h \cdot u_3 \partial_3 \mathbf{u}_h + \Delta \mathbf{u}_h \mathbf{c}_h(\mathbf{u}) \\
&+ \int_{\Omega} \epsilon^2 \Delta u_3 \mathbf{u}_h \nabla u_3 + \epsilon^2 \Delta u_3 u_3 \partial_3 u_3 + \Delta u_3 \epsilon \mathbf{c}_3(\mathbf{u})
\end{aligned} \tag{30}$$

One by one the terms on the right hand side can be bounded as follows, using the Holder, Ladyzhenskaya and Young inequalities.

$$\begin{aligned}
& \int_{\Omega} \Delta \mathbf{u}_h \cdot \mathbf{u}_h \nabla \mathbf{u}_h \leq \|\Delta \mathbf{u}_h\|_2 \|\mathbf{u}_h\|_6 \|\nabla \mathbf{u}_h\|_3 \\
& \leq \|\Delta \mathbf{u}_h\|_2 \|\mathbf{u}_h\|_6 \|\nabla \mathbf{u}_h\|_2^{1/2} \|\Delta \mathbf{u}_h\|_2^{1/2} \leq \|\Delta \mathbf{u}_h\|_2^{3/2} \|\mathbf{u}_h\|_6 \|\nabla \mathbf{u}_h\|_2^{1/2},
\end{aligned} \tag{31}$$

$$\begin{aligned}
& \int_{\Omega} \Delta \mathbf{u}_h \cdot u_3 \partial_3 \mathbf{u}_h = \int_{\Omega} \left(\int_0^z |\nabla \cdot \mathbf{u}_h| \right) \partial_3 \mathbf{u}_h \Delta \mathbf{u}_h \\
& \leq \|\nabla \mathbf{u}_h\|_2^{1/2} \|\partial_3 \mathbf{u}_h\|_2^{1/2} \|\nabla \partial_3 \mathbf{u}_h\|_2^{1/2} \|\Delta \mathbf{u}_h\|_2^{3/2},
\end{aligned} \tag{32}$$

using the same inequality obtained in [8],

$$\int_{\Omega} \Delta \mathbf{u}_h \mathbf{c}_h(\mathbf{u}) \leq \|\mathbf{u}_h\|_2 \|\mathbf{c}_h(\mathbf{u})\|_2 \tag{33}$$

$$\int_{\Omega} \epsilon^2 \Delta u_3 \mathbf{u}_h \nabla u_3 \leq \|\epsilon \Delta u_3\|_2 \|\mathbf{u}_h\|_6 \|\epsilon \nabla u_3\|_3, \tag{34}$$

$$\int_{\Omega} \epsilon^2 \Delta u_3 u_3 \partial_3 u_3 \leq \|\epsilon \Delta u_3\|_2 \|\epsilon u_3\|_6 \|\nabla \mathbf{u}_h\|_3, \tag{35}$$

and finally,

$$\int_{\Omega} \Delta u_3 \epsilon \mathbf{c}_3(\mathbf{u}) \leq \|\Delta u_3\|_2 \|\mathbf{c}_3(\mathbf{u})\|_2. \tag{36}$$

4.2 H_1 estimate for C

5 Concluding remarks

References

- [1] R. A. Adams and J. J. Fournier. *Sobolev spaces*, volume 140. Academic press, 2003.
- [2] I. Ali, S. Kalla, and H. Khajah. A partial differential equation related to a problem in atmospheric pollution. *Mathematical and computer modelling*, 28(12):1–6, 1998.
- [3] J. D. Anderson and J. Wendt. *Computational fluid dynamics*, volume 206. Springer, 1995.
- [4] G. Andria, A. Lay-Ekuakille, and M. Notarnicola. Mathematical models for atmospheric and industrial pollutant prediction. In *XVI IMEKO World Congress, Vienna, Austria*, 2000.
- [5] P. S. Arya. *Introduction to micrometeorology*, volume 79. Elsevier, 2001.
- [6] P. Azérad and F. Guillén. Mathematical justification of the hydrostatic approximation in the primitive equations of geophysical fluid dynamics. *SIAM Journal on Mathematical Analysis*, 33(4):847–859, 2001.
- [7] O. Besson and M. Laydi. Some estimates for the anisotropic navier-stokes equations and for the hydrostatic approximation. *ESAIM: Mathematical Modelling and Numerical Analysis*, 26(7):855–865, 1992.
- [8] C. Cao and E. S. Titi. Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. *Annals of Mathematics*, pages 245–267, 2007.
- [9] H. E. de Swart. The governing equations and the dominant balances of flow in the atmosphere and ocean. *USPC Summerschool, Utrecht*, 2016.
- [10] P. Goyal and A. Kumar. Mathematical modeling of air pollutants: An application to indian urban city. In *Air Quality-Models and Applications*. InTech, 2011.

- [11] B. Hosseini. Dispersion of pollutants in the atmosphere: a numerical study. Master's thesis, Simon Fraser University, 2013.
- [12] P. Kathirgamanathan, R. McKibbin, and R. McLachlan. Source term estimation of pollution from an instantaneous point source. *Research Letters in Information Mathematic Science*, Vol. 3, pp. 59-67., 2002.
- [13] P.-L. Lions. *Mathematical Topics in Fluid Mechanics: Volume 2: Compressible Models*, volume 2. Oxford University Press on Demand, 1996.
- [14] P. Marsaleix, F. Auclair, and C. Estournel. Considerations on open boundary conditions for regional and coastal ocean models. *Journal of Atmospheric and Oceanic Technology*, 23(11):1604–1613, 2006.
- [15] A. Monin. On the boundary condition on the earth surface for diffusing pollution. In *Advances in Geophysics*, volume 6, pages 435–436. Elsevier, 1959.
- [16] J. Pedlosky. *Geophysical fluid dynamics*. Springer Science & Business Media, 2013.
- [17] M. Prodanova, J. L. Perez, D. Syrakov, R. San Jose, K. Ganev, N. Miloshev, and S. Roglev. Application of mathematical models to simulate an extreme air pollution episode in the bulgarian city of stara zagora. *Applied Mathematical Modelling*, 32(8):1607–1619, 2008.
- [18] J. Simon. Compact sets in the space $l^p(0, t; b)$. *Annali di Matematica pura ed applicata*, 146(1):65–96, 1986.
- [19] R. Temam. *Navier-Stokes equations: theory and numerical analysis*, volume 343. American Mathematical Soc., 2001.
- [20] R. Temam and M. Ziane. Some mathematical problems in geophysical fluid dynamics. *Handbook of mathematical fluid dynamics*, 3:535–658, 2005.
- [21] Q. Wang, W. Zhou, D. Wang, and D. Dong. Ocean model open boundary conditions with volume, heat and salinity conservation constraints. *Advances in Atmospheric Sciences*, 31(1):188–196, 2014.
- [22] G.-T. Yeh and C.-H. Huang. Three-dimensional air pollutant modeling in the lower atmosphere. *Boundary-Layer Meteorology*, 9(4):381–390, 1975.

- [23] S. Zhuk, T. T. Tchrakian, S. Moore, R. Ordóñez-Hurtado, and R. Shorten. On source-term parameter estimation for linear advection-diffusion equations with uncertain coefficients. *SIAM Journal on Scientific Computing*, 38(4):A2334–A2356, 2016.