

Table S2

Characteristic	cluster 1:lower-intake/sugar-enriched N = 114 [†]	cluster2:higher-intake N = 59 [†]
Age	60 (14)	62 (18)
Sex		
Male	50 (44%)	45 (76%)
Female	64 (56%)	14 (24%)
Disease		
Acute myeloid leukemia	47 (41%)	21 (36%)
MDS/MPN	38 (33%)	19 (32%)
Non-Hodgkin's lymphoma	6 (5.3%)	11 (19%)
Acute lymphoid leukemia	11 (9.6%)	1 (1.7%)
Other	7 (6.1%)	4 (6.8%)
Myeloma	4 (3.5%)	1 (1.7%)
Chronic lymphocytic leukemia	1 (0.9%)	2 (3.4%)
Graft type		
Unmodified bone marrow or PBSC	47 (41%)	41 (69%)
T-cell depleted PBSC	57 (50%)	15 (25%)
Cord blood	10 (8.8%)	3 (5.1%)
Intensity of conditioning regimen		
Ablative	71 (62%)	29 (49%)
Reduced intensity	39 (34%)	13 (22%)
Nonmyeloablative	4 (3.5%)	17 (29%)
GvHD prophylaxis		
CNI-based	36 (32%)	23 (39%)
PTCy-based	21 (18%)	21 (36%)
T-cell depleted PBSC	57 (50%)	15 (25%)
Days exposed to broad-spectrum antibiotics	9.0 (11.0)	3.0 (12.0)
[†] Median (IQR); n (%)		