

Environmental variability across different scales of biological organization

Maggie Slein

Reed College, Summer 2020

Contents

Acknowledgements	3
Abstract	3
Main objective	3
Introduction	4
Methods	5
Results	6
Conclusions	10

Acknowledgements

This work would not have been possible without the ingenuity and dedication of Dr. Mary O'Connor, Dr. Joey Bernhardt, Dr. Jacob Usinowicz, and Dr. Sam Fey. Without their guidance and support, this project would have been impossible. I would also like to thank the CIEE Variability working group for their willingness to let a novice take on part of their project goals. Thank you to Matt and Ellen Cutts, for without their kindness in the midst of a pandemic, there would have been no place to complete this work. The biggest thank you to my sister, Elizabeth, and her patience in listening to be babble about this work. Lastly, thank you the Reed College Biology Department and the Arch and Fran Diack Student Field Research Award for supporting this research.

Abstract

Climate change continues to push the environment and its inhabitants to the brink of their limits, albeit thermally or spatially, highlighting the importance of organisms' ability to cope in a more variable, unreliable, and stochastic world. While recent studies have demonstrated that increasing the non-linearity of thermal patterns is more detrimental to organismal performance than simply increases in the temperature (Vasseur et al 2014), there still remains a lack of agreement in the field of ecology as to how both variation and variation type influences biological responses at all levels of biological organization. Here, we aim to describe patterns of variability study in the field of ecology across all levels of organization, from changes in amplitude to changes in the predictability of variation, and the contrast environmental variation studies between different levels of organization.

Main objective

To summarize current field of environmental variation across all levels of biological organization and the pitfalls of current inquiries

Introduction

Understanding the limits of performance for organisms, populations, communities, and ecosystems has been a pertinent field of study in ecology for the last several decades (cite). However, climate change has burgeoned a revival of those questions in the face of a rapidly changing world, particularly in an increasingly variable world. Environmental variation has appeared as several terms (alternating, fluctuating, varying) to describe a counter to constant conditions in a variety of performance and dynamics studies at varying levels of biological organization (Resilva et al 2014, etc). Variation treatments often feature a range of temperatures rather than a discrete constant temperature, however, detailed patterns of variation were sporadically reported (cite). Beyond serving as an important counter to constant conditions, environmental variation has been partitioned into three subfields: temporal variation, spatial variation, and spatiotemporal variation (the interaction between both) (Di Cecco and Gouhier et al 2017). Temporal variation manipulates an environmental variable over a period of time to understand performance dynamics at the scale of interest, while spatial variation manipulates access to environmental space to understand how it affects persistence (cite). More recently, several studies have investigated the interaction between both temporal and spatial variation to understand which of the two is the dominating factor in patterns of variation (Vasseur and Fox 2009, Gonzalez and Holt 2002, Matthews and Gonzalez 2007, Fontaine and Gonzalez 2005). While these different categories are key for deducing the effects of variation in both space and time, the manipulation of variation within those groups is of particular interest to uncovering its complete effects on the environment.

Altering the frequency of environmental variation is not a new concept in the field of ecology (cite). However, Vasseur and Yodzis' (2004) emphasis of the importance of environmental variation color in biological processes responded to and coincided with several community (cite-Descamps-Julien and Gonzalez 2005, Long et al 2007) and population (Orland and Lawlor 2004) studies focused explicitly on how the color of environmental variation causes significant shifts in response patterns both temporally and spatially. Broadly, frequency ($1/T$, T =period) is a measure of the number of occurrences of a repeating event per unit time (cite). With respect to environmental variation, longer periods correspond to lower frequencies and shorter periods correspond to higher frequencies. Colloquially, "reddened series" have become synonymous with lower frequencies while "whitened series" have become synonymous with higher frequencies. Vasseur and Yodzis (2004) underscore that "an important characteristic of environmental noise is its spectrum, which describes the variance as a sum of sinusoidal waves of different frequencies." Reddened series feature differing amounts of variance across time, whereas whitened series feature equal variance across time. Reddened series are also referred to as autocorrelated series, such that due to their periodic nature, organisms can track their periodicity accordingly. Autocorrelation is of particular importance to the field of ecology as over the last several decades, environmental variables (like temperature) have become more autocorrelated and are predicted to become increasingly correlated as a result of climate change (Matthews and Gonzalez 2006, Wiggley et al 1998). Several articles have cited the importance and dominance of autocorrelated variation in driving and environmental patterns, from inflationary population effects in conjunction with dispersal (cite) to population synchrony (cite). While it would seem that the color spectrum of variation is of pressing importance, these studies remain a limited area of study nearly a decade and a half later, with most studies continuing to focus on periodic, diurnal fluctuations in amplitude (cite).

Methods

To investigate variation type, duration, and relevance across all levels of biological organization, we framed our review around these guiding questions:

1. What studies have been conducted in which environmental variability has been the treatment variable (all else being controlled)?
2. Can we summarize what types of studies have been done and where there are key gaps?
3. What aspect environmental variability was varied (SD, autocorrelation)?
4. What biological organization of response variables?
5. Can this be augmented with less-controlled studies where environmental variation occurs naturally or as a by-product of other treatment?

To answer these questions we examined environmental variation in several papers from two particular bodies of literature: plankton population and communities studies and insect organismal level studies. The plankton population and community literature focused explicitly and exhaustively on the manipulation of environmental noise. There are few studies that focus on the explicit manipulation of environmental color. The insect

to pilot preliminary conclusions with respect to variation type and fluctuation period with respect to generation time. We then summarized our conclusions from pilot work below.

Results

Result 1: Generation time is not an accurate predictor for the period of fluctuation across all levels of organization

Warning: Removed 25 rows containing missing values (geom_point).




Figure 1. Generation time (hours) and period of fluctuation (hours) across different levels of biological organization (community, host-parasite, individual, population, predator-prey) and larger organisms groups (plants, plankton, fish, aquatic invertebrates, and amphibians).

Nearly all organismal and population level studies interested in thermal performance only consider diurnal patterns of fluctuations, often varying the range of temperatures or the amplitude over a daily cycle. Few explicitly reference their justification behind the period of the fluctuation, perhaps assuming a daily period is intuitive based current environmental patterns (circadian rhythm, diel vertical migration, etc). Ironically, the small population of ecological studies focused on environmental color (variation frequency) are some of the only studies to explicitly account for study organisms relative generation times to the periodicity of the fluctuations induced (Orland and Lawler 2004, Fontaine and Gonzalez 2005). These studies emphasize that there is likely to be little effect of variation on performance if the period of the fluctuations is less than the organisms generation time, which, it appears they often are. Orland and Lawler (2004) conclude that the longer period of their fluctuation regime was the driving factor in their autocorrelated treatment, suggesting that longer periodicity may have an important effect on performance. Similarly, Fontaine and Gonzalez (2005) justify the two periods of their variation treatments as they are relative to the

generation time of the predator's generation time and the life span. Both of these studies featured fluctuation periods much long than most studies, with periods fluctuating over more than 5 days.

The mismatch between generation time and period of fluctuation is apparent, with nearly all studies featuring generation times of less than 100 days and fluctuation periods of less than 2 days (figure 1). This pattern speaks to an emphasis on diurnal fluctuations over potentially longer periods of fluctuation, which appears to contradict predictions for increased autocorrelation in the environment and suggestions to look at longer periods of fluctuation (Orland and Lawlor).

Result 2: Periodic variation on a diurnal period is prioritized in organismal level studies, both periodic and colored variation are prioritized in population level studies, while community level studies prioritize stochastic and autocorrelated variation and do so on longer periods




Figure 2. Generation time (days) and period of fluctuation (days) across all levels of biological organization (individual, population, community, host-parasite, predator-prey) paneled by whether utilized colored variation (red or white colored noise) or neglected to do so (not considered)




Figure 3. Generation time (days) and period of fluctuation (days) across all levels of biological organization (individual, population, community, host-parasite, predator-prey) and larger organism grouping (amphibian, aquatic invert, fish, insect, plankton, plant) paneled by whether utilized colored variation (red or white colored noise) or neglected to do so (not considered)

In addition to lacking diversity in fluctuation period as well as organismal generation time, most of the studies featuring longer periods occurred at the population level and were exclusively planktonic population studies (figure 1). Most studies exclusively focused on variation type featured planktonic communities and were explicitly interested in both generation time and varying fluctuation periods (Orland and Lawlor 2004, Fontaine and Gonzalez 2005, etc).

Most studies that did not account for variation color were at the individual level exclusively and were almost exclusively insect based studies (Figure 3). The emphasis on more predictable variation, simply amplifying current patterns of variability is a hallmark of these studies (cite). This theme is common across many organismal level studies interested in non-rate responses (development size, shape, egg load, etc.) (cite), population level studies interested in rate responses (growth rate, development time, fecundity, longevity, etc.) (cite). Given that most TPC studies are most interested in the organismal or population level responses to short term variation in nature (as TPC are usually on a 24 hours cycle), it is not surprising that most are interested in variation in amplitude and less in the additional stochasticity present in the natural environment. Khelifa et al (2019) demonstrated that correcting for non-linearities with high resolution data when comparing two laboratory temperature treatments, one constant and one diurnally fluctuating, allows for harmonious thermal performance between the two treatment groups. However, when attempting to accomplish the same but with field observations, featuring two treatments, one constant and one ambient measurement of field conditions over time, their methods proved unsuccessful in

accurately predicting thermal performance. Khelifa et al highlights the importance of ambient, stochastic variation in accurately predicting thermal performance, as they conclude that variability studies under laboratory conditions may underestimate thermal performance. While an important conclusion, it is one that is at odds with what the collection of literature on environmental variation has advocated for (and disagrees on). It has been established and continually reinforced that environmental variation has become increasingly autocorrelated over the last several decades and is predicted to continue to do so under climate change. Though not explicitly referenced, one can imply that Khelifa et al's conclusions about field conditions needing to be prioritized over lab studies comment on the need for more explicit investigation of autocorrelated variation.

Result 3: Larger organismal groups and biological organization level feature delineations in study interests and design




Figure 4. Generation time (days) and period of fluctuation (days) across all levels of biological organization (individual, population, community, host-parasite, predator-prey) and larger organism grouping (amphibian, aquatic invert, fish, insect, plankton, plant)

Most studies were concentrated at a finer scale than the ranges of generation times and fluctuation periods than the range allowed for (figure 1). In focusing on the concentration of studies featuring generation times of less than 90 days and fluctuations periods of less than 2 days, patterns with respect to biological organization and larger organismal groups emerged (figure 4). Limited studies were conducted of plankton at the individual level and insects at the community level. Additionally, almost all planktonic studies utilized organisms with generations times of less than 10 days, while almost all insect based studies utilized organisms with a broader range of generation times, from 0 to less than 40 days.

Conclusions

The preliminary findings in these two particular sets of literature suggest disparities in how variation is pertinent to different scales of biological organization. While studies spanned from the organismal level to the community level, few studies featured organisms with generation times of less than <