

Journal Articles

- [1] M. Aaboud, “Measurement of inclusive jet and dijet cross-sections in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **05**, 195 (2018).
- [2] M. Aaboud, “Measurement of the production cross-section of a single top quark in association with a Z boson in proton-proton collisions at 13 TeV with the ATLAS detector”, Phys. Lett. **B780**, 557–577 (2018).
- [3] M. Aaboud, “Search for a Structure in the $B_s^0\pi^\pm$ Invariant Mass Spectrum with the ATLAS Experiment”, Phys. Rev. Lett. **120**, 202007 (2018).
- [4] M. Aaboud, “Search for an invisibly decaying Higgs boson or dark matter candidates produced in association with a Z boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B776**, 318–337 (2018).
- [5] M. Aaboud, “Search for heavy ZZ resonances in the $\ell^+\ell^-\ell^+\ell^-$ and $\ell^+\ell^-\nu\bar{\nu}$ final states using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 293 (2018).
- [6] M. Aaboud, “Search for the Decay of the Higgs Boson to Charm Quarks with the ATLAS Experiment”, Phys. Rev. Lett. **120**, 211802 (2018).
- [7] M. Aaboud, “A search for lepton-flavor-violating decays of the Z boson into a τ -lepton and a light lepton with the ATLAS detector”, Submitted to: Phys. Rev. (2018).
- [8] M. Aaboud, “A search for pair-produced resonances in four-jet final states at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 250 (2018).
- [9] M. Aaboud, “A search for resonances decaying into a Higgs boson and a new particle X in the $XH \rightarrow qqbb$ final state with the ATLAS detector”, Phys. Lett. **B779**, 24–45 (2018).
- [10] M. Aaboud, “Angular analysis of $B_d^0 \rightarrow K^*\mu^+\mu^-$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, (2018).
- [11] M. Aaboud, “Combination of inclusive and differential $t\bar{t}$ charge asymmetry measurements using ATLAS and CMS data at $\sqrt{s} = 7$ and 8 TeV”, JHEP **04**, 033 (2018).
- [12] M. Aaboud, “Combined measurement of differential and total cross sections in the $H \rightarrow \gamma\gamma$ and the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channels at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [13] M. Aaboud, “Direct top-quark decay width measurement in the $t\bar{t}$ lepton+jets channel at $\sqrt{s}=8$ TeV with the ATLAS experiment”, Eur. Phys. J. **C78**, 129 (2018).
- [14] M. Aaboud, “Evidence for the associated production of the Higgs boson and a top quark pair with the ATLAS detector”, Phys. Rev. **D97**, 072003 (2018).
- [15] M. Aaboud, “Measurement of colour flow using jet-pull observables in $t\bar{t}$ events with the ATLAS experiment at $\sqrt{s} = 13$ TeV”, (2018).
- [16] M. Aaboud, “Measurement of differential cross sections and W^+/W^- cross-section ratios for W boson production in association with jets at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **05**, 077 (2018).
- [17] M. Aaboud, “Measurement of differential cross sections of isolated-photon plus heavy-flavour jet production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Lett. **B776**, 295–317 (2018).
- [18] M. Aaboud, “Measurement of differential cross-sections of a single top quark produced in association with a W boson at $\sqrt{s} = 13$ TeV with ATLAS”, Eur. Phys. J. **C78**, 186 (2018).
- [19] M. Aaboud, “Measurement of dijet azimuthal decorrelations in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector and determination of the strong coupling”, (2018).
- [20] M. Aaboud, “Measurement of jet fragmentation in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, (2018).
- [21] M. Aaboud, “Measurement of longitudinal flow decorrelations in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ and 5.02 TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 142 (2018).
- [22] M. Aaboud, “Measurement of long-range multiparticle azimuthal correlations with the subevent cumulant method in pp and $p + Pb$ collisions with the ATLAS detector at the CERN Large Hadron Collider”, Phys. Rev. **C97**, 024904 (2018).
- [23] M. Aaboud, “Measurement of τ polarisation in $Z/\gamma^* \rightarrow \tau\tau$ decays in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 163 (2018).
- [24] M. Aaboud, “Measurement of quarkonium production in proton-lead and proton-proton collisions at 5.02 TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 171 (2018).

- [25] M. Aaboud, “Measurement of the cross section for isolated-photon plus jet production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, Phys. Lett. **B780**, 578–602 (2018).
- [26] M. Aaboud, “Measurement of the cross-section for producing a W boson in association with a single top quark in pp collisions at $\sqrt{s} = 13$ TeV with ATLAS”, JHEP **01**, 063 (2018).
- [27] M. Aaboud, “Measurement of the exclusive $\gamma\gamma \rightarrow \mu^+\mu^-$ process in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B777**, 303–323 (2018).
- [28] M. Aaboud, “Measurement of the Higgs boson coupling properties in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 095 (2018).
- [29] M. Aaboud, “Measurement of the Higgs boson mass in the $H \rightarrow ZZ^* \rightarrow 4\ell$ and $H \rightarrow \gamma\gamma$ channels with $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, (2018).
- [30] M. Aaboud, “Measurement of the inclusive and fiducial $t\bar{t}$ production cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 487 (2018).
- [31] M. Aaboud, “Measurement of the nuclear modification factor for inclusive jets in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, (2018).
- [32] M. Aaboud, “Measurement of the production cross section of three isolated photons in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Lett. **B781**, 55–76 (2018).
- [33] M. Aaboud, “Measurement of the suppression and azimuthal anisotropy of muons from heavy-flavor decays in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, (2018).
- [34] M. Aaboud, “Measurement of the W-boson mass in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 110 (2018).
- [35] M. Aaboud, “Measurements of b -jet tagging efficiency with the ATLAS detector using $t\bar{t}$ events at $\sqrt{s} = 13$ TeV”, (2018).
- [36] M. Aaboud, “Measurements of differential cross sections of top quark pair production in association with jets in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, (2018).
- [37] M. Aaboud, “Measurements of Higgs boson properties in the diphoton decay channel with 36 fb^{-1} of pp collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [38] M. Aaboud, “Measurements of $t\bar{t}$ differential cross-sections of highly boosted top quarks decaying to all-hadronic final states in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, (2018).
- [39] M. Aaboud, “Observation of centrality-dependent acoplanarity for muon pairs produced via two-photon scattering in Pb+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, (2018).
- [40] M. Aaboud, “Observation of Higgs boson production in association with a top quark pair at the LHC with the ATLAS detector”, (2018).
- [41] M. Aaboud, “Operation and performance of the ATLAS Tile Calorimeter in Run 1”, (2018).
- [42] M. Aaboud, “Performance of missing transverse momentum reconstruction with the ATLAS detector using proton-proton collisions at $\sqrt{s} = 13$ TeV”, (2018).
- [43] M. Aaboud, “Probing the quantum interference between singly and doubly resonant top-quark production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Submitted to: Phys. Rev. Lett. (2018).
- [44] M. Aaboud, “Prompt and non-prompt J/ψ and $\psi(2S)$ suppression at high transverse momentum in 5.02 TeV Pb+Pb collisions with the ATLAS experiment”, (2018).
- [45] M. Aaboud, “Search for a heavy Higgs boson decaying into a Z boson and another heavy Higgs boson in the $\ell\ell b\bar{b}$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [46] M. Aaboud, “Search for a new heavy gauge boson resonance decaying into a lepton and missing transverse momentum in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, Eur. Phys. J. **C78**, 401 (2018).
- [47] M. Aaboud, “Search for additional heavy neutral Higgs and gauge bosons in the ditau final state produced in 36 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **01**, 055 (2018).
- [48] M. Aaboud, “Search for B-L R -parity-violating top squarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS experiment”, Phys. Rev. **D97**, 032003 (2018).
- [49] M. Aaboud, “Search for chargino-neutralino production using recursive jigsaw reconstruction in final states with two or three charged leptons in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [50] M. Aaboud, “Search for dark matter and other new phenomena in events with an energetic jet and large missing transverse momentum using the ATLAS detector”, JHEP **01**, 126 (2018).
- [51] M. Aaboud, “Search for dark matter produced in association with bottom or top quarks in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C78**, 18 (2018).

- [52] M. Aaboud, “Search for diboson resonances with boson-tagged jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B777**, 91–113 (2018).
- [53] M. Aaboud, “Search for doubly charged Higgs boson production in multi-lepton final states with the ATLAS detector using proton–proton collisions at $\sqrt{s} = 13$ TeV”, Eur. Phys. J. **C78**, 199 (2018).
- [54] M. Aaboud, “Search for electroweak production of supersymmetric particles in final states with two or three leptons at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [55] M. Aaboud, “Search for electroweak production of supersymmetric states in scenarios with compressed mass spectra at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. **D97**, 052010 (2018).
- [56] M. Aaboud, “Search for flavor-changing neutral currents in top quark decays $t \rightarrow Hc$ and $t \rightarrow Hu$ in multilepton final states in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [57] M. Aaboud, “Search for flavour-changing neutral current top-quark decays $t \rightarrow qZ$ in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [58] M. Aaboud, “Search for heavy particles decaying into top-quark pairs using lepton-plus-jets events in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Submitted to: Eur. Phys. J. (2018).
- [59] M. Aaboud, “Search for heavy resonances decaying into a W or Z boson and a Higgs boson in final states with leptons and b -jets in 36 fb^{-1} of $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, JHEP **03**, 174 (2018).
- [60] M. Aaboud, “Search for heavy resonances decaying into WW in the $e\nu\mu\nu$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 24 (2018).
- [61] M. Aaboud, “Search for heavy resonances decaying to a photon and a hadronically decaying $Z/W/H$ boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [62] M. Aaboud, “Search for Higgs boson decays into pairs of light (pseudo)scalar particles in the $\gamma\gamma jj$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B782**, 750–767 (2018).
- [63] M. Aaboud, “Search for Higgs boson decays to beyond-the-Standard-Model light bosons in four-lepton events with the ATLAS detector at $\sqrt{s} = 13$ TeV”, (2018).
- [64] M. Aaboud, “Search for High-Mass Resonances Decaying to $\tau\nu$ in pp Collisions at $\sqrt{s}=13\text{TeV}$ with the ATLAS Detector”, Phys. Rev. Lett. **120**, 161802 (2018).
- [65] M. Aaboud, “Search for light resonances decaying to boosted quark pairs and produced in association with a photon or a jet in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [66] M. Aaboud, “Search for long-lived charginos based on a disappearing-track signature in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **06**, 022 (2018).
- [67] M. Aaboud, “Search for long-lived, massive particles in events with displaced vertices and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Phys. Rev. **D97**, 052012 (2018).
- [68] M. Aaboud, “Search for low-mass dijet resonances using trigger-level jets with the ATLAS detector in pp collisions at $\sqrt{s} = 13$ TeV”, (2018).
- [69] M. Aaboud, “Search for new phenomena in high-mass final states with a photon and a jet from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C78**, 102 (2018).
- [70] M. Aaboud, “Search for new phenomena using the invariant mass distribution of same-flavour opposite-sign dilepton pairs in events with missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, (2018).
- [71] M. Aaboud, “Search for pair- and single-production of vector-like quarks in final states with at least one Z boson decaying into a pair of electrons or muons in pp collision data collected with the ATLAS detector at $\sqrt{s} = 13$ TeV”, (2018).
- [72] M. Aaboud, “Search for pair production of heavy vector-like quarks decaying into high- p_T W bosons and top quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [73] M. Aaboud, “Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [74] M. Aaboud, “Search for pair production of higgsinos in final states with at least three b -tagged jets in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, Submitted to: Phys. Rev. (2018).
- [75] M. Aaboud, “Search for pair production of up-type vector-like quarks and for four-top-quark events in final states with multiple b -jets with the ATLAS detector”, (2018).
- [76] M. Aaboud, “Search for photonic signatures of gauge-mediated supersymmetry in 13 TeV pp collisions with the ATLAS detector”, Phys. Rev. **D97**, 092006 (2018).

- [77] M. Aaboud, “Search for resonances in the mass distribution of jet pairs with one or two jets identified as b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [78] M. Aaboud, “Search for resonant WZ production in the fully leptonic final state in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [79] M. Aaboud, “Search for R-parity-violating supersymmetric particles in multi-jet final states produced in p - p collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC”, (2018).
- [80] M. Aaboud, “Search for squarks and gluinos in final states with jets and missing transverse momentum using 36fb^{-1} of $\sqrt{s} = 13\text{TeV}$ pp collision data with the ATLAS detector”, Phys. Rev. **D97**, 112001 (2018).
- [81] M. Aaboud, “Search for supersymmetry in events with four or more leptons in $\sqrt{s} = 13$ TeV pp collisions with ATLAS”, (2018).
- [82] M. Aaboud, “Search for supersymmetry in final states with charm jets and missing transverse momentum in 13 TeV pp collisions with the ATLAS detector”, (2018).
- [83] M. Aaboud, “Search for the direct production of charginos and neutralinos in final states with tau leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C78**, 154 (2018).
- [84] M. Aaboud, “Search for the Higgs boson produced in association with a vector boson and decaying into two spin-zero particles in the $H \rightarrow aa \rightarrow 4b$ channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [85] M. Aaboud, “Search for the standard model Higgs boson produced in association with top quarks and decaying into a $b\bar{b}$ pair in pp collisions at $\sqrt{s} = 13\text{TeV}$ with the ATLAS detector”, Phys. Rev. **D97**, 072016 (2018).
- [86] M. Aaboud, “Search for top squarks decaying to tau sleptons in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2018).
- [87] M. Aaboud, “Search for $W' \rightarrow tb$ decays in the hadronic final state using pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B781**, 327–348 (2018).
- [88] M. Aaboud, “Search for WW/WZ resonance production in $\ell\nu qq$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 042 (2018).
- [89] M. Aaboud, “Searches for heavy ZZ and ZW resonances in the $\ell\ell qq$ and $\nu\nu qq$ final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 009 (2018).
- [90] M. Aaboud, “ $ZZ \rightarrow \ell^+\ell^-\ell'^+\ell'^-$ cross-section measurements and search for anomalous triple gauge couplings in 13 TeV pp collisions with the ATLAS detector”, Phys. Rev. **D97**, 032005 (2018).
- [91] L. Bauerdick, “HEP Software Foundation Community White Paper Working Group - Data Analysis and Interpretation”, edited by M. S. Neubauer (2018).
- [92] M. Aaboud, “Evidence for the $H \rightarrow b\bar{b}$ decay with the ATLAS detector”, JHEP **12**, 024 (2017).
- [93] M. Aaboud, “Jet energy scale measurements and their systematic uncertainties in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. **D96**, 072002 (2017).
- [94] M. Aaboud, “Measurement of b -hadron pair production with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **11**, 062 (2017).
- [95] M. Aaboud, “Measurement of lepton differential distributions and the top quark mass in $t\bar{t}$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 804 (2017).
- [96] M. Aaboud, “Measurement of the cross-section for electroweak production of dijets in association with a Z boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B775**, 206–228 (2017).
- [97] M. Aaboud, “Measurement of the Drell-Yan triple-differential cross section in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **12**, 059 (2017).
- [98] M. Aaboud, “Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, JHEP **11**, 191 (2017).
- [99] M. Aaboud, “Performance of the ATLAS Track Reconstruction Algorithms in Dense Environments in LHC Run 2”, Eur. Phys. J. **C77**, 673 (2017).
- [100] M. Aaboud, “Search for direct top squark pair production in final states with two leptons in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C77**, 898 (2017).
- [101] M. Aaboud, “Search for pair production of heavy vector-like quarks decaying to high- p_T W bosons and b quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 141 (2017).
- [102] M. Aaboud, “Searches for the $Z\gamma$ decay mode of the Higgs boson and for new high-mass resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 112 (2017).

- [103] M. Aaboud, “Study of ordered hadron chains with the ATLAS detector”, Phys. Rev. **D96**, 092008 (2017).
- [104] M. Aaboud, “Study of $WW\gamma$ and $WZ\gamma$ production in pp collisions at $\sqrt{s} = 8$ TeV and search for anomalous quartic gauge couplings with the ATLAS experiment”, Eur. Phys. J. **C77**, 646 (2017).
- [105] M. Aaboud, “A measurement of the calorimeter response to single hadrons and determination of the jet energy scale uncertainty using LHC Run-1 pp -collision data with the ATLAS detector”, Eur. Phys. J. **C77**, 26 (2017).
- [106] M. Aaboud, “A measurement of the soft-drop jet mass in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, (2017).
- [107] M. Aaboud, “Analysis of the Wtb vertex from the measurement of triple-differential angular decay rates of single top quarks produced in the t -channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **12**, 017 (2017).
- [108] M. Aaboud, “Determination of the strong coupling constant α_s from transverse energy–energy correlations in multijet events at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Eur. Phys. J. **C77**, 872 (2017).
- [109] M. Aaboud, “Electron efficiency measurements with the ATLAS detector using 2012 LHC proton–proton collision data”, Eur. Phys. J. **C77**, 195 (2017).
- [110] M. Aaboud, “Evidence for light-by-light scattering in heavy-ion collisions with the ATLAS detector at the LHC”, Nature Phys. **13**, 852–858 (2017).
- [111] M. Aaboud, “Femtoscopia with identified charged pions in proton-lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV with ATLAS”, Phys. Rev. **C96**, 064908 (2017).
- [112] M. Aaboud, “Fiducial, total and differential cross-section measurements of t -channel single top-quark production in pp collisions at 8 TeV using data collected by the ATLAS detector”, Eur. Phys. J. **C77**, 531 (2017).
- [113] M. Aaboud, “High- E_T isolated-photon plus jets production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Nucl. Phys. **B918**, 257–316 (2017).
- [114] M. Aaboud, “Identification and rejection of pile-up jets at high pseudorapidity with the ATLAS detector”, Eur. Phys. J. **C77**, [Erratum: Eur. Phys. J.C77,no.10,712(2017)], 580 (2017).
- [115] M. Aaboud, “Jet reconstruction and performance using particle flow with the ATLAS Detector”, Eur. Phys. J. **C77**, 466 (2017).
- [116] M. Aaboud, “Measurement of charged-particle distributions sensitive to the underlying event in $\sqrt{s} = 13$ TeV proton-proton collisions with the ATLAS detector at the LHC”, JHEP **03**, 157 (2017).
- [117] M. Aaboud, “Measurement of detector-corrected observables sensitive to the anomalous production of events with jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, Eur. Phys. J. **C77**, 765 (2017).
- [118] M. Aaboud, “Measurement of forward-backward multiplicity correlations in lead-lead, proton-lead, and proton-proton collisions with the ATLAS detector”, Phys. Rev. **C95**, 064914 (2017).
- [119] M. Aaboud, “Measurement of inclusive and differential cross sections in the $H \rightarrow ZZ^* \rightarrow 4\ell$ decay channel in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **10**, 132 (2017).
- [120] M. Aaboud, “Measurement of jet activity produced in top-quark events with an electron, a muon and two b -tagged jets in the final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 220 (2017).
- [121] M. Aaboud, “Measurement of jet fragmentation in 5.02 TeV proton–lead and proton–proton collisions with the ATLAS detector”, (2017).
- [122] M. Aaboud, “Measurement of jet fragmentation in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC”, Eur. Phys. J. **C77**, 379 (2017).
- [123] M. Aaboud, “Measurement of jet p_T correlations in Pb+Pb and pp collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B774**, 379–402 (2017).
- [124] M. Aaboud, “Measurement of multi-particle azimuthal correlations in pp , p +Pb and low-multiplicity Pb+Pb collisions with the ATLAS detector”, Eur. Phys. J. **C77**, 428 (2017).
- [125] M. Aaboud, “Measurement of the cross section for inclusive isolated-photon production in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, Phys. Lett. **B770**, 473–493 (2017).
- [126] M. Aaboud, “Measurement of the inclusive cross-sections of single top-quark and top-antiquark t -channel production in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **04**, 086 (2017).
- [127] M. Aaboud, “Measurement of the inclusive jet cross-sections in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 020 (2017).

- [128] M. Aaboud, “Measurement of the k_t splitting scales in $Z \rightarrow \ell\ell$ events in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 026 (2017).
- [129] M. Aaboud, “Measurement of the prompt J/ψ pair production cross-section in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 76 (2017).
- [130] M. Aaboud, “Measurement of the $t\bar{t}\gamma$ production cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **11**, 086 (2017).
- [131] M. Aaboud, “Measurement of the $t\bar{t}$ production cross section in the $\tau + \text{jets}$ final state in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Rev. **D95**, 072003 (2017).
- [132] M. Aaboud, “Measurement of the $t\bar{t}Z$ and $t\bar{t}W$ production cross sections in multilepton final states using 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 40 (2017).
- [133] M. Aaboud, “Measurement of the W boson polarisation in $t\bar{t}$ events from pp collisions at $\sqrt{s} = 8$ TeV in the lepton + jets channel with ATLAS”, Eur. Phys. J. **C77**, 264 (2017).
- [134] M. Aaboud, “Measurement of the W^+W^- production cross section in pp collisions at a centre-of-mass energy of $\sqrt{s} = 13$ TeV with the ATLAS experiment”, Phys. Lett. **B773**, 354–374 (2017).
- [135] M. Aaboud, “Measurement of the ZZ production cross section in proton-proton collisions at $\sqrt{s} = 8$ TeV using the $ZZ \rightarrow \ell^-\ell^+\ell'^-\ell'^+$ and $ZZ \rightarrow \ell^-\ell^+\nu\bar{\nu}$ channels with the ATLAS detector”, JHEP **01**, 099 (2017).
- [136] M. Aaboud, “Measurement of W boson angular distributions in events with high transverse momentum jets at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Lett. **B765**, 132–153 (2017).
- [137] M. Aaboud, “Measurement of $W^\pm W^\pm$ vector-boson scattering and limits on anomalous quartic gauge couplings with the ATLAS detector”, Phys. Rev. **D96**, 012007 (2017).
- [138] M. Aaboud, “Measurement of $WW/WZ \rightarrow \ell\nu qq'$ production with the hadronically decaying boson reconstructed as one or two jets in pp collisions at $\sqrt{s} = 8$ TeV with ATLAS, and constraints on anomalous gauge couplings”, Eur. Phys. J. **C77**, 563 (2017).
- [139] M. Aaboud, “Measurements of $\psi(2S)$ and $X(3872) \rightarrow J/\psi\pi^+\pi^-$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **01**, 117 (2017).
- [140] M. Aaboud, “Measurements of charge and CP asymmetries in b -hadron decays using top-quark events collected by the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **02**, 071 (2017).
- [141] M. Aaboud, “Measurements of electroweak Wjj production and constraints on anomalous gauge couplings with the ATLAS detector”, Eur. Phys. J. **C77**, 474 (2017).
- [142] M. Aaboud, “Measurements of integrated and differential cross sections for isolated photon pair production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D95**, 112005 (2017).
- [143] M. Aaboud, “Measurements of long-range azimuthal anisotropies and associated Fourier coefficients for pp collisions at $\sqrt{s} = 5.02$ and 13 TeV and $p+\text{Pb}$ collisions at $\sqrt{s_{\text{NN}}} = 5.02$ TeV with the ATLAS detector”, Phys. Rev. **C96**, 024908 (2017).
- [144] M. Aaboud, “Measurements of the production cross section of a Z boson in association with jets in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 361 (2017).
- [145] M. Aaboud, “Measurements of top quark spin observables in $t\bar{t}$ events using dilepton final states in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, JHEP **03**, 113 (2017).
- [146] M. Aaboud, “Measurements of top-quark pair differential cross-sections in the $e\mu$ channel in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, Eur. Phys. J. **C77**, 292 (2017).
- [147] M. Aaboud, “Measurements of top-quark pair to Z -boson cross-section ratios at $\sqrt{s} = 13, 8, 7$ TeV with the ATLAS detector”, JHEP **02**, 117 (2017).
- [148] M. Aaboud, “Performance of the ATLAS Transition Radiation Tracker in Run 1 of the LHC: tracker properties”, JINST **12**, P05002 (2017).
- [149] M. Aaboud, “Performance of the ATLAS Trigger System in 2015”, Eur. Phys. J. **C77**, 317 (2017).
- [150] M. Aaboud, “Precision measurement and interpretation of inclusive W^+ , W^- and Z/γ^* production cross sections with the ATLAS detector”, Eur. Phys. J. **C77**, 367 (2017).
- [151] M. Aaboud, “Probing the W $t\bar{b}$ vertex structure in t -channel single-top-quark production and decay in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **04**, 124 (2017).
- [152] M. Aaboud, “Reconstruction of primary vertices at the ATLAS experiment in Run 1 proton–proton collisions at the LHC”, Eur. Phys. J. **C77**, 332 (2017).
- [153] M. Aaboud, “Search for a scalar partner of the top quark in the jets plus missing transverse momentum final state at $\sqrt{s}=13$ TeV with the ATLAS detector”, JHEP **12**, 085 (2017).

- [154] M. Aaboud, “Search for anomalous electroweak production of WW/WZ in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, *Phys. Rev.* **D95**, 032001 (2017).
- [155] M. Aaboud, “Search for dark matter at $\sqrt{s} = 13$ TeV in final states containing an energetic photon and large missing transverse momentum with the ATLAS detector”, *Eur. Phys. J.* **C77**, 393 (2017).
- [156] M. Aaboud, “Search for dark matter in association with a Higgs boson decaying to b -quarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B765**, 11–31 (2017).
- [157] M. Aaboud, “Search for dark matter in association with a Higgs boson decaying to two photons at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev.* **D96**, 112004 (2017).
- [158] M. Aaboud, “Search for Dark Matter Produced in Association with a Higgs Boson Decaying to $b\bar{b}$ using 36 fb⁻¹ of pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **119**, 181804 (2017).
- [159] M. Aaboud, “Search for direct top squark pair production in events with a Higgs or Z boson, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *JHEP* **08**, 006 (2017).
- [160] M. Aaboud, “Search for exclusive Higgs and Z boson decays to $\phi\gamma$ and $\rho\gamma$ with the ATLAS detector”, (2017).
- [161] M. Aaboud, “Search for Heavy Higgs Bosons A/H Decaying to a Top Quark Pair in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, *Phys. Rev. Lett.* **119**, 191803 (2017).
- [162] M. Aaboud, “Search for heavy resonances decaying to a W or Z boson and a Higgs boson in the $q\bar{q}^{(\prime)}b\bar{b}$ final state in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B774**, 494–515 (2017).
- [163] M. Aaboud, “Search for heavy resonances decaying to a Z boson and a photon in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B764**, 11–30 (2017).
- [164] M. Aaboud, “Search for new high-mass phenomena in the dilepton final state using 36 fb¹ of proton-proton collision data at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **10**, 182 (2017).
- [165] M. Aaboud, “Search for new phenomena in a lepton plus high jet multiplicity final state with the ATLAS experiment using $\sqrt{s} = 13$ TeV proton-proton collision data”, *JHEP* **09**, 088 (2017).
- [166] M. Aaboud, “Search for new phenomena in dijet events using 37 fb⁻¹ of pp collision data collected at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev.* **D96**, 052004 (2017).
- [167] M. Aaboud, “Search for new phenomena in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 13$ pp collisions with the ATLAS detector”, *Eur. Phys. J.* **C77**, 144 (2017).
- [168] M. Aaboud, “Search for new phenomena in high-mass diphoton final states using 37 fb⁻¹ of proton-proton collisions collected at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B775**, 105–125 (2017).
- [169] M. Aaboud, “Search for new phenomena with large jet multiplicities and missing transverse momentum using large-radius jets and flavour-tagging at ATLAS in 13 TeV pp collisions”, *JHEP* **12**, 034 (2017).
- [170] M. Aaboud, “Search for new resonances decaying to a W or Z boson and a Higgs boson in the $\ell^+\ell^-b\bar{b}$, $\ell\nu b\bar{b}$, and $\nu\bar{\nu}b\bar{b}$ channels with pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B765**, 32–52 (2017).
- [171] M. Aaboud, “Search for pair production of vector-like top quarks in events with one lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *JHEP* **08**, 052 (2017).
- [172] M. Aaboud, “Search for squarks and gluinos in events with an isolated lepton, jets, and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev.* **D96**, 112010 (2017).
- [173] M. Aaboud, “Search for supersymmetry in events with b -tagged jets and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **11**, 195 (2017).
- [174] M. Aaboud, “Search for Supersymmetry in final states with missing transverse momentum and multiple b -jets in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 107 (2017).
- [175] M. Aaboud, “Search for supersymmetry in final states with two same-sign or three leptons and jets using 36 fb⁻¹ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, *JHEP* **09**, 084 (2017).
- [176] M. Aaboud, “Search for the dimuon decay of the Higgs boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev. Lett.* **119**, 051802 (2017).
- [177] M. Aaboud, “Search for top quark decays $t \rightarrow qH$, with $H \rightarrow \gamma\gamma$, in $\sqrt{s} = 13$ TeV pp collisions using the ATLAS detector”, *JHEP* **10**, 129 (2017).

- [178] M. Aaboud, “Search for top-squark pair production in final states with one lepton, jets, and missing transverse momentum using 36 fb⁻¹ of $\sqrt{s} = 13$ TeV pp collision data with the ATLAS detector”, (2017).
- [179] M. Aaboud, “Search for triboson $W^\pm W^\pm W^\mp$ production in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C77**, 141 (2017).
- [180] M. Aaboud, “Studies of $Z\gamma$ production in association with a high-mass dijet system in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **07**, 107 (2017).
- [181] M. Aaboud, “Study of the material of the ATLAS inner detector for Run 2 of the LHC”, JINST **12**, P12009 (2017).
- [182] M. Aaboud, “Top-quark mass measurement in the all-hadronic $t\bar{t}$ decay channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 118 (2017).
- [183] G. Aad, “Performance of algorithms that reconstruct missing transverse momentum in $\sqrt{s} = 8$ TeV proton-proton collisions in the ATLAS detector”, Eur. Phys. J. **C77**, 241 (2017).
- [184] G. Aad, “Search for lepton-flavour-violating decays of the Higgs and Z bosons with the ATLAS detector”, Eur. Phys. J. **C77**, 70 (2017).
- [185] G. Aad, “Topological cell clustering in the ATLAS calorimeters and its performance in LHC Run 1”, Eur. Phys. J. **C77**, 490 (2017).
- [186] T. Aaltonen, “Observation of the $Y(4140)$ structure in the $J/\psi\phi$ mass spectrum in $B^\pm \rightarrow J/\psi\phi K^\pm$ decays”, Mod. Phys. Lett. **A32**, 1750139 (2017).
- [187] A. A. Alves, “A Roadmap for HEP Software and Computing R&D for the 2020s”, (2017).
- [188] P. Elmer, **M. Neubauer**, M. D. Sokoloff, “Strategic Plan for a Scientific Software Innovation Institute (S2I2) for High Energy Physics”, (2017).
- [189] M. Aaboud, “Measurement of the Inelastic Proton-Proton Cross Section at $\sqrt{s} = 13$ TeV with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **117**, 182002 (2016).
- [190] M. Aaboud, “A measurement of material in the ATLAS tracker using secondary hadronic interactions in 7 TeV pp collisions”, JINST **11**, P11020 (2016).
- [191] M. Aaboud, “Charged-particle distributions at low transverse momentum in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC”, Eur. Phys. J. **C76**, 502 (2016).
- [192] M. Aaboud, “Dark matter interpretations of ATLAS searches for the electroweak production of supersymmetric particles in $\sqrt{s} = 8$ TeV proton-proton collisions”, JHEP **09**, 175 (2016).
- [193] M. Aaboud, “Luminosity determination in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector at the LHC”, Eur. Phys. J. **C76**, 653 (2016).
- [194] M. Aaboud, “Measurement of exclusive $\gamma\gamma \rightarrow W^+W^-$ production and search for exclusive Higgs boson production in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Phys. Rev. **D94**, 032011 (2016).
- [195] M. Aaboud, “Measurement of jet activity in top quark events using the $e\mu$ final state with two b -tagged jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 074 (2016).
- [196] M. Aaboud, “Measurement of the $b\bar{b}$ dijet cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 670 (2016).
- [197] M. Aaboud, “Measurement of the photon identification efficiencies with the ATLAS detector using LHC Run-1 data”, Eur. Phys. J. **C76**, 666 (2016).
- [198] M. Aaboud, “Measurement of the relative width difference of the $B^0\text{--}\bar{B}^0$ system with the ATLAS detector”, JHEP **06**, 081 (2016).
- [199] M. Aaboud, “Measurement of the top quark mass in the $t\bar{t} \rightarrow$ dilepton channel from $\sqrt{s} = 8$ TeV ATLAS data”, Phys. Lett. **B761**, 350–371 (2016).
- [200] M. Aaboud, “Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B761**, 158–178 (2016).
- [201] M. Aaboud, “Measurement of the $t\bar{t}$ production cross-section using $e\mu$ events with b -tagged jets in pp collisions at $\sqrt{s}=13$ TeV with the ATLAS detector”, Phys. Lett. **B761**, [Erratum: Phys. Lett.B772,879(2017)], 136–157 (2016).
- [202] M. Aaboud, “Measurement of the $W^\pm Z$ boson pair-production cross section in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, Phys. Lett. **B762**, 1–22 (2016).
- [203] M. Aaboud, “Measurement of top quark pair differential cross-sections in the dilepton channel in pp collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS”, Phys. Rev. **D94**, 092003 (2016).
- [204] M. Aaboud, “Measurement of W^+W^- production in association with one jet in proton–proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B763**, 114–133 (2016).

- [205] M. Aaboud, “Search for bottom squark pair production in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J.* **C76**, 547 (2016).
- [206] M. Aaboud, “Search for charged Higgs bosons produced in association with a top quark and decaying via $H^\pm \rightarrow \tau\nu$ using pp collision data recorded at $\sqrt{s} = 13$ TeV by the ATLAS detector”, *Phys. Lett.* **B759**, 555–574 (2016).
- [207] M. Aaboud, “Search for dark matter produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B763**, 251–268 (2016).
- [208] M. Aaboud, “Search for heavy long-lived charged R -hadrons with the ATLAS detector in 3.2 fb^{-1} of proton–proton collision data at $\sqrt{s} = 13$ TeV”, *Phys. Lett.* **B760**, 647–665 (2016).
- [209] M. Aaboud, “Search for Higgs and Z Boson Decays to $\phi\gamma$ with the ATLAS Detector”, *Phys. Rev. Lett.* **117**, 111802 (2016).
- [210] M. Aaboud, “Search for high-mass new phenomena in the dilepton final state using proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B761**, 372–392 (2016).
- [211] M. Aaboud, “Search for metastable heavy charged particles with large ionization energy loss in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS experiment”, *Phys. Rev.* **D93**, 112015 (2016).
- [212] M. Aaboud, “Search for Minimal Supersymmetric Standard Model Higgs bosons H/A and for a Z' boson in the $\tau\tau$ final state produced in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS Detector”, *Eur. Phys. J.* **C76**, 585 (2016).
- [213] M. Aaboud, “Search for new phenomena in different-flavour high-mass dilepton final states in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J.* **C76**, 541 (2016).
- [214] M. Aaboud, “Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **06**, 059 (2016).
- [215] M. Aaboud, “Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector”, *Phys. Rev.* **D94**, 032005 (2016).
- [216] M. Aaboud, “Search for new resonances in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B762**, 334–352 (2016).
- [217] M. Aaboud, “Search for pair production of Higgs bosons in the $b\bar{b}b\bar{b}$ final state using proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Rev.* **D94**, 052002 (2016).
- [218] M. Aaboud, “Search for resonances in diphoton events at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **09**, 001 (2016).
- [219] M. Aaboud, “Search for resonances in the mass distribution of jet pairs with one or two jets identified as b -jets in proton–proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Phys. Lett.* **B759**, 229–246 (2016).
- [220] M. Aaboud, “Search for scalar leptoquarks in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS experiment”, *New J. Phys.* **18**, 093016 (2016).
- [221] M. Aaboud, “Search for squarks and gluinos in events with hadronically decaying tau leptons, jets and missing transverse momentum in proton–proton collisions at $\sqrt{s} = 13$ TeV recorded with the ATLAS detector”, *Eur. Phys. J.* **C76**, 683 (2016).
- [222] M. Aaboud, “Search for squarks and gluinos in final states with jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *Eur. Phys. J.* **C76**, 392 (2016).
- [223] M. Aaboud, “Search for supersymmetry in a final state containing two photons and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions at the LHC using the ATLAS detector”, *Eur. Phys. J.* **C76**, 517 (2016).
- [224] M. Aaboud, “Search for TeV-scale gravity signatures in high-mass final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 13$ TeV”, *Phys. Lett.* **B760**, 520–537 (2016).
- [225] M. Aaboud, “Search for the Higgs boson produced in association with a W boson and decaying to four b -quarks via two spin-zero particles in pp collisions at 13 TeV with the ATLAS detector”, *Eur. Phys. J.* **C76**, 605 (2016).
- [226] M. Aaboud, “Search for the Standard Model Higgs boson produced by vector-boson fusion and decaying to bottom quarks in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, *JHEP* **11**, 112 (2016).
- [227] M. Aaboud, “Search for top squarks in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 13$ TeV pp collisions with the ATLAS detector”, *Phys. Rev.* **D94**, 052009 (2016).
- [228] M. Aaboud, “Searches for heavy diboson resonances in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, *JHEP* **09**, 173 (2016).

- [229] M. Aaboud, “Study of hard double-parton scattering in four-jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, JHEP **11**, 110 (2016).
- [230] M. Aaboud, “Study of the rare decays of B_s^0 and B^0 into muon pairs from data collected during the LHC Run 1 with the ATLAS detector”, Eur. Phys. J. **C76**, 513 (2016).
- [231] G. Aad, “A new method to distinguish hadronically decaying boosted Z bosons from W bosons using the ATLAS detector”, Eur. Phys. J. **C76**, 238 (2016).
- [232] G. Aad, “A search for an excited muon decaying to a muon and two jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, New J. Phys. **18**, 073021 (2016).
- [233] G. Aad, “A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **02**, 062 (2016).
- [234] G. Aad, “A search for top squarks with R-parity-violating decays to all-hadronic final states with the ATLAS detector in $\sqrt{s} = 8$ TeV proton-proton collisions”, JHEP **06**, 067 (2016).
- [235] G. Aad, “Beam-induced and cosmic-ray backgrounds observed in the ATLAS detector during the LHC 2012 proton-proton running period”, JINST **11**, P05013 (2016).
- [236] G. Aad, “Centrality, rapidity and transverse momentum dependence of isolated prompt photon production in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV measured with the ATLAS detector”, Phys. Rev. **C93**, 034914 (2016).
- [237] G. Aad, “Charged-particle distributions in pp interactions at $\sqrt{s} = 8$ TeV measured with the ATLAS detector”, Eur. Phys. J. **C76**, 403 (2016).
- [238] G. Aad, “Charged-particle distributions in $\sqrt{s} = 13$ TeV pp interactions measured with the ATLAS detector at the LHC”, Phys. Lett. **B758**, 67–88 (2016).
- [239] G. Aad, “Combination of searches for WW , WZ , and ZZ resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B755**, 285–305 (2016).
- [240] G. Aad, “Constraints on non-Standard Model Higgs boson interactions in an effective Lagrangian using differential cross sections measured in the $H \rightarrow \gamma\gamma$ decay channel at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B753**, 69–85 (2016).
- [241] G. Aad, “Dijet production in $\sqrt{s} = 7$ TeV pp collisions with large rapidity gaps at the ATLAS experiment”, Phys. Lett. **B754**, 214–234 (2016).
- [242] G. Aad, “Evidence for single top-quark production in the s -channel in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector using the Matrix Element Method”, Phys. Lett. **B756**, 228–246 (2016).
- [243] G. Aad, “Identification of boosted, hadronically decaying W bosons and comparisons with ATLAS data taken at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. **C76**, 154 (2016).
- [244] G. Aad, “Identification of high transverse momentum top quarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **06**, 093 (2016).
- [245] G. Aad, “Measurement of $D^{*\pm}$, D^\pm and D_s^\pm meson production cross sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. **B907**, 717–763 (2016).
- [246] G. Aad, “Measurement of event-shape observables in $Z \rightarrow \ell^+\ell^-$ events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC”, Eur. Phys. J. **C76**, 375 (2016).
- [247] G. Aad, “Measurement of fiducial differential cross sections of gluon-fusion production of Higgs bosons decaying to $WW \rightarrow e$ with the ATLAS detector at $\sqrt{s} = 8$ TeV”, JHEP **08**, 104 (2016).
- [248] G. Aad, “Measurement of jet charge in dijet events from $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Phys. Rev. **D93**, 052003 (2016).
- [249] G. Aad, “Measurement of the angular coefficients in Z -boson events using electron and muon pairs from data taken at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 159 (2016).
- [250] G. Aad, “Measurement of the centrality dependence of the charged-particle pseudorapidity distribution in proton–lead collisions at $\sqrt{s_{NN}} = 5.02$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 199 (2016).
- [251] G. Aad, “Measurement of the charge asymmetry in highly boosted top-quark pair production in $\sqrt{s} = 8$ TeV pp collision data collected by the ATLAS experiment”, Phys. Lett. **B756**, 52–71 (2016).
- [252] G. Aad, “Measurement of the charge asymmetry in top-quark pair production in the lepton-plus-jets final state in pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, [Erratum: Eur. Phys. J. **C77**, 564 (2017)], 87 (2016).
- [253] G. Aad, “Measurement of the charged-particle multiplicity inside jets from $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C76**, 322 (2016).

- [254] G. Aad, “Measurement of the correlations between the polar angles of leptons from top quark decays in the helicity basis at $\sqrt{s} = 7\text{TeV}$ using the ATLAS detector”, Phys. Rev. **D93**, 012002 (2016).
- [255] G. Aad, “Measurement of the CP-violating phase ϕ_s and the B_s^0 meson decay width difference with $B_s^0 \rightarrow J/\psi\phi$ decays in ATLAS”, JHEP **08**, 147 (2016).
- [256] G. Aad, “Measurement of the dependence of transverse energy production at large pseudorapidity on the hard-scattering kinematics of proton-proton collisions at $\sqrt{s} = 2.76\text{ TeV}$ with ATLAS”, Phys. Lett. **B756**, 10–28 (2016).
- [257] G. Aad, “Measurement of the differential cross-section of highly boosted top quarks as a function of their transverse momentum in $\sqrt{s} = 8\text{ TeV}$ proton-proton collisions using the ATLAS detector”, Phys. Rev. **D93**, 032009 (2016).
- [258] G. Aad, “Measurement of the differential cross-sections of prompt and non-prompt production of J/ψ and $\psi(2S)$ in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 283 (2016).
- [259] G. Aad, “Measurement of the double-differential high-mass Drell-Yan cross section in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **08**, 009 (2016).
- [260] G. Aad, “Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **08**, 005 (2016).
- [261] G. Aad, “Measurement of the production cross-section of a single top quark in association with a W boson at 8 TeV with the ATLAS experiment”, JHEP **01**, 064 (2016).
- [262] G. Aad, “Measurement of the transverse momentum and ϕ_η^* distributions of Drell-Yan lepton pairs in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. **C76**, 291 (2016).
- [263] G. Aad, “Measurement of the ZZ Production Cross Section in pp Collisions at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS Detector”, Phys. Rev. Lett. **116**, 101801 (2016).
- [264] G. Aad, “Measurement of total and differential W^+W^- production cross sections in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector and limits on anomalous triple-gauge-boson couplings”, JHEP **09**, 029 (2016).
- [265] G. Aad, “Measurement of W^\pm and Z -boson production cross sections in pp collisions at $\sqrt{s} = 13\text{ TeV}$ with the ATLAS detector”, Phys. Lett. **B759**, 601–621 (2016).
- [266] G. Aad, “Measurements of fiducial cross-sections for $t\bar{t}$ production with one or two additional b-jets in pp collisions at $\sqrt{s} = 8\text{ TeV}$ using the ATLAS detector”, Eur. Phys. J. **C76**, 11 (2016).
- [267] G. Aad, “Measurements of four-lepton production in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Phys. Lett. **B753**, 552–572 (2016).
- [268] G. Aad, “Measurements of the charge asymmetry in top-quark pair production in the dilepton final state at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Phys. Rev. **D94**, 032006 (2016).
- [269] G. Aad, “Measurements of the Higgs boson production and decay rates and constraints on its couplings from a combined ATLAS and CMS analysis of the LHC pp collision data at $\sqrt{s} = 7$ and 8 TeV ”, JHEP **08**, 045 (2016).
- [270] G. Aad, “Measurements of the Higgs boson production and decay rates and coupling strengths using pp collision data at $\sqrt{s} = 7$ and 8 TeV in the ATLAS experiment”, Eur. Phys. J. **C76**, 6 (2016).
- [271] G. Aad, “Measurements of top-quark pair differential cross-sections in the lepton+jets channel in pp collisions at $\sqrt{s} = 8\text{ TeV}$ using the ATLAS detector”, Eur. Phys. J. **C76**, 538 (2016).
- [272] G. Aad, “Measurements of $W^\pm Z$ production cross sections in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector and limits on anomalous gauge boson self-couplings”, Phys. Rev. **D93**, 092004 (2016).
- [273] G. Aad, “Measurements of $Z\gamma$ and $Z\gamma\gamma$ production in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Phys. Rev. **D93**, 112002 (2016).
- [274] G. Aad, “Muon reconstruction performance of the ATLAS detector in proton-proton collision data at $\sqrt{s} = 13\text{ TeV}$ ”, Eur. Phys. J. **C76**, 292 (2016).
- [275] G. Aad, “Observation of Long-Range Elliptic Azimuthal Anisotropies in $\sqrt{s} = 13$ and 2.76 TeV pp Collisions with the ATLAS Detector”, Phys. Rev. Lett. **116**, 172301 (2016).
- [276] G. Aad, “Performance of b -Jet Identification in the ATLAS Experiment”, JINST **11**, P04008 (2016).
- [277] G. Aad, “Performance of pile-up mitigation techniques for jets in pp collisions at $\sqrt{s} = 8\text{ TeV}$ using the ATLAS detector”, Eur. Phys. J. **C76**, 581 (2016).
- [278] G. Aad, “Probing lepton flavour violation via neutrinoless $\tau \rightarrow 3\mu$ decays with the ATLAS detector”, Eur. Phys. J. **C76**, 232 (2016).
- [279] G. Aad, “Reconstruction of hadronic decay products of tau leptons with the ATLAS experiment”, Eur. Phys. J. **C76**, 295 (2016).

- [280] G. Aad, “Search for a high-mass Higgs boson decaying to a W boson pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **01**, 032 (2016).
- [281] G. Aad, “Search for an additional, heavy Higgs boson in the $H \rightarrow ZZ$ decay channel at $\sqrt{s} = 8$ TeV in pp collision data with the ATLAS detector”, Eur. Phys. J. **C76**, 45 (2016).
- [282] G. Aad, “Search for anomalous couplings in the Wtb vertex from the measurement of double differential angular decay rates of single top quarks produced in the t -channel with the ATLAS detector”, JHEP **04**, 023 (2016).
- [283] G. Aad, “Search for charged Higgs bosons in the $H^\pm \rightarrow tb$ decay channel in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, JHEP **03**, 127 (2016).
- [284] G. Aad, “Search for dark matter produced in association with a Higgs boson decaying to two bottom quarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D93**, 072007 (2016).
- [285] G. Aad, “Search for direct top squark pair production in final states with two tau leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 81 (2016).
- [286] G. Aad, “Search for flavour-changing neutral current top-quark decays to qZ in pp collision data collected with the ATLAS detector at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. **C76**, 12 (2016).
- [287] G. Aad, “Search for gluinos in events with an isolated lepton, jets and missing transverse momentum at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 565 (2016).
- [288] G. Aad, “Search for invisible decays of a Higgs boson using vector-boson fusion in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **01**, 172 (2016).
- [289] G. Aad, “Search for magnetic monopoles and stable particles with high electric charges in 8 TeV pp collisions with the ATLAS detector”, Phys. Rev. **D93**, 052009 (2016).
- [290] G. Aad, “Search for new phenomena in dijet mass and angular distributions from pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Lett. **B754**, 302–322 (2016).
- [291] G. Aad, “Search for new phenomena in events with at least three photons collected in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 210 (2016).
- [292] G. Aad, “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum with ATLAS using $\sqrt{s} = 13$ TeV proton-proton collisions”, Phys. Lett. **B757**, 334–355 (2016).
- [293] G. Aad, “Search for new phenomena with photon+jet events in proton-proton collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, JHEP **03**, 041 (2016).
- [294] G. Aad, “Search for pair production of gluinos decaying via stop and sbottom in events with b -jets and large missing transverse momentum in pp collisions at $\sqrt{s} = 13$ TeV with the ATLAS detector”, Phys. Rev. **D94**, 032003 (2016).
- [295] G. Aad, “Search for single production of a vector-like quark via a heavy gluon in the $4b$ final state with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. **B758**, 249–268 (2016).
- [296] G. Aad, “Search for single production of vector-like quarks decaying into Wb in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 442 (2016).
- [297] G. Aad, “Search for single top-quark production via flavour-changing neutral currents at 8 TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 55 (2016).
- [298] G. Aad, “Search for strong gravity in multijet final states produced in pp collisions at $\sqrt{s} = 13$ TeV using the ATLAS detector at the LHC”, JHEP **03**, 026 (2016).
- [299] G. Aad, “Search for supersymmetry at $\sqrt{s} = 13$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector”, Eur. Phys. J. **C76**, 259 (2016).
- [300] G. Aad, “Search for the electroweak production of supersymmetric particles in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Phys. Rev. **D93**, 052002 (2016).
- [301] G. Aad, “Search for the production of single vector-like and excited quarks in the Wt final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **02**, 110 (2016).
- [302] G. Aad, “Search for the Standard Model Higgs boson decaying into $b\bar{b}$ produced in association with top quarks decaying hadronically in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **05**, 160 (2016).
- [303] G. Aad, “Search for the Standard Model Higgs boson produced in association with a vector boson and decaying into a tau pair in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D93**, 092005 (2016).
- [304] G. Aad, “Searches for scalar leptoquarks in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C76**, 5 (2016).

- [305] G. Aad, “Study of the $B_c^+ \rightarrow J/\psi D_s^+$ and $B_c^+ \rightarrow J/\psi D_s^{*+}$ decays with the ATLAS detector”, Eur. Phys. J. **C76**, 4 (2016).
- [306] G. Aad, “Test of CP Invariance in vector-boson fusion production of the Higgs boson using the Optimal Observable method in the ditau decay channel with the ATLAS detector”, Eur. Phys. J. **C76**, 658 (2016).
- [307] G. Aad, “The performance of the jet trigger for the ATLAS detector during 2011 data taking”, Eur. Phys. J. **C76**, 526 (2016).
- [308] G. Aad, “Transverse momentum, rapidity, and centrality dependence of inclusive charged-particle production in $\sqrt{s_{NN}} = 5.02$ TeV $p + \text{Pb}$ collisions measured by the ATLAS experiment”, Phys. Lett. **B763**, 313–336 (2016).
- [309] M. Abolins, “The ATLAS Data Acquisition and High Level Trigger system”, JINST **11**, P06008 (2016).
- [310] G. Aad, “A search for high-mass resonances decaying to $\tau^+\tau^-$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **07**, 157 (2015).
- [311] G. Aad, “A search for $t\bar{t}$ resonances using lepton-plus-jets events in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 148 (2015).
- [312] G. Aad, “Analysis of events with b -jets and a pair of leptons of the same charge in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10**, 150 (2015).
- [313] G. Aad, “ATLAS Run 1 searches for direct pair production of third-generation squarks at the Large Hadron Collider”, Eur. Phys. J. **C75**, [Erratum: Eur. Phys. J. C76,no.3,153(2016)], 510 (2015).
- [314] G. Aad, “Centrality and rapidity dependence of inclusive jet production in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector”, Phys. Lett. **B748**, 392–413 (2015).
- [315] G. Aad, “Combined Measurement of the Higgs Boson Mass in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS and CMS Experiments”, Phys. Rev. Lett. **114**, 191803 (2015).
- [316] G. Aad, “Constraints on new phenomena via Higgs boson couplings and invisible decays with the ATLAS detector”, JHEP **11**, 206 (2015).
- [317] G. Aad, “Constraints on the off-shell Higgs boson signal strength in the high-mass ZZ and WW final states with the ATLAS detector”, Eur. Phys. J. **C75**, 335 (2015).
- [318] G. Aad, “Determination of spin and parity of the Higgs boson in the $WW^* \rightarrow e\nu\mu\nu$ decay channel with the ATLAS detector”, Eur. Phys. J. **C75**, 231 (2015).
- [319] G. Aad, “Determination of the ratio of b -quark fragmentation fractions f_s/f_d in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **115**, 262001 (2015).
- [320] G. Aad, “Determination of the top-quark pole mass using $t\bar{t} + 1$ -jet events collected with the ATLAS experiment in 7 TeV pp collisions”, JHEP **10**, 121 (2015).
- [321] G. Aad, “Differential top-antitop cross-section measurements as a function of observables constructed from final-state particles using pp collisions at $\sqrt{s} = 7$ TeV in the ATLAS detector”, JHEP **06**, 100 (2015).
- [322] G. Aad, “Evidence for the Higgs-boson Yukawa coupling to tau leptons with the ATLAS detector”, JHEP **04**, 117 (2015).
- [323] G. Aad, “Evidence of W Production in pp Collisions at $s=8\text{TeV}$ and Limits on Anomalous Quartic Gauge Couplings with the ATLAS Detector”, Phys. Rev. Lett. **115**, 031802 (2015).
- [324] G. Aad, “Identification and energy calibration of hadronically decaying tau leptons with the ATLAS experiment in pp collisions at $\sqrt{s}=8$ TeV”, Eur. Phys. J. **C75**, 303 (2015).
- [325] G. Aad, “Jet energy measurement and its systematic uncertainty in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 17 (2015).
- [326] G. Aad, “Measurement of charged-particle spectra in $\text{Pb}+\text{Pb}$ collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector at the LHC”, JHEP **09**, 050 (2015).
- [327] G. Aad, “Measurement of colour flow with the jet pull angle in $t\bar{t}$ events using the ATLAS detector at $\sqrt{s} = 8$ TeV”, Phys. Lett. **B750**, 475–493 (2015).
- [328] G. Aad, “Measurement of differential J/ψ production cross sections and forward-backward ratios in $p + \text{Pb}$ collisions with the ATLAS detector”, Phys. Rev. **C92**, 034904 (2015).
- [329] G. Aad, “Measurement of exclusive $\gamma\gamma \rightarrow \ell^+\ell^-$ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B749**, 242–261 (2015).
- [330] G. Aad, “Measurement of four-jet differential cross sections in $\sqrt{s} = 8$ TeV proton-proton collisions using the ATLAS detector”, JHEP **12**, 105 (2015).

- [331] G. Aad, “Measurement of Spin Correlation in Top-Antitop Quark Events and Search for Top Squark Pair Production in pp Collisions at $\sqrt{s} = 8$ TeV Using the ATLAS Detector”, Phys. Rev. Lett. **114**, 142001 (2015).
- [332] G. Aad, “Measurement of the branching ratio $\Gamma(\Lambda_b^0 \rightarrow \psi(2S)\Lambda^0)/\Gamma(\Lambda_b^0 \rightarrow J/\psi\Lambda^0)$ with the ATLAS detector”, Phys. Lett. **B751**, 63–80 (2015).
- [333] G. Aad, “Measurement of the charge asymmetry in dileptonic decays of top quark pairs in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **05**, 061 (2015).
- [334] G. Aad, “Measurement of the correlation between flow harmonics of different order in lead-lead collisions at $\sqrt{s_{NN}}=2.76$ TeV with the ATLAS detector”, Phys. Rev. **C92**, 034903 (2015).
- [335] G. Aad, “Measurement of the forward-backward asymmetry of electron and muon pair-production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **09**, 049 (2015).
- [336] G. Aad, “Measurement of the inclusive jet cross-section in proton-proton collisions at $\sqrt{s} = 7$ TeV using 4.5 fb¹ of data with the ATLAS detector”, JHEP **02**, [Erratum: JHEP09,141(2015)], 153 (2015).
- [337] G. Aad, “Measurement of the production and lepton charge asymmetry of W bosons in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 23 (2015).
- [338] G. Aad, “Measurement of the production of neighbouring jets in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B751**, 376–395 (2015).
- [339] G. Aad, “Measurement of the top pair production cross section in 8 TeV proton-proton collisions using kinematic information in the lepton+jets final state with ATLAS”, Phys. Rev. **D91**, 112013 (2015).
- [340] G. Aad, “Measurement of the top quark mass in the $t\bar{t} \rightarrow$ lepton+jets and $t\bar{t} \rightarrow$ dilepton channels using $\sqrt{s} = 7$ TeV ATLAS data”, Eur. Phys. J. **C75**, 330 (2015).
- [341] G. Aad, “Measurement of the top-quark mass in the fully hadronic decay channel from ATLAS data at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C75**, 158 (2015).
- [342] G. Aad, “Measurement of the transverse polarization of Λ and $\bar{\Lambda}$ hyperons produced in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. **D91**, 032004 (2015).
- [343] G. Aad, “Measurement of the $t\bar{t}$ production cross-section as a function of jet multiplicity and jet transverse momentum in 7 TeV proton-proton collisions with the ATLAS detector”, JHEP **01**, 020 (2015).
- [344] G. Aad, “Measurement of the $t\bar{t}W$ and $t\bar{t}Z$ production cross sections in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **11**, 172 (2015).
- [345] G. Aad, “Measurement of the $WW + WZ$ cross section and limits on anomalous triple gauge couplings using final states with one lepton, missing transverse momentum, and two jets with the ATLAS detector at $\sqrt{s} = 7$ TeV”, JHEP **01**, 049 (2015).
- [346] G. Aad, “Measurement of three-jet production cross-sections in pp collisions at 7 TeV centre-of-mass energy using the ATLAS detector”, Eur. Phys. J. **C75**, 228 (2015).
- [347] G. Aad, “Measurement of transverse energy-energy correlations in multi-jet events in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector and determination of the strong coupling constant $\alpha_s(m_Z)$ ”, Phys. Lett. **B750**, 427–447 (2015).
- [348] G. Aad, “Measurements of Higgs boson production and couplings in the four-lepton channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, Phys. Rev. **D91**, 012006 (2015).
- [349] G. Aad, “Measurements of the Nuclear Modification Factor for Jets in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **114**, 072302 (2015).
- [350] G. Aad, “Measurements of the top quark branching ratios into channels with leptons and quarks with the ATLAS detector”, Phys. Rev. **D92**, 072005 (2015).
- [351] G. Aad, “Measurements of the Total and Differential Higgs Boson Production Cross Sections Combining the $H \rightarrow$ and $H \rightarrow ZZ^* \rightarrow 4$ Decay Channels at $\sqrt{s}=8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 091801 (2015).
- [352] G. Aad, “Measurements of the W production cross sections in association with jets with the ATLAS detector”, Eur. Phys. J. **C75**, 82 (2015).
- [353] G. Aad, “Modelling $Z \rightarrow \tau\tau$ processes in ATLAS with τ -embedded $Z \rightarrow \mu\mu$ data”, JINST **10**, P09018 (2015).
- [354] G. Aad, “Observation and measurement of Higgs boson decays to WW^* with the ATLAS detector”, Phys. Rev. **D92**, 012006 (2015).

- [355] G. Aad, “Observation and measurements of the production of prompt and non-prompt J/ψ mesons in association with a Z boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 229 (2015).
- [356] G. Aad, “Observation of top-quark pair production in association with a photon and measurement of the $t\bar{t}\gamma$ production cross section in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. **D91**, 072007 (2015).
- [357] G. Aad, “Performance of the ATLAS muon trigger in pp collisions at $\sqrt{s} = 8$ TeV”, Eur. Phys. J. **C75**, 120 (2015).
- [358] G. Aad, “Search for a Charged Higgs Boson Produced in the Vector-Boson Fusion Mode with Decay $H^\pm \rightarrow W^\pm Z$ using pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Experiment”, Phys. Rev. Lett. **114**, 231801 (2015).
- [359] G. Aad, “Search for a CP-odd Higgs boson decaying to Zh in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B744**, 163–183 (2015).
- [360] G. Aad, “Search for a Heavy Neutral Particle Decaying to $e\mu$, $e\tau$, or $\mu\tau$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 031801 (2015).
- [361] G. Aad, “Search for a new resonance decaying to a W or Z boson and a Higgs boson in the $\ell\ell/\ell\nu/\nu\nu + b\bar{b}$ final states with the ATLAS detector”, Eur. Phys. J. **C75**, 263 (2015).
- [362] G. Aad, “Search for anomalous production of prompt same-sign lepton pairs and pair-produced doubly charged Higgs bosons with $\sqrt{s} = 8$ TeV pp collisions using the ATLAS detector”, JHEP **03**, 041 (2015).
- [363] G. Aad, “Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau^\pm\nu$ in fully hadronic final states using pp collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **03**, 088 (2015).
- [364] G. Aad, “Search for dark matter in events with heavy quarks and missing transverse momentum in pp collisions with the ATLAS detector”, Eur. Phys. J. **C75**, 92 (2015).
- [365] G. Aad, “Search for Dark Matter in Events with Missing Transverse Momentum and a Higgs Boson Decaying to Two Photons in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **115**, 131801 (2015).
- [366] G. Aad, “Search for direct pair production of a chargino and a neutralino decaying to the 125 GeV Higgs boson in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C75**, 208 (2015).
- [367] G. Aad, “Search for flavour-changing neutral current top quark decays $t \rightarrow Hq$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **12**, 061 (2015).
- [368] G. Aad, “Search for $H \rightarrow \gamma\gamma$ produced in association with top quarks and constraints on the Yukawa coupling between the top quark and the Higgs boson using data taken at 7 TeV and 8 TeV with the ATLAS detector”, Phys. Lett. **B740**, 222–242 (2015).
- [369] G. Aad, “Search for heavy lepton resonances decaying to a Z boson and a lepton in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **09**, 108 (2015).
- [370] G. Aad, “Search for heavy long-lived multi-charged particles in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, Eur. Phys. J. **C75**, 362 (2015).
- [371] G. Aad, “Search for heavy Majorana neutrinos with the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, JHEP **07**, 162 (2015).
- [372] G. Aad, “Search for Higgs and Z Boson Decays to J/ψ and (nS) with the ATLAS Detector”, Phys. Rev. Lett. **114**, 121801 (2015).
- [373] G. Aad, “Search For Higgs Boson Pair Production in the $\gamma\gamma b\bar{b}$ Final State using pp Collision Data at $\sqrt{s} = 8$ TeV from the ATLAS Detector”, Phys. Rev. Lett. **114**, 081802 (2015).
- [374] G. Aad, “Search for Higgs boson pair production in the $b\bar{b}b\bar{b}$ final state from pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 412 (2015).
- [375] G. Aad, “Search for Higgs bosons decaying to aa in the $\mu\mu\tau\tau$ final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS experiment”, Phys. Rev. **D92**, 052002 (2015).
- [376] G. Aad, “Search for high-mass diboson resonances with boson-tagged jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **12**, 055 (2015).
- [377] G. Aad, “Search for high-mass diphoton resonances in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D92**, 032004 (2015).
- [378] G. Aad, “Search for invisible decays of the Higgs boson produced in association with a hadronically decaying vector boson in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 337 (2015).
- [379] G. Aad, “Search for invisible particles produced in association with single-top-quarks in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 79 (2015).

- [380] G. Aad, “Search for lepton-flavour-violating $H \rightarrow$ decays of the Higgs boson with the ATLAS detector”, JHEP **11**, 211 (2015).
- [381] G. Aad, “Search for long-lived, weakly interacting particles that decay to displaced hadronic jets in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D92**, 012010 (2015).
- [382] G. Aad, “Search for low-scale gravity signatures in multi-jet final states with the ATLAS detector at $\sqrt{s} = 8$ TeV”, JHEP **07**, 032 (2015).
- [383] G. Aad, “Search for massive, long-lived particles using multitrack displaced vertices or displaced lepton pairs in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D92**, 072004 (2015).
- [384] G. Aad, “Search for massive supersymmetric particles decaying to many jets using the ATLAS detector in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Rev. **D91**, [Erratum: Phys. Rev.D93,no.3,039901(2016)], 112016 (2015).
- [385] G. Aad, “Search for metastable heavy charged particles with large ionisation energy loss in pp collisions at $\sqrt{s} = 8$ TeV using the ATLAS experiment”, Eur. Phys. J. **C75**, 407 (2015).
- [386] G. Aad, “Search for new light gauge bosons in Higgs boson decays to four-lepton final states in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector at the LHC”, Phys. Rev. **D92**, 092001 (2015).
- [387] G. Aad, “Search for New Phenomena in Dijet Angular Distributions in Proton-Proton Collisions at $\sqrt{s} = 8$ TeV Measured with the ATLAS Detector”, Phys. Rev. Lett. **114**, 221802 (2015).
- [388] G. Aad, “Search for new phenomena in events with a photon and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D91**, [Erratum: Phys. Rev.D92,no.5,059903(2015)], 012008 (2015).
- [389] G. Aad, “Search for new phenomena in events with three or more charged leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 138 (2015).
- [390] G. Aad, “Search for new phenomena in final states with an energetic jet and large missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, [Erratum: Eur. Phys. J.C75,no.9,408(2015)], 299 (2015).
- [391] G. Aad, “Search for new phenomena in the dijet mass distribution using $p - p$ collision data at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D91**, 052007 (2015).
- [392] G. Aad, “Search for pair production of a new heavy quark that decays into a W boson and a light quark in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D92**, 112007 (2015).
- [393] G. Aad, “Search for pair-produced long-lived neutral particles decaying in the ATLAS hadronic calorimeter in pp collisions at $\sqrt{s} = 8$ TeV”, Phys. Lett. **B743**, 15–34 (2015).
- [394] G. Aad, “Search for photonic signatures of gauge-mediated supersymmetry in 8 TeV pp collisions with the ATLAS detector”, Phys. Rev. **D92**, 072001 (2015).
- [395] G. Aad, “Search for production of vector-like quark pairs and of four top quarks in the lepton-plus-jets final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **08**, 105 (2015).
- [396] G. Aad, “Search for production of WW/WZ resonances decaying to a lepton, neutrino and jets in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, [Erratum: Eur. Phys. J.C75,370(2015)], 209 (2015).
- [397] G. Aad, “Search for resonant diboson production in the $\ell\ell q\bar{q}$ final state in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 69 (2015).
- [398] G. Aad, “Search for Scalar Charm Quark Pair Production in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **114**, 161801 (2015).
- [399] G. Aad, “Search for s -channel single top-quark production in proton-proton collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B740**, 118–136 (2015).
- [400] G. Aad, “Search for squarks and gluinos in events with isolated leptons, jets and missing transverse momentum at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **04**, 116 (2015).
- [401] G. Aad, “Search for supersymmetry in events containing a same-flavour opposite-sign dilepton pair, jets, and large missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C75**, [Erratum: Eur. Phys. J.C75,no.10,463(2015)], 318 (2015).
- [402] G. Aad, “Search for the associated production of the Higgs boson with a top quark pair in multilepton final states with the ATLAS detector”, Phys. Lett. **B749**, 519–541 (2015).
- [403] G. Aad, “Search for the $b\bar{b}$ decay of the Standard Model Higgs boson in associated $(W/Z)H$ production with the ATLAS detector”, JHEP **01**, 069 (2015).
- [404] G. Aad, “Search for the Standard Model Higgs boson produced in association with top quarks and decaying into $b\bar{b}$ in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 349 (2015).

- [405] G. Aad, “Search for the X_b and other hidden-beauty states in the $\pi^+\pi^-\Upsilon(1S)$ channel at ATLAS”, Phys. Lett. **B740**, 199–217 (2015).
- [406] G. Aad, “Search for type-III Seesaw heavy leptons in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. **D92**, 032001 (2015).
- [407] G. Aad, “Search for vector-like B quarks in events with one isolated lepton, missing transverse momentum and jets at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D91**, 112011 (2015).
- [408] G. Aad, “Search for $W' \rightarrow t\bar{b}$ in the lepton plus jets final state in proton-proton collisions at a centre-of-mass energy of $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B743**, 235–255 (2015).
- [409] G. Aad, “Search for $W' \rightarrow tb \rightarrow qqbb$ decays in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Eur. Phys. J. **C75**, 165 (2015).
- [410] G. Aad, “Searches for heavy long-lived charged particles with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 8$ TeV”, JHEP **01**, 068 (2015).
- [411] G. Aad, “Searches for Higgs boson pair production in the $hh \rightarrow bb\tau\tau, \gamma\gamma WW^*, \gamma\gamma bb, bbbb$ channels with the ATLAS detector”, Phys. Rev. **D92**, 092004 (2015).
- [412] G. Aad, “Simultaneous measurements of the $t\bar{t}$, W^+W^- , and $Z/\gamma^* \rightarrow \tau\tau$ production cross-sections in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D91**, 052005 (2015).
- [413] G. Aad, “Study of the spin and parity of the Higgs boson in diboson decays with the ATLAS detector”, Eur. Phys. J. **C75**, [Erratum: Eur. Phys. J. C76,no.3,152(2016)], 476 (2015).
- [414] G. Aad, “Study of (W/Z)H production and Higgs boson couplings using $H \rightarrow WW^*$ decays with the ATLAS detector”, JHEP **08**, 137 (2015).
- [415] G. Aad, “Summary of the ATLAS experiment’s sensitivity to supersymmetry after LHC Run 1 — interpreted in the phenomenological MSSM”, JHEP **10**, 134 (2015).
- [416] G. Aad, “Summary of the searches for squarks and gluinos using $\sqrt{s} = 8$ TeV pp collisions with the ATLAS experiment at the LHC”, JHEP **10**, 054 (2015).
- [417] G. Aad, “Two-particle Bose–Einstein correlations in pp collisions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector”, Eur. Phys. J. **C75**, 466 (2015).
- [418] G. Aad, “Z boson production in p +Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV measured with the ATLAS detector”, Phys. Rev. **C92**, 044915 (2015).
- [419] G. Volpi, “The ATLAS fast tracker processor design”, PoS **VERTEX2015**, 040 (2015).
- [420] G. Aad, “A measurement of the ratio of the production cross sections for W and Z bosons in association with jets with the ATLAS detector”, Eur. Phys. J. **C74**, 3168 (2014).
- [421] G. Aad, “A neural network clustering algorithm for the ATLAS silicon pixel detector”, JINST **9**, P09009 (2014).
- [422] G. Aad, “Comprehensive measurements of t -channel single top-quark production cross sections at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D90**, 112006 (2014).
- [423] G. Aad, “Electron and photon energy calibration with the ATLAS detector using LHC Run 1 data”, Eur. Phys. J. **C74**, 3071 (2014).
- [424] G. Aad, “Electron reconstruction and identification efficiency measurements with the ATLAS detector using the 2011 LHC proton-proton collision data”, Eur. Phys. J. **C74**, 2941 (2014).
- [425] G. Aad, “Evidence for Electroweak Production of $W^\pm W^\pm jj$ in pp Collisions at $\sqrt{s} = 8$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **113**, 141803 (2014).
- [426] G. Aad, “Fiducial and differential cross sections of Higgs boson production measured in the four-lepton decay channel in pp collisions at $\sqrt{s}=8$ TeV with the ATLAS detector”, Phys. Lett. **B738**, 234–253 (2014).
- [427] G. Aad, “Flavor tagged time-dependent angular analysis of the $B_s \rightarrow J/\psi\phi$ decay and extraction of $\Delta\Gamma$ s and the weak phase ϕ_s in ATLAS”, Phys. Rev. **D90**, 052007 (2014).
- [428] G. Aad, “Light-quark and gluon jet discrimination in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C74**, 3023 (2014).
- [429] G. Aad, “Measurement of χ_{c1} and χ_{c2} production with $\sqrt{s} = 7$ TeV pp collisions at ATLAS”, JHEP **07**, 154 (2014).
- [430] G. Aad, “Measurement of differential production cross-sections for a Z boson in association with b -jets in 7 TeV proton-proton collisions with the ATLAS detector”, JHEP **10**, 141 (2014).
- [431] G. Aad, “Measurement of dijet cross sections in pp collisions at 7 TeV centre-of-mass energy using the ATLAS detector”, JHEP **05**, 059 (2014).
- [432] G. Aad, “Measurement of distributions sensitive to the underlying event in inclusive Z-boson production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C74**, 3195 (2014).

- [433] G. Aad, “Measurement of event-plane correlations in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector”, Phys. Rev. **C90**, 024905 (2014).
- [434] G. Aad, “Measurement of flow harmonics with multi-particle cumulants in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. **C74**, 3157 (2014).
- [435] G. Aad, “Measurement of Higgs boson production in the diphoton decay channel in pp collisions at center-of-mass energies of 7 and 8 TeV with the ATLAS detector”, Phys. Rev. **D90**, 112015 (2014).
- [436] G. Aad, “Measurement of inclusive jet charged-particle fragmentation functions in Pb+Pb collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B739**, 320–342 (2014).
- [437] G. Aad, “Measurement of long-range pseudorapidity correlations and azimuthal harmonics in $\sqrt{s_{NN}} = 5.02$ TeV proton-lead collisions with the ATLAS detector”, Phys. Rev. **C90**, 044906 (2014).
- [438] G. Aad, “Measurement of the centrality and pseudorapidity dependence of the integrated elliptic flow in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Eur. Phys. J. **C74**, 2982 (2014).
- [439] G. Aad, “Measurement of the cross section of high transverse momentum $Z \rightarrow b\bar{b}$ production in proton–proton collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS Detector”, Phys. Lett. **B738**, 25–43 (2014).
- [440] G. Aad, “Measurement of the cross-section of high transverse momentum vector bosons reconstructed as single jets and studies of jet substructure in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, New J. Phys. **16**, 113013 (2014).
- [441] G. Aad, “Measurement of the electroweak production of dijets in association with a Z-boson and distributions sensitive to vector boson fusion in proton-proton collisions at $\sqrt{s} = 8$ TeV using the ATLAS detector”, JHEP **04**, 031 (2014).
- [442] G. Aad, “Measurement of the Higgs boson mass from the $H \rightarrow \gamma\gamma$ and $H \rightarrow ZZ^* \rightarrow 4\ell$ channels with the ATLAS detector using 25 fb^{-1} of pp collision data”, Phys. Rev. **D90**, 052004 (2014).
- [443] G. Aad, “Measurement of the inclusive isolated prompt photons cross section in pp collisions at $\sqrt{s} = 7\text{TeV}$ with the ATLAS detector using 4.6fb^{-1} ”, Phys. Rev. **D89**, 052004 (2014).
- [444] G. Aad, “Measurement of the low-mass Drell-Yan differential cross section at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **06**, 112 (2014).
- [445] G. Aad, “Measurement of the mass difference between top and anti-top quarks in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Lett. **B728**, 363–379 (2014).
- [446] G. Aad, “Measurement of the muon reconstruction performance of the ATLAS detector using 2011 and 2012 LHC proton–proton collision data”, Eur. Phys. J. **C74**, 3130 (2014).
- [447] G. Aad, “Measurement of the parity-violating asymmetry parameter α_b and the helicity amplitudes for the decay $\Lambda_b^0 \rightarrow J/\psi + \Lambda^0$ with the ATLAS detector”, Phys. Rev. **D89**, 092009 (2014).
- [448] G. Aad, “Measurement of the production cross section of prompt J/ψ mesons in association with a W^\pm boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **04**, 172 (2014).
- [449] G. Aad, “Measurement of the production cross-section of $\psi(2S) \rightarrow J/\psi(\rightarrow \mu^+\mu^-)\pi^+\pi^-$ in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS”, JHEP **09**, 079 (2014).
- [450] G. Aad, “Measurement of the production of a W boson in association with a charm quark in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **05**, 068 (2014).
- [451] G. Aad, “Measurement of the top quark pair production charge asymmetry in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **02**, 107 (2014).
- [452] G. Aad, “Measurement of the total cross section from elastic scattering in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. **B889**, 486–548 (2014).
- [453] G. Aad, “Measurement of the $t\bar{t}$ production cross-section using $e\mu$ events with b-tagged jets in pp collisions at $\sqrt{s} = 7$ and 8 TeV with the ATLAS detector”, Eur. Phys. J. **C74**, [Addendum: Eur. Phys. J.C76,no.11,642(2016)], 3109 (2014).
- [454] G. Aad, “Measurement of the underlying event in jet events from 7 TeV proton-proton collisions with the ATLAS detector”, Eur. Phys. J. **C74**, 2965 (2014).
- [455] G. Aad, “Measurement of the Z/γ^* boson transverse momentum distribution in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **09**, 145 (2014).
- [456] G. Aad, “Measurements of fiducial and differential cross sections for Higgs boson production in the diphoton decay channel at $\sqrt{s} = 8$ TeV with ATLAS”, JHEP **09**, 112 (2014).
- [457] G. Aad, “Measurements of Four-Lepton Production at the Z Resonance in pp Collisions at $\sqrt{s} = 7$ and 8 TeV with ATLAS”, Phys. Rev. Lett. **112**, 231806 (2014).
- [458] G. Aad, “Measurements of jet vetoes and azimuthal decorrelations in dijet events produced in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. **C74**, 3117 (2014).

- [459] G. Aad, “Measurements of normalized differential cross sections for $t\bar{t}$ production in pp collisions at $\sqrt{s} = 7\text{TeV}$ using the ATLAS detector”, Phys. Rev. **D90**, 072004 (2014).
- [460] G. Aad, “Measurements of spin correlation in top-antitop quark events from proton-proton collisions at $\sqrt{s} = 7\text{ TeV}$ using the ATLAS detector”, Phys. Rev. **D90**, 112016 (2014).
- [461] G. Aad, “Monitoring and data quality assessment of the ATLAS liquid argon calorimeter”, JINST **9**, P07024 (2014).
- [462] G. Aad, “Muon reconstruction efficiency and momentum resolution of the ATLAS experiment in proton-proton collisions at $\sqrt{s} = 7\text{ TeV}$ in 2010”, Eur. Phys. J. **C74**, 3034 (2014).
- [463] G. Aad, “Observation of an Excited B_c^\pm Meson State with the ATLAS Detector”, Phys. Rev. Lett. **113**, 212004 (2014).
- [464] G. Aad, “Operation and performance of the ATLAS semiconductor tracker”, JINST **9**, P08009 (2014).
- [465] G. Aad, “Search for a multi-Higgs-boson cascade in $W^+W^b\bar{b}$ events with the ATLAS detector in pp collisions at $\sqrt{s} = 8\text{TeV}$ ”, Phys. Rev. **D89**, 032002 (2014).
- [466] G. Aad, “Search for contact interactions and large extra dimensions in the dilepton channel using proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. **C74**, 3134 (2014).
- [467] G. Aad, “Search for dark matter in events with a hadronically decaying W or Z boson and missing transverse momentum in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Phys. Rev. Lett. **112**, 041802 (2014).
- [468] G. Aad, “Search for dark matter in events with a Z boson and missing transverse momentum in pp collisions at $\sqrt{s}=8\text{ TeV}$ with the ATLAS detector”, Phys. Rev. **D90**, 012004 (2014).
- [469] G. Aad, “Search for direct pair production of the top squark in all-hadronic final states in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **09**, 015 (2014).
- [470] G. Aad, “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 8\text{TeV}$ pp collisions with the ATLAS detector”, JHEP **04**, 169 (2014).
- [471] G. Aad, “Search for direct production of charginos, neutralinos and sleptons in final states with two leptons and missing transverse momentum in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **05**, 071 (2014).
- [472] G. Aad, “Search for direct top squark pair production in events with a Z boson, b-jets and missing transverse momentum in sqrt(s)=8 TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C74**, 2883 (2014).
- [473] G. Aad, “Search for direct top-squark pair production in final states with two leptons in pp collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS detector”, JHEP **06**, 124 (2014).
- [474] G. Aad, “Search for Higgs boson decays to a photon and a Z boson in pp collisions at $\sqrt{s}=7$ and 8 TeV with the ATLAS detector”, Phys. Lett. **B732**, 8–27 (2014).
- [475] G. Aad, “Search for high-mass dilepton resonances in pp collisions at $\sqrt{s} = 8\text{TeV}$ with the ATLAS detector”, Phys. Rev. **D90**, 052005 (2014).
- [476] G. Aad, “Search for Invisible Decays of a Higgs Boson Produced in Association with a Z Boson in ATLAS”, Phys. Rev. Lett. **112**, 201802 (2014).
- [477] G. Aad, “Search for long-lived neutral particles decaying into lepton jets in proton-proton collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **11**, 088 (2014).
- [478] G. Aad, “Search for microscopic black holes and string balls in final states with leptons and jets with the ATLAS detector at sqrt(s) = 8 TeV”, JHEP **08**, 103 (2014).
- [479] G. Aad, “Search for neutral Higgs bosons of the minimal supersymmetric standard model in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **11**, 056 (2014).
- [480] G. Aad, “Search for new particles in events with one lepton and missing transverse momentum in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **09**, 037 (2014).
- [481] G. Aad, “Search for new phenomena in photon+jet events collected in proton–proton collisions at sqrt(s) = 8 TeV with the ATLAS detector”, Phys. Lett. **B728**, 562–578 (2014).
- [482] G. Aad, “Search for new resonances in $W\gamma$ and $Z\gamma$ final states in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, Phys. Lett. **B738**, 428–447 (2014).
- [483] G. Aad, “Search for nonpointing and delayed photons in the diphoton and missing transverse momentum final state in 8 TeV pp collisions at the LHC using the ATLAS detector”, Phys. Rev. **D90**, 112005 (2014).
- [484] G. Aad, “Search for pair and single production of new heavy quarks that decay to a Z boson and a third-generation quark in pp collisions at $\sqrt{s} = 8\text{ TeV}$ with the ATLAS detector”, JHEP **11**, 104 (2014).

- [485] G. Aad, “Search for pair-produced third-generation squarks decaying via charm quarks or in compressed supersymmetric scenarios in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Rev. **D90**, 052008 (2014).
- [486] G. Aad, “Search for Quantum Black Hole Production in High-Invariant-Mass Lepton+Jet Final States Using pp Collisions at $\sqrt{s} = 8$ TeV and the ATLAS Detector”, Phys. Rev. Lett. **112**, 091804 (2014).
- [487] G. Aad, “Search for Scalar Diphoton Resonances in the Mass Range 65 – 600 GeV with the ATLAS Detector in pp Collision Data at $\sqrt{s} = 8$ TeV”, Phys. Rev. Lett. **113**, 171801 (2014).
- [488] G. Aad, “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using $\sqrt{s} = 8$ TeV proton–proton collision data”, JHEP **09**, 176 (2014).
- [489] G. Aad, “Search for strong production of supersymmetric particles in final states with missing transverse momentum and at least three b -jets at $\sqrt{s} = 8$ TeV proton-proton collisions with the ATLAS detector”, JHEP **10**, 024 (2014).
- [490] G. Aad, “Search for supersymmetry at $\sqrt{s} = 8$ TeV in final states with jets and two same-sign leptons or three leptons with the ATLAS detector”, JHEP **06**, 035 (2014).
- [491] G. Aad, “Search for supersymmetry in events with four or more leptons in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, Phys. Rev. **D90**, 052001 (2014).
- [492] G. Aad, “Search for supersymmetry in events with large missing transverse momentum, jets, and at least one tau lepton in 20 fb⁻¹ of $\sqrt{s} = 8$ TeV proton-proton collision data with the ATLAS detector”, JHEP **09**, 103 (2014).
- [493] G. Aad, “Search for the direct production of charginos, neutralinos and staus in final states with at least two hadronically decaying taus and missing transverse momentum in pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, JHEP **10**, 096 (2014).
- [494] G. Aad, “Search for the lepton flavor violating decay $Z \rightarrow e$ in pp collisions at \sqrt{s} TeV with the ATLAS detector”, Phys. Rev. **D90**, 072010 (2014).
- [495] G. Aad, “Search for the Standard Model Higgs boson decay to $\mu^+ \mu^-$ with the ATLAS detector”, Phys. Lett. **B738**, 68–86 (2014).
- [496] G. Aad, “Search for top quark decays $t \rightarrow qH$ with $H \rightarrow \gamma\gamma$ using the ATLAS detector”, JHEP **06**, 008 (2014).
- [497] G. Aad, “Search for top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 8$ TeV pp collisions with the ATLAS detector”, JHEP **11**, 118 (2014).
- [498] G. Aad, “Search for WZ resonances in the fully leptonic channel using pp collisions at $\sqrt{s} = 8$ TeV with the ATLAS detector”, Phys. Lett. **B737**, 223–243 (2014).
- [499] G. Aad, “Standalone vertex finding in the ATLAS muon spectrometer”, JINST **9**, P02001 (2014).
- [500] G. Aad, “Study of heavy-flavor quarks produced in association with top-quark pairs at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. **D89**, 072012 (2014).
- [501] G. Aad, “The differential production cross section of the ϕ (1020) meson in $\sqrt{s} = 7$ TeV pp collisions measured with the ATLAS detector”, Eur. Phys. J. **C74**, 2895 (2014).
- [502] G. Aad, “A search for high-mass resonances decaying to $\tau^+ \tau^-$ in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B719**, 242–260 (2013).
- [503] G. Aad, “A search for prompt lepton-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B719**, 299–317 (2013).
- [504] G. Aad, “ATLAS search for new phenomena in dijet mass and angular distributions using pp collisions at $\sqrt{s} = 7$ TeV”, JHEP **01**, 029 (2013).
- [505] G. Aad, “Characterisation and mitigation of beam-induced backgrounds observed in the ATLAS detector during the 2011 proton-proton run”, JINST **8**, P07004 (2013).
- [506] G. Aad, “Dynamics of isolated-photon plus jet production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. **B875**, 483–535 (2013).
- [507] G. Aad, “Evidence for the spin-0 nature of the Higgs boson using ATLAS data”, Phys. Lett. **B726**, 120–144 (2013).
- [508] G. Aad, “Improved luminosity determination in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC”, Eur. Phys. J. **C73**, 2518 (2013).
- [509] G. Aad, “Jet energy measurement with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C73**, 2304 (2013).
- [510] G. Aad, “Jet energy resolution in proton-proton collisions at $\sqrt{s} = 7$ TeV recorded in 2010 with the ATLAS detector”, Eur. Phys. J. **C73**, 2306 (2013).

- [511] G. Aad, “Measurement of angular correlations in Drell-Yan lepton pairs to probe Z/γ^* boson transverse momentum at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. **B720**, 32–51 (2013).
- [512] G. Aad, “Measurement of charged-particle event shape variables in $\sqrt{s} = 7$ TeV proton-proton interactions with the ATLAS detector”, Phys. Rev. **D88**, 032004 (2013).
- [513] G. Aad, “Measurement of hard double-parton interactions in $W(\rightarrow l\nu) + 2$ jet events at $\sqrt{s}=7$ TeV with the ATLAS detector”, New J. Phys. **15**, 033038 (2013).
- [514] G. Aad, “Measurement of isolated-photon pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **01**, 086 (2013).
- [515] G. Aad, “Measurement of jet shapes in top-quark pair events at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. **C73**, 2676 (2013).
- [516] G. Aad, “Measurement of k_T splitting scales in $W \rightarrow l\nu$ events at $\sqrt{s}=7$ TeV with the ATLAS detector”, Eur. Phys. J. **C73**, 2432 (2013).
- [517] G. Aad, “Measurement of the Azimuthal Angle Dependence of Inclusive Jet Yields in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Rev. Lett. **111**, 152301 (2013).
- [518] G. Aad, “Measurement of the cross-section for W boson production in association with b-jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **06**, 084 (2013).
- [519] G. Aad, “Measurement of the differential cross-section of B^+ meson production in pp collisions at $\sqrt{s} = 7$ TeV at ATLAS”, JHEP **10**, 042 (2013).
- [520] G. Aad, “Measurement of the distributions of event-by-event flow harmonics in lead-lead collisions at $= 2.76$ TeV with the ATLAS detector at the LHC”, JHEP **11**, 183 (2013).
- [521] G. Aad, “Measurement of the flavour composition of dijet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C73**, 2301 (2013).
- [522] G. Aad, “Measurement of the high-mass Drell-Yan differential cross-section in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. **B725**, 223–242 (2013).
- [523] G. Aad, “Measurement of the inclusive jet cross section in pp collisions at $\sqrt{s}=2.76$ TeV and comparison to the inclusive jet cross section at $\sqrt{s}=7$ TeV using the ATLAS detector”, Eur. Phys. J. **C73**, 2509 (2013).
- [524] G. Aad, “Measurement of the jet radius and transverse momentum dependence of inclusive jet suppression in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B719**, 220–241 (2013).
- [525] G. Aad, “Measurement of the production cross section of jets in association with a Z boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **07**, 032 (2013).
- [526] G. Aad, “Measurement of the top quark charge in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **11**, 031 (2013).
- [527] G. Aad, “Measurement of the $t\bar{t}$ production cross section in the tau+jets channel using the ATLAS detector”, Eur. Phys. J. **C73**, 2328 (2013).
- [528] G. Aad, “Measurement of the Λ_b^0 lifetime and mass in the ATLAS experiment”, Phys. Rev. **D87**, 032002 (2013).
- [529] G. Aad, “Measurement of Top Quark Polarization in Top-Antitop Events from Proton-Proton Collisions at $\sqrt{s} = 7$ TeV Using the ATLAS Detector”, Phys. Rev. Lett. **111**, 232002 (2013).
- [530] G. Aad, “Measurement of Upsilon production in 7 TeV pp collisions at ATLAS”, Phys. Rev. **D87**, 052004 (2013).
- [531] G. Aad, “Measurement of W^+W^- production in pp collisions at $\sqrt{s}=7$ TeV with the ATLAS detector and limits on anomalous WWZ and WW couplings”, Phys. Rev. **D87**, [Erratum: Phys. Rev.D88,no.7,079906(2013)], 112001 (2013).
- [532] G. Aad, “Measurement of Z boson Production in Pb+Pb Collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **110**, 022301 (2013).
- [533] G. Aad, “Measurement of ZZ production in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous ZZZ and $ZZ\gamma$ couplings with the ATLAS detector”, JHEP **03**, 128 (2013).
- [534] G. Aad, “Measurement with the ATLAS detector of multi-particle azimuthal correlations in p+Pb collisions at $\sqrt{s_{NN}} = 5.02$ TeV”, Phys. Lett. **B725**, 60–78 (2013).
- [535] G. Aad, “Measurements of Higgs boson production and couplings in diboson final states with the ATLAS detector at the LHC”, Phys. Lett. **B726**, [Erratum: Phys. Lett.B734,406(2014)], 88–119 (2013).
- [536] G. Aad, “Measurements of top quark pair relative differential cross-sections with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C73**, 2261 (2013).

- [537] G. Aad, “Measurements of $W\gamma$ and $Z\gamma$ production in pp collisions at $\sqrt{s}=7\text{TeV}$ with the ATLAS detector at the LHC”, Phys. Rev. **D87**, [Erratum: Phys. Rev.D91,no.11,119901(2015)], 112003 (2013).
- [538] G. Aad, “Multi-channel search for squarks and gluinos in $\sqrt{s} = 7 \text{ TeV}$ pp collisions with the ATLAS detector”, Eur. Phys. J. **C73**, 2362 (2013).
- [539] G. Aad, “Observation of Associated Near-Side and Away-Side Long-Range Correlations in $\sqrt{s_{NN}}=5.02\text{TeV}$ Proton-Lead Collisions with the ATLAS Detector”, Phys. Rev. Lett. **110**, 182302 (2013).
- [540] G. Aad, “Performance of jet substructure techniques for large- R jets in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$ using the ATLAS detector”, JHEP **09**, 076 (2013).
- [541] G. Aad, “Search for a heavy narrow resonance decaying to $e\mu$, $e\tau$, or $\mu\tau$ with the ATLAS detector in $\sqrt{s} = 7 \text{ TeV}$ pp collisions at the LHC”, Phys. Lett. **B723**, 15–32 (2013).
- [542] G. Aad, “Search for a light charged Higgs boson in the decay channel $H^+ \rightarrow c\bar{s}$ in $t\bar{t}$ events using pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Eur. Phys. J. **C73**, 2465 (2013).
- [543] G. Aad, “Search for charged Higgs bosons through the violation of lepton universality in $t\bar{t}$ events using pp collision data at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS experiment”, JHEP **03**, 076 (2013).
- [544] G. Aad, “Search for charginos nearly mass degenerate with the lightest neutralino based on a disappearing-track signature in pp collisions at $\sqrt{s}=8\text{TeV}$ with the ATLAS detector”, Phys. Rev. **D88**, 112006 (2013).
- [545] G. Aad, “Search for contact interactions and large extra dimensions in dilepton events from pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. **D87**, 015010 (2013).
- [546] G. Aad, “Search for dark matter candidates and large extra dimensions in events with a jet and missing transverse momentum with the ATLAS detector”, JHEP **04**, 075 (2013).
- [547] G. Aad, “Search for dark matter candidates and large extra dimensions in events with a photon and missing transverse momentum in pp collision data at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. Lett. **110**, 011802 (2013).
- [548] G. Aad, “Search for direct chargino production in anomaly-mediated supersymmetry breaking models based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, JHEP **01**, 131 (2013).
- [549] G. Aad, “Search for direct production of charginos and neutralinos in events with three leptons and missing transverse momentum in $\sqrt{s} = 7 \text{ TeV}$ pp collisions with the ATLAS detector”, Phys. Lett. **B718**, 841–859 (2013).
- [550] G. Aad, “Search for direct slepton and gaugino production in final states with two leptons and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7 \text{ TeV}$ ”, Phys. Lett. **B718**, 879–901 (2013).
- [551] G. Aad, “Search for direct third-generation squark pair production in final states with missing transverse momentum and two b -jets in $\sqrt{s} = 8 \text{ TeV}$ pp collisions with the ATLAS detector”, JHEP **10**, 189 (2013).
- [552] G. Aad, “Search for displaced muonic lepton jets from light Higgs boson decay in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. **B721**, 32–50 (2013).
- [553] G. Aad, “Search for excited electrons and muons in $\sqrt{s}=8 \text{ TeV}$ proton-proton collisions with the ATLAS detector”, New J. Phys. **15**, 093011 (2013).
- [554] G. Aad, “Search for Extra Dimensions in diphoton events using proton-proton collisions recorded at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector at the LHC”, New J. Phys. **15**, 043007 (2013).
- [555] G. Aad, “Search for light top squark pair production in final states with leptons and b^- jets with the ATLAS detector in $\sqrt{s} = 7 \text{ TeV}$ proton-proton collisions”, Phys. Lett. **B720**, 13–31 (2013).
- [556] G. Aad, “Search for long-lived, heavy particles in final states with a muon and multi-track displaced vertex in proton-proton collisions at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Lett. **B719**, 280–298 (2013).
- [557] G. Aad, “Search for long-lived, multi-charged particles in pp collisions at $\sqrt{s}=7 \text{ TeV}$ using the ATLAS detector”, Phys. Lett. **B722**, 305–323 (2013).
- [558] G. Aad, “Search for long-lived stopped R-hadrons decaying out-of-time with pp collisions using the ATLAS detector”, Phys. Rev. **D88**, 112003 (2013).
- [559] G. Aad, “Search for microscopic black holes in a like-sign dimuon final state using large track multiplicity with the ATLAS detector”, Phys. Rev. **D88**, 072001 (2013).
- [560] G. Aad, “Search for new phenomena in events with three charged leptons at $\sqrt{s} = 7 \text{ TeV}$ with the ATLAS detector”, Phys. Rev. **D87**, 052002 (2013).

- [561] G. Aad, “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum at $\sqrt{s}=8$ TeV proton-proton collisions using the ATLAS experiment”, JHEP **10**, [Erratum: JHEP01,109(2014)], 130 (2013).
- [562] G. Aad, “Search for new phenomena in the WW to $\ell\nu\ell'\nu'$ final state in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B718**, 860–878 (2013).
- [563] G. Aad, “Search for nonpointing photons in the diphoton and E_T^{miss} final state in $\sqrt{s}=7$ TeV proton-proton collisions using the ATLAS detector”, Phys. Rev. **D88**, 012001 (2013).
- [564] G. Aad, “Search for pair production of heavy top-like quarks decaying to a high-pT W boson and a b quark in the lepton plus jets final state at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. **B718**, 1284–1302 (2013).
- [565] G. Aad, “Search for pair-produced massive coloured scalars in four-jet final states with the ATLAS detector in proton-proton collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C73**, 2263 (2013).
- [566] G. Aad, “Search for resonances decaying into top-quark pairs using fully hadronic decays in pp collisions with ATLAS at $\sqrt{s} = 7$ TeV”, JHEP **01**, 116 (2013).
- [567] G. Aad, “Search for resonant diboson production in the $WW/WZ \rightarrow jj$ decay channels with the ATLAS detector at $\sqrt{s} = 7$ TeV”, Phys. Rev. **D87**, 112006 (2013).
- [568] G. Aad, “Search for single b^* -quark production with the ATLAS detector at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B721**, 171–189 (2013).
- [569] G. Aad, “Search for squarks and gluinos with the ATLAS detector in final states with jets and missing transverse momentum using 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collision data”, Phys. Rev. **D87**, 012008 (2013).
- [570] G. Aad, “Search for supersymmetry in events with photons, bottom quarks, and missing transverse momentum in proton-proton collisions at a centre-of-mass energy of 7 TeV with the ATLAS detector”, Phys. Lett. **B719**, 261–279 (2013).
- [571] G. Aad, “Search for $t\bar{t}$ resonances in the lepton plus jets final state with ATLAS using 4.7 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. **D88**, 012004 (2013).
- [572] G. Aad, “Search for the neutral Higgs bosons of the Minimal Supersymmetric Standard Model in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **02**, 095 (2013).
- [573] G. Aad, “Search for third generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **06**, 033 (2013).
- [574] G. Aad, “Search for WH production with a light Higgs boson decaying to prompt electron-jets in proton-proton collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, New J. Phys. **15**, 043009 (2013).
- [575] G. Aad, “Searches for heavy long-lived sleptons and R-Hadrons with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B720**, 277–308 (2013).
- [576] G. Aad, “Single hadron response measurement and calorimeter jet energy scale uncertainty with the ATLAS detector at the LHC”, Eur. Phys. J. **C73**, 2305 (2013).
- [577] G. Aad, “Technical Design Report for the Phase-I Upgrade of the ATLAS TDAQ System”, (2013).
- [578] G. Aad, “Triggers for displaced decays of long-lived neutral particles in the ATLAS detector”, JINST **8**, P07015 (2013).
- [579] T. Aaltonen, “Measurement of the B_c^- meson lifetime in the decay $B_c^- \rightarrow J/\psi \pi^-$ ”, Phys. Rev. **D87**, 011101 (2013).
- [580] T. Aaltonen, “Measurement of the mass difference between top and antitop quarks”, Phys. Rev. **D87**, 052013 (2013).
- [581] T. Aaltonen, “Observation of the Production of a W Boson in Association with a Single Charm Quark”, Phys. Rev. Lett. **110**, 071801 (2013).
- [582] T. Aaltonen, “Production of $K_S^0, K^{*\pm}(892)$ and $\phi^0(1020)$ in minimum bias events and K_S^0 and Λ^0 in jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96 \text{ TeV}$ ”, Phys. Rev. **D88**, 092005 (2013).
- [583] T. Aaltonen, “Search for a Two-Higgs-Boson Doublet Using a Simplified Model in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96 \text{ TeV}$ ”, Phys. Rev. Lett. **110**, 121801 (2013).
- [584] T. Aaltonen, “Search for the Higgs boson in the all-hadronic final state using the full CDF data set”, JHEP **02**, 004 (2013).
- [585] T. Aaltonen, “Searches for the Higgs boson decaying to $W^+W^- \rightarrow \ell^+\nu\ell^-\bar{\nu}$ with the CDF II detector”, Phys. Rev. **D88**, 052012 (2013).
- [586] T. Aaltonen, “ W boson polarization measurement in the $t\bar{t}$ dilepton channel using the CDF II Detector”, Phys. Lett. **B722**, 48–54 (2013).

- [587] J. Anderson, “A fast hardware tracker for the ATLAS trigger system”, Nucl. Instrum. Meth. **A718**, 258–259 (2013).
- [588] G. Aad, “A measurement of the ratio of the W and Z cross sections with exactly one associated jet in pp collisions at $\sqrt{s} = 7$ TeV with ATLAS”, Phys. Lett. **B708**, 221–240 (2012).
- [589] G. Aad, “A particle consistent with the Higgs Boson observed with the ATLAS Detector at the Large Hadron Collider”, Science **338**, 1576–1582 (2012).
- [590] G. Aad, “A search for flavour changing neutral currents in top-quark decays in pp collision data collected with the ATLAS detector at $\sqrt{s} = 7$ TeV”, JHEP **09**, 139 (2012).
- [591] G. Aad, “A search for $t\bar{t}$ resonances in lepton+jets events with highly boosted top quarks collected in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **09**, 041 (2012).
- [592] G. Aad, “A search for $t\bar{t}$ resonances with the ATLAS detector in 2.05 fb^{-1} of proton-proton collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 2083 (2012).
- [593] G. Aad, “A study of the material in the ATLAS inner detector using secondary hadronic interactions”, JINST **7**, P01013 (2012).
- [594] G. Aad, “ATLAS Measurements of the Properties of Jets for Boosted Particle Searches”, Phys. Rev. **D86**, 072006 (2012).
- [595] G. Aad, “ATLAS search for a heavy gauge boson decaying to a charged lepton and a neutrino in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 2241 (2012).
- [596] G. Aad, “Combined search for the Standard Model Higgs boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D86**, 032003 (2012).
- [597] G. Aad, “Combined search for the Standard Model Higgs boson using up to 4.9 fb^{-1} of pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector at the LHC”, Phys. Lett. **B710**, 49–66 (2012).
- [598] G. Aad, “Determination of the strange quark density of the proton from ATLAS measurements of the $W \rightarrow \ell\nu$ and $Z \rightarrow \ell\ell$ cross sections”, Phys. Rev. Lett. **109**, 012001 (2012).
- [599] G. Aad, “Electron performance measurements with the ATLAS detector using the 2010 LHC proton-proton collision data”, Eur. Phys. J. **C72**, 1909 (2012).
- [600] G. Aad, “Evidence for the associated production of a W boson and a top quark in ATLAS at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B716**, 142–159 (2012).
- [601] G. Aad, “Forward-backward correlations and charged-particle azimuthal distributions in pp interactions using the ATLAS detector”, JHEP **07**, 019 (2012).
- [602] G. Aad, “Further search for supersymmetry at $\sqrt{s} = 7$ TeV in final states with jets, missing transverse momentum and isolated leptons with the ATLAS detector”, Phys. Rev. **D86**, 092002 (2012).
- [603] G. Aad, “Hunt for new phenomena using large jet multiplicities and missing transverse momentum with ATLAS in 4.7 fb^{-1} of $\sqrt{s} = 7$ TeV proton-proton collisions”, JHEP **07**, 167 (2012).
- [604] G. Aad, “Jet mass and substructure of inclusive jets in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS experiment”, JHEP **05**, 128 (2012).
- [605] G. Aad, “K-short and Λ production in pp interactions at $\sqrt{s} = 0.9$ and 7 TeV measured with the ATLAS detector at the LHC”, Phys. Rev. **D85**, 012001 (2012).
- [606] G. Aad, “Measurement of $D^{*+/-}$ meson production in jets from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 052005 (2012).
- [607] G. Aad, “Measurement of event shapes at large momentum transfer with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 2211 (2012).
- [608] G. Aad, “Measurement of inclusive jet and dijet production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. **D86**, 014022 (2012).
- [609] G. Aad, “Measurement of inclusive two-particle angular correlations in pp collisions with the ATLAS detector at the LHC”, JHEP **05**, 157 (2012).
- [610] G. Aad, “Measurement of τ polarization in $W^- \rightarrow \tau\nu$ decays with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 2062 (2012).
- [611] G. Aad, “Measurement of $t\bar{t}$ production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. **C72**, 2043 (2012).
- [612] G. Aad, “Measurement of the azimuthal anisotropy for charged particle production in $\sqrt{s_{NN}} = 2.76$ TeV lead-lead collisions with the ATLAS detector”, Phys. Rev. **C86**, 014907 (2012).
- [613] G. Aad, “Measurement of the azimuthal ordering of charged hadrons with the ATLAS detector”, Phys. Rev. **D86**, 052005 (2012).
- [614] G. Aad, “Measurement of the b-hadron production cross section using decays to $D^*\mu^- X$ final states in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Nucl. Phys. **B864**, 341–381 (2012).

- [615] G. Aad, “Measurement of the centrality dependence of the charged particle pseudorapidity distribution in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B710**, 363–382 (2012).
- [616] G. Aad, “Measurement of the charge asymmetry in top quark pair production in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Eur. Phys. J. **C72**, 2039 (2012).
- [617] G. Aad, “Measurement of the cross section for the production of a W boson in association with b^- jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B707**, 418–437 (2012).
- [618] G. Aad, “Measurement of the cross section for top-quark pair production in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector using final states with two high-pt leptons”, JHEP **05**, 059 (2012).
- [619] G. Aad, “Measurement of the cross-section for b -jets produced in association with a Z boson at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B706**, 295–313 (2012).
- [620] G. Aad, “Measurement of the inclusive W^\pm and Z/γ cross sections in the electron and muon decay channels in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 072004 (2012).
- [621] G. Aad, “Measurement of the isolated di-photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 012003 (2012).
- [622] G. Aad, “Measurement of the polarisation of W bosons produced with large transverse momentum in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Eur. Phys. J. **C72**, 2001 (2012).
- [623] G. Aad, “Measurement of the production cross section for Z/γ in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 032009 (2012).
- [624] G. Aad, “Measurement of the production cross section of an isolated photon associated with jets in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 092014 (2012).
- [625] G. Aad, “Measurement of the pseudorapidity and transverse momentum dependence of the elliptic flow of charged particles in lead-lead collisions at $\sqrt{s_{NN}} = 2.76$ TeV with the ATLAS detector”, Phys. Lett. **B707**, 330–348 (2012).
- [626] G. Aad, “Measurement of the t -channel single top-quark production cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B717**, 330–350 (2012).
- [627] G. Aad, “Measurement of the top quark mass with the template method in the $t\bar{t} \rightarrow$ lepton + jets channel using ATLAS data”, Eur. Phys. J. **C72**, 2046 (2012).
- [628] G. Aad, “Measurement of the top quark pair cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV using final states with an electron or a muon and a hadronically decaying τ lepton”, Phys. Lett. **B717**, 89–108 (2012).
- [629] G. Aad, “Measurement of the top quark pair production cross section in pp collisions at $\sqrt{s} = 7$ TeV in dilepton final states with ATLAS”, Phys. Lett. **B707**, 459–477 (2012).
- [630] G. Aad, “Measurement of the top quark pair production cross-section with ATLAS in the single lepton channel”, Phys. Lett. **B711**, 244–263 (2012).
- [631] G. Aad, “Measurement of the Transverse Momentum Distribution of W Bosons in pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Phys. Rev. **D85**, 012005 (2012).
- [632] G. Aad, “Measurement of the W boson polarization in top quark decays with the ATLAS detector”, JHEP **06**, 088 (2012).
- [633] G. Aad, “Measurement of the $W \rightarrow \tau\nu_\tau$ cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Phys. Lett. **B706**, 276–294 (2012).
- [634] G. Aad, “Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector and limits on anomalous gauge couplings”, Phys. Lett. **B712**, 289–308 (2012).
- [635] G. Aad, “Measurement of the $W^\pm Z$ production cross section and limits on anomalous triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B709**, 341–357 (2012).
- [636] G. Aad, “Measurement of the ZZ production cross section and limits on anomalous neutral triple gauge couplings in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 041804 (2012).
- [637] G. Aad, “Measurement of $W\gamma$ and $Z\gamma$ production cross sections in pp collisions at $\sqrt{s} = 7$ TeV and limits on anomalous triple gauge couplings with the ATLAS detector”, Phys. Lett. **B717**, 49–69 (2012).
- [638] G. Aad, “Measurement of WZ production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C72**, 2173 (2012).
- [639] G. Aad, “Measurements of the electron and muon inclusive cross-sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B707**, 438–458 (2012).

- [640] G. Aad, “Measurements of the pseudorapidity dependence of the total transverse energy in proton-proton collisions at $\sqrt{s} = 7$ TeV with ATLAS”, JHEP **11**, 033 (2012).
- [641] G. Aad, “Observation of a new χ_b state in radiative transitions to $\Upsilon(1S)$ and $\Upsilon(2S)$ at ATLAS”, Phys. Rev. Lett. **108**, 152001 (2012).
- [642] G. Aad, “Observation of a new particle in the search for the Standard Model Higgs boson with the ATLAS detector at the LHC”, Phys. Lett. **B716**, 1–29 (2012).
- [643] G. Aad, “Observation of spin correlation in $t\bar{t}$ events from pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Rev. Lett. **108**, 212001 (2012).
- [644] G. Aad, “Performance of Missing Transverse Momentum Reconstruction in Proton-Proton Collisions at 7 TeV with ATLAS”, Eur. Phys. J. **C72**, 1844 (2012).
- [645] G. Aad, “Performance of the ATLAS Trigger System in 2010”, Eur. Phys. J. **C72**, 1849 (2012).
- [646] G. Aad, “Rapidity gap cross sections measured with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 1926 (2012).
- [647] G. Aad, “Search for a fermiophobic Higgs boson in the diphoton decay channel with the ATLAS detector”, Eur. Phys. J. **C72**, 2157 (2012).
- [648] G. Aad, “Search for a heavy Standard Model Higgs boson in the channel $H \rightarrow ZZ \rightarrow \ell^+ \ell'^q \bar{q}$ using the ATLAS detector”, Phys. Lett. **B707**, 27–45 (2012).
- [649] G. Aad, “Search for a heavy top-quark partner in final states with two leptons with the ATLAS detector at the LHC”, JHEP **11**, 094 (2012).
- [650] G. Aad, “Search for a light Higgs boson decaying to long-lived weakly-interacting particles in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 251801 (2012).
- [651] G. Aad, “Search for a standard model Higgs boson in the mass range 200–600 GeV in the $H \rightarrow ZZ \rightarrow \ell^+ \ell^- q \bar{q}$ decay channel with the ATLAS detector”, Phys. Lett. **B717**, 70–88 (2012).
- [652] G. Aad, “Search for a standard model Higgs boson in the $H \rightarrow ZZ \rightarrow \ell^+ \ell^- \nu \bar{\nu}$ decay channel using 4.7 fb $^{-1}$ of $\sqrt{s} = 7$ TeV data with the ATLAS detector”, Phys. Lett. **B717**, 29–48 (2012).
- [653] G. Aad, “Search for a supersymmetric partner to the top quark in final states with jets and missing transverse momentum at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **109**, 211802 (2012).
- [654] G. Aad, “Search for anomalous production of prompt like-sign lepton pairs at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **12**, 007 (2012).
- [655] G. Aad, “Search for anomalous production of prompt like-sign muon pairs and constraints on physics beyond the Standard Model with the ATLAS detector”, Phys. Rev. **D85**, 032004 (2012).
- [656] G. Aad, “Search for anomaly-mediated supersymmetry breaking with the ATLAS detector based on a disappearing-track signature in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C72**, 1993 (2012).
- [657] G. Aad, “Search for charged Higgs bosons decaying via $H^\pm \rightarrow \tau \nu$ in top quark pair events using pp collision data at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **06**, 039 (2012).
- [658] G. Aad, “Search for contact interactions in dilepton events from pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B712**, 40–58 (2012).
- [659] G. Aad, “Search for decays of stopped, long-lived particles from 7 TeV pp collisions with the ATLAS detector”, Eur. Phys. J. **C72**, 1965 (2012).
- [660] G. Aad, “Search for diphoton events with large missing transverse momentum in 7 TeV proton-proton collision data with the ATLAS detector”, Phys. Lett. **B718**, 411–430 (2012).
- [661] G. Aad, “Search for diphoton events with large missing transverse momentum in fb $^{-1}$ of 7 TeV proton-proton collision data with the ATLAS detector”, Phys. Lett. **B710**, 519–537 (2012).
- [662] G. Aad, “Search for direct top squark pair production in final states with one isolated lepton, jets, and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions using 4.7 fb $^{-1}$ of ATLAS data”, Phys. Rev. Lett. **109**, 211803 (2012).
- [663] G. Aad, “Search for displaced vertices arising from decays of new heavy particles in 7 TeV pp collisions at ATLAS”, Phys. Lett. **B707**, 478–496 (2012).
- [664] G. Aad, “Search for doubly-charged Higgs bosons in like-sign dilepton final states at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C72**, 2244 (2012).
- [665] G. Aad, “Search for down-type fourth generation quarks with the ATLAS detector in events with one lepton and hadronically decaying W bosons”, Phys. Rev. Lett. **109**, 032001 (2012).
- [666] G. Aad, “Search for events with large missing transverse momentum, jets, and at least two tau leptons in 7 TeV proton-proton collision data with the ATLAS detector”, Phys. Lett. **B714**, 180–196 (2012).

- [667] G. Aad, “Search for excited leptons in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 072003 (2012).
- [668] G. Aad, “Search for extra dimensions using diphoton events in 7 TeV proton-proton collisions with the ATLAS detector”, Phys. Lett. **B710**, 538–556 (2012).
- [669] G. Aad, “Search for FCNC single top-quark production at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B712**, 351–369 (2012).
- [670] G. Aad, “Search for first generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B709**, [Erratum: Phys. Lett. **B711**, 442(2012)], 158–176 (2012).
- [671] G. Aad, “Search for gluinos in events with two same-sign leptons, jets and missing transverse momentum with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. Lett. **108**, 241802 (2012).
- [672] G. Aad, “Search for heavy neutrinos and right-handed W bosons in events with two leptons and jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C72**, 2056 (2012).
- [673] G. Aad, “Search for heavy vector-like quarks coupling to light quarks in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B712**, 22–39 (2012).
- [674] G. Aad, “Search for high-mass resonances decaying to dilepton final states in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **11**, 138 (2012).
- [675] G. Aad, “Search for lepton flavour violation in the $e\mu$ continuum with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, Eur. Phys. J. **C72**, 2040 (2012).
- [676] G. Aad, “Search for light scalar top quark pair production in final states with two leptons with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Eur. Phys. J. **C72**, 2237 (2012).
- [677] G. Aad, “Search for magnetic monopoles in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Rev. Lett. **109**, 261803 (2012).
- [678] G. Aad, “Search for new particles decaying to ZZ using final states with leptons and jets with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. **B712**, 331–350 (2012).
- [679] G. Aad, “Search for New Phenomena in $t\bar{t}$ Events With Large Missing Transverse Momentum in Proton-Proton Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Phys. Rev. Lett. **108**, 041805 (2012).
- [680] G. Aad, “Search for New Physics in the Dijet Mass Distribution using 1 fb⁻¹ of pp Collision Data at $\sqrt{s} = 7$ TeV collected by the ATLAS Detector”, Phys. Lett. **B708**, 37–54 (2012).
- [681] G. Aad, “Search for pair production of a heavy up-type quark decaying to a W boson and a b quark in the lepton+jets channel with the ATLAS detector”, Phys. Rev. Lett. **108**, 261802 (2012).
- [682] G. Aad, “Search for pair production of a new quark that decays to a Z boson and a bottom quark with the ATLAS detector”, Phys. Rev. Lett. **109**, 071801 (2012).
- [683] G. Aad, “Search for pair production of massive particles decaying into three quarks with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, JHEP **12**, 086 (2012).
- [684] G. Aad, “Search for pair-produced heavy quarks decaying to Wq in the two-lepton channel at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D86**, 012007 (2012).
- [685] G. Aad, “Search for production of resonant states in the photon-jet mass distribution using pp collisions at $\sqrt{s} = 7$ TeV collected by the ATLAS detector”, Phys. Rev. Lett. **108**, 211802 (2012).
- [686] G. Aad, “Search for resonant top plus jet production in $t\bar{t} + \text{jets}$ events with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. **D86**, 091103 (2012).
- [687] G. Aad, “Search for resonant WZ production in the $WZ \rightarrow \ell\nu\ell'\ell'$ channel in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Rev. **D85**, 112012 (2012).
- [688] G. Aad, “Search for R-parity-violating supersymmetry in events with four or more leptons in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, JHEP **12**, 124 (2012).
- [689] G. Aad, “Search for same-sign top-quark production and fourth-generation down-type quarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **04**, 069 (2012).
- [690] G. Aad, “Search for scalar bottom quark pair production with the ATLAS detector in pp Collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. Lett. **108**, 181802 (2012).
- [691] G. Aad, “Search for scalar top quark pair production in natural gauge mediated supersymmetry models with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B715**, 44–60 (2012).
- [692] G. Aad, “Search for second generation scalar leptoquarks in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C72**, 2151 (2012).
- [693] G. Aad, “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. **B710**, 67–85 (2012).

- [694] G. Aad, “Search for strong gravity signatures in same-sign dimuon final states using the ATLAS detector at the LHC”, Phys. Lett. **B709**, 322–340 (2012).
- [695] G. Aad, “Search for Supersymmetry in Events with Large Missing Transverse Momentum, Jets, and at Least One Tau Lepton in 7 TeV Proton-Proton Collision Data with the ATLAS Detector”, Eur. Phys. J. **C72**, 2215 (2012).
- [696] G. Aad, “Search for supersymmetry in events with three leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, Phys. Rev. Lett. **108**, 261804 (2012).
- [697] G. Aad, “Search for supersymmetry in final states with jets, missing transverse momentum and one isolated lepton in $\sqrt{s} = 7$ TeV pp collisions using 1 fb^{-1} of ATLAS data”, Phys. Rev. **D85**, [Erratum: Phys. Rev. **D87**, 099903(2013)], 012006 (2012).
- [698] G. Aad, “Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b^- jets with the ATLAS detector”, Phys. Rev. **D85**, 112006 (2012).
- [699] G. Aad, “Search for supersymmetry with jets, missing transverse momentum and at least one hadronically decaying τ lepton in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B714**, 197–214 (2012).
- [700] G. Aad, “Search for tb resonances in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **109**, 081801 (2012).
- [701] G. Aad, “Search for TeV-scale gravity signatures in final states with leptons and jets with the ATLAS detector at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B716**, 122–141 (2012).
- [702] G. Aad, “Search for the decay $B_s^0 \rightarrow \mu\mu$ with the ATLAS detector”, Phys. Lett. **B713**, 387–407 (2012).
- [703] G. Aad, “Search for the Higgs boson in the $H \rightarrow WW \rightarrow l\nu jj$ decay channel at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B718**, 391–410 (2012).
- [704] G. Aad, “Search for the Higgs boson in the $H \rightarrow WW(*) \rightarrow \ell_\nu \ell_\nu$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **108**, 111802 (2012).
- [705] G. Aad, “Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ(*) \rightarrow 4\ell$ with 4.8 fb^{-1} of pp collision data at $\sqrt{s} = 7$ TeV with ATLAS”, Phys. Lett. **B710**, 383–402 (2012).
- [706] G. Aad, “Search for the Standard Model Higgs boson in the diphoton decay channel with 4.9 fb^{-1} of pp collisions at $\sqrt{s} = 7$ TeV with ATLAS”, Phys. Rev. Lett. **108**, 111803 (2012).
- [707] G. Aad, “Search for the Standard Model Higgs boson in the H to $\tau^+\tau^-$ decay mode in $\sqrt{s} = 7$ TeV pp collisions with ATLAS”, JHEP **09**, 070 (2012).
- [708] G. Aad, “Search for the Standard Model Higgs boson in the $H \rightarrow WW(*) \rightarrow \ell\nu\ell\nu$ decay mode with 4.7 $/fb$ of ATLAS data at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B716**, 62–81 (2012).
- [709] G. Aad, “Search for the Standard Model Higgs boson produced in association with a vector boson and decaying to a b -quark pair with the ATLAS detector”, Phys. Lett. **B718**, 369–390 (2012).
- [710] G. Aad, “Search for top and bottom squarks from gluino pair production in final states with missing transverse energy and at least three b -jets with the ATLAS detector”, Eur. Phys. J. **C72**, 2174 (2012).
- [711] G. Aad, “Searches for supersymmetry with the ATLAS detector using final states with two leptons and missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. **B709**, 137–157 (2012).
- [712] G. Aad, “Study of jets produced in association with a W boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D85**, 092002 (2012).
- [713] G. Aad, “Time-dependent angular analysis of the decay $B_s^0 \rightarrow J/\psi\phi$ and extraction of $\Delta\Gamma_s$ and the CP-violating weak phase ϕ_s by ATLAS”, JHEP **12**, 072 (2012).
- [714] G. Aad, “Underlying event characteristics and their dependence on jet size of charged-particle jet events in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D86**, 072004 (2012).
- [715] T. Aaltonen, “A Search for dark matter in events with one jet and missing transverse energy in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 211804 (2012).
- [716] T. Aaltonen, “An additional study of multi-muon events produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. **B710**, 278–283 (2012).
- [717] T. Aaltonen, “Combination of CDF and D0 measurements of the W boson helicity in top quark decays”, Phys. Rev. **D85**, 071106 (2012).
- [718] T. Aaltonen, “Combination of the top-quark mass measurements from the Tevatron collider”, Phys. Rev. **D86**, 092003 (2012).
- [719] T. Aaltonen, “Combined search for the standard model Higgs boson decaying to a $b\bar{b}$ pair using the full CDF data set”, Phys. Rev. Lett. **109**, 111802 (2012).

- [720] T. Aaltonen, “Diffractive Dijet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D86**, 032009 (2012).
- [721] T. Aaltonen, “Evidence for a particle produced in association with weak bosons and decaying to a bottom-antibottom quark pair in Higgs boson searches at the Tevatron”, Phys. Rev. Lett. **109**, 071804 (2012).
- [722] T. Aaltonen, “Evidence for the charmless annihilation decay mode $B_s^0 \rightarrow \pi^+\pi^-$ ”, Phys. Rev. Lett. **108**, 211803 (2012).
- [723] T. Aaltonen, “Measurement of $B_s^0 \rightarrow D_s^{(*)+}D_s^{(*)-}$ Branching Ratios”, Phys. Rev. Lett. **108**, 201801 (2012).
- [724] T. Aaltonen, “Measurement of CP-violating asymmetries in $D^0 \rightarrow \pi^+\pi^-$ and $D^0 \rightarrow K^+K^-$ decays at CDF”, Phys. Rev. **D85**, 012009 (2012).
- [725] T. Aaltonen, “Measurement of CP-violation asymmetries in $D^0 \rightarrow K_S\pi^+\pi^-$ ”, Phys. Rev. **D86**, 032007 (2012).
- [726] T. Aaltonen, “Measurement of the Bottom-Strange Meson Mixing Phase in the Full CDF Data Set”, Phys. Rev. Lett. **109**, 171802 (2012).
- [727] T. Aaltonen, “Measurement of the branching fraction $\mathcal{B}(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-\pi^+\pi^-)$ at CDF”, Phys. Rev. **D85**, 032003 (2012).
- [728] T. Aaltonen, “Measurement of the CP-Violating Phase $\beta_s^{J/\psi\phi}$ in $B_s^0 \rightarrow J/\psi\phi$ Decays with the CDF II Detector”, Phys. Rev. **D85**, 072002 (2012).
- [729] T. Aaltonen, “Measurement of the difference of CP-violating asymmetries in $D^0 \rightarrow K^+K^-$ and $D^0 \rightarrow \pi^+\pi^-$ decays at CDF”, Phys. Rev. Lett. **109**, 111801 (2012).
- [730] T. Aaltonen, “Measurement of the masses and widths of the bottom baryons Σ_b^{+-} and Σ_b^{*+-} ”, Phys. Rev. **D85**, 092011 (2012).
- [731] T. Aaltonen, “Measurement of the Top Quark Mass in the All-Hadronic Mode at CDF”, Phys. Lett. **B714**, 24–31 (2012).
- [732] T. Aaltonen, “Measurement of the WZ Cross Section and Triple Gauge Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D86**, 031104 (2012).
- [733] T. Aaltonen, “Measurement of ZZ production in leptonic final states at \sqrt{s} of 1.96 TeV at CDF”, Phys. Rev. Lett. **108**, 101801 (2012).
- [734] T. Aaltonen, “Measurements of Angular Distributions of Muons From Υ Meson Decays in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 151802 (2012).
- [735] T. Aaltonen, “Measurements of the Angular Distributions in the Decays $B \rightarrow K^{(*)}\mu^+\mu^-$ at CDF”, Phys. Rev. Lett. **108**, 081807 (2012).
- [736] T. Aaltonen, “Measurements of the Top-quark Mass and the $t\bar{t}$ Cross Section in the Hadronic τ + Jets Decay Channel at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **109**, 192001 (2012).
- [737] T. Aaltonen, “Novel inclusive search for the Higgs boson in the four-lepton final state at CDF”, Phys. Rev. **D86**, 072012 (2012).
- [738] T. Aaltonen, “Observation of Exclusive Gamma Gamma Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 081801 (2012).
- [739] T. Aaltonen, “Precise measurement of the W -boson mass with the CDF II detector”, Phys. Rev. Lett. **108**, 151803 (2012).
- [740] T. Aaltonen, “Precision Top-Quark Mass Measurements at CDF”, Phys. Rev. Lett. **109**, 152003 (2012).
- [741] T. Aaltonen, “Production of Λ , $\bar{\Lambda}^0$, Ξ^\pm and Ω^\pm Hyperons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D86**, 012002 (2012).
- [742] T. Aaltonen, “Search for a dark matter candidate produced in association with a single top quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 201802 (2012).
- [743] T. Aaltonen, “Search for a heavy particle decaying to a top quark and a light quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 211805 (2012).
- [744] T. Aaltonen, “Search for a Heavy Vector Boson Decaying to Two Gluons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D86**, 112002 (2012).
- [745] T. Aaltonen, “Search for a Higgs Boson in the Diphoton Final State in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 011801 (2012).
- [746] T. Aaltonen, “Search for a Higgs boson in the diphoton final state using the full CDF data set from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. **B717**, 173–181 (2012).
- [747] T. Aaltonen, “Search for a low mass Standard Model Higgs boson in the $\tau - \tau$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **108**, 181804 (2012).

- [748] T. Aaltonen, “Search for anomalous production of multiple leptons in association with W and Z bosons at CDF”, Phys. Rev. **D85**, 092001 (2012).
- [749] T. Aaltonen, “Search for heavy metastable particles decaying to jet pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 012007 (2012).
- [750] T. Aaltonen, “Search for Higgs Bosons Produced in Association with b -quarks”, Phys. Rev. **D85**, 032005 (2012).
- [751] T. Aaltonen, “Search for high-mass resonances decaying into ZZ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 012008 (2012).
- [752] T. Aaltonen, “Search for Neutral Higgs Bosons in Events with Multiple Bottom Quarks at the Tevatron”, Phys. Rev. **D86**, 091101 (2012).
- [753] T. Aaltonen, “Search for new phenomena in events with two Z bosons and missing transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 011104 (2012).
- [754] T. Aaltonen, “Search for Scalar Top Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, JHEP **10**, 158 (2012).
- [755] T. Aaltonen, “Search for Standard Model Higgs Boson Production in Association with a W Boson at CDF”, Phys. Rev. **D85**, 052002 (2012).
- [756] T. Aaltonen, “Search for Standard Model Higgs Boson Production in Association with a W Boson Using a Matrix Element Technique at CDF in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 072001 (2012).
- [757] T. Aaltonen, “Search for the Rare Radiative Decay: $W \rightarrow \pi\gamma$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 032001 (2012).
- [758] T. Aaltonen, “Search for the standard model Higgs boson decaying to a $b\bar{b}$ pair in events with no charged leptons and large missing transverse energy using the full CDF data set”, Phys. Rev. Lett. **109**, 111805 (2012).
- [759] T. Aaltonen, “Search for the standard model Higgs boson decaying to a bb pair in events with one charged lepton and large missing transverse energy using the full CDF data set”, Phys. Rev. Lett. **109**, 111804 (2012).
- [760] T. Aaltonen, “Search for the standard model Higgs boson decaying to a bb pair in events with two oppositely-charged leptons using the full CDF data set”, Phys. Rev. Lett. **109**, 111803 (2012).
- [761] T. Aaltonen, “Search for the standard model Higgs boson produced in association with a W^\pm boson with 7.5 fb^{-1} integrated luminosity at CDF”, Phys. Rev. **D86**, 032011 (2012).
- [762] T. Aaltonen, “Search for the standard model Higgs boson produced in association with a Z Boson in 7.9 fb^{-1} of p anti- p collisions at $\sqrt{s} = 1.96$ TeV using the CDF II detector”, Phys. Lett. **B715**, 98–104 (2012).
- [763] T. Aaltonen, “Search for the standard model Higgs boson produced in association with top quarks using the full CDF data set”, Phys. Rev. Lett. **109**, 181802 (2012).
- [764] T. Aaltonen, “Search for $WZ + ZZ$ production with MET + jets with b enhancement at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 012002 (2012).
- [765] T. Aaltonen, “Study of Substructure of High Transverse Momentum Jets Produced in Proton-Antiproton Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D85**, 091101 (2012).
- [766] T. Aaltonen, “Transverse momentum cross section of e^+e^- pairs in the Z -boson region from $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D86**, 052010 (2012).
- [767] J. Anderson, “FTK: A fast track trigger for ATLAS”, JINST **7**, C10002 (2012).
- [768] A. Andreani, “The Fast Tracker real time processor and its impact on muon isolation, tau and b-Jet online selections at ATLAS”, IEEE Trans. Nucl. Sci. **59**, 348–357 (2012).
- [769] G. Aad, “Charged-particle multiplicities in pp interactions measured with the ATLAS detector at the LHC”, New J. Phys. **13**, 053033 (2011).
- [770] G. Aad, “Measurement of inclusive jet and dijet cross sections in proton-proton collisions at 7 TeV centre-of-mass energy with the ATLAS detector”, Eur. Phys. J. **C71**, 1512 (2011).
- [771] G. Aad, “Measurement of the centrality dependence of J/ψ yields and observation of Z production in lead-lead collisions with the ATLAS detector at the LHC”, Phys. Lett. **B697**, 294–312 (2011).
- [772] G. Aad, “Measurement of the inclusive isolated prompt photon cross section in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D83**, 052005 (2011).
- [773] G. Aad, “Search for quark contact interactions in dijet angular distributions in pp collisions at $\sqrt{s} = 7$ TeV measured with the ATLAS detector”, Phys. Lett. **B694**, 327–345 (2011).

- [774] G. Aad, “Studies of the performance of the ATLAS detector using cosmic-ray muons”, Eur. Phys. J. **C71**, 1593 (2011).
- [775] G. Aad, “Study of Jet Shapes in Inclusive Jet Production in pp Collisions at $\sqrt{s} = 7$ TeV using the ATLAS Detector”, Phys. Rev. **D83**, 052003 (2011).
- [776] G. Aad, “Inclusive search for same-sign dilepton signatures in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **10**, 107 (2011).
- [777] G. Aad, “Limits on the production of the Standard Model Higgs Boson in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C71**, 1728 (2011).
- [778] G. Aad, “Luminosity Determination in pp Collisions at $\sqrt{s} = 7$ TeV Using the ATLAS Detector at the LHC”, Eur. Phys. J. **C71**, 1630 (2011).
- [779] G. Aad, “Measurement of Dijet Azimuthal Decorrelations in pp Collisions at $\sqrt{s} = 7$ TeV”, Phys. Rev. Lett. **106**, 172002 (2011).
- [780] G. Aad, “Measurement of dijet production with a veto on additional central jet activity in pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, JHEP **09**, 053 (2011).
- [781] G. Aad, “Measurement of multi-jet cross sections in proton-proton collisions at a 7 TeV center-of-mass energy”, Eur. Phys. J. **C71**, 1763 (2011).
- [782] G. Aad, “Measurement of the $\Upsilon(1S)$ production cross-section in pp collisions at $\sqrt{s} = 7$ TeV in ATLAS”, Phys. Lett. **B705**, 9–27 (2011).
- [783] G. Aad, “Measurement of the differential cross-sections of inclusive, prompt and non-prompt J/ψ production in proton-proton collisions at $\sqrt{s} = 7$ TeV”, Nucl. Phys. **B850**, 387–444 (2011).
- [784] G. Aad, “Measurement of the inclusive and dijet cross-sections of b^- jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Eur. Phys. J. **C71**, 1846 (2011).
- [785] G. Aad, “Measurement of the inclusive isolated prompt photon cross-section in pp collisions at $\sqrt{s} = 7$ TeV using 35 pb⁻¹ of ATLAS data”, Phys. Lett. **B706**, 150–167 (2011).
- [786] G. Aad, “Measurement of the Inelastic Proton-Proton Cross-Section at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Nature Commun. **2**, 463 (2011).
- [787] G. Aad, “Measurement of the jet fragmentation function and transverse profile in proton-proton collisions at a center-of-mass energy of 7 TeV with the ATLAS detector”, Eur. Phys. J. **C71**, 1795 (2011).
- [788] G. Aad, “Measurement of the production cross section for W-bosons in association with jets in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B698**, 325–345 (2011).
- [789] G. Aad, “Measurement of the top quark-pair production cross section with ATLAS in pp collisions at $\sqrt{s} = 7$ TeV”, Eur. Phys. J. **C71**, 1577 (2011).
- [790] G. Aad, “Measurement of the transverse momentum distribution of Z/* bosons in proton-proton collisions at $\sqrt{s}=7$ TeV with the ATLAS detector”, Phys. Lett. **B705**, 415–434 (2011).
- [791] G. Aad, “Measurement of the W charge asymmetry in the $W \rightarrow \mu\nu$ decay mode in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B701**, 31–49 (2011).
- [792] G. Aad, “Measurement of the WW cross section in $\sqrt{s} = 7$ TeV pp collisions with ATLAS”, Phys. Rev. Lett. **107**, 041802 (2011).
- [793] G. Aad, “Measurement of the Z to tau tau Cross Section with the ATLAS Detector”, Phys. Rev. **D84**, 112006 (2011).
- [794] G. Aad, “Measurement of underlying event characteristics using charged particles in pp collisions at $\sqrt{s} = 900\text{GeV}$ and 7 TeV with the ATLAS detector”, Phys. Rev. **D83**, 112001 (2011).
- [795] G. Aad, “Measurement of Wgamma and Zgamma production in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, JHEP **09**, 072 (2011).
- [796] G. Aad, “Measurements of underlying-event properties using neutral and charged particles in pp collisions at 900 GeV and 7 TeV with the ATLAS detector at the LHC”, Eur. Phys. J. **C71**, 1636 (2011).
- [797] G. Aad, “Properties of jets measured from tracks in proton-proton collisions at center-of-mass energy $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. **D84**, 054001 (2011).
- [798] G. Aad, “Search for a heavy gauge boson decaying to a charged lepton and a neutrino in 1 fb⁻¹ of pp collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector”, Phys. Lett. **B705**, 28–46 (2011).
- [799] G. Aad, “Search for a heavy neutral particle decaying into an electron and a muon using 1 fb⁻¹ of ATLAS data”, Eur. Phys. J. **C71**, 1809 (2011).
- [800] G. Aad, “Search for a heavy particle decaying into an electron and a muon with the ATLAS detector in $\sqrt{s} = 7$ TeV pp collisions at the LHC”, Phys. Rev. Lett. **106**, 251801 (2011).

- [801] G. Aad, “Search for a Standard Model Higgs boson in the $H \rightarrow ZZ \rightarrow l^+l^- \nu \nu^-$ decay channel with the ATLAS detector”, Phys. Rev. Lett. **107**, 221802 (2011).
- [802] G. Aad, “Search for an excess of events with an identical flavour lepton pair and significant missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS detector”, Eur. Phys. J. **C71**, 1647 (2011).
- [803] G. Aad, “Search for Contact Interactions in Dimuon Events from pp Collisions at $\sqrt{s} = 7$ TeV with the ATLAS Detector”, Phys. Rev. **D84**, 011101 (2011).
- [804] G. Aad, “Search for dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **107**, 272002 (2011).
- [805] G. Aad, “Search for Diphoton Events with Large Missing Transverse Energy in 7 TeV Proton-Proton Collisions with the ATLAS Detector”, Phys. Rev. Lett. **106**, 121803 (2011).
- [806] G. Aad, “Search for Diphoton Events with Large Missing Transverse Energy with 36 pb⁻¹ of 7 TeV Proton-Proton Collision Data with the ATLAS Detector”, Eur. Phys. J. **C71**, 1744 (2011).
- [807] G. Aad, “Search for heavy long-lived charged particles with the ATLAS detector in pp collisions at $\sqrt{s} = 7$ TeV”, Phys. Lett. **B703**, 428–446 (2011).
- [808] G. Aad, “Search for high mass dilepton resonances in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS experiment”, Phys. Lett. **B700**, 163–180 (2011).
- [809] G. Aad, “Search for high-mass states with one lepton plus missing transverse momentum in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B701**, 50–69 (2011).
- [810] G. Aad, “Search for Massive Colored Scalars in Four-Jet Final States in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS Detector”, Eur. Phys. J. **C71**, 1828 (2011).
- [811] G. Aad, “Search for Massive Long-lived Highly Ionising Particles with the ATLAS Detector at the LHC”, Phys. Lett. **B698**, 353–370 (2011).
- [812] G. Aad, “Search for neutral MSSM Higgs bosons decaying to $\tau^+\tau^-$ pairs in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Lett. **B705**, 174–192 (2011).
- [813] G. Aad, “Search for new phenomena in final states with large jet multiplicities and missing transverse momentum using $\sqrt{s} = 7$ TeV pp collisions with the ATLAS detector”, JHEP **11**, 099 (2011).
- [814] G. Aad, “Search for new phenomena with the monojet and missing transverse momentum signature using the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. **B705**, 294–312 (2011).
- [815] G. Aad, “Search for New Physics in Dijet Mass and Angular Distributions in pp Collisions at $\sqrt{s} = 7$ TeV Measured with the ATLAS Detector”, New J. Phys. **13**, 053044 (2011).
- [816] G. Aad, “Search for pair production of first or second generation leptoquarks in proton-proton collisions at $\sqrt{s} = 7$ TeV using the ATLAS detector at the LHC”, Phys. Rev. **D83**, 112006 (2011).
- [817] G. Aad, “Search for squarks and gluinos using final states with jets and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV proton-proton collisions”, Phys. Lett. **B701**, 186–203 (2011).
- [818] G. Aad, “Search for stable hadronising squarks and gluinos with the ATLAS experiment at the LHC”, Phys. Lett. **B701**, 1–19 (2011).
- [819] G. Aad, “Search for supersymmetric particles in events with lepton pairs and large missing transverse momentum in $\sqrt{s} = 7$ TeV proton-proton collisions with the ATLAS experiment”, Eur. Phys. J. **C71**, 1682 (2011).
- [820] G. Aad, “Search for supersymmetry in pp collisions at $\sqrt{s} = 7$ TeV in final states with missing transverse momentum and b -jets”, Phys. Lett. **B701**, 398–416 (2011).
- [821] G. Aad, “Search for supersymmetry using final states with one lepton, jets, and missing transverse momentum with the ATLAS detector in $\sqrt{s} = 7$ TeV pp ”, Phys. Rev. Lett. **106**, 131802 (2011).
- [822] G. Aad, “Search for the Higgs boson in the $H \rightarrow WW \rightarrow lvjj$ decay channel in pp collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, Phys. Rev. Lett. **107**, 231801 (2011).
- [823] G. Aad, “Search for the Standard Model Higgs boson in the decay channel $H \rightarrow ZZ^{(*)} \rightarrow 4$ with the ATLAS detector”, Phys. Lett. **B705**, 435–451 (2011).
- [824] G. Aad, “Search for the Standard Model Higgs boson in the two photon decay channel with the ATLAS detector at the LHC”, Phys. Lett. **B705**, 452–470 (2011).
- [825] T. Aaltonen, “A Search for resonant production of $t\bar{t}$ pairs in 4.8 fb⁻¹ of integrated luminosity of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D84**, 072004 (2011).
- [826] T. Aaltonen, “Evidence for a Mass Dependent Forward-Backward Asymmetry in Top Quark Pair Production”, Phys. Rev. **D83**, 112003 (2011).
- [827] T. Aaltonen, “Evidence for $t\bar{t}\gamma$ Production and Measurement of $\sigma_t\bar{t}\gamma/\sigma_t\bar{t}$ ”, Phys. Rev. **D84**, 031104 (2011).

- [828] T. Aaltonen, “First Measurement of the Angular Coefficients of Drell-Yan e^+e^- pairs in the Z Mass Region from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 241801 (2011).
- [829] T. Aaltonen, “First Search for Multijet Resonances in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **107**, 042001 (2011).
- [830] T. Aaltonen, “Improved determination of the sample composition of dimuon events produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Eur. Phys. J. **C71**, 1720 (2011).
- [831] T. Aaltonen, “Invariant Mass Distribution of Jet Pairs Produced in Association with a W boson in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 171801 (2011).
- [832] T. Aaltonen, “Limits on Anomalous Trilinear Gauge Couplings in $Z\gamma$ Events from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 051802 (2011).
- [833] T. Aaltonen, “Measurement of b hadron lifetimes in exclusive decays containing a J/ψ in p^- pbar collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 121804 (2011).
- [834] T. Aaltonen, “Measurement of branching ratio and B_s^0 lifetime in the decay $B_s^0 \rightarrow J/\psi f_0(980)$ at CDF”, Phys. Rev. **D84**, 052012 (2011).
- [835] T. Aaltonen, “Measurement of Event Shapes in Proton-Antiproton Collisions at Center-of-Mass Energy 1.96 TeV”, Phys. Rev. **D83**, 112007 (2011).
- [836] T. Aaltonen, “Measurement of Polarization and Search for CP-Violation in $B_s^0 \rightarrow \phi\phi$ Decays”, Phys. Rev. Lett. **107**, 261802 (2011).
- [837] T. Aaltonen, “Measurement of the B^- lifetime using a simulation free approach for trigger bias correction”, Phys. Rev. **D83**, 032008 (2011).
- [838] T. Aaltonen, “Measurement of the B_s Lifetime in Fully and Partially Reconstructed $B_s \rightarrow D_s^-(\phi\pi^-)X$ Decays in $\bar{p} - p$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 272001 (2011).
- [839] T. Aaltonen, “Measurement of the Cross Section for Prompt Isolated Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 102003 (2011).
- [840] T. Aaltonen, “Measurement of the Cross Section for Prompt Isolated Diphoton Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D84**, 052006 (2011).
- [841] T. Aaltonen, “Measurement of the Forward-Backward Asymmetry in the $B \rightarrow K^{(*)}\mu^+\mu^-$ Decay and First Observation of the $B_s^0 \rightarrow \phi\mu^+\mu^-$ Decay”, Phys. Rev. Lett. **106**, 161801 (2011).
- [842] T. Aaltonen, “Measurement of the mass difference between t and \bar{t} quarks”, Phys. Rev. Lett. **106**, 152001 (2011).
- [843] T. Aaltonen, “Measurement of the Top Pair Production Cross Section in the Lepton + Jets Channel Using a Jet Flavor Discriminant”, Phys. Rev. **D84**, 031101 (2011).
- [844] T. Aaltonen, “Measurement of the Top Quark Mass in the Lepton+Jets Channel Using the Lepton Transverse Momentum”, Phys. Lett. **B698**, 371–379 (2011).
- [845] T. Aaltonen, “Measurement of the top-quark mass in the lepton+jets channel using a matrix element technique with the CDF II detector”, Phys. Rev. **D84**, 071105 (2011).
- [846] T. Aaltonen, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using events with large Missing E_T and jets”, Phys. Rev. **D84**, 032003 (2011).
- [847] T. Aaltonen, “Measurement of the $t\bar{t}$ Production Cross Section with an in situ Calibration of b -jet Identification Efficiency”, Phys. Rev. **D83**, 071102 (2011).
- [848] T. Aaltonen, “Measurement of $t\bar{t}$ Spin Correlation in $p\bar{p}$ Collisions Using the CDF II Detector at the Tevatron”, Phys. Rev. **D83**, 031104 (2011).
- [849] T. Aaltonen, “Measurements of branching fraction ratios and CP-asymmetries in suppressed $B^- \rightarrow D(\rightarrow K^+\pi^-)K^-$ and $B^- \rightarrow D(\rightarrow K^+\pi^-)\pi^-$ decays”, Phys. Rev. **D84**, 091504 (2011).
- [850] T. Aaltonen, “Measurements of Direct CP Violating Asymmetries in Charmless Decays of Strange Bottom Mesons and Bottom Baryons”, Phys. Rev. Lett. **106**, 181802 (2011).
- [851] T. Aaltonen, “Measurements of the properties of $\Lambda_c(2595)$, $\Lambda_c(2625)$, $\Sigma_c(2455)$, and $\Sigma_c(2520)$ baryons”, Phys. Rev. **D84**, 012003 (2011).
- [852] T. Aaltonen, “Observation of $B_s^0 \rightarrow J/\psi K^{*0}(892)$ and $B_s^0 \rightarrow J/\psi K_S^0$ Decays”, Phys. Rev. **D83**, 052012 (2011).
- [853] T. Aaltonen, “Observation of the Ξ_b^0 Baryon”, Phys. Rev. Lett. **107**, 102001 (2011).
- [854] T. Aaltonen, “Observation of the Baryonic Flavor-Changing Neutral Current Decay $\Lambda_b \rightarrow \Lambda\mu^+\mu^-$ ”, Phys. Rev. Lett. **107**, 201802 (2011).
- [855] T. Aaltonen, “Search for a Heavy Top-Like Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 261801 (2011).

- [856] T. Aaltonen, “Search for a New Heavy Gauge Boson W' with Electron + missing ET Event Signature in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D83**, 031102 (2011).
- [857] T. Aaltonen, “Search for a Very Light CP-Odd Higgs Boson in Top Quark Decays from p^-p Collisions at 1.96 TeV”, Phys. Rev. Lett. **107**, 031801 (2011).
- [858] T. Aaltonen, “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B^0 \rightarrow \mu^+\mu^-$ Decays with CDF II”, Phys. Rev. Lett. **107**, [Addendum: Phys. Rev. Lett.107,no.23,239903(2011)], 191801 (2011).
- [859] T. Aaltonen, “Search for heavy bottom-like quarks decaying to an electron or muon and jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 141803 (2011).
- [860] T. Aaltonen, “Search for High Mass Resonances Decaying to Muon Pairs in $\sqrt{s} = 1.96$ TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **106**, 121801 (2011).
- [861] T. Aaltonen, “Search for New Dielectron Resonances and Randall-Sundrum Gravitons at the Collider Detector at Fermilab”, Phys. Rev. Lett. **107**, 051801 (2011).
- [862] T. Aaltonen, “Search for New Heavy Particles Decaying to $ZZ \rightarrow \ell\ell\ell\ell, \ell\ell jj$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D83**, 112008 (2011).
- [863] T. Aaltonen, “Search for new physics in high p_T like-sign dilepton events at CDF II”, Phys. Rev. Lett. **107**, 181801 (2011).
- [864] T. Aaltonen, “Search for New T' Particles in Final States with Large Jet Multiplicities and Missing Transverse Energy in $ppbar$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **107**, 191803 (2011).
- [865] T. Aaltonen, “Search for Production of Heavy Particles Decaying to Top Quarks and Invisible Particles in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **106**, 191801 (2011).
- [866] T. Aaltonen, “Search for Randall-Sundrum Gravitons in the Diphoton Channel at CDF”, Phys. Rev. **D83**, 011102 (2011).
- [867] T. Aaltonen, “Search for resonant production of $t\bar{t}$ decaying to jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D84**, 072003 (2011).
- [868] T. Aaltonen, “Search for the Higgs boson in the all-hadronic final state using the CDF II detector”, Phys. Rev. **D84**, 052010 (2011).
- [869] T. Aaltonen, “Top quark mass measurement using the template method at CDF”, Phys. Rev. **D83**, 111101 (2011).
- [870] T. Aaltonen, “Top-quark mass measurement using events with missing transverse energy and jets at CDF”, Phys. Rev. Lett. **107**, 232002 (2011).
- [871] S. Amerio, “ATLAS FTK: Fast track trigger”, PoS **VERTEX2011**, 040 (2011).
- [872] **M. S. Neubauer**, “Diboson production at colliders”, Ann. Rev. Nucl. Part. Sci. **61**, 223–250 (2011).
- [873] **M. S. Neubauer**, “Search for the Standard Model Higgs Boson in the Lepton + Missing Transverse Energy + Jets Final State in ATLAS”, (2011).
- [874] G. Aad, “Charged-particle multiplicities in pp interactions at $\sqrt{s} = 900$ GeV measured with the ATLAS detector at the LHC”, Phys. Lett. **B688**, 21–42 (2010).
- [875] G. Aad, “Commissioning of the ATLAS Muon Spectrometer with Cosmic Rays”, Eur. Phys. J. **C70**, 875–916 (2010).
- [876] G. Aad, “Drift Time Measurement in the ATLAS Liquid Argon Electromagnetic Calorimeter using Cosmic Muons”, Eur. Phys. J. **C70**, 755–785 (2010).
- [877] G. Aad, “Measurement of the $W \rightarrow \ell\nu$ and $Z/\gamma^* \rightarrow \ell\ell$ production cross sections in proton-proton collisions at $\sqrt{s} = 7$ TeV with the ATLAS detector”, JHEP **12**, 060 (2010).
- [878] G. Aad, “Performance of the ATLAS Detector using First Collision Data”, JHEP **09**, 056 (2010).
- [879] G. Aad, “Readiness of the ATLAS Liquid Argon Calorimeter for LHC Collisions”, Eur. Phys. J. **C70**, 723–753 (2010).
- [880] G. Aad, “Readiness of the ATLAS Tile Calorimeter for LHC collisions”, Eur. Phys. J. **C70**, 1193–1236 (2010).
- [881] G. Aad, “Search for New Particles in Two-Jet Final States in 7 TeV Proton-Proton Collisions with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **105**, 161801 (2010).
- [882] G. Aad, “The ATLAS Inner Detector commissioning and calibration”, Eur. Phys. J. **C70**, 787–821 (2010).
- [883] G. Aad, “The ATLAS Simulation Infrastructure”, Eur. Phys. J. **C70**, 823–874 (2010).
- [884] G. Aad, “Observation of a Centrality-Dependent Dijet Asymmetry in Lead-Lead Collisions at $\sqrt{s_{NN}} = 2.77$ TeV with the ATLAS Detector at the LHC”, Phys. Rev. Lett. **105**, 252303 (2010).
- [885] T. Aaltonen, “A Search for the Higgs Boson Using Neural Networks in Events with Missing Energy and b-quark Jets in p anti- p Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **104**, 141801 (2010).

- [886] T. Aaltonen, “A Study of the associated production of photons and b-quark jets in p p-bar collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. **D81**, 052006 (2010).
- [887] T. Aaltonen, “Combination of Tevatron searches for the standard model Higgs boson in the W+W-decay mode”, Phys. Rev. Lett. **104**, 061802 (2010).
- [888] T. Aaltonen, “Combined Tevatron upper limit on $gg \rightarrow H \rightarrow W^+W^-$ and constraints on the Higgs boson mass in fourth-generation fermion models”, Phys. Rev. **D82**, 011102 (2010).
- [889] T. Aaltonen, “Diffractive W and Z Production at the Fermilab Tevatron”, Phys. Rev. **D82**, 112004 (2010).
- [890] T. Aaltonen, “Direct Top-Quark Width Measurement CDF”, Phys. Rev. Lett. **105**, 232003 (2010).
- [891] T. Aaltonen, “Exclusion of an Exotic Top Quark with -4/3 Electric Charge Using Soft Lepton Tagging”, Phys. Rev. Lett. **105**, 101801 (2010).
- [892] T. Aaltonen, “First Measurement of the b-jet Cross Section in Events with a W Boson in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **104**, 131801 (2010).
- [893] T. Aaltonen, “First Measurement of the Ratio $\sigma(t\bar{t})/\sigma(Z/\gamma^{**} \rightarrow \ell\bar{\ell})$ and Precise Extraction of the $t\bar{t}$ Cross Section”, Phys. Rev. Lett. **105**, 012001 (2010).
- [894] T. Aaltonen, “Improved Search for a Higgs Boson Produced in Association with $Z \rightarrow \ell^+\ell^-$ in proton antiproton Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 251802 (2010).
- [895] T. Aaltonen, “Inclusive Search for Standard Model Higgs Boson Production in the WW Decay Channel using the CDF II Detector”, Phys. Rev. Lett. **104**, 061803 (2010).
- [896] T. Aaltonen, “Measurement of the Λ_b Lifetime in $\Lambda_b \rightarrow \Lambda_c^+ \pi^-$ Decays in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 102002 (2010).
- [897] T. Aaltonen, “Measurement of the Top Pair Production Cross Section in the Dilepton Decay Channel in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. **D82**, 052002 (2010).
- [898] T. Aaltonen, “Measurement of the Top Quark Mass and $p\bar{p} \rightarrow t\bar{t}$ Cross Section in the All-Hadronic Mode with the CDFII Detector”, Phys. Rev. **D81**, 052011 (2010).
- [899] T. Aaltonen, “Measurement of the Top Quark Mass in the Dilepton Channel Using m_{T2} at CDF”, Phys. Rev. **D81**, 031102 (2010).
- [900] T. Aaltonen, “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s}=1.96\text{ TeV}$ using Soft Electron b-Tagging”, Phys. Rev. **D81**, 092002 (2010).
- [901] T. Aaltonen, “Measurement of the W^+W^- Production Cross Section and Search for Anomalous $WW\gamma$ and WWZ Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 201801 (2010).
- [902] T. Aaltonen, “Measurement of the $WW + WZ$ Production Cross Section Using a Matrix Element Technique in Lepton + Jets Events”, Phys. Rev. **D82**, 112001 (2010).
- [903] T. Aaltonen, “Measurement of the $WW+WZ$ Production Cross Section Using the Lepton+Jets Final State at CDF II”, Phys. Rev. Lett. **104**, 101801 (2010).
- [904] T. Aaltonen, “Measurement of W-Boson Polarization in Top-quark Decay in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 042002 (2010).
- [905] T. Aaltonen, “Measurement of Z gamma Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. **D82**, 031103 (2010).
- [906] T. Aaltonen, “Measurements of Branching Fraction Ratios and CP Asymmetries in $B^\pm \rightarrow D_{CP} K^\pm$ Decays in Hadron Collisions”, Phys. Rev. **D81**, 031105 (2010).
- [907] T. Aaltonen, “Measurements of the top-quark mass using charged particle tracking”, Phys. Rev. **D81**, 032002 (2010).
- [908] T. Aaltonen, “Observation of Single Top Quark Production and Measurement of $|V_{tb}|$ with CDF”, Phys. Rev. **D82**, 112005 (2010).
- [909] T. Aaltonen, “Search for Anomalous Production of Events with Two Photons and Additional Energetic Objects at CDF”, Phys. Rev. **D82**, 052005 (2010).
- [910] T. Aaltonen, “Search for New Bottomlike Quark Pair Decays $Q\bar{Q} \rightarrow (tW)(\bar{t}W)$ in Same-Charge Dilepton Events”, Phys. Rev. Lett. **104**, 091801 (2010).
- [911] T. Aaltonen, “Search for New Color-Octet Vector Particle Decaying to t anti-t in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Lett. **B691**, 183–190 (2010).
- [912] T. Aaltonen, “Search for New Physics with a Dijet plus Missing Transverse Energy Signature in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **105**, 131801 (2010).
- [913] T. Aaltonen, “Search for Pair Production of Supersymmetric Top Quarks in Dilepton Events from p anti-p Collisions at $S^{**}(1/2) = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **104**, 251801 (2010).

- [914] T. Aaltonen, “Search for R-parity Violating Decays of τ sneutrinos to $e\mu$, $\mu\tau$, and $e\tau$ Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **105**, 191801 (2010).
- [915] T. Aaltonen, “Search for single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV in the missing transverse energy plus jets topology”, Phys. Rev. **D81**, 072003 (2010).
- [916] T. Aaltonen, “Search for Supersymmetry with Gauge-Mediated Breaking in Diphoton Events with Missing Transverse Energy at CDF II”, Phys. Rev. Lett. **104**, 011801 (2010).
- [917] T. Aaltonen, “Search for Technicolor Particles Produced in Association with a W Boson at CDF”, Phys. Rev. Lett. **104**, 111802 (2010).
- [918] T. Aaltonen, “Search for the Production of Scalar Bottom Quarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **105**, 081802 (2010).
- [919] T. Aaltonen, “Search for the supersymmetric partner of the top quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D82**, 092001 (2010).
- [920] T. Aaltonen, “Search for WW and WZ resonances decaying to electron, missing E_T , and two jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **104**, 241801 (2010).
- [921] T. Aaltonen, “Studying the Underlying Event in Drell-Yan and High Transverse Momentum Jet Production at the Tevatron”, Phys. Rev. **D82**, 034001 (2010).
- [922] T. Aaltonen, “Top Quark Mass Measurement in the Lepton + Jets Channel Using a Matrix Element Method and *in situ* Jet Energy Calibration”, Phys. Rev. Lett. **105**, 252001 (2010).
- [923] T. Aaltonen, “Updated Search for the Flavor-Changing Neutral-Current Decay $D^0 \rightarrow \mu^+\mu^-$ ”, Phys. Rev. **D82**, 091105 (2010).
- [924] T. A. Aaltonen, “Measurement of $d\sigma/dy$ of Drell-Yan e^+e^- pairs in the Z Mass Region from $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Lett. **B692**, 232–239 (2010).
- [925] M. Abolins, “The ATLAS Trigger/DAQ Authorlist, version 4.0”, (2010).
- [926] A. Andreani, “Enhancement of the ATLAS trigger system with a hardware tracker finder FTK”, JINST **5**, C12037 (2010).
- [927] J. D. Hobbs, **M. S. Neubauer**, S. Willenbrock, “Tests of the Standard Electroweak Model at the Energy Frontier”, (2010).
- [928] G. Aad, “Expected Performance of the ATLAS Experiment - Detector, Trigger and Physics”, (2009).
- [929] G. Aad, “Triggering on Long-Lived Neutral Particles in the ATLAS Detector”, (2009).
- [930] T. Aaltonen, “A Measurement of the t anti-t Cross Section in p anti-p Collisions at $\sqrt{s} = 1.96$ -TeV using Dilepton Events with a Lepton plus Track Selection”, Phys. Rev. **D79**, 112007 (2009).
- [931] T. Aaltonen, “A Search for high-mass resonances decaying to dimuons at CDF”, Phys. Rev. Lett. **102**, 091805 (2009).
- [932] T. Aaltonen, “A Search for the Associated Production of the Standard-Model Higgs Boson in the All-Hadronic Channel”, Phys. Rev. Lett. **103**, 221801 (2009).
- [933] T. Aaltonen, “A Search for the Higgs Boson Produced in Association with $Z \rightarrow \ell^+\ell^-$ Using the Matrix Element Method at CDF II”, Phys. Rev. **D80**, 071101 (2009).
- [934] T. Aaltonen, “Direct Measurement of the W Production Charge Asymmetry in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **102**, 181801 (2009).
- [935] T. Aaltonen, “Evidence for a Narrow Near-Threshold Structure in the $J/\psi\phi$ Mass Spectrum in $B^+ \rightarrow J/\psi\phi K^+$ Decays”, Phys. Rev. Lett. **102**, 242002 (2009).
- [936] T. Aaltonen, “First Direct Bound on the Total Width of the Top Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **102**, 042001 (2009).
- [937] T. Aaltonen, “First Measurement of the Ratio of Branching Fractions $B(\Lambda_b^0 \rightarrow \Lambda_c^+\mu^-\bar{\nu}_\mu)/B(\Lambda_b^0 \rightarrow \Lambda_c^+\pi^-)$ ”, Phys. Rev. **D79**, 032001 (2009).
- [938] T. Aaltonen, “First Measurement of the t anti-t Differential Cross Section $d\sigma/dM(t \text{ anti-}t)$ in p anti-p Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **102**, 222003 (2009).
- [939] T. Aaltonen, “First observation of $\bar{B}_s^0 \rightarrow D_s^\pm K^\mp$ and measurement of the ratio of branching fractions $B(\bar{B}_s^0 \rightarrow D_s^\pm K^\mp)/B(\bar{B}_s^0 \rightarrow D_s^\pm \pi^\mp)$ ”, Phys. Rev. Lett. **103**, 191802 (2009).
- [940] T. Aaltonen, “First Observation of Electroweak Single Top Quark Production”, Phys. Rev. Lett. **103**, 092002 (2009).
- [941] T. Aaltonen, “First Observation of Vector Boson Pairs in a Hadronic Final State at the Tevatron Collider”, Phys. Rev. Lett. **103**, 091803 (2009).
- [942] T. Aaltonen, “First simultaneous measurement of the top quark mass in the lepton + jets and dilepton channels at CDF”, Phys. Rev. **D79**, 092005 (2009).
- [943] T. Aaltonen, “Global Search for New Physics with 2.0 fb^{-1} at CDF”, Phys. Rev. **D79**, 011101 (2009).

- [944] T. Aaltonen, “Inclusive Search for Squark and Gluino Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 121801 (2009).
- [945] T. Aaltonen, “Measurement of Cross Sections for b Jet Production in Events with a Z Boson in p^- anti- p Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, 052008 (2009).
- [946] T. Aaltonen, “Measurement of Particle Production and Inclusive Differential Cross Sections in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, [Erratum: Phys. Rev.D82,119903(2010)], 112005 (2009).
- [947] T. Aaltonen, “Measurement of Resonance Parameters of Orbitally Excited Narrow B^0 Mesons”, Phys. Rev. Lett. **102**, 102003 (2009).
- [948] T. Aaltonen, “Measurement of the b-Hadron Production Cross Section Using Decays to MU- d0 X Final States in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, 092003 (2009).
- [949] T. Aaltonen, “Measurement of the fraction of $t\bar{t}$ production via gluon-gluon fusion in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, 031101 (2009).
- [950] T. Aaltonen, “Measurement of the Inclusive Isolated Prompt Photon Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ using the CDF Detector”, Phys. Rev. **D80**, 111106 (2009).
- [951] T. Aaltonen, “Measurement of the k_T Distribution of Particles in Jets Produced in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 232002 (2009).
- [952] T. Aaltonen, “Measurement of the top quark mass at CDF using the ‘neutrino ϕ weighting’ template method on a lepton plus isolated track sample”, Phys. Rev. **D79**, 072005 (2009).
- [953] T. Aaltonen, “Measurement of the Top Quark Mass Using the Invariant Mass of Lepton Pairs in Soft Muon b-tagged Events”, Phys. Rev. **D80**, 051104 (2009).
- [954] T. Aaltonen, “Measurement of the top quark mass with dilepton events selected using neuroevolution at CDF”, Phys. Rev. Lett. **102**, 152001 (2009).
- [955] T. Aaltonen, “Measurement of the $t\bar{t}$ Production Cross Section in 2 fb^{-1} of $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ Using Lepton Plus Jets Events with Soft Muon b-Tagging”, Phys. Rev. **D79**, 052007 (2009).
- [956] T. Aaltonen, “Measurement of W-Boson Helicity Fractions in Top-Quark Decays Using $\cos\theta^*$ ”, Phys. Lett. **B674**, 160–167 (2009).
- [957] T. Aaltonen, “Observation of exclusive charmonium production and $\gamma + \gamma$ to $\mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **102**, 242001 (2009).
- [958] T. Aaltonen, “Observation of New Charmless Decays of Bottom Hadrons”, Phys. Rev. Lett. **103**, 031801 (2009).
- [959] T. Aaltonen, “Observation of the Ω_b^- and Measurement of the Properties of the Ξ_b^- and Ω_b^- ”, Phys. Rev. **D80**, 072003 (2009).
- [960] T. Aaltonen, “Precision Measurement of the $X(3872)$ Mass in $J/\psi\pi^+\pi^-$ Decays”, Phys. Rev. Lett. **103**, 152001 (2009).
- [961] T. Aaltonen, “Production of $\psi(2S)$ Mesons in p anti-p Collisions at 1.96-TeV ”, Phys. Rev. **D80**, 031103 (2009).
- [962] T. Aaltonen, “Search for a Fermiophobic Higgs Boson Decaying into Diphotons in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **103**, 061803 (2009).
- [963] T. Aaltonen, “Search for a Higgs Boson Decaying to Two W Bosons at CDF”, Phys. Rev. Lett. **102**, 021802 (2009).
- [964] T. Aaltonen, “Search for a Higgs Boson in $WH \rightarrow \ell\nu b\bar{b}$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **103**, 101802 (2009).
- [965] T. Aaltonen, “Search for Anomalous Production of Events with a Photon, Jet, b-quark Jet, and Missing Transverse Energy”, Phys. Rev. **D80**, 052003 (2009).
- [966] T. Aaltonen, “Search for charged Higgs bosons in decays of top quarks in p anti-p collisions at $s^{**}(1/2) = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **103**, 101803 (2009).
- [967] T. Aaltonen, “Search for exclusive Z boson production and observation of high mass $p\bar{p} \rightarrow \gamma\gamma \rightarrow p + \ell\ell + \bar{p}$ events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **102**, 222002 (2009).
- [968] T. Aaltonen, “Search for Gluino-Mediated Sbottom Production in p anti-p Collisions at $s^{**}(1/2) = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 221801 (2009).
- [969] T. Aaltonen, “Search for Hadronic Decays of W and Z Bosons in Photon Events in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-GeV}$ ”, Phys. Rev. **D80**, 052011 (2009).
- [970] T. Aaltonen, “Search for Higgs bosons predicted in two-Higgs-doublet models via decays to tau lepton pairs in 1.96-TeV p anti-p collisions”, Phys. Rev. Lett. **103**, 201801 (2009).
- [971] T. Aaltonen, “Search for High-Mass e^+e^- Resonances in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 031801 (2009).

- [972] T. Aaltonen, “Search for Long-Lived Massive Charged Particles in 1.96 TeV $\bar{p}p$ Collisions”, Phys. Rev. Lett. **103**, 021802 (2009).
- [973] T. Aaltonen, “Search for Maximal Flavor Violating Scalars in Same-Charge Lepton Pairs in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **102**, 041801 (2009).
- [974] T. Aaltonen, “Search for narrow resonances lighter than Υ mesons”, Eur. Phys. J. **C62**, 319–326 (2009).
- [975] T. Aaltonen, “Search for new particles decaying into dijets in proton-antiproton collisions at $s^{*(1/2)} = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, 112002 (2009).
- [976] T. Aaltonen, “Search for new physics in the $\mu\mu + e/\mu + \cancel{E}_T$ channel with a low- p_T lepton threshold at the Collider Detector at Fermilab”, Phys. Rev. **D79**, 052004 (2009).
- [977] T. Aaltonen, “Search for Standard Model Higgs Boson Production in Association with a W Boson using a Neural Network Discriminant at CDF”, Phys. Rev. **D80**, 012002 (2009).
- [978] T. Aaltonen, “Search for the Decays $B_s^0 \rightarrow e^+\mu^-$ and $B_s^0 \rightarrow e^+e^-$ in CDF Run II”, Phys. Rev. Lett. **102**, 201801 (2009).
- [979] T. Aaltonen, “Search for the Neutral Current Top Quark Decay $t \rightarrow Zc$ Using Ratio of Z-Boson + 4 Jets to W-Boson + 4 Jets Production”, Phys. Rev. **D80**, 052001 (2009).
- [980] T. Aaltonen, “Search for the Production of Narrow t anti-b Resonances in 1.9 fb-1 of p anti-p Collisions at $s^{*(1/2)} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **103**, 041801 (2009).
- [981] T. Aaltonen, “Search for the Rare Decays $B^+ \rightarrow \mu^+\mu^-K^+$, $B^0 \rightarrow \mu^+\mu^-K^{*0}(892)$, and $B_s^0 \rightarrow \mu^+\mu^-\phi$ at CDF”, Phys. Rev. **D79**, 011104 (2009).
- [982] T. Aaltonen, “Search for top-quark production via flavor-changing neutral currents in W+1 jet events at CDF”, Phys. Rev. Lett. **102**, 151801 (2009).
- [983] T. Aaltonen, “Search for WW and WZ Production in Lepton Plus Jets Final State at CDF”, Phys. Rev. **D79**, 112011 (2009).
- [984] T. Aaltonen, “Searching the Inclusive Lepton + Photon + Missing E(T) + b-quark Signature for Radiative Top Quark Decay and Non-Standard-Model Processes”, Phys. Rev. **D80**, 011102 (2009).
- [985] T. Aaltonen, “Top Quark Mass Measurement in the Lepton plus Jets Channel Using a Modified Matrix Element Method”, Phys. Rev. **D79**, 072001 (2009).
- [986] T. Aaltonen, “Top Quark Mass Measurement in the $t\bar{t}$ All Hadronic Channel using a Matrix Element Technique in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D79**, 072010 (2009).
- [987] M. Abolins, “The ATLAS Trigger/DAQ Authorlist, version 3.0”, (2009).
- [988] M. Abolins, “The ATLAS Trigger/DAQ Authorlist, version 3.1”, (2009).
- [989] A. Abulencia, “Measurement of the Ratio of Branching Fractions $B(B^+ \rightarrow J/\psi\pi^+)/B(B^+ \rightarrow J/\psi K^+)$ ”, Phys. Rev. **D79**, 112003 (2009).
- [990] A. Annovi, “Development of FTK architecture: a fast hardware track trigger for the ATLAS detector”, (2009).
- [991] A. Annovi, “The fast tracker architecture for the LHC baseline luminosity”, PoS **EPS-HEP2009**, 136 (2009).
- [992] T. Aaltonen, “A Direct measurement of the W boson width in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 071801 (2008).
- [993] T. Aaltonen, “Cross-section constrained top quark mass measurement from dilepton events at the Tevatron”, Phys. Rev. Lett. **100**, 062005 (2008).
- [994] T. Aaltonen, “Evidence for $D^0 - \bar{D}^0$ mixing using the CDF II Detector”, Phys. Rev. Lett. **100**, 121802 (2008).
- [995] T. Aaltonen, “First Flavor-Tagged Determination of Bounds on Mixing-Induced CP Violation in $B_s^0 \rightarrow J/\psi\phi$ Decays”, Phys. Rev. Lett. **100**, 161802 (2008).
- [996] T. Aaltonen, “First measurement of the fraction of top quark pair production through gluon-gluon fusion”, Phys. Rev. **D78**, 111101 (2008).
- [997] T. Aaltonen, “First measurement of the production of a W boson in association with a single charm quark in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 091803 (2008).
- [998] T. Aaltonen, “First Measurement of ZZ Production in panti-p Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 201801 (2008).
- [999] T. Aaltonen, “First Observation of the Decay $B_s^0 \rightarrow D_s^- D_s^+$ and Measurement of Its Branching Ratio”, Phys. Rev. Lett. **100**, 021803 (2008).
- [1000] T. Aaltonen, “First Run II Measurement of the W Boson Mass”, Phys. Rev. **D77**, 112001 (2008).
- [1001] T. Aaltonen, “Forward-Backward Asymmetry in Top Quark Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 202001 (2008).

- [1002] T. Aaltonen, “Limits on the production of narrow $t\bar{t}$ resonances in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D77**, 051102 (2008).
- [1003] T. Aaltonen, “Measurement of b -jet Shapes in Inclusive Jet Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D78**, 072005 (2008).
- [1004] T. Aaltonen, “Measurement of correlated $b^-\bar{b}$ production in $p^-\bar{p}$ collisions at $\sqrt{s} = 1960\text{ GeV}$ ”, Phys. Rev. **D77**, 072004 (2008).
- [1005] T. Aaltonen, “Measurement of inclusive jet cross-sections in $Z/\gamma^* \rightarrow e^+e^- + \text{jets}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 102001 (2008).
- [1006] T. Aaltonen, “Measurement of lifetime and decay-width difference in $B_s^0 \rightarrow J/\psi\phi$ decays”, Phys. Rev. Lett. **100**, 121803 (2008).
- [1007] T. Aaltonen, “Measurement of Ratios of Fragmentation Fractions for Bottom Hadrons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D77**, 072003 (2008).
- [1008] T. Aaltonen, “Measurement of the cross section for W^- boson production in association with jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D77**, 011108 (2008).
- [1009] T. Aaltonen, “Measurement of the Inclusive Jet Cross Section at the Fermilab Tevatron p anti- p Collider Using a Cone-Based Jet Algorithm”, Phys. Rev. **D78**, [Erratum: Phys. Rev. **D79**, 119902(2009)], 052006 (2008).
- [1010] T. Aaltonen, “Measurement of the Single Top Quark Production Cross Section at CDF”, Phys. Rev. Lett. **101**, 252001 (2008).
- [1011] T. Aaltonen, “Model-Independent and Quasi-Model-Independent Search for New Physics at CDF”, Phys. Rev. **D78**, 012002 (2008).
- [1012] T. Aaltonen, “Observation of Exclusive Dijet Production at the Fermilab Tevatron $p^-\bar{p}$ Collider”, Phys. Rev. **D77**, 052004 (2008).
- [1013] T. Aaltonen, “Observation of orbitally excited B_s mesons”, Phys. Rev. Lett. **100**, 082001 (2008).
- [1014] T. Aaltonen, “Observation of the Decay $B^+ \rightarrow J/\psi\pi^+$ and Measurement of the B^+ Mass”, Phys. Rev. Lett. **100**, 182002 (2008).
- [1015] T. Aaltonen, “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ decays with $2fb^{-1}$ of $p\bar{p}$ collisions”, Phys. Rev. Lett. **100**, 101802 (2008).
- [1016] T. Aaltonen, “Search for chargino-neutralino production in $p\bar{p}$ collisions at 1.96-TeV with high- $p(T)$ leptons”, Phys. Rev. **D77**, 052002 (2008).
- [1017] T. Aaltonen, “Search for Doubly Charged Higgs Bosons with Lepton-Flavor-Violating Decays involving Tau Leptons”, Phys. Rev. Lett. **101**, 121801 (2008).
- [1018] T. Aaltonen, “Search for Heavy, Long-Lived Neutralinos that Decay to Photons at CDF II Using Photon Timing”, Phys. Rev. **D78**, 032015 (2008).
- [1019] T. Aaltonen, “Search for Heavy Top-like Quarks Using Lepton Plus Jets Events in 1.96-TeV $p\bar{p}$ Collisions”, Phys. Rev. Lett. **100**, 161803 (2008).
- [1020] T. Aaltonen, “Search for large extra dimensions in final states containing one photon or jet and large missing transverse energy produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **101**, 181602 (2008).
- [1021] T. Aaltonen, “Search for New Heavy Particles Decaying to $Z^0Z^0 \rightarrow eeee$ in $p - \bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. **D78**, 012008 (2008).
- [1022] T. Aaltonen, “Search for Pair Production of Scalar Top Quarks Decaying to a τ Lepton and a b Quark in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 071802 (2008).
- [1023] T. Aaltonen, “Search for resonant $t\bar{t}$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96\text{-TeV}$ ”, Phys. Rev. Lett. **100**, 231801 (2008).
- [1024] T. Aaltonen, “Search for Standard Model Higgs Boson Production in Association with a W Boson at CDF”, Phys. Rev. **D78**, 032008 (2008).
- [1025] T. Aaltonen, “Search for Standard Model Higgs Bosons Produced in Association with W Bosons”, Phys. Rev. Lett. **100**, 041801 (2008).
- [1026] T. Aaltonen, “Search for Supersymmetry in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{-TeV}$ Using the Tripleton Signature of Chargino-Neutralino Production”, Phys. Rev. Lett. **101**, 251801 (2008).
- [1027] T. Aaltonen, “Search for the Flavor Changing Neutral Current Decay $t \rightarrow Zq$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **101**, 192002 (2008).
- [1028] T. Aaltonen, “Search for the Higgs boson in events with missing transverse energy and b quark jets produced in proton-antiproton collisions at $\sqrt{s} = 1.96\text{ TeV}$ ”, Phys. Rev. Lett. **100**, 211801 (2008).

- [1029] T. Aaltonen, “Search for the Higgs boson produced with $Z \rightarrow \ell^+ \ell^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **101**, 251803 (2008).
- [1030] T. Aaltonen, “Search for Third Generation Vector Leptoquarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D77**, 091105 (2008).
- [1031] T. Aaltonen, “Two-Particle Momentum Correlations in Jets Produced in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D77**, 092001 (2008).
- [1032] A. Abulencia, “Cross Section Measurements of High-p(T) Dilepton Final-State Processes Using a Global Fitting Method”, Phys. Rev. **D78**, 012003 (2008).
- [1033] A. Annovi, “A hardware track finder for the ATLAS trigger”, Nuovo Cim. **B123**, 981–983 (2008).
- [1034] T. Aaltonen, “First measurement of the W boson mass in Run II of the Tevatron”, Phys. Rev. Lett. **99**, 151801 (2007).
- [1035] T. Aaltonen, “First observation of heavy baryons Σ_b and Σ_b^{*} ”, Phys. Rev. Lett. **99**, 202001 (2007).
- [1036] T. Aaltonen, “Limits on Anomalous Triple Gauge Couplings in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D76**, 111103 (2007).
- [1037] T. Aaltonen, “Measurement of the $p\bar{p} \rightarrow t\bar{t}$ production cross-section and the top quark mass at $\sqrt{s} = 1.96$ -TeV in the all-hadronic decay mode”, Phys. Rev. **D76**, 072009 (2007).
- [1038] T. Aaltonen, “Measurement of the top-quark mass in all-hadronic decays in p anti-p collisions at CDF II”, Phys. Rev. Lett. **98**, 142001 (2007).
- [1039] T. Aaltonen, “Measurement of the top-quark mass using missing E_T + jets events with secondary vertex b -tagging at CDF II”, Phys. Rev. **D75**, 111103 (2007).
- [1040] T. Aaltonen, “Model-Independent Global Search for New High-p(T) Physics at CDF”, (2007) 10.2172/922303.
- [1041] T. Aaltonen, “Observation and mass measurement of the baryon Xi_b^- ”, Phys. Rev. Lett. **99**, 052002 (2007).
- [1042] T. Aaltonen, “Search for a high-mass diphoton state and limits on Randall-Sundrum gravitons at CDF”, Phys. Rev. Lett. **99**, 171801 (2007).
- [1043] T. Aaltonen, “Search for chargino-neutralino production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 191806 (2007).
- [1044] T. Aaltonen, “Search for Direct Pair Production of Supersymmetric Top and Supersymmetric Bottom Quarks in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D76**, 072010 (2007).
- [1045] T. Aaltonen, “Search for exclusive $\gamma\gamma$ production in hadron-hadron collisions”, Phys. Rev. Lett. **99**, 242002 (2007).
- [1046] T. Aaltonen, “Search for New Particles Leading to Z + jets Final States in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D76**, 072006 (2007).
- [1047] T. Aaltonen, “Search for new physics in high mass electron-positron events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 171802 (2007).
- [1048] G. Aarons, “ILC Reference Design Report Volume 3 - Accelerator”, edited by N. Phinney, N. Toge, and N. J. Walker (2007).
- [1049] A. Abulencia, “Analysis of the quantum numbers J^{PC} of the $X(3872)$ ”, Phys. Rev. Lett. **98**, 132002 (2007).
- [1050] A. Abulencia, “First Measurement of the Ratio of Central-Electron to Forward-Electron W Partial Cross Sections in p anti-p Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **98**, 251801 (2007).
- [1051] A. Abulencia, “Inclusive search for new physics with like-sign dilepton events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 221803 (2007).
- [1052] A. Abulencia, “Measurement of $\sigma(\Lambda_b^0)/\sigma(\bar{B}^0) \times \text{BR}(\Lambda_b^0 \rightarrow \Lambda_c^+ \pi^-)/\text{BR}(\bar{B}^0 \rightarrow D^+ \pi^-)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 122002 (2007).
- [1053] A. Abulencia, “Measurement of $\sigma_{\chi_{c2}} \mathcal{B}(\chi_{c2} \rightarrow J/\psi \gamma)/\sigma_{\chi_{c1}} \mathcal{B}(\chi_{c1} \rightarrow J/\psi \gamma)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 232001 (2007).
- [1054] A. Abulencia, “Measurement of $\sigma(p\bar{p} \rightarrow Z) \cdot \text{Br}(Z \rightarrow 2\tau)$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D75**, 092004 (2007).
- [1055] A. Abulencia, “Measurement of the Λ_b^0 Lifetime in $\Lambda_b^0 \rightarrow J/\psi \Lambda^0$ in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 122001 (2007).
- [1056] A. Abulencia, “Measurement of the B^+ production cross-section in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D75**, 012010 (2007).
- [1057] A. Abulencia, “Measurement of the Helicity Fractions of W Bosons from Top Quark Decays using Fully Reconstructed $t\bar{t}$ Events with CDF II”, Phys. Rev. **D75**, 052001 (2007).

- [1058] A. Abulencia, “Measurement of the Inclusive Jet Cross Section using the k_T algorithm in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV with the CDF II Detector”, Phys. Rev. **D75**, [Erratum: Phys. Rev. **D75**, 119901(2007)], 092006 (2007).
- [1059] A. Abulencia, “Measurement of the Ratios of Branching Fractions $B(B_0(s) \rightarrow D(s) \pi^+ \pi^+ \pi^-) / B(B_0 \rightarrow D^- \pi^+ \pi^+ \pi^-)$ and $B(B_0(s) \rightarrow D(s) \pi^+) / B(B_0 \rightarrow D^- \pi^+)$ ”, Phys. Rev. Lett. **98**, 061802 (2007).
- [1060] A. Abulencia, “Measurement of the Top Quark Mass in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV using the Decay Length Technique”, Phys. Rev. **D75**, 071102 (2007).
- [1061] A. Abulencia, “Measurements of inclusive W and Z cross sections in p anti-p collisions at $s^{*(1/2)} = 1.96$ -TeV”, J. Phys. **G34**, 2457–2544 (2007).
- [1062] A. Abulencia, “Observation of Exclusive Electron-Positron Production in Hadron-Hadron Collisions”, Phys. Rev. Lett. **98**, 112001 (2007).
- [1063] A. Abulencia, “Observation of WZ Production”, Phys. Rev. Lett. **98**, 161801 (2007).
- [1064] A. Abulencia, “Polarization of J/ψ and ψ_{2S} mesons produced in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **99**, 132001 (2007).
- [1065] A. Abulencia, “Precise measurement of the top quark mass in the lepton+jets topology at CDF II”, Phys. Rev. Lett. **99**, 182002 (2007).
- [1066] A. Abulencia, “Precision measurement of the top quark mass from dilepton events at CDF II”, Phys. Rev. **D75**, 031105 (2007).
- [1067] A. Abulencia, “Search for anomalous production of multi-lepton events in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 131804 (2007).
- [1068] A. Abulencia, “Search for Exotic $S=-2$ Baryons in proton-antiproton Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D75**, 032003 (2007).
- [1069] A. Abulencia, “Search for heavy, long-lived particles that decay to photons at CDF II”, Phys. Rev. Lett. **99**, 121801 (2007).
- [1070] A. Abulencia, “Search for new physics in lepton + photon + X events with 929 pb⁽⁻¹⁾ of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D75**, 112001 (2007).
- [1071] A. Abulencia, “Search for V+A current in top quark decay in p anti-p collisions at $s^{*(1/2)} = 1.96$ -TeV”, Phys. Rev. Lett. **98**, 072001 (2007).
- [1072] A. Abulencia, “Search for W-prime boson decaying to electron-neutrino pairs in p anti-p collisions at $s^{*(1/2)} = 1.96$ -TeV”, Phys. Rev. **D75**, 091101 (2007).
- [1073] B. Aharmim, “Determination of the ν_e and total ^8B solar neutrino fluxes with the Sudbury neutrino observatory phase I data set”, Phys. Rev. **C75**, 045502 (2007).
- [1074] A. Abulencia, “A search for scalar bottom quarks from gluino decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 171802 (2006).
- [1075] A. Abulencia, “A search for $t \rightarrow \tau \nu q$ in $t\bar{t}$ production.”, Phys. Lett. **B639**, 172–178 (2006).
- [1076] A. Abulencia, “Direct search for Dirac magnetic monopoles in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **96**, 201801 (2006).
- [1077] A. Abulencia, “Evidence for the exclusive decay $B_c^\pm \rightarrow J/\psi \pi^\pm$ and measurement of the mass of the B_c meson”, Phys. Rev. Lett. **96**, 082002 (2006).
- [1078] A. Abulencia, “Measurement of mass and width of the excited charmed meson states $D_0(1)$ and $D^*_0(2)$ at CDF”, Phys. Rev. **D73**, 051104 (2006).
- [1079] A. Abulencia, “Measurement of the b jet cross-section in events with a Z boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. **D74**, 032008 (2006).
- [1080] A. Abulencia, “Measurement of the $B_s^0 - \bar{B}_s^0$ Oscillation Frequency”, Phys. Rev. Lett. **97**, 062003 (2006).
- [1081] A. Abulencia, “Measurement of the $B(c)^+$ meson lifetime using $B(c)^+ \rightarrow J/\psi e^+ \nu(e)$ ”, Phys. Rev. Lett. **97**, 012002 (2006).
- [1082] A. Abulencia, “Measurement of the dipion mass spectrum in $X(3872) \rightarrow J/\psi \pi^+ \pi^-$ decays.”, Phys. Rev. Lett. **96**, 102002 (2006).
- [1083] A. Abulencia, “Measurement of the helicity of W bosons in top-quark decays”, Phys. Rev. **D73**, 111103 (2006).
- [1084] A. Abulencia, “Measurement of the inclusive jet cross section in $p\bar{p}$ interactions at $\sqrt{s} = 1.96$ -TeV using a cone-based jet algorithm.”, Phys. Rev. **D74**, 071103 (2006).
- [1085] A. Abulencia, “Measurement of the inclusive jet cross section using the $k(t)$ algorithm in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 122001 (2006).

- [1086] A. Abulencia, “Measurement of the ratio of branching fractions $B(D0 \rightarrow K^+\pi^-) / B(D0 \rightarrow K^-\pi^+)$ using the CDF II Detector”, Phys. Rev. **D74**, 031109 (2006).
- [1087] A. Abulencia, “Measurement of the ratios of branching fractions $B(B_s^0 \rightarrow D_s^-\pi^+)/B(B^0 \rightarrow D^-\pi^+)$ and $B(B^+ \rightarrow \bar{D}^0\pi^+)/B(B^0 \rightarrow D^-\pi^+)$ ”, Phys. Rev. Lett. **96**, 191801 (2006).
- [1088] A. Abulencia, “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **97**, 082004 (2006).
- [1089] A. Abulencia, “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV in the All Hadronic Decay Mode”, Phys. Rev. **D74**, 072005 (2006).
- [1090] A. Abulencia, “Measurement of the $t\bar{t}$ Production Cross Section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV using Lepton + Jets Events with Jet Probability b^- tagging”, Phys. Rev. **D74**, 072006 (2006).
- [1091] A. Abulencia, “Measurement of the top quark mass using template methods on dilepton events in proton antiproton collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. **D73**, 112006 (2006).
- [1092] A. Abulencia, “Measurement of the top quark mass with the dynamical likelihood method using lepton plus jets events with b-tags in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. **D73**, 092002 (2006).
- [1093] A. Abulencia, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV using missing E_T + jets events with secondary vertex b -tagging”, Phys. Rev. Lett. **96**, 202002 (2006).
- [1094] A. Abulencia, “Observation of $B^0 (s) \rightarrow K^+K^-$ and Measurements of Branching Fractions of Charmless Two-body Decays of B^0 and B_s^0 Mesons in $p\bar{p}$ Collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 211802 (2006).
- [1095] A. Abulencia, “Observation of $B_s^0 - \bar{B}_s^0$ Oscillations”, Phys. Rev. Lett. **97**, 242003 (2006).
- [1096] A. Abulencia, “Observation of $B_s^0 \rightarrow \psi(2S)\phi$ and measurement of ratio of branching fractions $B(B_s^0 \rightarrow \psi(2S)\phi)/B(B_s^0 \rightarrow J/\psi\phi)$ ”, Phys. Rev. Lett. **96**, 231801 (2006).
- [1097] A. Abulencia, “Precision top quark mass measurement in the lepton + jets topology in p anti-p collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 022004 (2006).
- [1098] A. Abulencia, “Search for a neutral Higgs boson decaying to a W boson pair in p antip collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 081802 (2006).
- [1099] A. Abulencia, “Search for anomalous semileptonic decay of heavy flavor hadrons produced in association with a W boson at CDF II”, Phys. Rev. **D73**, 051101 (2006).
- [1100] A. Abulencia, “Search for charged Higgs bosons from top quark decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 042003 (2006).
- [1101] A. Abulencia, “Search for excited and exotic muons in the $\mu\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 191802 (2006).
- [1102] A. Abulencia, “Search for H to b anti- b produced in association with W bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 081803 (2006).
- [1103] A. Abulencia, “Search for high-mass resonances decaying to $e\mu$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. Lett. **96**, 211802 (2006).
- [1104] A. Abulencia, “Search for Large Extra Dimensions in the Production of Jets and Missing Transverse Energy in p anti-p Collisions at $s^{**}(1/2) = 1.96$ TeV”, Phys. Rev. Lett. **97**, 171802 (2006).
- [1105] A. Abulencia, “Search for neutral MSSM Higgs bosons decaying to tau pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **96**, 011802 (2006).
- [1106] A. Abulencia, “Search for new physics in lepton + photon + X events with 305 pb^{-1} of $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV”, Phys. Rev. Lett. **97**, 031801 (2006).
- [1107] A. Abulencia, “Search for second-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ -TeV.”, Phys. Rev. **D73**, 051102 (2006).
- [1108] A. Abulencia, “Search for $Z' \rightarrow e^+e^-$ using dielectron mass and angular distribution.”, Phys. Rev. Lett. **96**, 211801 (2006).
- [1109] A. Abulencia, “Top quark mass measurement from dilepton events at CDF II”, Phys. Rev. Lett. **96**, 152002 (2006).
- [1110] A. Abulencia, “Top quark mass measurement from dilepton events at CDF II with the matrix-element method”, Phys. Rev. **D74**, 032009 (2006).
- [1111] A. Abulencia, “Top quark mass measurement using the template method in the lepton + jets channel at CDF II”, Phys. Rev. **D73**, 032003 (2006).
- [1112] D. Acosta, “Measurement of b hadron masses in exclusive J/ψ decays with the CDF detector”, Phys. Rev. Lett. **96**, 202001 (2006).

- [1113] D. Acosta, “Measurement of the azimuthal angle distribution of leptons from W boson decays as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D73**, 052002 (2006).
- [1114] D. Acosta, “Search for W and Z bosons in the reaction $p\bar{p} \rightarrow 2\text{jets} + \gamma$ at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D73**, 012001 (2006).
- [1115] A. Abulencia, “Search for $B_s \rightarrow \mu^+\mu^-$ and $B_d \rightarrow \mu^+\mu^-$ decays in $p\bar{p}$ collisions with CDF II”, Phys. Rev. Lett. **95**, [Erratum: Phys. Rev. Lett.95,249905(2005)], 221805 (2005).
- [1116] A. Abulencia, “Search for new high mass particles decaying to lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 252001 (2005).
- [1117] D. Acosta, “A search for supersymmetric Higgs bosons in the di-tau decay mode in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D72**, 072004 (2005).
- [1118] D. Acosta, “Comparison of three-jet events in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV to predictions from a next-to-leading order QCD calculation”, Phys. Rev. **D71**, 032002 (2005).
- [1119] D. Acosta, “First evidence for $B_s^0 \rightarrow \phi\phi$ decay and measurements of branching ratio and A_{CP} for $B^+ \rightarrow \phi K^+$ ”, Phys. Rev. Lett. **95**, 031801 (2005).
- [1120] D. Acosta, “First measurements of inclusive W and Z cross sections from Run II of the Tevatron collider”, Phys. Rev. Lett. **94**, 091803 (2005).
- [1121] D. Acosta, “ K_S^0 and Λ^0 production studies in $p\bar{p}$ collisions at $\sqrt{s} = 1800$ -GeV and 630-GeV”, Phys. Rev. **D72**, 052001 (2005).
- [1122] D. Acosta, “Measurement of $B(t \rightarrow Wb)/B(t \rightarrow Wq)$ at the Collider Detector at Fermilab”, Phys. Rev. Lett. **95**, 102002 (2005).
- [1123] D. Acosta, “Measurement of charged particle multiplicities in gluon and quark jets in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **94**, 171802 (2005).
- [1124] D. Acosta, “Measurement of partial widths and search for direct CP violation in D^0 meson decays to K^-K^+ and $\pi^-\pi^+$ ”, Phys. Rev. Lett. **94**, 122001 (2005).
- [1125] D. Acosta, “Measurement of the cross section for prompt diphoton production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 022003 (2005).
- [1126] D. Acosta, “Measurement of the cross section for $t\bar{t}$ production in $p\bar{p}$ collisions using the kinematics of lepton + jets events”, Phys. Rev. **D72**, 052003 (2005).
- [1127] D. Acosta, “Measurement of the forward-backward charge asymmetry from $W \rightarrow e\nu$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, 051104 (2005).
- [1128] D. Acosta, “Measurement of the forward-backward charge asymmetry of electron positron pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, 052002 (2005).
- [1129] D. Acosta, “Measurement of the J/ψ meson and b -hadron production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1960$ GeV”, Phys. Rev. **D71**, 032001 (2005).
- [1130] D. Acosta, “Measurement of the lifetime difference between B_s mass eigenstates”, Phys. Rev. Lett. **94**, 101803 (2005).
- [1131] D. Acosta, “Measurement of the moments of the hadronic invariant mass distribution in semileptonic B decays”, Phys. Rev. **D71**, 051103 (2005).
- [1132] D. Acosta, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using kinematic fitting of b -tagged lepton + jet events”, Phys. Rev. **D71**, 072005 (2005).
- [1133] D. Acosta, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using lepton + jets events with secondary vertex b -tagging”, Phys. Rev. **D71**, 052003 (2005).
- [1134] D. Acosta, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using lepton plus jets events with semileptonic B decays to muons”, Phys. Rev. **D72**, 032002 (2005).
- [1135] D. Acosta, “Measurement of the W boson polarization in top decay at CDF at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D71**, [Erratum: Phys. Rev.D71,059901(2005)], 031101 (2005).
- [1136] D. Acosta, “Measurement of the W^+W^- production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using dilepton events”, Phys. Rev. Lett. **94**, 211801 (2005).
- [1137] D. Acosta, “Measurement of $W\gamma$ and $Z\gamma$ production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **94**, 041803 (2005).
- [1138] D. Acosta, “Measurements of $b\bar{b}$ azimuthal production correlations in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D71**, 092001 (2005).
- [1139] D. Acosta, “Search for anomalous kinematics in $t\bar{t}$ dilepton events at CDF II”, Phys. Rev. Lett. **95**, 022001 (2005).

- [1140] D. Acosta, “Search for anomalous production of diphoton events with missing transverse energy at CDF and limits on gauge-mediated supersymmetry-breaking models”, Phys. Rev. **D71**, 031104 (2005).
- [1141] D. Acosta, “Search for $\Lambda_b \rightarrow p\pi$ and $\Lambda_b \rightarrow pK$ decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D72**, 051104 (2005).
- [1142] D. Acosta, “Search for electroweak single top quark production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, 012005 (2005).
- [1143] D. Acosta, “Search for excited and exotic electrons in the $e\gamma$ decay channel in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **94**, 101802 (2005).
- [1144] D. Acosta, “Search for first-generation scalar leptoquarks in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D72**, 051107 (2005).
- [1145] D. Acosta, “Search for Higgs bosons decaying into $b\bar{b}$ and produced in association with a vector boson in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **95**, 051801 (2005).
- [1146] D. Acosta, “Search for long-lived doubly-charged Higgs bosons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **95**, 071801 (2005).
- [1147] D. Acosta, “Search for new physics using high mass tau pairs from 1.96 TeV $p\bar{p}$ collisions”, Phys. Rev. Lett. **95**, 131801 (2005).
- [1148] D. Acosta, “Search for scalar leptoquark pairs decaying to $\nu\bar{\nu}q\bar{q}$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, [Erratum: Phys. Rev.D71,119901(2005)], 112001 (2005).
- [1149] D. Acosta, “Search for ZZ and ZW production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, 091105 (2005).
- [1150] D. Acosta, “Study of jet shapes in inclusive jet production in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. **D71**, 112002 (2005).
- [1151] B. Aharmim, “Electron energy spectra, fluxes, and day-night asymmetries of B-8 solar neutrinos from measurements with NaCl dissolved in the heavy-water detector at the Sudbury Neutrino Observatory”, Phys. Rev. **C72**, 055502 (2005).
- [1152] M. Casarsa, S. C. Hsu, E. D. Lipeles, **M. Neubauer**, S. Sarkar, I. Sfiligoi, F. K. Wuerthwein, F. Wuerthwein, “The CDF analysis farm”, AIP Conf. Proc. **794**, 275–278 (2005).
- [1153] V. M. Abazov, “Combination of CDF and D0 Results on W Boson Mass and Width”, Phys. Rev. **D70**, 092008 (2004).
- [1154] D. Acosta, “Direct photon cross section with conversions at CDF”, Phys. Rev. **D70**, 074008 (2004).
- [1155] D. Acosta, “Inclusive double pomeron exchange at the Fermilab Tevatron $\bar{p}p$ collider”, Phys. Rev. Lett. **93**, 141601 (2004).
- [1156] D. Acosta, “Inclusive search for anomalous production of high p_T like-sign lepton pairs in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **93**, 061802 (2004).
- [1157] D. Acosta, “Measurement of the average time-integrated mixing probability of b -flavored hadrons produced at the Tevatron”, Phys. Rev. **D69**, 012002 (2004).
- [1158] D. Acosta, “Measurement of the polar-angle distribution of leptons from W boson decay as a function of the W transverse momentum in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. **D70**, 032004 (2004).
- [1159] D. Acosta, “Measurement of the $t\bar{t}$ production cross section in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV using dilepton events”, Phys. Rev. Lett. **93**, 142001 (2004).
- [1160] D. Acosta, “Observation of the narrow state $X(3872) \rightarrow J/\psi\pi^+\pi^-$ in $\bar{p}p$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 072001 (2004).
- [1161] D. Acosta, “Optimized search for single top quark production at the Fermilab tevatron”, Phys. Rev. **D69**, 052003 (2004).
- [1162] D. Acosta, “Search for doubly-charged Higgs bosons decaying to dileptons in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 221802 (2004).
- [1163] D. Acosta, “Search for Kaluza-Klein graviton emission in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ -TeV using the missing energy signature”, Phys. Rev. Lett. **92**, 121802 (2004).
- [1164] D. Acosta, “Search for pair production of scalar top quarks in R -parity violating decay modes in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, Phys. Rev. Lett. **92**, 051803 (2004).
- [1165] D. Acosta, “The underlying event in hard interactions at the Tevatron $\bar{p}p$ collider”, Phys. Rev. **D70**, 072002 (2004).
- [1166] D. Acosta, “Search for $B_s^0 \rightarrow \mu^+\mu^-$ and $B_d^0 \rightarrow \mu^+\mu^-$ decays in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, Phys. Rev. Lett. **93**, 032001 (2004).
- [1167] B. Aharmim, “Electron antineutrino search at the Sudbury Neutrino Observatory”, Phys. Rev. **D70**, 093014 (2004).

- [1168] S. N. Ahmed, “Constraints on nucleon decay via ‘invisible’ modes from the Sudbury Neutrino Observatory”, *Phys. Rev. Lett.* **92**, 102004 (2004).
- [1169] T. H. Kim, **M. Neubauer**, I. Sfiligoi, L. Weems, F. K. Wuerthwein, “The CDF Central Analysis Farm”, *IEEE Trans. Nucl. Sci.* **51**, 892–896 (2004).
- [1170] D. Acosta, “Central pseudorapidity gaps in events with a leading antiproton at the Fermilab Tevatron $\bar{p}p$ collider”, *Phys. Rev. Lett.* **91**, 011802 (2003).
- [1171] D. Acosta, “Measurement of prompt charm meson production cross sections in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev. Lett.* **91**, 241804 (2003).
- [1172] D. Acosta, “Measurement of the mass difference $m(D_s^+) - m(D^+)$ at CDF II”, *Phys. Rev.* **D68**, 072004 (2003).
- [1173] D. Acosta, “Search for a W' Boson Decaying to a Top and Bottom Quark Pair in 1.8 TeV $p\bar{p}$ Collisions”, *Phys. Rev. Lett.* **90**, 081802 (2003).
- [1174] D. Acosta, “Search for Associated Production of Υ and Vector Boson in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV”, *Phys. Rev. Lett.* **90**, 221803 (2003).
- [1175] D. Acosta, “Search for lepton flavor violating decays of a heavy neutral particle in $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, *Phys. Rev. Lett.* **91**, 171602 (2003).
- [1176] D. Acosta, “Search for Long-Lived Charged Massive Particles in $\bar{p}p$ Collisions at $\sqrt{s} = 1.8$ TeV”, *Phys. Rev. Lett.* **90**, 131801 (2003).
- [1177] D. Acosta, “Search for the flavor-changing neutral current decay $D^0 \rightarrow \mu^+\mu^-$ in $p\bar{p}$ collisions at $\sqrt{s} = 1.96$ TeV”, *Phys. Rev.* **D68**, 091101 (2003).
- [1178] D. Acosta, “Search for the supersymmetric partner of the top quark in dilepton events from $p\bar{p}$ collisions at $\sqrt{s} = 1.8$ TeV”, *Phys. Rev. Lett.* **90**, 251801 (2003).
- [1179] **M. S. Neubauer**, “Computing for Run II at CDF”, *Nucl. Instrum. Meth.* **A502**, 386–390 (2003).
- [1180] D. Acosta, “Cross Section for Forward J/ψ Production in $p\bar{p}$ Collisions at $\sqrt{s} = 1.8$ TeV”, *Phys. Rev.* **D66**, 092001 (2002).
- [1181] Q. R. Ahmad, “Direct evidence for neutrino flavor transformation from neutral current interactions in the Sudbury Neutrino Observatory”, *Phys. Rev. Lett.* **89**, 011301 (2002).
- [1182] Q. R. Ahmad, “Measurement of day and night neutrino energy spectra at SNO and constraints on neutrino mixing parameters”, *Phys. Rev. Lett.* **89**, 011302 (2002).
- [1183] A. B. McDonald, “Direct evidence for neutrino flavor transformation from neutral-current interactions in SNO”, *AIP Conf. Proc.* **646**, 43–58 (2002).
- [1184] A. W. P. Poon, “Neutrino observations from the Sudbury Neutrino Observatory”, *AIP Conf. Proc.* **610**, 218–230 (2002).
- [1185] A. W. P. Poon, “Solar neutrino observations at the Sudbury Neutrino Observatory”, *eConf* **C020805**, TTH01 (2002).
- [1186] Q. R. Ahmad, “Measurement of the rate of $\nu_e + d \rightarrow p + p + e^-$ interactions produced by 8B solar neutrinos at the Sudbury Neutrino Observatory”, *Phys. Rev. Lett.* **87**, 071301 (2001).
- [1187] A. B. McDonald, “Measurement of CC interactions produced by 8B solar neutrinos at SNO”, *PoS hep2001*, 186 (2001).
- [1188] **M. S. Neubauer**, “First solar neutrino observations from SNO”, *Int. J. Mod. Phys.* **A16S1B**, 715–717 (2001).
- [1189] J. Boger, “The Sudbury neutrino observatory”, *Nucl. Instrum. Meth.* **A449**, 172–207 (2000).

Conference Proceedings

- [1190] G. Kazakevich, R. Johnson, **M. Neubauer**, V. Lebedev, W. Schappert, V. Yakovlev, “Methods of phase and power control in magnetron transmitters for superconducting accelerators”, *Proceedings, 8th International Particle Accelerator Conference (IPAC 2017): Copenhagen, Denmark, May 14-19, 2017* (2017), THPIK122.
- [1191] T. Aaltonen, “Combined CDF and D0 Upper Limits on Standard Model Higgs Boson Production with up to 8.2 fb^{-1} of Data”, *Proceedings, 46th Rencontres de Moriond on Electroweak Interactions and Unified Theories: La Thuile, Italy, March 13-20, 2011* (2011).
- [1192] **M. S. Neubauer**, “A Fast Hardware Tracker for the ATLAS Trigger System”, *Particles and fields. Proceedings, Meeting of the Division of the American Physical Society, DPF 2011, Providence, USA, August 9-13, 2011* (2011).

- [1193] A. Annovi, “The Evolution of FTK, a Real-Time Tracker for Hadron Collider Experiments”, Astroparticle, particle and space physics, detectors and medical physics applications. Proceedings, 11th International Conference on Advanced Technology and Particle Physics, ICATPP 11, Como, Italy, October 5-9, 2009 (2010), pp. 653–657.
- [1194] T. Ciodaro, “Muon Detection Based on a Hadronic Calorimeter”, Proceedings, 2010 IEEE Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC 2010): Knoxville, Tennessee, October 30-November 6, 2010 (2010), pp. 772–776.
- [1195] M. Neubauer, “A Fast Hardware Tracker for the ATLAS Trigger System”, (2009).
- [1196] M. S. Neubauer, “A fast hardware tracker for the ATLAS trigger system”, Proceedings, 2009 IEEE Nuclear Science Symposium and Medical Imaging Conference (NSS/MIC 2009): Orlando, Florida, October 25-31, 2009 (2009), pp. 1876–1879.
- [1197] M. S. Neubauer, “Diboson physics at the Tevatron”, Proceedings, 41st Rencontres de Moriond, 2006 QCD and High Energy Hadronic Interactions: La Thuile, Val d’Aoste, Italy, Mar 18-25, 2006 (2006), pp. 113–116.
- [1198] A. Sill, “Globally distributed user analysis computing at CDF”, Computing in high energy physics and nuclear physics. Proceedings, Conference, CHEP’04, Interlaken, Switzerland, September 27-October 1, 2004 (2005), pp. 1139–1142.
- [1199] F. K. Wuerthwein, E. D. Lipeles, M. Neubauer, I. Sfiligoi, “The Condor based CDF CAF”, Computing in high energy physics and nuclear physics. Proceedings, Conference, CHEP’04, Interlaken, Switzerland, September 27-October 1, 2004 (2005), pp. 978–981.
- [1200] A. McDonald, “First results from the Sudbury Neutrino Observatory”, Neutrino oscillations. Proceedings, NO-VE International Workshop, Venice, Italy, July 24-26, 2001. c: C01/07/24.2 (2001), pp. 1–16.