

# Omega Cross-Section

---

Martin Sobotzik

Institute for Nuclear Physics  
Johannes Gutenberg-University of Mainz

Mainz, March 2019



2019-03-21



$\omega$  Cross-Section Studies

Omega Cross-Section

Martin Sobotzik  
Institute for Nuclear Physics  
Johannes Gutenberg University of Mainz

Mainz, March 2019





Ols Cross Section; Dip at about  $\cos(\theta) = -0.3$


# Increasing the Number of Bins




$\omega$  Cross-Section Studies

2019-03-21

Increasing the Number of Bins



Increased number of bins to 40; now there is still a dip at  $\cos(\theta) = -0.3$  but also a peak at  $\cos(\theta) = -0.4$



Increased number of bins to 40; now there is still a dip at  $\cos(\theta) = -0.3$  but also a peak at  $\cos(\theta) = -0.4$



# Comparing 10 Bin to 40 Bin Cross Section



$\omega$  Cross-Section Studies




2019-03-21

Comparing 10 Bin to 40 Bin Cross Section



Both Cross Sections are shown.

## Efficiency



Cross Section and efficiency. There is an efficiency drop at  $\cos(\theta) \approx -0.3$

# Taking a closer Look





## $\omega$ Cross-Section Studies

2019-03-21

### Taking a closer Look

$$\gamma p \rightarrow \omega \rightarrow \gamma \pi^0 p \rightarrow \gamma \gamma p$$

There is a 1:1 correlation between the polar angle of  $p$  and  $\omega$  for fixed  $E(\gamma)$ !



# Taking a closer Look





## $\omega$ Cross-Section Studies

2019-03-21

### Taking a closer Look

$$\gamma p \rightarrow \omega \rightarrow \gamma \pi^0 p \rightarrow \gamma \gamma p$$

There is a 1:1 correlation between the polar angle of  $p$  and  $\omega$  for fixed  $E(\gamma)$ !






Energy of Proton for  $\cos(\theta_\omega) = [-0.45, -0.35]$  (Peak)

A2

 $\omega$  Cross-Section Studies


2019-03-21

The energy of protons looks similar for MC and Data



Energy of protons for  $\cos(\theta_\omega) = [-0.45, -0.35]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

Energy of Proton for  $\cos(\theta_\omega) = [-0.45, -0.35]$  (Peak)



Energy of protons for  $\cos(\theta_\omega) = [-0.45, -0.35]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.




# $\theta$ of Proton for $\cos(\theta_\omega) = [-0.45, -0.35]$ (Peak)




A2

 $\omega$  Cross-Section Studies

2019-03-21

 $\theta$  of Proton for  $\cos(\theta_\omega) = [-0.45, -0.35]$  (Peak)


$\theta$  of protons for  $\cos(\theta_\omega) = [-0.45, -0.35]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



 $\theta_p$  (lab) $\theta_p$  (lab)

$\theta$  of protons for  $\cos(\theta_\omega) = [-0.45, -0.35]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.


Energy of Protons for  $\cos(\theta_\omega) = [-0.35, -0.25]$  (Dip) $\omega$  Cross-Section Studies

2019-03-21


Energy of Protons for  $\cos(\theta_\omega) = [-0.35, -0.25]$   
(Dip)



E (lab) Proton



E (lab) Proton






Energy of protons for  $\cos(\theta_\omega) = [-0.35, -0.25]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

$\theta$  of Proton for  $\cos(\theta_\omega) = [-0.35, -0.25]$  (Dip)


A2

 $\omega$  Cross-Section Studies

2019-03-21

 $\theta$  of Proton for  $\cos(\theta_\omega) = [-0.35, -0.25]$  (Dip) $\theta_p$  (lab) $\theta_p$  (lab)

$\theta$  of protons for  $\cos(\theta_\omega) = [-0.35, -0.25]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



$\theta$  of protons for  $\cos(\theta_\omega) = [-0.35, -0.25]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



Energy of Protons for  $\cos(\theta_\omega) = [0.35, 0.45]$  (Good)

A2

 $\omega$  Cross-Section Studies



2019-03-21

Energy of Protons for  $\cos(\theta_\omega) = [0.35, 0.45]$   
(Good)



E (lab) Proton

E (lab) Proton




Energy of protons for  $\cos(\theta_\omega) = [0.35, 0.45]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

$\theta$  of Proton for  $\cos(\theta_\omega) = [0.35, 0.45]$  (Good)




A2

 $\omega$  Cross-Section Studies

2019-03-21

 $\theta$  of Proton for  $\cos(\theta_\omega) = [0.35, 0.45]$  (Good)



$\theta$  of protons for  $\cos(\theta_\omega) = [0.35, 0.45]$ . Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

 $\theta_p$  (lab) $\theta_p$  (lab)

$\theta$  of protons for  $\cos(\theta_\omega) = [0.35, 0.45]$ . Red are MC and blue are beamtime data.



Left side is just measured, right side is after KFit.

Proton:  $\theta_{fit}$  vs.  $\theta_{gen}$



$\cos(\theta_{fit})$  vs.  $\cos(\theta_{gen})$  for all protons.

2019-03-21

 $\omega: \theta_{fit} \text{ vs. } \theta_{gen}$  $\cos(\theta_{fit}) \text{ vs. } \cos(\theta_{gen}) \text{ for all } \omega.$

# Unfolding

---

2019-03-21

ω Cross-Section Studies  
└ Unfolding

Unfolding

---



- $\mu$  is the *true* distribution given by nature
- detector effects are then described by the migration matrix  $R$ . (inefficiencies, bias and smearing)
- This results in the distribution  $\nu$ .

$$\nu_i = \sum_{j=1}^M R_{ij} \mu_j$$

- With infinite statistics, it would be possible to recover the original distribution by inverting the migration matrix

$$\mu = R^{-1} \nu$$

- Use numerical methods to invert the migration matrix



- $\mu$  is the *true* distribution given by nature
- detector effects are then described by the migration matrix  $R$ . (inefficiencies, bias and smearing)
- This results in the distribution  $\nu$ .

$$\nu_i = \sum_{j=1}^M R_{ij} \mu_j$$

- With infinite statistics, it would be possible to recover the original distribution by inverting the migration matrix

$$\mu = R^{-1} \nu$$

- Use numerical methods to invert the migration matrix



Example for Unfolding. Blue is true distribution. Red is measured distribution. Black Dots are the unfolded distribution. Green is the fit of the unfolded distribution



In region of  $\cos\theta(\omega)_{gen} \approx -0.35$ , see broader distribution of  $\cos\theta(\omega)_{fit} - i$   
simple 1D efficiency correction may not work well enough

Flat  $\omega$ ; Folded MC Data $\omega$  Cross-Section Studies

Unfolding

Flat  $\omega$ ; Folded MC Data

2019-03-21

Flat  $\omega$  was used. MC fitted data were unfolded.Flat  $\omega$  was used. MC fitted data were unfolded.

- Drop in the measured cross section at  $\cos(\theta_\omega) \approx -0.35$  is caused by a drop in the efficiency at that region
- Inefficiency is caused by the protons hitting the edge of the CB  
→ They are not reconstructed properly
- There are differences between MC and Data
- The differences are too big to make the Unfolding work