

Omega Cross-Section

Martin Sobotzik

Institute for Nuclear Physics
Johannes Gutenberg-University of Mainz

Mainz, March 2019



2019-03-21


ω Cross-Section Studies

Omega Cross-Section


Martin Sobotzik
Institute for Nuclear Physics
Johannes Gutenberg University of Mainz

Mainz, March 2019





ω Cross Section; Dip at about $\cos(\theta) = -0.3$




Increasing the Number of Bins




ω Cross-Section Studies

2019-03-21

Increasing the Number of Bins



Increased number of bins to 40; now there is still a dip at $\cos(\theta) = -0.3$ but also a peak at $\cos(\theta) = -0.4$



Increased number of bins to 40; now there is still a dip at $\cos(\theta) = -0.3$ but also a peak at $\cos(\theta) = -0.4$



Comparing 10 Bin to 40 Bin Cross Section



ω Cross-Section Studies




2019-03-21

Comparing 10 Bin to 40 Bin Cross Section



Both Cross Sections are shown.

Efficiency



Cross Section and efficiency. There is an efficiency drop at $\cos(\theta) \approx -0.3$

Taking a closer Look





ω Cross-Section Studies

2019-03-21

Taking a closer Look

$$\gamma p \rightarrow \omega \rightarrow \gamma \pi^0 p \rightarrow \gamma \gamma p$$

There is a 1:1 correlation between the polar angle of p and ω for fixed $E(\gamma)$!



Taking a closer Look





ω Cross-Section Studies

2019-03-21

Taking a closer Look

$$\gamma p \rightarrow \omega \rightarrow \gamma \pi^0 p \rightarrow \gamma \gamma p$$

There is a 1:1 correlation between the polar angle of p and ω for fixed $E(\gamma)$!






Energy of Proton for $\cos(\theta_\omega) = [-0.45, -0.35]$ (Peak)

A2


 ω Cross-Section Studies

2019-03-21

The energy of protons looks similar for MC and Data







Energy of protons for $\cos(\theta_\omega) = [-0.45, -0.35]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



Energy of protons for $\cos(\theta_\omega) = [-0.45, -0.35]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

ω Cross-Section Studies

2019-03-21

 θ of Proton for $\cos(\theta_\omega) = [-0.45, -0.35]$ (Peak) θ_p (lab) θ_p (lab)

θ of protons for $\cos(\theta_\omega) = [-0.45, -0.35]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.


Energy of Protons for $\cos(\theta_\omega) = [-0.35, -0.25]$ (Dip)

A2

 ω Cross-Section Studies



2019-03-21

Energy of Protons for $\cos(\theta_\omega) = [-0.35, -0.25]$
(Dip)




E (lab) Proton


E (lab) Proton



Energy of protons for $\cos(\theta_\omega) = [-0.35, -0.25]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



Energy of protons for $\cos(\theta_\omega) = [-0.35, -0.25]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



θ of protons for $\cos(\theta_\omega) = [-0.35, -0.25]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.




ω Cross-Section Studies

2019-03-21

θ of Proton for $\cos(\theta_\omega) = [-0.35, -0.25]$ (Dip)

θ of Proton for $\cos(\theta_\omega) = [-0.35, -0.25]$ (Dip)

A2

 θ_p (lab) θ_p (lab)

θ of protons for $\cos(\theta_\omega) = [-0.35, -0.25]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.



Energy of Protons for $\cos(\theta_\omega) = [0.35, 0.45]$ (Good)

A2

 ω Cross-Section Studies



2019-03-21

Energy of Protons for $\cos(\theta_\omega) = [0.35, 0.45]$
(Good)



E (lab) Proton

E (lab) Proton




Energy of protons for $\cos(\theta_\omega) = [0.35, 0.45]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

θ of Proton for $\cos(\theta_\omega) = [0.35, 0.45]$ (Good)




A2

 ω Cross-Section Studies

2019-03-21

 θ of Proton for $\cos(\theta_\omega) = [0.35, 0.45]$ (Good)



θ of protons for $\cos(\theta_\omega) = [0.35, 0.45]$. Red are MC and blue are beamtime data. Left side is just measured, right side is after KFit.

 θ_p (lab) θ_p (lab)

θ of protons for $\cos(\theta_\omega) = [0.35, 0.45]$. Red are MC and blue are beamtime data.



Left side is just measured, right side is after KFit.

Proton: θ_{fit} vs. θ_{gen}



$\cos(\theta_{fit})$ vs. $\cos(\theta_{gen})$ for all protons.

2019-03-21

 $\omega: \theta_{fit} \text{ vs. } \theta_{gen}$  $\cos(\theta_{fit}) \text{ vs. } \cos(\theta_{gen}) \text{ for all } \omega.$

Unfolding

2019-03-21

ω Cross-Section Studies
└ Unfolding

Unfolding

- μ is the *true* distribution given by nature
- detector effects are then described by the migration matrix R .
(inefficiencies, bias and smearing)
- This results in the distribution ν .

$$\nu_i = \sum_{j=1}^M R_{ij} \mu_j$$

- With infinite statistics, it would be possible to recover the original distribution by inverting the migration matrix

$$\mu = R^{-1} \nu$$

- Use numerical methods to invert the migration matrix


- μ is the *true* distribution given by nature
- detector effects are then described by the migration matrix R .
(inefficiencies, bias and smearing)
- This results in the distribution ν .

$$\nu_i = \sum_{j=1}^M R_{ij} \mu_j$$


- With infinite statistics, it would be possible to recover the original distribution by inverting the migration matrix

$$\mu = R^{-1} \nu$$

- Use numerical methods to invert the migration matrix



Example for Unfolding. Blue is true distribution. Red is measured distribution. Black Dots are the unfolded distribution. Green is the fit of the unfolded distribution



Example for Unfolding. Blue is true distribution. Red is measured distribution. Black Dots are the unfolded distribution. Green is the fit of the unfolded distribution

In region of $\cos\theta(\omega)_{gen} \approx -0.35$, see broader distribution of $\cos\theta(\omega)_{fit} - i$
simple 1D efficiency correction may not work well enough

Flat ω ; Folded MC Data



A2

 ω Cross-Section Studies


└ Unfolding

└ Flat ω ; Folded MC Data

2019-03-21

Flat ω was used. MC fitted data were unfolded.Flat ω was used. MC fitted data were unfolded.

Flat ω ; Folded MC Data



Flat ω was used. MC fitted data were unfolded.

- ω are generated with a flat phase space
- A migration matrix is calculated
- This migration matrix is then used to unfold realistic MC
- Unfortunately, the unfolded cross section does not have the desired shape
→ Unfolding was unsuccessful

ω Cross-Section Studies

Unfolding

Flat ω ; Folded MC Data


2019-03-21

- Drop in the measured cross section at $\cos(\theta_\omega) \approx -0.35$ is caused by a drop in the efficiency at that region
- Inefficiency is caused by the protons hitting the edge of the CB
→ They are not reconstructed properly
- There are differences between MC and Data
- The differences are too big to make the Unfolding work

- Drop in the measured cross section at $\cos(\theta_\omega) \approx -0.35$ is caused by a drop in the efficiency at that region
- Inefficiency is caused by the protons hitting the edge of the CB
→ They are not reconstructed properly
- There are differences between MC and Data
- The differences are too big to make the Unfolding work

2019-03-21

Backup

Flat generated ω 