

# Systematic Studies for the $\pi^0$ Calibration of the Crystal-Ball Detector

Martin Sobotzik

Johannes-Gutenberg Universität Mainz

xx.yy.2017

# Inhaltsverzeichnis

- 1 Motivation
- 2 Preparation
- 3 Studies
- 4 Further Results

# The Process

$$\gamma + p \rightarrow \pi^0 + p \rightarrow p + \gamma_1 \gamma_2 \quad (1)$$

$$m_{\pi^0} = \sqrt{2E_1 E_2 (1 - \cos(\alpha))} \quad (2)$$

# The Process

$$\gamma + p \rightarrow \pi^0 + p \rightarrow p + \gamma_1 \gamma_2 \quad (1)$$

$$m_{\pi^0} = \sqrt{2E_1 E_2 (1 - \cos(\alpha))} \quad (2)$$

- Is there an energy dependency in the CB-Detector and how can it be checked?

# The Process

$$\gamma + p \rightarrow \pi^0 + p \rightarrow p + \gamma_1 \gamma_2 \quad (1)$$

$$m_{\pi^0} = \sqrt{2E_1 E_2 (1 - \cos(\alpha))} \quad (2)$$

- Is there an energy dependency in the CB-Detector and how can it be checked?  
 $\rightarrow |E_1 - E_2| < 25 \text{ MeV}$

# The Process

$$\gamma + p \rightarrow \pi^0 + p \rightarrow p + \gamma_1 \gamma_2 \quad (1)$$

$$m_{\pi^0} = \sqrt{2E_1 E_2 (1 - \cos(\alpha))} \quad (2)$$

- Is there an energy dependency in the CB-Detector and how can it be checked?  
 $\rightarrow |E_1 - E_2| < 25 \text{ MeV}$
- What are the reasons for the dependency?

# Crystal-Ball-Function / Reduction of the Underground

- Gaussian Fit Function  $\rightarrow$  Crystal-Ball Fit Function

# Crystal-Ball-Function / Reduction of the Underground

- Gaussian Fit Function  $\rightarrow$  Crystal-Ball Fit Function
- Check if the registered particles are charged  
 $\rightarrow$  Reduction of the underground



# Crystal-Ball-Function / Reduction of the Underground

- Gaussian Fit Function  $\rightarrow$  Crystal-Ball Fit Function
- Check if the registered particles are charged  
 $\rightarrow$  Reduction of the underground




Abbildung: Beamtime: Example for not reduced and reduced underground

# Event Generator

Reasons for a new event generator:

# Event Generator

Reasons for a new event generator:

- The condition  $|E_1 - E_2| < 25 \text{ MeV}$  is a strong cut  
→ There is no package with enough events

# Event Generator

Reasons for a new event generator:

- The condition  $|E_1 - E_2| < 25 \text{ MeV}$  is a strong cut  
→ There is no package with enough events
- Creating a new package with enough events would take too much time (multiple days on blaster)

# Event Generator

Reasons for a new event generator:

- The condition  $|E_1 - E_2| < 25 \text{ MeV}$  is a strong cut  
→ There is no package with enough events
- Creating a new package with enough events would take to much time (multiple days on blaster)
- It would be better if the same generator is used for all studies  
→ The generator should be able to simulate MAMI-Beam and isotropic decay

# Event Generator

# No Additional Cut

- Beamtime October 2014
- No additional cut





Abbildung: Beamtime: No additional cut

# Detectors On At The Edge

- Beamtime October 2014
- Neglect the detectors at the edge



**Abbildung:** Beamtime: With and without considerations of the detectors on the edge of the beam entrance and exit



# Simulation

# Hot and Dead Crystals