Bhabha Tracking Efficiencies

Martin Sobotzik

01.03.2019

Johannes Gutenberg Universität Mainz

Motivation


- I want to study the tracking efficiency of Bhabha processes
- For this I first take a look at electrons which are only detected by the ECL
- They are therefor labeled and treated as gammas
- In the next step I only take information from the ECL and count the electrons there without the information from the tracking detectors
- Afterwards the number of tracks in the tracking detectors is count to get the efficiency
- This idea comes from some plots presented by Sam in previous tracking and ECL meeting.


Getting Started

- All cuts were taken from Sam's studies:
 - gamma:probe '(E > 0.1)'
 - gamma:tag '(clusterE > 3.0)'
 - vpho:cand 'reconstructed from gamma:probe and gamma:tag'
 - $0.296706 < \theta < 2.61799 \rightarrow$ It has to hit the ECL
 - nCleanedTracks[abs(dz) < 2.0 and abs(dr) < 0.5 and nCDCHits > 0 and pt > 0.15] < 1 \rightarrow bad quality hits
 - $M(vpho) > 8.0\,{\rm GeV} \to {\sf To}$ cut away background (not from his email but surely he is using something like that)

Reproducing Plots


- Sam used Prod6 on all of his plots, and since I try to reproduce his plots I will use it as well
- Sam used all Prod6 data. I am only using the following: /hsm/belle2/bdata/Data/release-02-01-00/DB00000438 /prod00000006/e0003/4S/r02*/all/mdst.sub00/*.root
- Sam's plots are on the left.
- \bullet Three lines. The middle one is $\gamma\gamma$ the two others are ee






Reproducing Plots


Same plots but zoomed in:






Reproducing Plots


- ullet The middle peak is $\gamma\gamma$ the two other peaks are ee
- My $\gamma\gamma$ peak is way higher (Maybe there are some cuts that I am not considering?)






Some more plots


Here is no Mass cut on the vpho



Some more plots



Some more plots



The next Steps

- Investigate the best cuts for our purposes and apply them
- Select only the ee candidates and study the tracking efficiency using that sample
- Producing the same plots shown here using MC sample (from a preliminary study using MC10 it looks like very few events survive applying the descibed selection)