

INVOLUTIVE PARTIALLY-ORDERED SEMILATTICES

PETER JIPSEN AND MELISSA SUGIMOTO

ABSTRACT. We give a full description of the structure of finite posets with an involutive residuated binary operation that is associative, commutative and idempotent. A duality between Plonka sums of generalized Boolean algebras and semilattice direct systems of sets and partial functions is used to give a logarithmically smaller description of these partially ordered algebras. The results specialize to commutative idempotent involutive residuated lattices.

1. INTRODUCTION

2. BACKGROUND ON INVOLUTIVE PO-SEMIGROUPS

A *residuated po-semigroup* $(A, \leq, \cdot, \backslash, /)$ is a partially-ordered set (A, \leq) with an associative binary operation \cdot and two *residuals* that satisfy

$$(\text{res}) \quad xy \leq z \iff x \leq z/y \iff y \leq x \backslash z$$

for all $x, y, z \in A$. The first equivalence ensures that \cdot is order-preserving in the first argument ($x \leq y \leq yz/z \implies xz \leq yz$), and likewise the second equivalence implies order-preservation in the second argument. A *residuated lattice-ordered semigroup* $(A, \wedge, \vee, \cdot, \backslash, /)$ (or *rl-semigroup* for short) is a residuated po-semigroup for which the partial order is a lattice order. If A contains a constant 1 that is an identity element for \cdot then A is called a *residuated lattice*.

A *residuated po-semilattice* is a residuated po-semigroup in which the multiplication is commutative and idempotent.

An *involutive partially-ordered semigroup* or *ipo-semigroup* is of the form $(A, \leq, \cdot, \sim, -)$ such that (A, \leq) is a poset, \cdot is an associative operation and the *left and right linear negations* $\sim, -$ are an *involutive pair*, i.e., $\sim \sim x = x = - -x$, $x \leq \sim y \iff y \leq -x$ and

$$(\text{ires}) \quad xy \leq z \iff x \leq -(y \cdot \sim z) \iff y \leq \sim(-z \cdot x)$$

holds for all $x, y, z \in A$. It follows that the operation \cdot is order-preserving in each argument, and that $\sim, -$ are both order-reversing.

The axiom (ires) shows that $z/y = -(y \cdot \sim z \text{ and } x \setminus z = \sim(-z \cdot x))$, but it is often convenient to use the equivalent formulation

$$(\text{ires}') \quad xy \leq z \iff y \cdot \sim z \leq \sim x \iff -z \cdot x \leq -y$$

which says that the three variables can be rotated left (resp. right) cyclically as long as the left (resp. right) linear negation is applied to the variables that move across the inequality symbol.

If (A, \leq) is a lattice, with join \vee and meet \wedge , then $(A, \vee, \wedge, \cdot, \sim, -)$ is an *involutive lattice-ordered semigroup* or *il-semigroup*, and in this case it follows that $x(y \vee z) = xy \vee xz$ and $(x \vee y)z = xz \vee yz$.

An operation \cdot is *idempotent* if it satisfies $x \cdot x = x$ and *commutative* if $xy = yx$.

Lemma 1. *If \cdot is idempotent, then (ires) or (ires') imply $\sim -x = x = -\sim x$ and $x \leq \sim y \iff y \leq -x$.*

Proof. (to be inserted) □

In a commutative idempotent ipo-semigroup there is another semi-lattice order, called the *multiplicative order*, that is defined by

$$x \sqsubseteq y \iff xy = x.$$

Therefore these algebras are called *ipo-semilattices*. Note that commutativity implies $\sim x = -x$, hence by Lemma 1 ipo-semilattices can be defined succinctly as partially ordered algebras $(A, \leq, \cdot, -)$ such that (A, \cdot) is a semilattice and

$$(\text{cres}) \quad xy \leq z \iff x \leq -(y \cdot -z) \quad \text{for all } x, y, z \in A$$

or equivalently

$$(\text{cres}') \quad xy \leq z \iff y \cdot -z \leq -x \quad \text{for all } x, y, z \in A.$$

It then follows that $-$ is an order-reversing bijection and $--x = x$, hence it is natural to define the operation $x + y = -(-x \cdot -y)$.

The most prominent examples of ipo-semilattices are Boolean algebras. They arise as the case when the partial order \leq and the semi-lattice order \sqsubseteq coincide. Usually Boolean algebras are defined with constant operations 0 and 1, but in this special case the terms $x \cdot -x$ and $-(x \cdot -x)$ are constant unary operations that fulfill the same role.

An *involutive residuated poset* is an ipo-semigroup with a multiplicative identity element 1, in which case $\sim 1 = -1$, and this element is denoted 0. Similarly, an *involutive residuated lattice* is an il-semigroup with a multiplicative identity 1. Note that an ipo-semilattice has a multiplicative identity if and only if it has a top element in the multiplicative order.


# of elem. $n =$	1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16
CIdInRL	1	1	1	2	2	4	4	9	10	21	22	49	52	114	121	270
CIdInRP	1	1	1	2	2	4	4	9	10	22	24	53	61	134	157	343
$i\ell$ -semilattices	1	1	1	3	4	10	17	42	80	191						
ipo-semilattices	1	1	1	3	4	10	17	43	82	201						

TABLE 1. Number of algebras up to isomorphism

The aim of this paper is to give a full description of the structure of finite commutative idempotent $i\ell$ -semigroups ($i\ell$ -semilattices for short). For finite commutative idempotent involutive residuated lattices such a description has been given in [3].

The number of algebras with n elements (up to isomorphism) for the classes mentioned above have been calculated with Mace4 [4] and are given in Table 1.

The smallest $i\ell$ -semilattice that is not a commutative idempotent involutive residuated lattice has 4 elements. Its lattice and multiplicative order are show in Figure 1.

FIGURE 1. The smallest $i\ell$ -semilattice that does not have an identity.

The smallest ipo-semilattice that is not lattice-ordered has 8 elements. Its poset and multiplicative order are show in Figure 2.

3. STRUCTURAL PROPERTIES OF IPO-SEMILATTICES

It is an interesting project to adapt the arguments from [3] to the more general setting of $i\ell$ -semilattices.

The first step is to prove that ipo-semilattices are disjoint unions of Boolean algebras.

Lemma 2. *Let \mathbf{A} be a residuated po-semilattice and suppose $x \setminus x = y \setminus y$ for $x, y \in A$. Then*

- (1) $x \sqsubseteq y \iff x \leq y$,
- (2) $x \setminus x = xy \setminus xy$,




FIGURE 2. The smallest ipo-semilattice that is not lattice-ordered.

- (3) if $y \setminus y = z \setminus z$ then $x \setminus x = yz \setminus yz$, and
- (4) if $y \sqsubseteq z \sqsubseteq x \setminus x$ then $x \setminus x = z \setminus z$.

Proof. (1) Assume $x \setminus x = y \setminus y$ and $x \sqsubseteq y$. Then $xy = x$, and by idempotence $xx = x$ implies $x \leq x \setminus x$. Hence $x \leq y \setminus y$, and it follows that $xy \leq y$. Since $xy = x$ we deduce $x \leq y$.

Conversely, assume $x \setminus x = y \setminus y$ and $x \leq y$. By idempotence $yy = y$, hence $y \leq y \setminus y = x \setminus x$, and it follows that $xy \leq x$. Again by idempotence $x = xx \leq xy$ since \cdot is order-preserving and $x \leq y$. Therefore $xy = x$, or equivalently, $x \sqsubseteq y$.

(2) Assume $x \setminus x = y \setminus y$. Then $x \setminus x \leq y \setminus y$. Equivalently, $y(x \setminus x) \leq y$, and since we have order-preservation and associativity, $xy(x \setminus x) \leq xy$. Hence $x \setminus x \leq xy \setminus xy$.

For the reverse inequality, $xy \setminus y \leq xy \setminus y$ implies $xy(xy \setminus y) \leq y$. Using associativity and commutativity, we obtain $x(xy \setminus y) \leq y \setminus y = x \setminus x$. Hence $xx(xy \setminus y) \leq x$. Using idempotence and substituting xy for y , it follows that $x(xxy \setminus xy) \leq x$ and therefore $xy \setminus xy \leq x \setminus x$.

(3) holds since (2) implies that $y \setminus y = yz \setminus yz$.

(4) We will need that $z(z \setminus z) = z$. By idempotence, we obtain that $z \leq z \setminus z$, and then that $z \leq z(z \setminus z)$. For the reverse inequality, $z \setminus z \leq z \setminus z$, and equivalently, $z(z \setminus z) \leq z$. Thus we have the desired equality. Now assume $y \sqsubseteq z \sqsubseteq x \setminus x$, hence $yz = y$ and $z(x \setminus x) = z$, and therefore $x \setminus x \leq z \setminus z$. Furthermore, $yz(z \setminus z) = yz$, so $y(z \setminus z) = y$, and consequently $z \setminus z \leq y \setminus y = x \setminus x$. It follows that $x \setminus x = z \setminus z$. \square

Define an equivalence relation \equiv on A by $x \equiv y \iff x \setminus x = y \setminus y$. Part (1) of the previous lemma shows that the partial order \leq and the semilattice order \sqsubseteq agree on each equivalence class of \equiv . Define the notation

$$1_x = x \setminus x.$$

Lemma 3. *Let \mathbf{A} be a residuated po-semilattice and let $x \in A$. Then*

- (1) $x(x \setminus x) = x$,
- (2) $(x \setminus x) \setminus (x \setminus x) = x \setminus x$,
- (3) *each equivalence class of \equiv is a meet-semilattice $([x]_{\equiv}, \cdot)$ with identity element 1_x as top element of $[x]_{\equiv}$,*
- (4) $x \sqsubseteq y$ *implies* $1_y \leq 1_x$,
- (5) $1_x \cdot 1_y \leq 1_{xy}$.

Proof. (1) We have $x(x \setminus x) \leq x$ and $x \leq x \setminus x$ from (res) and idempotence. Hence $x = xx \leq x(x \setminus x)$ and thus equality holds.

(2) From $y(y \setminus y) \leq y$ and associativity we deduce $xy(y \setminus y) \leq xy$ and therefore $y \setminus y \leq xy \setminus xy$. Replacing y by $x \setminus x$ and making use of (1) produces $(x \setminus x) \setminus (x \setminus x) \leq x \setminus x$. The reverse inequality follows from idempotence.

(3) We first note that (2) implies 1_x is a member of $[x]_{\equiv}$. It is the top element of the equivalence class of x since $x \equiv y$ implies $y \leq y \setminus y = x \setminus x = 1_x$. Hence $y \sqsubseteq 1_x$ and therefore $y1_x = y$ for all $y \in [x]_{\equiv}$.

Part (3) of the previous lemma shows that each equivalence class is closed under multiplication ($y, z \in [x]_{\equiv}$ implies $yz \in [x]_{\equiv}$).

(4) In a residuated po-semigroup we always have $y(y \setminus y) \leq y$, hence $xy(y \setminus y) \leq xy$. If we assume $xy = x$ then it follows that $x(y \setminus y) \leq x$ and therefore $1_y = y \setminus y \leq x \setminus x = 1_x$.

(5) Since $xy \sqsubseteq x, y$ we obtain $1_x, 1_y \leq 1_{xy}$ from (4). Hence $1_x \cdot 1_y \leq 1_{xy} \cdot 1_{xy} = 1_{xy}$. \square

Hence if A is finite, each class $[x]_{\equiv}$ also has a least element, denoted by 0_x . Lemma 2(4) shows that each class is an interval, i.e., $[x]_{\equiv} = \{y \in A \mid 0_x \sqsubseteq y \sqsubseteq 1_x\} = \{y \in A \mid 0_x \leq y \leq 1_x\}$.

We now specialize to *involutive* po-semilattices. In this case $1_x = x \setminus x = -(x \cdot -x)$ and we define

$$0_x = -1_x = x \cdot -x.$$

By Lemma 3(3) 1_x is the top element of $[x]_{\equiv}$, and the lemma below shows that 0_x is the bottom element, hence each equivalence class has a bottom element even without the assumption of finiteness.

For an ipo-semilattice A , define the term $x + y = -(-x \cdot -y)$, so $1_x = x + -x$. Let $\mathbb{B}_x = \{a \in A \mid 0_x \sqsubseteq a \sqsubseteq 1_x\}$. By the previous lemma $y \in \mathbb{B}_x \iff 0_x = 0_y$, and if $0_x = 0_y$ then $x \sqsubseteq y \iff x \leq y$.

The next lemma shows that ipo-semilattices have many interesting structural properties.

Lemma 4. *Let \mathbf{A} be an ipo-semilattice. Then for all $x, y, z \in A$*

- (1) $\mathbb{B}_x = [x]$ is closed under linear negation, i.e., $y \in \mathbb{B}_x \implies -y \in \mathbb{B}_x$, hence 0_x is the bottom element of \mathbb{B}_x ,
- (2) $x \sqsubseteq y$ implies $0_x \sqsubseteq 0_y$,
- (3) $0_x \sqsubseteq 0_y \iff 0_x \leq 0_y \iff 1_y \leq 1_x$,
- (4) $x \leq x \cdot 1_y$,
- (5) $x \sqsubseteq y$ implies $1_x \sqsubseteq 1_y$,
- (6) $x = 1_x$ and $y = 1_y$ imply $xy = 1_{xy}$,
- (7) $1_x \cdot y = 0_x$ implies $y = 0_x$,
- (8) $x(y + z) \leq xy + xz$ and $(x + y)(x + z) \leq x + yz$,
- (9) $0_{xy} \leq 0_x \cdot 0_y$,
- (10) $x, y \sqsubseteq z$ implies $0_x \leq 1_y$, and
- (11) $0_x \leq 1_y$ implies $0_x \cdot 0_y = 0_{xy}$.

Proof. (1) To see that 0_x is a member of $[x]$, note that $0_x \setminus 0_x = -(x \cdot -x \cdot (x \setminus x)) = -(x(x \setminus x) \cdot -x) = -(x \cdot -x)$ by commutativity and Lemma 3(1), hence $0_x \setminus 0_x = x \setminus x$.

Now Lemma 2(4) implies $\mathbb{B}_x \subseteq [x]$. Since 1_x is the largest element of $[x]$ with respect to \leq , it follows that 0_x is the smallest, hence $[x] \subseteq \mathbb{B}_x$.

If $y \in [x]$ then $x \setminus x = y \setminus y$, hence $-(x \cdot -x) = -(y \cdot -y)$. This equation still holds if we replace y by $-y$ since commutativity and $--y = y$ hold. Therefore $x \setminus x = -y \setminus -y$, hence $-y \in [x]$.

For (2), we have that $-(x \cdot y) \leq -(x \cdot y)$ by reflexivity. Using commutativity and the residuation property, $x \cdot -(x \cdot y) \leq -y$. But, $x \cdot y = x$, so it follows that $x \cdot -x \leq -y$, and by idempotence, $x \cdot -x \leq x \cdot -x \cdot -y$. Again, using commutativity, associativity and $x \cdot y = x$, we obtain $x \cdot -x \leq x \cdot -x \cdot y \cdot -y$, or equivalently $0_x \leq 0_x \cdot 0_y$.

To show the reverse inequality, we note that $0_y \leq y$ since \leq and \sqsubseteq coincide on \mathbb{B}_y . Then $y \cdot -y \leq y$, and by order preservation, $y \cdot -y \cdot (x \cdot -x) \leq y \cdot (x \cdot -x)$. By associativity and commutativity, $y \cdot -y \cdot x \cdot -x \leq x \cdot y \cdot -x$, and hence $y \cdot -y \cdot x \cdot -x \leq x \cdot -x$. Consequently $0_x \cdot 0_y = 0_x$.

(3) First note that $0_x \cdot -0_x = 0_x \cdot 1_x = 0_x$, hence $0_{0_x} = 0_x$. Now the forward direction follows from (2) since $0_x \sqsubseteq 0_y$ implies $0_x = 0_{0_x} \leq 0_{0_y} = 0_y$.

For the reverse direction, assume $0_x \leq 0_y$. Since $0_y \leq y$ and $0_y \leq -y$, we have $0_x \leq y, -y$. By idempotence $0_x \leq y0_x$ and $0_x \leq -y0_x$. Involution implies $-y, y \leq 1_x$, hence $-y0_x \leq 1_x0_x = 0_x$ and similarly $y0_x \leq 0_x$. Thus equality holds in both cases and we get $0_x0_y = 0_x \cdot y \cdot -y = 0_x \cdot -y = 0_x$, i.e., $0_x \sqsubseteq 0_y$.

(4) First we observe that $x \cdot 1_y \leq x \cdot 1_y$ implies $x \cdot -(x \cdot 1_y) \leq -1_y = 0_y \leq y$, hence by idempotence $x \cdot -(x \cdot 1_y) \leq xy$. Replacing x by $x \cdot -(x \cdot 1_y)$ we obtain

$$x \cdot -(x \cdot 1_y) \cdot -(x \cdot -(x \cdot 1_y) \cdot 1_y) \leq x \cdot -(x \cdot 1_y)y \leq -x.$$

where the last inequality follows from (cres'), idempotence and $y \leq 1_y$. The term on the left is $x \cdot -(x \cdot 1_y) \cdot 1_{x1_y} = x \cdot -(x \cdot 1_y)$, hence it follows that $x \cdot -(x \cdot 1_y) \leq -x$. Applying (cres') and idempotence once more, we get $x \leq x \cdot 1$.

(5) From $x \sqsubseteq y$ it follows that $1_y \leq 1_x$ by (2) and (3). Multiplying by 1_x on both sides and using idempotence produces $1_x 1_y \leq 1_x$. The reverse inequality always holds by (4).

(6) Assume $x = 1_x$ and $y = 1_y$. By Lemma 3(5) we have $xy \leq 1_{xy}$. To prove $1_{xy} \leq xy$ it suffices to show $-(xy) \leq xy \cdot -(xy)$. By (4) $z \leq z1_x = xz$ and $z \leq z1_y = yz$, hence $-(xy) \leq y \cdot -(xy) \leq xy \cdot -(xy)$.

(7) Assume $1_x \cdot y = 0_x$. By (cres') we have $1_x \cdot 1_x \leq -y$, hence $y \leq 0_x$. It follows that $0_y \leq y \leq 0_x \leq x, -x$. From $y \cdot -y \leq -x$ we deduce $xy \leq y$, and from $y \cdot -y \leq x$ we $-x \cdot y \leq y$. Now multiplying the assumption by y on both sides gives

$$1_x \cdot y = 0_x \cdot y = x \cdot -x \cdot y \leq xy \leq y.$$

Therefore $0_x = 1_x \cdot y \leq y$, and by antisymmetry we have $y = 0_x$.

(8) Using the definition of $+$, we need to show that $x \cdot -(-y \cdot -z) \leq -(-(xy) \cdot -(xz))$, or equivalently by (cres') $x \cdot -(xy) \cdot -(xz) \leq -y \cdot -z$.

From $xy \leq xy$ we deduce that $x \cdot -(xy) \leq -y$, hence left-multiplication by x and idempotence give $x \cdot -(xy) \leq x \cdot -y$. Multiplying on the right by $-(xz)$ produces $x \cdot -(xy) \cdot -(xz) \leq x \cdot -y \cdot -(xz) = -y \cdot x \cdot -(xz) \leq -y \cdot -z$.

The second inequality follows by De Morgan's laws: $-(xy + xz) \leq -(x(y + z))$, hence $(-x + -y)(-x + -z) = -x + -y \cdot -z$. Finally replace x, y, z by $-x, -y, -z$ and simplify.

(9) First we use (8) to see that $xy(x+y) \leq xyx + xyy = xy + xy = xy$. Hence $-(xy) \leq -(xy(x+y)) = -(xy - (-x \cdot -y))$. Multiplying by xy on both sides we obtain

$$xy \cdot -(xy) \leq xy \cdot -(xy - (-x \cdot -y)) \leq xy(-x \cdot -y),$$

where the last inequality holds because $xy \cdot -(xy - (-x \cdot -y)) \leq -x \cdot -y$ by (cres'). Therefore $0_{xy} \leq 0_x 0_y$.

(10) Assume $x, y \sqsubseteq z$, i.e., $xz = x$ and $yz = y$. Since $z \cdot -z = 0_z \leq z$ we have $xz \cdot -z \leq xz$ and hence $x \cdot -z \leq x$. From (cres') we obtain $x \cdot -x \leq z$.

Using (cres') with $yz \leq yz$ gives $y \cdot -(yz) \leq -z$, and using $yz = y$ this simplifies to $y \cdot -y \leq -z$ or equivalently $z \leq -(y \cdot -y)$. By transitivity we have $0_x \leq z \leq 1_y$.

(11) Assume $0_x \leq 1_y$, i.e., $x \cdot -x \leq -(y \cdot -y)$. By two applications of (cres') this is equivalent to $xy \cdot -y \leq x$ and $-x \cdot -y \leq -(xy)$.

Multiplying by xy on both sides produces $0_x \cdot 0_y \leq 0_{xy}$. The reverse inequality follows from (9). \square

The example in Figure 1 shows that the assumptions in Lemma 4(10), (11) are needed. In this algebra, $0_a = 1_a = a$, $0_b = 1_b = b$, $0_\perp = 0_\top = \perp$ and $1_\perp = 1_\top = \top$. Hence $0_a \not\leq 1_b$ and $0_a \cdot 0_b = ab = \top \neq 0_{ab} = 0_\top = \perp$.

Theorem 5. *The semilattice intervals $(\mathbb{B}_x, \cdot, +, -, 0_x, 1_x)$ are Boolean algebras and they partition A .*

Proof. Recall that the monoid order \sqsubseteq and the partial order \leq agree on all elements of \mathbb{B}_x for fixed x , so for all $y, z \in \mathbb{B}_x$,

$$y \leq z \iff y \sqsubseteq z \iff yz = y \iff y + z = z.$$

The involution $-$ shows that (\mathbb{B}_x, \cdot) and $(\mathbb{B}_x, +)$ are dually isomorphic semilattices with respect to the same underlying partial order \leq , hence $(\mathbb{B}_x, \cdot, +)$ is a lattice. It is complemented since $0_x = y \cdot -y$ and $1_x = y + -y$ for all y . To show it is a Boolean algebra, it remains to prove the distributive law $w(y+z) = wy + wz$. By Lemma 4(8) the inequality $w(y+z) \leq wy + wz$ holds for all $w, y, z \in A$. The reverse inequality is true for $w, y, z \in \mathbb{B}_x$ since in that case we have $wy, wz \sqsubseteq w(y+z)$, hence $wy + wz \sqsubseteq w(y+z)$, or equivalently $wy + wz \leq w(y+z)$. \square

It is an open problem to characterize the posets that are reducts of involutive po-semigroups, even for idempotent and/or finite involutive residuated posets.

Note that any downward-closed set in the multiplicative order of an ipo-semilattice that is a union of Boolean components is again the multiplicative order of some ipo-semilattice. However the diagram in Figure 2 above shows that the same result does not hold for il -semilattices. In this example the multiplicative order \sqsubseteq is a downset of the multiplicative order of $S_3 \times S_3$, where S_3 is the (unique) 3-element il -semilattice (also known as the 3-element Sugihara monoid). However the 8-element example is only an ipo-semilattice rather than an ie -semilattice.

4. SEMILATTICE DIRECT SYSTEMS AND PŁONKA SUMS

In this section we prove that the multiplicative order \sqsubseteq of a finite ipo-semilattice is a Płonka sum of a semilattice direct system of generalized Boolean algebras. In the next section we show that the partial order \leq is determined by the multiplicative order. Hence the structure of finite ipo-semilattices is in one-one correspondence with semilattice direct systems of generalized Boolean algebras. We first recall that

generalized Boolean algebras are the variety **GBA** of residuated lattices generated by the 2-element residuated lattice $\mathbf{2}_r$. An equational basis for this variety is given by the identities for residuated lattices together with $xy = x \wedge y$ and $(y \setminus x)/y = x \vee y$ ([2, Prop. 3.23]. Finite members of **GBA** are reducts of Boolean algebras without complementation or a constant designating the bottom element. Homomorphisms between GBAs are residuated lattice homomorphisms, hence they preserve the top element 1, join, meet, and residuals, but not necessarily complementation or 0.

A *semilattice direct system* (or *sd-system* for short) is a triple $\mathbf{B} = (\mathbf{B}_i, h_{ij}, I)$ such that

- I is a semilattice,
- $\{\mathbf{B}_i : i \in I\}$ is a family of algebras of the same type with disjoint universes,
- $h_{ij} : \mathbf{B}_i \rightarrow \mathbf{B}_j$ is a homomorphism for all $i \geq j \in I$ such that h_{ii} is the identity on \mathbf{B}_i and for all $i \geq j \geq k$, $h_{jk} \circ h_{ij} = h_{ik}$.

Note that I here is considered as a meet-semilattice, whereas in the literature on sd-systems it is usually taken to be a join-semilattice (hence we use \geq in the above definition rather than \leq).

The *Plonka sum* over \mathbf{B} is the algebra $P_1(\mathbf{B}) = \bigcup_{i \in I} \mathbf{B}_i$ with each fundamental operation $g^{\mathbf{B}}$ defined by

$$g^{\mathbf{B}}(b_{i_1}, \dots, b_{i_n}) = g^{\mathbf{B}_j}(h_{i_1 j}(b_{i_1}), \dots, h_{i_n j}(b_{i_n}))$$

where $b_{i_k} \in \mathbf{B}_{i_k}$ and $j = i_1 \cdots i_n$ is the semilattice meet of $i_1, \dots, i_n \in I$.

Theorem 6. *Let \mathbf{A} be an ipo-semilattice, and define $I = (\{1_x \mid x \in A\}, \cdot)$. Then*

- (1) $\mathbf{B} = (\mathbb{B}_i, h_{ij}, I)$ is a sd-system of Boolean algebras, where each $h_{ij} : \mathbb{B}_i \rightarrow \mathbb{B}_j$ is a generalized Boolean algebra homomorphism defined by $h_{ij}(x) = x \cdot j$,
- (2) if \mathbf{A} is lattice-ordered, then the image $h_{ij}[\mathbb{B}_i]$ is a filter,
- (3) the Plonka sum $P_1(\mathbf{B})$ reconstructs the reduct algebra $(A, \cdot, +, -)$.

Proof. (1) The set I is closed under the semilattice operation \cdot of \mathbf{A} by Lemma 4(6). The maps h_{ij} satisfy $h_{ij}(xy) = xyj = xjyj = h_{ij}(x)h_{ij}(y)$ since multiplication is commutative and idempotent, and $h_{ij}(x + y) = (x + y)j \leq xj + yj = h_{ij}(x) + h_{ij}(y)$ by Lemma 4(8). However they should be distributive lattice homomorphisms SO HERE IS STILL A GAP. Finally for $x \in \mathbb{B}_i$, $h_{ii}(x) = xi = x1_x = x$, and if $i \geq j \geq k$ then $h_{jk}(h_{ij}(x)) = xjk = xk = h_{ik}(x)$. Since $1_i = i$ and $1_j = j$ it follows that $i \geq j$ implies $h_{ij}(1_i) = ij = j = 1_j$.

(2), (3) TBD

□

The representation of ipo-semilattices by certain Płonka sums of Boolean algebras gives a good structural description of the multiplication operation of the ipo-semilattice. Reconstructing the partial order \leq is not so simple, but it is possible.

First we note that a dual representation of sd-systems of Boolean algebras gives a much more compact way of drawing finite ipo-semilattices. Every finite Boolean algebra \mathbb{B}_i is isomorphic to the powerset Boolean algebra of its finite set X_i of atoms. For $i \leq j$, the distributive lattice homomorphism h_{ji} corresponds to the partial map $f_{ij} : X_i \rightarrow X_j$ defined by $f_{ij}(a) = b \iff a \leq h_{ji}(b)$ and $a \not\leq h_{ji}(0_j)$. Given an atom $a \in X_i$ there exists at most one such atom $b \in X_j$ since if $a \leq h_{ji}(b')$ for an atom $b' \neq b$ then $a \leq h_{ji}(b)h_{ji}(b') = h_{ji}(bb') = h_{ji}(0_j)$.

A *sd-system of partial maps* is a triple $\mathbf{X} = (X_i, f_{ij}, I)$ such that

- I is a semilattice,
- $\{X_i : i \in I\}$ is a family of disjoint sets, and
- $f_{ij} : X_i \rightarrow X_j$ is a partial map for all $i \leq j \in I$ such that $f_{ii} = id_{X_i}$ and for all $i \leq j \leq k$, $f_{jk} \circ f_{ij} = f_{ik}$.

If I and all X_i are finite, then \mathbf{X} can be displayed as a Hasse diagram of I with nodes of I replaced by the sets X_i and their elements. For a pair $i \leq j$ such that i is covered by j , the partial map f_{ij} is given by edges from elements of X_i to elements of X_j (see Figure ??). The empty set is labeled by \emptyset and the empty partial map is indicated by a dotted line between the sets. This graphical representation and the fact that systems of partial maps are logarithmically smaller than the corresponding finite ipo-semilattices make them a useful tool for investigating finite commutative idempotent involutive residuated lattices.

The systems of partial maps that arise from ipo-semilattices have two additional properties. The system \mathbf{X} is *proper* if for $i < j$, f_{ij} is a proper partial map, i.e., $\text{dom}(f_{ij})$ is a proper subset of X_i . It is *disjoint* if for all $i, j \leq k$, $\text{dom}(f_{ij,i}) \cap \text{dom}(f_{ij,j}) = \emptyset$.

For a sd-system $\mathbf{B} = (\mathbf{B}_i, h_{ij}, I)$ of complete and atomic Boolean algebras with generalized Boolean algebra homomorphisms, define a sd-system $\text{At}(\mathbf{B}) = (X_i, f_{ij}, I)$ of partial maps by $X_i = \text{At}(B_i)$ = the set of atoms of B_i , and for $i \leq j$ define $f_{ij} : X_i \rightarrow X_j$ by $f_{ij}(a) = b \iff a \leq h_{ji}(b)$ and $a \not\leq h_{ji}(0_j)$.

Conversely, given a sd-system $\mathbf{X} = (X_i, f_{ij}, I)$ of partial maps, define a sd-system $P(\mathbf{X}) = (\mathcal{P}(X_i), h_{ij}, I)$ of Boolean algebras with distributive lattice homomorphisms given by $h_{ij}(S) = f_{ji}^{-1}[S] \cup (X_j \setminus \text{dom}(f_{ji}))$.

Theorem 7. (1) *For every sd-system \mathbf{B} of complete and atomic Boolean algebras, $\mathcal{P}(\text{At}(\mathbf{B}))$ is isomorphic to \mathbf{B} .*

- (2) For every sd-system \mathbf{X} of partial maps, $At(\mathcal{P}(\mathbf{X}))$ is isomorphic to \mathbf{X} .
- (3) Every a sd-system of partial maps obtained from an ipo-semilattice is proper and disjoint.

Proof. (1) and (2) follow from a slight generalization of Tarski's duality between the category of complete and atomic Boolean algebras and the category of sets. (3) Let $\mathbf{B} = (\mathbb{B}_i, h_{ij}, I)$ be an sd-system obtained from an ipo-semilattice. Then $At(\mathbf{B})$ is proper by Lemma 4(7) and disjoint by Lemma 4(8), (9). \square

We want to show that certain sd-systems of Boolean algebras can be used to construct ipo-semilattices. This means we have to construct the partial order \leq and prove that the residuation property holds. We first consider the case of a 2-element semilattice.

For a Boolean algebra $\mathbf{A} = (A, +, 0, \cdot, 1, -)$ and any element $a \in A$, the principal filter $\uparrow a = \{x \in A : a \leq x\}$ is again a Boolean algebra $\mathbf{A}+a = (\uparrow a, +, a, \cdot, 1, -^a)$ where $+, \cdot$ are the operations of \mathbf{A} restricted to $\uparrow a$ and $-^a x = -x + a$.

Let \mathbf{A} and \mathbf{B} be Boolean algebras, and for an element $a \in A$ let $h : \mathbf{B} \rightarrow \mathbf{A}+a$ be a homomorphism. We will assume that $A \cap B = \emptyset$ and define a semilattice operation \cdot on $C = A \cup B$ as follows:

$$x \cdot^C y = \begin{cases} xy & \text{if } x, y \in A \text{ or } x, y \in B \\ xh(y) & \text{if } x \in A, y \in B \\ h(x)y & \text{if } x \in B, y \in A. \end{cases}$$

This operation is clearly idempotent and commutative. Checking associativity consists of x, y, z is in A and the other two are in B (the other cases are similar).

$$\begin{aligned} (x \cdot^C y) \cdot^C z &= (xh(y))h(z) = xh(yz) = x \cdot^C (y \cdot^C z) \\ (x \cdot^C y) \cdot^C z &= (h(x)y)h(z) = h(x)(yh(z)) = x \cdot^C (y \cdot^C z) \\ (x \cdot^C y) \cdot^C z &= h(xy)z = h(x)(h(y)z) = x \cdot^C (y \cdot^C z) \end{aligned}$$

We also have an involution on C given by the union of $-^A$ and $-^B$.

5. FIGURES (TO BE PLACED LATER)

These are some figures of ipo-semilattices and their dual representations that ought to be placed appropriately amongst the sections above. We also have an eight element $i\ell$ -semilattice with both its lattice and multiplicative orders.




FIGURE 3. A fourteen-element ipo-semilattice with identity and its dual representation.




FIGURE 4. Another fourteen-element ipo-semilattice with identity and its dual representation.




FIGURE 5. Another fourteen-element ipo-semilattice with identity and its dual representation.




FIGURE 6. Another fourteen-element ipo-semilattice with identity and its dual representation.




FIGURE 7. An 8-element il -semilattice. Its multiplicative order shows its unital sub- il -semilattices.

REFERENCES

- [1] Bonzio, S., Loi, A., Peruzzi, L.: A duality for involutive bisemilattices, *Studia Logica*, (2019) 107: 423–444.
- [2] Galatos, N., Jipsen, P., Kowalski, T., Ono, H.: “Residuated lattices: an algebraic glimpse at substructural logics”, *Studies in Logic and the Foundations of Mathematics*, Vol. 151, Elsevier, 2007.
- [3] Jipsen, P., Tuyt, O., Valota, D.: Structural characterization of commutative idempotent involutive residuated lattices, preprint, 2020.
- [4] McCune, W.: Prover9 and Mace4. www.cs.unm.edu/~mccune/prover9/, 2005–2010.