

Limited evidence of test-retest reliability in infant-directed speech preference in a large
pre-registered infant experiment

Melanie S. Schreiner^{1,2}, Martin Zettersten^{3,4}, Christina Bergmann⁵, Michael C. Frank⁶,
Tom Fritzsche⁷, Nayeli Gonzalez-Gomez⁸, Kiley Hamlin⁹, Natalia Kartushina¹⁰, Danielle J.
Kellier¹¹, Nivedita Mani^{1,2}, Julien Mayor¹⁰, Jenny Saffran³, Mohinish Shukla¹², Priya
Silverstein^{13, 14}, Melanie Soderstrom¹⁵, & Matthias Lippold^{1,2}

¹ University of Goettingen

² Leibniz Science Campus PrimateCognition

³ University of Wisconsin-Madison

⁴ Princeton University

⁵ Max Planck Institute for Psycholinguistics

⁶ Stanford University

⁷ University of Potsdam

⁸ Oxford Brookes University

⁹ University of British Columbia

¹⁰ University of Oslo

¹¹ University of Pennsylvania

¹² Università di Padova

¹³ Institute for Globally Distributed Open Research

¹⁴ Ashland University

¹⁵ University of Manitoba

Author Note

Acknowledgements. This work was supported in part by a Leibniz ScienceCampus Primate Cognition seed fund awarded to MSc and ML, a grant from the Research Council of Norway (project number 301625) and its Centres of Excellence funding scheme (project number 223265) awarded to NK, an ERC Grant (agreement number 773202 – ERC 2017, “BabyRhythm”) awarded to MSh, a ManyBabies SSHRC Partnership Development Grant awarded to MSo, and a grant from the NSF awarded to MZ (NSF DGE-1747503).

Conflict of Interest Statement. The authors declare that there are no conflicts of interest for this work.

Data Availability Statement. All code for reproducing the paper is available at <https://github.com/msschreiner/MB1T>. Data and materials are available on OSF (https://osf.io/zeqka/?view_only=e027502f4e7f49408cfb2cba38f7b506).

CRedit author statement. Outside of the position of the first, the second, and the last author, authorship position was determined by sorting authors’ last names in alphabetical order. An overview of authorship contributions following the CRediT taxonomy can be viewed here: <https://docs.google.com/spreadsheets/d/1jDvb0xL1U6YbXrpPZ1UyfyQ7yYK9aXo002UaArqy35U/edit?usp=sharing>.

Correspondence concerning this article should be addressed to Melanie S. Schreiner, Gosslerstr. 14, 37073 Göttingen. E-mail: melanie.schreiner@psych.uni-goettingen.de

Abstract

Test-retest reliability — establishing that measurements remain consistent across multiple testing sessions — is critical to measuring, understanding, and predicting individual differences in infant language development. However, previous attempts to establish measurement reliability in infant speech perception tasks are limited, and reliability of frequently-used infant measures is largely unknown. The current study investigated the test-retest reliability of infants' preference for infant-directed speech (hereafter, IDS) over adult-directed speech (hereafter, ADS) in a large sample ($N=158$) in the context of the ManyBabies1 collaborative research project (hereafter, MB1; Frank et al., 2017; ManyBabies Consortium, 2020). Labs of the original MB1 study were asked to bring in participating infants for a second appointment retesting infants on their IDS preference. This approach allows us to estimate test-retest reliability across three different methods used to investigate preferential listening in infancy: the head-turn preference procedure, central fixation, and eye-tracking. Overall, we find no consistent evidence of test-retest reliability in measures of infants' speech preference (overall $r = .09$, 95% CI $[-.06, .25]$). While increasing the number of trials that infants needed to contribute for inclusion in the analysis revealed a numeric growth in test-retest reliability, it also considerably reduced the study's effective sample size. Therefore, future research on infant development should take into account that not all experimental measures may be appropriate for assessing individual differences between infants.

Keywords: language acquisition; speech perception; infant-directed speech; adult-directed speech; test-retest reliability

Word count: 3998

Limited evidence of test-retest reliability in infant-directed speech preference in a large pre-registered infant experiment

Obtaining a quantitative measure of infants' cognitive abilities is an extraordinarily difficult endeavor. The most frequent way to assess what infants know or prefer is to track overt behavior. However, measuring overt behavior at early ages presents many challenges: participants' attention span is short, they do not follow instructions, their mood can change instantly, and their behavior is often inconsistent. Therefore, most measurements are noisy and the typical sample size of an infant study is small (around 20 infants per group), resulting in low power (Oakes, 2017). In addition, there is individual and environmental variation that may add even more noise to the data (e.g., Johnson & Zamuner, 2010). Despite these demanding conditions, reliable and robust methods for assessing infants' behavior are critical to understanding development.

In order to address these challenges, the ManyBabies collaborative research consortium was formed to conduct large-scale, conceptual, consensus-based replications of seminal findings to identify sources of variability and establish best practices for experimental studies in infancy (Frank et al., 2017). The first ManyBabies collaborative research project (hereafter, MB1, ManyBabies Consortium, 2020) explored the reproducibility of the well-studied phenomenon that infants prefer infant-directed speech (hereafter, IDS) over adult-directed speech (hereafter, ADS, Cooper & Aslin, 1990). Across many different cultures, infants are commonly addressed in IDS, which typically is characterized by higher pitch, greater pitch range, and shorter utterances, compared to the language used between interacting adults (Fernald et al., 1989). A large body of behavioral studies finds that infants show increased looking times when hearing IDS compared to ADS stimuli across ages and methods (Cooper & Aslin, 1990; see Dunst, Gorman, & Hamby, 2012 for a meta-analysis). This attentional enhancement is also documented in neurophysiological studies showing increased neural activation during IDS compared to

ADS exposure (Naoi et al., 2012; Zangl & Mills, 2007). IDS has also been identified as facilitating early word learning. In particular, infants' word segmentation abilities (Flocchia et al., 2016; Schreiner & Mani, 2017; Singh, Nestor, Parikh, & Yull, 2009; Thiessen, Hill, & Saffran, 2005) and their learning of word-object associations (Graf Estes & Hurley, 2013; Ma, Golinkoff, Houston, & Hirsh-Pasek, 2011) are enhanced in the context of IDS. In sum, several lines of evidence suggest that IDS is beneficial for early language development.

Within MB1, 67 labs contributed data from 2,329 infants showing that babies generally prefer to listen to IDS over ADS. Nevertheless, the overall effect size of $d = 0.35$ was smaller than a previously reported meta-analytic effect size of $d = 0.67$ (Dunst et al., 2012). The results revealed several additional factors that influenced the effect size. First, older infants showed a larger preference of IDS over ADS. Second, the stimulus language was linked to IDS preference, with North American English learning infants showing a larger IDS preference than infants learning other languages. Third, comparing the different methods employed, the head-turn preference procedure yielded the highest effect size, while the central fixation paradigm and eye-tracking methods revealed smaller effects. Finally, exploratory analyses assessed the effect of different inclusion criteria. Across methods, using stricter inclusion criteria led to an increase in effect sizes despite the larger proportion of excluded participants (see also Byers-Heinlein, Bergmann, & Savalei, 2021).

However, there is a difference between a result being reliable in a large sample of infants and the measurement of an individual infant being reliable. In studies tracking individual differences, the measured behavior during an experimental setting is often used to predict a cognitive function or specific skill later in life. Individual differences research of this kind often has substantial implications for theoretical and applied work. For example, research showing that infants' behavior in speech perception tasks can be linked to later language development (see Cristia, Seidl, Junge, Soderstrom, & Hagoort, 2014 for a meta-analysis) has the potential to identify infants at risk for later language delays or disorders. However, a necessary precondition for this link to be observable is that

individual differences between infants can be measured with high reliability at these earlier stages, in order to ensure that measured inter-individual variation mainly reflects differences in children's abilities rather than measurement error. How reliable are the measures used in infancy research?

Previous attempts to address the reliability of measurements have typically been limited to adult populations (Hedge, Powell, & Sumner, 2018), or have been conducted with small sample sizes (e.g., Houston, Horn, Qi, Ting, & Gao, 2007). For example, Colombo, Mitchell, and Horowitz (1988) used a paired-comparison task, in which infants were familiarized with a stimulus and presented with the familiarized and a novel stimulus side-by-side at test. Results indicated that infants' novelty preference was extremely variable from task to task. Assessing infants' performance from one week to another revealed that infants' attention measures were moderately reliable. However, reliability seemed to increase with the number of tasks infants completed in the younger age group, suggesting that reliability is influenced by the number of assessments. In addition, infants' performance from 4 to 7 months was longitudinally stable but somewhat smaller than week-to-week reliability. Cristia, Seidl, Singh, and Houston (2016) also retested infant populations by independently conducting 12 different experiments on infant speech perception at three different labs with different implementations of the individual studies. Hence, it was only after completed data collection that the data was pooled together by the different labs revealing potential confounds. Nevertheless, the results showed that reliability was extremely variable across the different experiments and labs and low overall (meta-analytic $r = .07$).

Against this background, the current study investigates test-retest reliability of infants' performance in a speech preference task. Within MB1, a multi-lab collaboration, we examine whether infants' preferential listening behavior to IDS and ADS is reliable across two different test sessions. We also investigate the influence of various moderators on the reliability of IDS preference (e.g., time between test and retest; infants' language

background).

Our study was faced with a critical design choice: what stimuli to use to assess test-retest reliability. One constraint on our study was that, since it was a follow-on to MB1, any stimulus we used would always be presented after the MB1 stimuli. One option would be simply to bring back infants and have them hear exactly the same stimulus materials. A weakness of this design would be the potential for stimulus familiarity effects, however, since infants would have heard the materials before. Further complicating matters, infants might show a preference for or against a familiar stimulus depending on their age (Hunter & Ames, 1988). The ideal solution then would be to create a brand new stimulus set with the same characteristics. Unfortunately, because of the process of how MB1 stimuli were created, we did not have enough normed raw recordings available to make brand new stimulus items that conformed to the same standards as the MB1 stimuli. We therefore chose an intermediate path: we reversed the ordering of MB1 stimuli. Average looking times in MB1 were always lower than 9s per trial, even for the youngest children on the earliest trials (the group who looked the longest on average), so most children in MB1 did not hear the second half of most trials. Thus, by reversing the order, we had a perfectly matched stimulus set that was relatively unfamiliar to most infants. The disadvantage of this design was that infants who looked longer might be more likely to hear a familiar clip heard in the previous study. If infants then showed a familiarity preference — an assumption which might not be true — the end result could be to inflate our estimates of test-retest reliability slightly, since longer lookers would on average look longer at retest due to their familiarity preference. We view this risk as relatively low, but do note that it is a limitation of our design.

The current study also explores whether there are any differences in test-retest reliability between three widely used methods: central fixation (CF), eye-tracking (ET), and the head-turn preference procedure (HPP). Exploring differences in CF, ET, and HPP, Junge et al. (2020) provide experimental and meta-analytic evidence in favor of using the

HPP in speech segmentation tasks. Similarly, the MB1 project reported an increase in the effect size for HPP compared to CF and ET (ManyBabies Consortium, 2020). HPP requires gross motor movements relative to other methods, such as CF and ET paradigms, for which subtle eye movements towards a monitor located in front of the child are sufficient. One possible explanation for the stronger effects with HPP may be a higher sensitivity to the contingency of the presentation of auditory stimuli and infants' head turns away from the typical forward-facing position. While these findings suggest that HPP may be a more sensitive index of infant preference, they do not necessarily imply higher reliability for individual infants' performance using HPP. For example, Marimon and Höhle (2022) found no evidence for test-retest reliability when testing infants' prosodic preferences using the HPP method. It remains an open question whether the same measures that produce larger effect sizes at the group-level also have higher test-retest reliability for individual infants (Byers-Heinlein, Bergmann, et al., 2021). Therefore, assessing the test-retest reliability of the different preference measures is crucial, so that researchers can make informed decisions about the appropriate methods for their particular research question. Critically, only measures with high test-retest reliability should be used for studies of individual differences.

Method

Preregistration

We preregistered the current study on the Open Science Framework (<https://osf.io/v5f8t>). Section S1 in the Supplementary Materials contains additional notes on the preregistration decisions and any deviations from the preregistered analytic plan.

193 Data Collection

194 A call was issued to all labs participating in the original MB1 study on January 24th,
 195 2018 (ManyBabies Consortium, 2020). The collection of retest session data was initially set
 196 to end on May 31st, 2018, one month after the end date of the original MB1 project. Due
 197 to the fact that the original MB1 project extended the time frame for data collection and
 198 the late start of data collection for the MB1 test-retest study, we also allowed participating
 199 labs to continue data collection past the scheduled end date.

200 Participants

201 Contributing labs were asked to re-recruit their monolingual participants between the
 202 ages of 6 to 12 months who had already participated in the MB1 project. If participating
 203 labs had not committed to testing either of these age groups, they were also allowed to
 204 re-recruit participants from the youngest age group of 3- to 6-month-olds and/or the oldest
 205 age group of 12- to 15-month-olds. Labs were asked to contribute half ($n=16$) or full
 206 samples ($n=32$); however, a lab's data was included in the study regardless of the number
 207 of included infants. The study was approved by each lab's respective ethics committee and
 208 parental consent was obtained for each infant prior to participation in the study.

209 Our final sample consisted of 158 monolingual infants from 7 different labs (Table 1).
 210 In order to be included in the study, infants needed a minimum of 90% first language
 211 exposure, to be born full term with no known developmental disorders, and normal hearing
 212 and vision. We excluded 18 additional participants (see Data Exclusion section for details).
 213 The mean age of infants included in the study was 245 days (range: 108 – 373 days;
 214 approximately 8.06 months).

Materials

Visual stimuli. The visual stimuli and instructions were identical to MB1. For the CF paradigm and ET, labs used a multicolored static checkerboard as the fixation stimulus as well as a multicolored moving circle with a ringing sound as an attention-getter between trials. For the HPP method, labs used their standard procedure, as in MB1.

Speech stimuli. We used the identical training stimuli of piano music from MB1. A second set of naturalistic IDS and ADS recordings of mothers either talking to their infant or to an experimenter was created for the retest session by reversing the order of clips within each sequence of the original study. This resulted in eight new sequences of natural IDS and eight new sequences of natural ADS with a length of 18 seconds each.

Procedure. Infants were retested using the identical procedure as during the first testing day: CF, HPP, or ET. Participating labs were asked to schedule test and retest sessions 7 days apart with a minimum number of 1 day and a maximum number of 31 days. However, infants whose time between test and retest exceeded 31 days were still included in the analyses ($n = 3$). The mean number of days between test and retest was 10 (range: 1 - 49).

A total of 18 trials, including two training, eight IDS, and eight ADS trials, were presented in one of four pseudo-randomized orders. Trial length was either infant-controlled or fixed depending on the lab's standard procedure: a trial stopped either if the infant looked away for 2 seconds or after the total trial duration of 18 seconds. The online coding experimenter and the parent listened to music masked with the stimuli of the study via noise-cancelling headphones. If the experimenter was in an adjacent room separate from the testing location, listening to masking music was optional for the experimenter.

Data exclusion. In total, 18 participants were excluded from the analysis. 4 participants were excluded for being preterm (defined as a gestation time of less than 37 weeks). 6 participants were excluded due to session errors involving an experimenter error

Table 1

Statistics of the included labs. n refers to the number of infants included in the final analysis.

Lab	Method	Language	Mean age (days)	N
InfantCog-UBC	central fixation	English	147	7
babylab-potsdam	HPP	German	227	22
babyling-oslo	eye-tracking	Norwegian	249	10
brookes-babylab	central fixation	English	267	18
infantll-madison	HPP	English	230	30
lancslab	eye-tracking	English	236	16
wsi-goettingen	HPP	German	242	16
wsi-goettingen	central fixation	German	280	39

(e.g., inaccurate coding or presentation of retest stimuli on the first test session). Individual trials were excluded if they were marked as trial errors (5.45% of remaining trials), i.e., if the infant was reported as fussy, an experimental or equipment error occurred, or there was parental interference during the task (e.g., if the parent spoke with the infant during the trial). Trials were also excluded if the minimum looking time of 2 s was not met (12.60% of the remaining trials). If a participant was unable to contribute at least one IDS and one ADS trial for either test or retest after trial-level exclusions, all data of that participant was excluded from the test-retest analyses (12 additional participants).

Table 2

Average looking times (in seconds) for each session and condition

Trial type	Session 1 Mean	Session 1 <i>SD</i>	Session 2 Mean	Session 2 <i>SD</i>
ADS	7.71	2.77	6.96	2.92
IDS	8.76	2.84	7.75	2.75

Results

IDS preference

First, we conducted confirmatory analyses examining infants' preference for IDS in both sessions. Two-samples t-tests comparing the difference in average looking time between IDS and ADS to zero revealed that infants showed a preference of IDS over ADS in Session 1, $t(157) = 6.47$, $p < .001$, and Session 2, $t(157) = 4.19$, $p < .001$, replicating the main finding from MB1 (Table 2). 68.35% of infants in Session 1 and 63.29% of infants in Session 2 showed a preference for IDS. In order to test whether there was a difference in the strength of the preference effect across sessions, we fit a linear mixed-effects model predicting infants' average difference in looking time between IDS and ADS from test session (1 vs. 2), including by-lab and by-participant random intercepts. There was no significant difference in the magnitude of infants' preference between the two sessions, $\beta = -0.30$, $SE = 0.24$, $p = .208$.

Reliability

We assessed test-retest reliability in two planned, confirmatory analyses. First, we fit a linear mixed-effects model predicting IDS preference in Session 2 from IDS preference in Session 1, including a by-lab random intercept. The results revealed no significant relationship between IDS preference in Session 1 and 2 (Table 3). Second, we calculated

Table 3

Coefficient estimates from a linear mixed effects model predicting IDS preference in Session 2.

	Estimate	SE	t	p
Intercept	0.87	0.46	1.92	0.10
IDS Preference Session 1	0.04	0.09	0.41	0.68




Figure 1. Correlation between IDS Preference in Session 1 and Session 2 in each lab and method. Dots indicate individual participants. Error bands represent 95 percent confidence intervals. The dashed line indicates no preference (i.e., a value of zero) for the first and second session, respectively.

Table 4

Coefficient estimates from a linear mixed effects model predicting IDS preference in Session 2 and Pearson correlation coefficient for each method separately.

Method	beta	SE	p	Pearson r
central fixation	-0.20	0.12	0.12	0.08
HPP	0.15	0.14	0.28	0.13
eye-tracking	0.03	0.16	0.84	0.02

the Pearson correlation coefficient. While a simple correlation coefficient might overestimate the test-retest reliability in our sample because it does not control for the differences between different labs and methods (HPP, CF, and ET), we felt it was important to also conduct a Pearson correlation as it is commonly used to assess reliability. The size of the correlation coefficient was not statistically different from zero and the estimate was small, $r = .09$, 95% CI $[-.06, .25]$, $t(156) = 1.19$, $p = .237$. Moreover, no significant correlations emerged in each sample considered separately (Figure 1; see Supplementary Materials S3 for a meta-analytic approach). 41.77% of the infants reversed their direction of preference for IDS versus ADS from the test to the retest session.

To investigate the test-retest reliability of each specific method, we computed Pearson correlation coefficients and the same mixed-effects model described above for HPP, CF, and ET separately (Table 4) in additional exploratory analyses. None of the three methods showed evidence of test-retest reliability. Neither the Pearson correlation coefficients nor the coefficients of the multilevel analysis were significant, all p -values > 0.12 . In planned secondary analyses, we found that time between test sessions, participant age, and language background did not moderate the relationship between IDS preference in session

1 and session 2 (see Supplementary Materials S2). Taken together, we find no significant evidence of test-retest reliability across our preregistered analyses.

Results with different inclusion criteria

To this point, all analyses were performed using the inclusion criteria from MB1, which required only that infants contribute at least one trial per condition for inclusion (i.e., one IDS and one ADS trial). However, more stringent inclusion criteria yielded larger effect sizes in MB1. We therefore conducted exploratory analyses assessing test-retest reliability after applying progressively stricter inclusion criteria, requiring two, four, six, and eight valid trials per condition. Applying stricter criteria — and thereby increasing the number of test trials — increased reliability numerically from $r = 0.07$ to $r = 0.34$ (Figure 2). In part due to a decrease in sample size, only one of these correlations was statistically significant (when requiring six trial pairs): two valid trial pairs, $t(152) = 0.90$, $p = .367$; four valid trial pairs, $t(143) = 1.03$, $p = .306$; six valid trial pairs, $t(98) = 2.23$, $p = .028$; eight valid trial pairs — all trials in both sessions — $t(22) = 1.68$, $p = .108$. The analyses provide tentative evidence that stricter inclusion criteria may lead to higher test-retest reliability, but at the cost of substantial decreases in sample size (see Supplementary Materials S5 for additional analyses).

General Discussion

The current study investigated the test-retest reliability of infants' preference for IDS over ADS. We retested the IDS preference of infants participating in the original MB1 project to assess the extent to which their pattern of preference would remain consistent across multiple testing sessions. While we replicated the original effect of infants' speech preference for IDS over ADS for both the test and retest session on the group-level, we found that infants' speech preference measures showed no evidence of test-retest reliability. In other words, we were unable to detect stable individual differences in infants' preference




Figure 2. IDS preferences of both sessions plotted against each other for each inclusion criterion. n indicates the number of included infants, r is the Pearson correlation coefficient as the indicator for reliability.

for IDS. This finding is consistent with past research suggesting low test-retest reliability in other infant paradigms (Cristia et al., 2016). Given that most experimental procedures conducted in infant research are interested in the comparison of groups, individual differences between participants within a specific condition are usually minimized by the experimental procedure while differences between conditions are maximized. Therefore, infant preference measures may be a good approach for capturing group-level phenomena, but may be less appropriate for examining individual differences in development.

Consistent with general psychometric theory (e.g., DeBolt, Rhemtulla, & Oakes, 2020), stricter inclusion criteria — and consequently a larger number of included test trials per participant — tended to increase the magnitude of the correlation between test sessions. However, this association was based on exploratory analyses and was in part only observed descriptively, and hence should be interpreted with caution. A similar effect on the group-level was found in the MB1 project, where a stricter inclusion criterion led to bigger effect sizes (ManyBabies Consortium, 2020). As in MB1, higher reliability through strict exclusions came at a high cost. In particular, with the strictest criterion, only a small portion of the original sample size (24 out of 158 infants) could be included in the final sample. In other words, applying stricter criteria leads to a higher drop-out rate and can dramatically reduce the sample size. In the case of studies in the field of developmental science, where there are many practical restrictions in collecting large samples of infants (e.g., birth rate in the area, restricted lab capacities, budget restrictions), a strict drop-out criterion may often be difficult to implement. Note that studies in developmental science already have above-average drop-out rates (Miller, 2017). In addition, drop out may not be random, and so having high drop-out rates can further limit the generalizability of a study. In fact, the number of trials individual infants contributed was highly correlated between test sessions in the current study (see Supplementary Materials S6). Particularly in the context of turning individual differences measures into diagnostic tools, high drop-out rates have an additional limitation of not being broadly usable.

An alternative approach to increasing the number of valid trials is to increase the number of experimental trials. This approach seeks to increase the likelihood that participants will contribute sufficient trials (after trial-level exclusions) to allow for precise individual-level estimates (DeBolt et al., 2020; see also Silverstein, Feng, Westermann, Parise, & Twomey, 2021). While this approach is promising, it may not always be feasible, because the attention span of a typical infant participant is limited. Therefore, prolonging the experimental procedure to maximize the absolute number of trials is often challenging in practice. Other avenues for obtaining higher numbers of valid trials may include changes in the procedure (e.g., Egger, Rowland, & Bergmann, 2020) or implementing multi-day test sessions (Fernald & Marchman, 2012).

As our results are only based on the phenomenon of IDS preference (albeit, with three widely used methods: HPP, CF, ET) it is essential to further assess the underlying reliability of preferential looking measures within other areas of speech perception (Marimon & Höhle, 2022). While most infants prefer IDS over ADS (Dunst et al., 2012), patterns of preferential looking in other tasks (e.g., speech segmentation) are often inconsistent and difficult to predict (Bergmann & Cristia, 2016). These inconsistencies in looking behavior are especially important to consider in the context of relating a direction of preference to later language development, and can sometimes lead to seemingly contradictory findings. That is, both familiarity and novelty responses have been suggested to be predictive of infants' later linguistic abilities (DePaolis, Vihman, & Keren-Portnoy, 2014; Newman, Ratner, Jusczyk, Jusczyk, & Dow, 2006; Newman, Rowe, & Ratner, 2016). In light of our findings, researchers conducting longitudinal studies with experimental data from young infants predicting future outcomes should be cautious, as there may be large intra-individual variability affecting preference measurement.

Limitations

While we had an above-average sample size for a study in infant research, we were unable to approach the number of participants collected within the original MB1 study. In addition to a delayed call, the extra effort of having to schedule a second lab visit for each participant and the fact that there were already other collaborative studies taking place simultaneously (MB1B, Byers-Heinlein, Tsui, Bergmann, et al., 2021; MB1G, Byers-Heinlein, Tsui, Van Renswoude, et al., 2021), might have contributed to a low participation rate. A higher sample size and a larger number of participating labs from different countries would have enabled us to conduct a more highly-powered test of differences in test-retest reliability across different methods, language backgrounds, and participant age.

A further limitation concerns the stimuli. While the order of the audio recording clips presented to infants within a given trial differed between the first and second session, the exact same stimulus material as in MB1 was used in both sessions. In particular, all children heard the exact same voices in Session 1 and in Session 2. From a practical point of view, this was the most straightforward solution for coordinating the experiment within the larger MB1 project. However, familiarity effects might have influenced infants' looking behavior. Infants with longer looking times in their first session might have had more opportunity to recognize familiar audio clips in their second session. For infants with short looking times, familiar audio clips would only occur towards the end of second-session trials, thus offering infants less opportunity to recognize voices from their first session. Therefore, inconsistent familiarity with the stimulus material in the second session across infants might have artificially lowered test-retest reliability. Moreover, infants' experience with a testing paradigm has been found to systematically affect looking time to familiar stimuli (Santolin, Garcia-Castro, Zettersten, Sebastian-Galles, & Saffran, 2021), further complicating the interpretation of infant familiarity preferences in retest sessions. On the

other hand, one factor that mitigates this concern is that infants' looking times generally declined in session 2 compared to session 1 (consistent with past work, e.g. Marimon & Höhle, 2022), limiting opportunities for infants to encounter previously experienced stimulus material.

Conclusion

Following the MB1 protocol, the current study could not detect test-retest reliability in measures of infants' preference for IDS over ADS. Subsequent analyses provided tentative evidence that stricter criteria for the inclusion of participants may enhance test-retest reliability at the cost of high drop-out rates. Developmental studies relying on stable individual differences between their participants need to consider the underlying reliability of their measures, and we recommend a broader assessment of test-retest reliability in infant research.

References

- Bergmann, C., & Cristia, A. (2016). Development of infants' segmentation of words from native speech: A meta-analytic approach. *Developmental Science*, 19(6), 901–917.
- Byers-Heinlein, K., Bergmann, C., & Savalei, V. (2021). Six solutions for more reliable infant research. *Infant and Child Development*, e2296.
- Byers-Heinlein, K., Tsui, A. S. M., Bergmann, C., Black, A. K., Brown, A., Carbajal, M. J., ... Wermelinger, S. (2021). A multilab study of bilingual infants: Exploring the preference for infant-directed speech. *Advances in Methods and Practices in Psychological Science*, 4(1), 2515245920974622.
- Byers-Heinlein, K., Tsui, R. K.-Y., Van Renswoude, D., Black, A. K., Barr, R., Brown, A., ... Singh, L. (2021). The development of gaze following in monolingual and bilingual infants: A multi-laboratory study. *Infancy*, 26(1), 4–38.
- Colombo, J., Mitchell, D. W., & Horowitz, F. D. (1988). Infant visual attention in the paired-comparison paradigm: Test-retest and attention-performance relations. *Child Development*, 1198–1210.
- Cooper, R. P., & Aslin, R. N. (1990). Preference for infant-directed speech in the first month after birth. *Child Development*, 61(5), 1584–1595.
- Cristia, A., Seidl, A., Junge, C., Soderstrom, M., & Hagoort, P. (2014). Predicting individual variation in language from infant speech perception measures. *Child Development*, 85(4), 1330–1345.
- Cristia, A., Seidl, A., Singh, L., & Houston, D. (2016). Test-retest reliability in infant speech perception tasks. *Infancy*, 21(5), 648–667.
- DeBolt, M. C., Rhemtulla, M., & Oakes, L. M. (2020). Robust data and power in infant research: A case study of the effect of number of infants and number of trials in visual preference procedures. *Infancy*, 25(4), 393–419.

- DePaolis, R. A., Vihman, M. M., & Keren-Portnoy, T. (2014). When do infants begin recognizing familiar words in sentences? *Journal of Child Language*, *41*(1), 226–239.
- Dunst, C., Gorman, E., & Hamby, D. (2012). Preference for infant-directed speech in preverbal young children. *Center for Early Literacy Learning*, *5*(1), 1–13. Retrieved from http://www.earlyliteracylearning.org/cellreviews/cellreviews_v5_n1.pdf
- Egger, J., Rowland, C. F., & Bergmann, C. (2020). Improving the robustness of infant lexical processing speed measures. *Behavior Research Methods*, *52*(5), 2188–2201.
- Fernald, A., & Marchman, V. A. (2012). Individual differences in lexical processing at 18 months predict vocabulary growth in typically developing and late-talking toddlers. *Child Development*, *83*(1), 203–222.
- Fernald, A., Taeschner, T., Dunn, J., Papousek, M., Boysson-Bardies, B. de, & Fukui, I. (1989). A cross-language study of prosodic modifications in mothers' and fathers' speech to preverbal infants. *Journal of Child Language*, *16*(3), 477–501.
- Floccia, C., Keren-Portnoy, T., DePaolis, R., Duffy, H., Delle Luche, C., Durrant, S., . . . Vihman, M. (2016). British english infants segment words only with exaggerated infant-directed speech stimuli. *Cognition*, *148*, 1–9.
- Frank, M. C., Bergelson, E., Bergmann, C., Cristia, A., Floccia, C., Gervain, J., . . . Yurovsky, D. (2017). A collaborative approach to infant research: Promoting reproducibility, best practices, and theory-building. *Infancy*, *22*(4), 421–435. <https://doi.org/10.1111/infa.12182>
- Graf Estes, K., & Hurley, K. (2013). Infant-directed prosody helps infants map sounds to meanings. *Infancy*, *18*(5), 797–824. <https://doi.org/10.1111/infa.12006>

- Hedge, C., Powell, G., & Sumner, P. (2018). The reliability paradox: Why robust cognitive tasks do not produce reliable individual differences. *Behavior Research Methods*, 50(3), 1166–1186.
- Houston, D. M., Horn, D. L., Qi, R., Ting, J. Y., & Gao, S. (2007). Assessing speech discrimination in individual infants. *Infancy*, 12(2), 119–145.
- Hunter, M. A., & Ames, E. W. (1988). A multifactor model of infant preferences for novel and familiar stimuli. *Advances in Infancy Research*, 5, 69–95.
- Johnson, E., & Zamuner, T. (2010). Using infant and toddler testing methods in language acquisition research. In E. Blom & S. Unsworth (Eds.), *Experimental methods in language acquisition research* (pp. 73–93). Amsterdam: John Benjamins Publishing Company.
- Junge, C., Everaert, E., Porto, L., Fikkert, P., Klerk, M. de, Keij, B., & Benders, T. (2020). Contrasting behavioral looking procedures: A case study on infant speech segmentation. *Infant Behavior and Development*, 60, 101448.
- Ma, W., Golinkoff, R. M., Houston, D. M., & Hirsh-Pasek, K. (2011). Word learning in infant-and adult-directed speech. *Language Learning and Development*, 7(3), 185–201.
- ManyBabies Consortium. (2020). Quantifying sources of variability in infancy research using the infant-directed-speech preference. *Advances in Methods and Practices in Psychological Science*, 3(1), 24–52.
- Marimon, M., & Höhle, B. (2022). Testing prosodic development with the headturn preference procedure: A test-retest reliability study. *Infant and Child Development*, e2362.
- Miller, S. A. (2017). *Developmental research methods*. Sage publications.
- Naoi, N., Minagawa-Kawai, Y., Kobayashi, A., Takeuchi, K., Nakamura, K., Yamamoto, J., & Shozo, K. (2012). Cerebral responses to infant-directed speech and the effect of talker familiarity. *Neuroimage*, 59(2), 1735–1744.

- Newman, R., Ratner, N. B., Jusczyk, A. M., Jusczyk, P. W., & Dow, K. A. (2006).
 Infants' early ability to segment the conversational speech signal predicts later
 language development: A retrospective analysis. *Developmental Psychology*,
42(4), 643.
- Newman, R., Rowe, M. L., & Ratner, N. B. (2016). Input and uptake at 7 months
 predicts toddler vocabulary: The role of child-directed speech and infant
 processing skills in language development. *Journal of Child Language*, *43*(5),
 1158–1173.
- Oakes, L. M. (2017). Sample size, statistical power, and false conclusions in infant
 looking-time research. *Infancy*, *22*(4), 436–469.
- Santolin, C., Garcia-Castro, G., Zettersten, M., Sebastian-Galles, N., & Saffran, J.
 R. (2021). Experience with research paradigms relates to infants' direction of
 preference. *Infancy*, *26*(1), 39–46. <https://doi.org/10.1111/infa.12372>
- Schreiner, M. S., & Mani, N. (2017). Listen up! Developmental differences in the
 impact of IDS on speech segmentation. *Cognition*, *160*, 98–102.
- Silverstein, P., Feng, J., Westermann, G., Parise, E., & Twomey, K. E. (2021).
 Infants learn to follow gaze in stages: Evidence confirming a robotic prediction.
Open Mind, 1–15.
- Singh, L., Nestor, S., Parikh, C., & Yull, A. (2009). Influences of infant-directed
 speech on early word recognition. *Infancy*, *14*(6), 654–666.
- Thiessen, E. D., Hill, E. A., & Saffran, J. R. (2005). Infant-directed speech
 facilitates word segmentation. *Infancy*, *7*(1), 53–71.
https://doi.org/10.1207/s15327078in0701_5
- Zangl, R., & Mills, D. L. (2007). Increased brain activity to infant-directed speech
 in 6-and 13-month-old infants. *Infancy*, *11*(1), 31–62.
https://doi.org/10.1207/s15327078in1101_2