Manybabies1 Test-Retest Supplementary Information

10

11

12

13

14

15

16

25

26

27

28

29

4 S1: Notes on and deviations from the preregistration

Below, we have compiled a list of notes on and deviations from the preregistered methods and analyses https://osf.io/v5f8t.

- All infants with usable data for both test and retest session were included in the
 analyses, regardless of the number of total of infants a lab was able to contribute after
 exclusion. This decision is consistent with past decisions in ManyBabies projects to
 be as inclusive about data inclusion as possible (ManyBabies Consortium, 2020).
 - A small number of infants with a time between sessions above 31 days were also included in the analyses (n=2).
- Consistent with analytic decisions in ManyBabies 1 (ManyBabies Consortium, 2020), total looking times were truncated at 18 seconds (the maximum trial time) in the small number of cases where recorded looking times were slightly greater than 18s (presumably due to small measurement error in recording infant looking times).
- In assessing differences in IDS preference between test and retest sessions, we 17 preregistered an additional linear mixed-effects model including a by-lab random 18 slope for session. This model yielded qualitatively equivalent results (see R 19 markdown analysis script for the main manuscript). However, the model resulted in a 20 singular fit, suggesting that the model specification may be overly complex and that 21 its estimates should be interpreted with caution. We therefore focused only on the 22 first preregistered model (including only by-lab and by-participant random 23 intercepts) in reporting the analyses in the main manuscript. 24
 - In assessing the reliability of IDS using a linear-mixed-effects model predicting IDS preference in session 2 from IDS preference in session 1, we also assessed the robustness of the results by fitting a second preregistered model with more complex random effects structure, including a by-lab random slope for IDS preference in session 1. This model is included in the main R markdown script and yields

- qualitatively equivalent results to the model reported in the manuscript that includes
 a by-lab random intercept only.
- We report a series of secondary planned analyses in the Supplementary Materials
 exploring potential moderating variables of time between test sessions (S2.1),
 participant age (S2.2.), and the language background of the participants (S2.3.).

55 S2: Secondary Analyses Investigating Possible Moderating Variables

S2.1. Time between test sessions. The number of days between the first and 36 second testing session varied widely across participants (mean: 10 days; range: 1 - 49 days). 37 We therefore tested for the possibility that the time between sessions might have an impact 38 on the reliability. We fit a linear mixed-effects model predicting IDS preference in session 2 39 from IDS preference in session 1 (mean-centered), number of days between testing sessions (mean-centered), and their interaction, including a by-lab random intercept and random slope for IDS preference in test session 1 (more complex random effects structure including additional random slopes for number of days between test sessions and its interaction with IDS preference in session 1 did not converge). We found no evidence that number of days between test sessions moderated the relationship between IDS preference at test session 1 and 2. Neither the main effect of time between sessions, β =-0.01, SE=0.03, p=.684, nor the interaction term, β =-0.01, SE=0.02, p=.465, showed significant effects.

S2.2. Participant Age.

48

S2.3. Language Background. NAE-learning infants showed greater IDS preferences than their non-NAE counterparts in MB1. We therefore also assessed if test-retest reliability interacted with children's language background. A multilevel analysis with Lab as random intercept, predicting the IDS preference in Session 2 based on the IDS preference in Session 1, NAE and the interaction of these two variables, revealed no interaction, β =0.29, SE=0.18, p=.115 (see Figure S1).




Figure 1. Infants' preference in Session 1 and Session 2 with individual data points and regression lines color-coded by method (central fixation, eye-tracking, or HPP). Results are plotted separately for North American English-learning infants (right panel) and infants learning other languages and dialects (right panel).

- 55 S3: Meta-analytic estimate
- 56 S4: Alternative Dependent Variables
- 57 S4.1. Log-transformed looking times.
- S4.2. Proportion novelty preference.
- 59 S5: Patterns of preference across sessions
- We also conducted analyses to explore whether there were any patterns of preference reversal across test sessions. While there was no strong correlation in the magnitude of IDS

preference between test session 1 and test session 2, here we asked whether infants consistently expressed the same preference across test sessions. Overall, 58.20% of the infants had a consistent preference from test to retest session, indicating that infants were not more likely than chance to maintain their preference from test session 1 to test session 2 (exact binomial test; p = 0.05). Of the 158 total infants, 44.90% of infants showed a consistent infant-directed speech preference and 13.30% showed a consistent adult-directed speech preference. 23.40% of infants switched from an infant-directed speech preference at test session 2 and 18.40% switched from an adult-directed speech preference to an infant-directed speech preference.

Next, we explored whether we could detect any systematic clustering of infants with distinct patterns of preference across the test and retest session. We took a bottom-up approach and conducted a k-means clustering of the test-retest difference data. We found little evidence of distinct clusters emerging from these groupings: the clusterings ranging from k=2 (2 clusters) to k=4 (4 clusters) appear to simply track whether participants are approximately above or below the mean looking time difference for test session 1 and test session 2, and the diagnostic elbow plot shows little evidence of a qualitative improvement as the number of clusters is increased.

₇₉ S6: Relationship between the number of trials infants contribute in each session

Are there stable individual differences in how likely an infant is to contribute a high
number of trials? To answer this question, we conducted an exploratory analysis
investigating whether there is a relationship between the number of trials an infant
contributed in session 1 and session 2. Do infants who contribute a higher number of trials
during their first testing session also tend to contribute more trials during their second
testing session? A positive correlation between trial numbers during the first and second
session would indicate that their is some stability in a given infants' likelihood of remaining
attentive throughout the experiment. On the other hand, the absence of a correlation




Figure 2. (A) Results from the k-means clustering analysis of IDS preference in session 1 and 2 for different numbers of k and (B) the corresponding elbow plot of the total within-cluster sum of squares. In (A), points represent indvidual participants' magnitude of looking time difference at test sessions 1 (x-axis) and 2 (y-axis). The solid line indicates no preference for IDS vs. ADS, the dotted lines indicate mean IDS preference at test session 1 and 2, respectively. Colors indicate clusters from the k-means clustering for different values of k.




Figure 3. Correlation between the number of trials contributed in session 1 and session 2. Each data point represents one infant. Colored lines represent linear fits for each method.

would indicate that the number of trials a given infant contributes is not predictive of how many trials they might contribute during their next session.

We found a strong positive correlation between number of trials contributed during the first and the second session r = .58, 95% CI [.47, .68], t(159) = 9.05, p < .001 (see Figure 1). This result suggests that if infants contribute a higher number of trials in one session, compared to other infants, they are likely to contribute a higher number of trials in their next session. This finding is consistent with the hypothesis that how attentive infants are throughout an experiment (and hence how many trials they contribute) is a stable individual difference, at least for some infant looking time tasks. Researchers should therefore be mindful of the fact that decisions about including or excluding infants based on trials contributed may selectively sample a specific sub-set of the infant population they are studying (Byers-Heinlein, Bergmann, & Savalei, 2021; DeBolt, Rhemtulla, & Oakes, 2020).

S7: Correlations in average looking times between sessions

S7.1: Relations between overall looking time in session 1 and 2. There is a strong relationship between average overall looking in the first test session and the second test session, even after controlling for number of trials in the first and second session.






Table 1 Average Looking during session 1 predicted from average looking at session 2, controlling for trial number for each session.

Predictor	b	95% CI	t	df	p
Intercept	2.55	[0.38, 4.73]	2.32	154	.022
Mean lt 1	0.42	[0.27, 0.58]	5.52	154	< .001
N 1	-0.08	[-0.24, 0.08]	-0.96	154	.338
N 2	0.18	[0.04, 0.32]	2.52	154	.013





##

##

134

```
## Call:
108
   ## lm(formula = LT_Retest_IDS ~ LT_Test_IDS + LT_Test_ADS, data = agg_by_subj_condition_
109
   ##
110
   ## Residuals:
111
   ##
           Min
                    1Q
                        Median
                                      3Q
                                             Max
112
   ## -4.2721 -1.7567 -0.2799 1.4822
                                          6.4805
113
   ##
114
   ## Coefficients:
115
                   Estimate Std. Error t value Pr(>|t|)
   ##
116
   ## (Intercept)
                     3.9749
                                 0.6902
                                           5.759 4.41e-08 ***
117
   ## LT_Test_IDS
                     0.2123
                                 0.1008
                                           2.105
                                                   0.0369 *
118
   ## LT Test ADS
                     0.2467
                                           2.362
                                                   0.0194 *
                                 0.1044
   ## ---
120
   ## Signif. codes: 0 '***' 0.001 '**' 0.05 '.' 0.1 ' ' 1
   ##
122
   ## Residual standard error: 2.52 on 155 degrees of freedom
123
         (7 observations deleted due to missingness)
   ##
124
   ## Multiple R-squared: 0.1771, Adjusted R-squared: 0.1665
125
   ## F-statistic: 16.68 on 2 and 155 DF, p-value: 2.751e-07
126
   ##
127
   ## Call:
128
   ## lm(formula = LT_Retest_ADS ~ LT_Test_IDS + LT_Test_ADS, data = agg_by_subj_condition_
   ##
130
   ## Residuals:
131
   ##
         Min
                  1Q Median
                                 3Q
                                       Max
132
   ## -5.556 -1.771 -0.489 1.254
                                     8.901
133
```

147

Coefficients: 135 Estimate Std. Error t value Pr(>|t|) 136 ## (Intercept) 3.2374 0.7356 4.401 2e-05 *** 137 ## LT Test IDS 0.1103 0.1075 1.026 0.30641 138 ## LT_Test_ADS 0.3563 0.1113 3.201 0.00166 ** 139 140 0 '*** 0.001 '** 0.01 '*' 0.05 '.' 0.1 ' '1 ## Signif. codes: 141 142 ## Residual standard error: 2.686 on 155 degrees of freedom 143 (7 observations deleted due to missingness) 144 ## Multiple R-squared: 0.1677, Adjusted R-squared: 0.157 145 ## F-statistic: 15.62 on 2 and 155 DF, p-value: 6.619e-07

S7.3: Relations for specific ADS and IDS stimuli between the first and





Table 2

Mixed-effects model results predicting looking time
during session 1 from looking time at session 2 at the
stimulus level.

Term	\hat{eta}	95% CI	t	df	p
Intercept	6.04	[4.99, 7.08]	11.35	6.88	< .001
LT Test	0.13	[0.05, 0.20]	3.46	25.38	.002

S8: By-item-pair preference scores across sessions



151

153

154

155

156

157

158

159

160

Table 3

Mixed-effects model results predicting IDS preference
during session 1 from IDS preference at session 2 at
the stimulus level.

Term	\hat{eta}	95% CI	t	df	p
Intercept	0.87	[0.45, 1.30]	4.04	122.79	< .001
Diff 1	0.10	[-0.02, 0.22]	1.63	6.31	.151

152 References

Byers-Heinlein, K., Bergmann, C., & Savalei, V. (2021). Six solutions for more reliable infant research. *Infant and Child Development*, e2296.

DeBolt, M. C., Rhemtulla, M., & Oakes, L. M. (2020). Robust data and power in infant research: A case study of the effect of number of infants and number of trials in visual preference procedures. *Infancy*, 25(4), 393–419.

ManyBabies Consortium. (2020). Quantifying sources of variability in infancy research using the infant-directed-speech preference. Advances in Methods and Practices in Psychological Science, 3(1), 24–52.