SABR Simulation for Risk Management

(Working Paper)

Maged Tawfik

January 12, 2026

1 Introduction

The SABR (Acronym for Stochastic alpha-beta-rho) model is a stochastic volatility model first studied by Hagan *et al*, 2002[6]. Due to its ability to replicate certain observed market phenomena, such as implied volatility smiles, frowns and skews, it has gained wide interest for the modeling of options. With its modeling power that can cover underlying assets that display normal, log-normal or CEV (Constant Elasticity of Variance) behavior, SABR is currently in extensive use in the area of fixed income derivatives.

Formally, assuming a filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$, SABR is given by the following equations

$$dF_t = F_t^{\beta}\sigma_t dW_t \qquad \beta \in [0,1],\ F_0 > 0 \tag{1.1a}$$

$$d\sigma_t = \nu\sigma_t dZ_t \qquad \nu \geq 0,\ \sigma_0 > 0 \tag{1.1b}$$

$$[dZ_t, dW_t] = \rho dt \qquad \rho \in [-1,1] \tag{1.1c}$$

where $F_t, \sigma_t \in \mathcal{F}_t$, $[.,.]$ is the quadratic variation operator and dW_t, dZ_t are standard Wiener increments.

Pricing derivatives based on the SABR model is done using either a PDE (Partial Differential Equation) solver based on the two-dimensional discretization of the Kolmogorov backward equation, or a Monte Carlo simulation approach. PDE solvers tend to have much faster convergence. On the other hand, each run is effective in pricing a single option. Conversely, a set of Monte Carlo paths, once generated, can be used to price any number of derivatives as long as they share the same underlying asset and expiry date. This makes Monte Carlo simulation a more favorable approach in risk management, where risk measures associated with very large fixed income derivative portfolios need to be evaluated.

[*]The opinions expressed in the paper are those of the author and do not necessarily reflect the view of Citibank.

[†]email: maged.tawfik@citi.com

2 Difficulties in SABR Modeling

To properly simulate the SABR model, we need to counter two essential difficulties. To explain the first difficulty, we use the following discretized version of (1.1)

$$\begin{aligned} F_t - F_{t-\Delta t} =& \bar{F}_t^{\beta} \sigma_? W_{\Delta t} \\ \sigma_t - \sigma_{t-\Delta t} =& \nu \bar{\sigma}_t Z_{\Delta t} \\ [W_{\Delta t}, Z_{\Delta t}] =& \rho \Delta t \end{aligned} \tag{2.1}$$

where $\bar{F}_t$ is an average value of F_s between time $t-\Delta t$ and t, $\bar{\sigma}_t$ is an average value of σ_s between time $t-\Delta t$ and t, and $\sigma_?$ is a representative value of σ between $t-\Delta t$ and t. The difficulty in choosing an appropriate value $\sigma_?$, when $\nu > 0$, stems from the fact that the continuous change of the value of σ is accompanied by a continuous change in F dependent on σ. Islah, 2009[9], showed that when $\beta \in \{0, 1\}$ or when $\rho = 0$, $\sigma_?$ can be exactly represented by the conditional average variance given as

$$\sigma_0 \sqrt{\Delta t \; I_{t-\Delta t}^{t}(\sigma_t)} \tag{2.2}$$

where

$$I_{t-\Delta t}^{t}(\sigma_t) := \frac{1}{\sigma_0^2 \, \Delta t} \int_{t-\Delta t}^{t} \sigma_s^2 \, ds \, \Big|_{\sigma_t} \tag{2.3}$$

Choi *et al*, 2025[2], give the following equivalent functional

$$I_{t-\Delta t}^{t}(\hat{Z}_1) \sim \int_0^1 \exp(2\nu \Delta t \, \hat{Z}_s) \, ds \, \big|_{\hat{Z}_1} \tag{2.4}$$

where $\hat{Z}_s \sim \mathcal{N}\left(-s\,\nu\sqrt{\Delta t}, \sqrt{s}\right)$. Although $I_{t-\Delta t}^{t}(\sigma_t)$ has a closed form PDF, as shown by Cai *et al*, 2017[4], there is no closed form for the CDF and its numerical integration is very difficult and highly error-prone. In their paper, Cai *et al* suggested the use of Laplace transform inversion to obtain an accurate and efficient representation of the CDF. We will be using a modified version of their Laplace inversion algorithm for the purposes of this paper.

The second difficulty in simulating a SABR process, is that the proxy for $\sigma_?$ given by (2.2) is no longer valid. No other exact proxy is currently available. As a result, one has to resort to plausible approximations such as those presented in [4],[2] and[6].

3 Sampling σ_t

In the process of simulating the SABR process, one may need to directly sample the volatility process itself. As per Equation (1.1b), σ_t follows a log-normal process. As a result, it can be sampled according

to

$$\begin{aligned}\sigma_t =& \sigma_0 \, \exp(\nu \sqrt{t} \, \hat{Z}_t) \\ \hat{Z}_t \sim& \mathcal{N}\left(-\tfrac{1}{2}\nu \sqrt{t}, \, 1\right) \; .\end{aligned} \tag{3.1}$$

Thus, σ_t is sampled *via* a single Gaussian variate.

4 Sampling the Average Variance

As explained in Section 2, the average volatility variance $I_0^t(\sigma_t)$ is needed for the simulation of the price process. As discussed earlier, the sampling of $I_0^t(\sigma_t)$ is fraught with difficulty, due to the lack of a closed form representation of its CDF. For the sake of presentational simplicity, we may notationally drop the σ_t dependency of I_0^h, in the remainder of the text, without changing the fact.

In[2], Choi *et al* proposed an approach for the sampling of I_0^t by matching its CDF to that of shifted log-normal ($\mathcal{SLN}(\mu, \sigma^2, \lambda)$) process with matching first four moments. The $\mathcal{SLN}$ process is defined as

$$\begin{aligned}Y \sim& \mathcal{SLN}(\mu, \sigma^2, \lambda) := \\ Y \sim& \mu \left[(1-\lambda) + \lambda \exp(\sigma X - \frac{\sigma^2}{2})\right] \qquad \mu > 0, \sigma > 0, 0 < \lambda \leq 1 \, ,\end{aligned} \tag{4.1}$$

where X is a standard Normal variate, where μ and σ^2 are the mean and variance of the LN process and λ is the shift parameter. In their work, Choi *et al* suggested setting $\lambda = \frac{5}{6}$ and sampling I_0^t according to

$$\begin{aligned}I_0^t \sim& \mathcal{SLN}\left(\mu, \sigma^2, \frac{5}{6}\right) \\ \sigma =& \sqrt{\ln\left(1 + \frac{36}{25} v^2\right)} \, ,\end{aligned} \tag{4.2}$$

where μ and v are drived from

$$\begin{aligned}\hat{Z} =& \frac{\ln(\sigma_t) - \ln(\sigma_0)}{\nu\sqrt{t}} \\ m_k(\hat{Z}) =& \frac{\Phi(\hat{Z} + k\nu\sqrt{t}) - \Phi(\hat{Z} - k\nu\sqrt{t})}{2k\nu\sqrt{t} \, \phi(\sqrt{\hat{Z}^2 + k^2\nu^2 t})} \\ c(\hat{Z}) =& \tfrac{1}{2}\left(\frac{\sigma_t}{\sigma_0} + \frac{\sigma_0}{\sigma_t}\right) \\ \mu =& \frac{\sigma_t}{\sigma_0} \, m_1(\hat{Z}) \\ v =& \left(\frac{\sigma_t}{\sigma_0}\right) \frac{\sqrt{m_2(\hat{Z}) - c(\hat{Z}) \, m_1(\hat{Z})}}{\mu \, \nu\sqrt{t}} \; - \mu \, ,\end{aligned} \tag{4.3}$$

where $\Phi(.)$ and $\phi(.)$ are standard normal CDF and PDF, respectively. Significant testing showed that while the approximation approach of (4.2) works reasonably well for t less than three years, it tends to diverge for larger time steps.

A more robust approach for the sampling of I_0^t is the exact approach of Cai *et al*[4], which is based on Laplace transform technique. In their paper they showed that the Laplace transform $L(\theta)$ of the CDF of $1/I_0^t$ is given by

$$\begin{aligned} L(\theta) &= \mathcal{L}(\theta; F_{1/I_0^t}(u)) = \int_0^\infty F_{1/I_0^t}(u)\ \exp(-\theta\, u)\, du \\ L(\theta) &= \frac{\sigma_0^2 t}{\theta}\ \exp\left\{\frac{[\ln(\sigma_t/\sigma_0)]^2 - [\psi(\theta\nu^2/\sigma_0^2)]^2}{2\nu^2\, t}\right\} \\ \psi(x) &= \operatorname{arcosh}\left(\frac{x\sigma_0}{\sigma_t} + \tfrac{1}{2}(\frac{\sigma_t}{\sigma_0} + \frac{\sigma_0}{\sigma_t})\right)\ . \end{aligned} \tag{4.4}$$

Several powerful numerical algorithms have been proposed in the literature for the inversion of Laplace transforms. We present here the Euler algorithm proposed by Abate and Whitt (2006)[1] for the inversion of (4.4), which reads

$$\begin{aligned} F_{1/I_0^t}(t) &= \frac{10^{\frac{M}{3}}}{t}\ \sum_{k=0}^{2M} (-1)^k \eta_k\ \Re\left(L(\frac{\beta_k}{t})\right) \\ \beta_k &= \frac{1}{3}\, M \ln(10)\ +\ \pi i k \\ \eta_0 &= \frac{1}{2}, \qquad \eta_k = 1,\ \ 1 \le k \le M \qquad \eta_{2M} = \ 2^{-M} \\ \eta_{2M-k} &= \eta_{2M-k+1}\ +\ 2^{-M}\binom{M}{k},\ \ 0 < k < M\ , \end{aligned} \tag{4.5}$$

where $i = \sqrt{-1}$, $\Re(.)$ is the real part of a complex number and M is chosen according to the desired level of accuracy. The number of correct significant digits is greater than $0.6M$, for well-behaved transforms. Notice that the CDF of I_0^t is equal to $1 - F_{1/I_0^t}$, since $I_0^t > 0$, and hence, the relationship between the two is a bijection and

$$\operatorname{sign}(\frac{d(1/I_0^t)}{dt}) = -\operatorname{sign}(\frac{dI_0^t}{dt})\ .$$

.

5 Sampling the CEV Process

Since the SABR model is a stochastic volatility generalization of the Constant Elasticity of Variance (CEV) model, sampling CEV variates is usually a needed step in the SABR simulation.

Definition 1 (CEV) *A CEV distribution which is denoted by*

$$F_t \sim \mathcal{CEV}_\beta(F_0, \sigma_0^2 t)\,, \tag{5.1}$$

is the distribution of F_t, which follows the stochastic process

$$dF_t = \sigma_0 F_t^\beta dW_t \quad \sigma \geq 0,\ F_0 \geq 0,\ 0 \leq \beta \leq 1\,, \tag{5.2}$$

where dW_t is a standard Wiener process.

Islah, 2009[9], showed that a SABR model with $\rho = 0$ is reducible to a CEV model by replacing $\sigma_0^2 t$ by I_0^t, and that the CEV CDF follows the following probability law

$$\begin{aligned}
\mathbb{P}(F_t = 0 \big| F_0, \sigma_0, \sigma_t, I_0^t) &= 1 \,-\, Q_{\chi^2}\left(A_0; \frac{1}{1-\beta}\right) \\
\mathbb{P}(F_t \leq u \big| F_0, \sigma_0, \sigma_t, I_0^t) &= 1 \,-\, Q_{\chi'^2}\left(A_0; \frac{1}{1-\beta}; C_0(u)\right) \quad u > 0\,,
\end{aligned} \tag{5.3}$$

where

$$A_0 = \frac{1}{I_0^t}\left(\frac{F_0^{1-\beta}}{1-\beta}\right)^2 \text{ and}$$

$$C_0(u) = \frac{1}{I_0^t}\left(\frac{u^{1-\beta}}{1-\beta}\right)^2,$$

and where $\chi^2(\delta)$ is the chi-squared distribution with δ degrees of freedom, and $\chi'^2(\delta, \gamma)$ is the non-central chi-squared distribution with δ degrees of freedom and γ non-centrality parameter. The sampling of F_t requires the numerical inversion of Equation (5.3), a process fraught with difficulty due to wide range of the parameters involved which lead to high degree of numerical instability.

A novel approach proposed by Choi *et al*,[2] based on the work of Makarov and Glew[12] and Kang[10] allows for the direct sampling of a CEV variate through the drawing of two gamma samples and one Poisson sample. Using their approach, we define

$$\begin{aligned}
z_t &:= z(F_t) \\
z(y) &:= \frac{u^{2(1-\beta)}}{(1-\beta)^2 \sigma_0^2 t I_0^t}\,.
\end{aligned}$$

A sample of z_t can be drawn according to

$$\begin{aligned}
Z_t &\sim 2\mathcal{G}(N+1) \\
N &\sim \mathcal{POIS}(\frac{z_0}{2} - X) \\
X &\sim \mathcal{G}\left(\frac{1}{2(1-\beta)}\right) \\
F_t &= \begin{cases} 0 & X \geq 0.5\, z_0 \\ \left(\sigma_0(1-\beta)\sqrt{t\, Z_t\, I_0^t}\right)^{\frac{1}{1-\beta}} & X < 0.5\, z_0\,, \end{cases}
\end{aligned} \tag{5.4}$$

where $\mathcal{G}(\alpha)$ is the gamma distribution with shape parameter α and unit scale parameter, and $\mathcal{POIS}(\lambda)$ is the Poisson distribution with intensity λ.

6 Approximating the SABR Process

As already mentioned in Section 2, the SABR process can be reduced to a CEV process, with $\sigma_0\sqrt{t\,I_0^t}$ playing the rôle of σ_t, whenever $\rho = 0$. As of the time of this writing, however, there is no known analytic expression of the general case of SABR dynamics with $\rho \neq 0$. In this section we present an approximation, based on an operator splitting technique, as suggested by Choi em et al[2]. According to this approach we rewrite Equation (1.1) as

$$\frac{dF_t}{F_t^\beta} = \sigma_t\,(\rho\,dZ_t + \sqrt{1-\rho^2}\,dZ_t^\perp) \quad \text{and} \quad \frac{d\sigma_t}{\sigma_t} = \nu\,dZ_t\,, \tag{6.1}$$

where dZ_t and $dZ_t^\perp$ are uncorrelated Wiener increments. Applying the operator splitting technique Equation (6.1) becomes

$$\frac{dF_t^{(1)}}{[F_t^{(1)}]^\beta} = \rho\,\sigma_t\,dZ_t \quad \text{and} \quad \frac{d\sigma_t}{\sigma_t} = \nu\,dZ_t \tag{6.2a}$$

$$\frac{dF_t^{(2)}}{[F_t^{(2)}]^\beta} = \sqrt{1-\rho^2}\,\sigma_t\,dZ_t^\perp \quad \text{and} \quad \frac{d\sigma_t}{\sigma_t} = \nu\,dZ_t\,. \tag{6.2b}$$

The operator splitting technique approximates the solution F_t of (6.1), with initial condition F_0, by $F_t \approx F_t^{(2)}\left(F_t^{(1)}(F_0)\right)$, where $F_t^{(1)}(F_0)$ is the solution $F_0^{(1)}$ of (6.2a), with initial condition $F_0^{(1)}=F_0$, and $F_t^{(2)}(F_t^{(1)}(F_0))$ is the solution $F_t^{(2)}$ of Equation (6.2b), with initial condition $F_0^{(2)} = F_t^{(1)}(F_0)$.

According to Section 2, Equation (6.2b) which obeys the SABR dynamics with zero correlation, is essentially a CEV process whose solution is

$$F_t^{(2)}|\sigma_t, I_0^t \sim \mathcal{CEV}_\beta\left(F_0^{(2)}, (1-\rho^2)\sigma_0^2 t I_0^t\right)\,. \tag{6.3}$$

This leaves us with Equation (6.2a), which has no known analytic solution. Choi em et al[2] suggested a frozen coefficient approximation in which they approximated $[F_t^{(1)}]^{(1-\beta)}$ by $F_0^{(1-\beta)}$, accordingly,

$$\begin{aligned}\frac{dF_t^{(1)}}{F_t^{(1)}} &= \frac{\rho\sigma_t}{[F_t^{(1)}]^{(1-\beta)}}\,dZ_t \\ &\approx \frac{\rho\sigma_t}{F_0^{(1-\beta)}}\,dZ_t\,.\end{aligned} \tag{6.4}$$

The solution $\tilde{F}_t^{(1)}$ of Equation (6.4) is

$$\tilde{F}_t^{(1)}(F_0) = F_0\,\exp\left(\frac{\rho}{\nu F_0^{(1-\beta)}}(\sigma_t - \sigma_0) - \frac{\rho^2\sigma_0^2 t I_0^t}{2F_0^{2(1-\beta)}}\right)\,. \tag{6.5}$$

Finally, conditional on σ_t and I_0^t, the CEV approximation of the general SABR process results in a terminal forward price F_t according to

$$F_t|\,\sigma_t, I_0^t \sim \mathcal{CEV}\left(\tilde{F}_t^{(1)},\, (1-\rho^2)\sigma_0^2\, t\, I_0^t\right) \tag{6.6}$$

As is the case with almost all approximation approaches of this nature, the accuracy of the final result increases in inverse proportion to the time step t.

7 Implied Volatility

Given an option pricing model, the *implied volatility* is a value, reflective of the variance of the stochastic process, that, when plugged into the model, gives the observed option price. Derivatives traders prefer implied volatility as a price quotation mechanism. This is due to the fact that it can provide stable option quotation that does not need to change with every movement of the underlying. Moreover, implied volatility reveals the nature of option prices as a function of option strike—generally referred to as the volatility smile.

Let $B(\sigma; u, k, \tau)$ be the price of an option of strike k, on an underlying of forward price u and price volatility σ, and which expires at time τ. Given a specific tuple (v, u_0, k, τ), where v the price of the option at time $t = 0$, the implied volatility of the option is

$$\sigma^{imp} = V(v; u_0, k, \tau)\,, \tag{7.1a}$$

that satisfies

$$B(V(v; u_0, k, \tau); u, k, \tau) = v\,, \tag{7.1b}$$

i.e. , $V(.) = B(.)^{-1}$.

When the dynamics of the underlying are believed to be governed by a geometric Brownian motion, the pricing model is log-normal and the implied volatility is often referred to as the Black or Black-Scholes volatility, σ^{BS}. For instance, in the case of a call option, σ^{BS} can be obtained from

$$B(\sigma; u_0, k, \tau) = u_0\, N(d_1) \;-\; k\, N(d2) \qquad \text{with} \qquad d_{1,2} = \frac{\log(u_0/k)}{\sigma^{BS}\sqrt{\tau}} \pm \frac{\sigma^{BS}\sqrt{\tau}}{2}\,, \tag{7.2}$$

where $N(.)$ is the CDF of the standard Gaussian distribution.

Similarly, when the dynamics are Gaussian in nature, the pricing model is Bachelier and the implied volatility is known as the normal volatility, σ^N, which in the case of a call option should satisfy

$$B(\sigma; u_0, k, \tau) = (u_0 - k)\, N(d_N) \;+\; \sigma^N\sqrt{\tau}\, n(d_N) \qquad \text{with} \qquad d_n = \frac{u_0 \;-\; k}{\sigma^N\sqrt{\tau}}\,, \tag{7.3}$$

where $n(.)$ is the PDF of the standard Gaussian distribution.

More relevant to the SABR model is the notion of the CEV implied volatility, σ^{CEV}, which is the volatility associated with the constant elasticity of variance model. In the case of a call option, σ^{CEV} obeys (see [5])

$$\begin{aligned} B(\sigma; u_0, k, \tau) = \bar{F}_{\chi^2}\left(\frac{k^{2\beta_*}}{[\sigma^{CEV}\beta_*]^2\,\tau}; 2 + \frac{1}{\beta_*}, \frac{1}{[\sigma^{CEV}\beta_*]^2\,\tau}\right) - \\ k\,F_{\chi^2}\left(\frac{1}{[\sigma^{CEV}\beta_*]^2\,\tau}; \frac{1}{\beta_*}, \frac{k^{2\beta_*}}{[\sigma^{CEV}\beta_*]^2\,\tau}\right), \end{aligned} \tag{7.4}$$

where $\beta_* = 1-\beta$, $\bar{F}_{\chi^2}(.) = 1 - F_{\chi^2}(.)$ and $F_{\chi^2}(x; r; x_0)$ is CDF of the non-central chi-squared distribution of r degrees of freedom and x_0 non-centrality parameter.

It goes without saying that any implied volatility can be converted to any other implied volatility. This is done by plugging the given implied volatility into the corresponding option pricing model. Once the option price is obtained, it can be used to obtain the desired implied volatility by, usually numerically, solving the relevant inverse pricing model $B^{-1}(.)$.

8 Asymptotic Representations

Since deriving closed form analytic formulas for derivatives pricing is, most of the time, impossible or impractical, a significant amount of research effort has been directed towards finding effective semi-analytic approximations. These are usually formulas for expressing implied volatility in terms of model parameters. In this section, we focus on semi-analytic representations of the SABR model. All formulas presented in this section will use the following normalized quantities,

$$\begin{aligned} f_t &= \frac{F_t}{F_0} \quad (f_0 = 1) \quad k = \frac{K}{F_0} \\ \hat{\sigma}_t &= \frac{\sigma_t}{\sigma_0} \quad (\hat{\sigma}_0 = 1) \quad \alpha = \frac{\sigma_0}{F_0^{\beta_*}}, \end{aligned} \tag{8.1}$$

where K is the option strike price. We will denote option expiry by T.

8.1 Hagan *et al*'s (HKLW) Formula

The Hagan approximation (2002)[6] is one of the most widely used semi-analytic approximations of the SABR model. It is very effective near the money and for small times to maturity. Hagan *et all* presented $O(T)$ approximations for both the normal and the Black-Scholes volatilities. Their normal approximation is more accurate. In a later paper, Hagan *el al* (2014)[7], derived a modified more accurate version of their earlier approximation. In the modified Hagan model, the normal implied volatility of the SABR model is given by the following chain of definitions:

$$
\begin{aligned}
\sigma^N =& \alpha \frac{q_N}{q} H(z)\,(1 + h_N T), \qquad z = \frac{\nu}{\alpha} q \\
q =& \int_1^k k^{-\beta}\, dk = \begin{cases} \frac{k^{\beta_*}-1}{\beta_*} & \text{if} \quad 0 \le \beta < 1 \\ \log k & \text{if} \quad \beta = 1 \end{cases} \\
q_N =& k - 1 \qquad (z_N = \frac{\nu}{\alpha} q_N) \\
H(z) =& \frac{z}{x(z)}, \quad x(z) = \log\left(\frac{V(z) + z + \rho}{1+\rho}\right), \quad V(x) = \sqrt{1 + 2\rho z + z^2} \\
h_N =& \log\left(k^{\beta/2} \frac{q}{q_N}\right) \frac{\alpha^2}{q^2} + \frac{\rho}{4} \frac{k^\beta - 1}{k-1} \alpha\nu + \frac{2 - 3\rho^2}{24} \nu^2 .
\end{aligned}
\tag{8.2}
$$

There are three special cases concerning Equation (8.2). First, when $z = 0$, $H(z)$ converges to 1. Secondly, when $\rho = 0$, $H(z)$ simplifies to $H(z) = z/arcsinh(z)$. Finally, the third case is the case of $\rho = -1$. In such a case, the value of $H(z)$ becomes

$$
H(z) = \begin{cases} 0 & \text{if} \quad z = 0 \\ \infty & \text{otherwise} . \end{cases} \tag{8.3}
$$

8.2 Choi and Wu's CEV Formula

In 2021 Choi and Wu[5] presented a powerful formula for the approximation of σ^{CEV}, based on Paulot's (2015) work[14]. This formula is better at handling out of the money options, large initial volatility and large vol-of-vol.

9 Option Portfolio

In the preceding sections we developed a framework for simulating the correlated movement of forward prices and volatilities at future time horizons. Such framework is useful for pricing options written at time t_0 with expiry that matches any of the simulation horizon. This assumes that the starting values of σ, ν, ρ, β are known at time t_0 and either, remain constant, or are given as known function in time. Moreover, options written on different tenors of the same interest rate curve have to reflect the stochastic dependence of the forward prices of the various tenors. Otherwise, simulation results may be totally unrealistic or even non-arbitrage-free. In this section we will address these two issues.

9.1 Forward Swap Volatility

Let us start by assuming the following risk neutral dynamics for the instantaneous forward rate $f(t,T)$, which is the rate at time T as observed at time t,

$$\begin{aligned} df(t,T) &= \sum_{i=1}^{n} \sigma_i(t,T)\,\bar{\sigma}_i(t,T)\,dt \;+\; \sum_{i=1}^{n} \sigma_i(t,T)\,dW_i(t) \\ \bar{\sigma}_i(t,T) &= \int_t^T \sigma_i(t,u)\,du\,, \end{aligned} \tag{9.1}$$

where $dW_i(t)$ are n standard uncorrelated Wiener increments.

Employing Ito's lemma and the arguments of HJM (see [8]), we get

$$\frac{dP(t,T)}{P(t,T)} = r(t)\,dt \;-\; \sum_{i=1}^{n} \bar{\sigma}_i(t,T)\,dW_i(t)\,, \tag{9.2}$$

where $r(t) = f(t,t)$.

The interest rate forward swap rate $S(t_x,t_s,t_e)$ is the coupon rate of the fixed leg of the swap, which is an $\mathcal{F}_t$-martingale under the annuity measure. The parameters t_s , t_e are the start and end dates of the swap, respectively, and t_x is the rate set date. Let the swap-corresponding annuity be defined as

$$A(t) = \sum_{i=0}^{n} \pi_i\,P(t,T_i) \qquad t \le t_s\,, \tag{9.3}$$

where n is the number of payments, T_i is the i-th payment date, $T_0 = t_s$, $T_n = t_e$ and π_i is a calendar factor. The value of $S(t_x,t_s,t_e)$, assuming simple calendar conventions, can be expressed as

$$S(t_x,t_s,t_e) = \frac{1-P(t_x,t_e)}{A(t_x)}\,. \tag{9.4}$$

Let us write the risk neutral dynamics of $S(t_x,t_s,t_e)$ as

$$dS(t,t_s,t_e) = \mu(t)\,dt \;+\; \sigma_{s,e,i}(t)\,dW_i(t)\,. \tag{9.5}$$

Employing Ito's lemma, we have, for $t \le t_s$

$$\begin{aligned} \sigma_{s,e,i}(t) &= \frac{1-P(t,t_e)}{A^2} \sum_{j=0}^{n} \pi_i P(t,T_j)\,\bar{\sigma}_i(t,T_j) \;+\; \frac{P(t,t_e)}{A(t)}\,\bar{\sigma}_n(t,t_e) \\ \sigma_{s,e,i}(t) &= \frac{1}{A(t)} \left(\frac{1-P(t,t_e)}{A(t)} \sum_{j=0}^{n} \pi_i P(t,T_j)\,\bar{\sigma}_i(t,T_j) \;+\; P(t,t_e)\,\bar{\sigma}_n(t,t_e) \right) \\ \sigma_{s,e,i}(t) &= \frac{1}{A(t)} \left(S(t,t_s,t_e) \sum_{j=0}^{n} \pi_i P(t,T_j)\,\bar{\sigma}_i(t,T_j) \;+\; P(t,t_e)\,\bar{\sigma}_n(t,t_e) \right)\,. \end{aligned} \tag{9.6}$$

Substituting from Equation (9.1), we get

$$\sigma_{s,e,i}(t) = \frac{1}{A(t)} \left(S(t, t_s, t_e) \sum_{j=0}^{n} \pi_i \, P(t, T_j) \int_t^{T_j} \sigma_i(t, u) \, du \; + \; P(t, t_e) \int_t^{t_e} \sigma_i(t, u) \, du \right) \; . \quad (9.7)$$

Equation (9.7) represents the forward swap volatility in terms of the terms structure volatilities of the instantaneous forward rates. Using this equation one can construct a set of arbitrage-free swap volatilities, given a set of instant forward volatilities. Similarly, given a set of *consistent* swap volatilities, one should be able to deduce, at least numerically, the instantaneous forward volatilities. The consistency condition is needed due to the fact that the nonlinear function of Equation (9.7) is necessarily invertible.

9.2 Historical or Implied Volatility

Since true volatilities are not market observable, a proxy is always needed for their estimation. Practitioners either, resort to the liquid options market and use the implied volatility as a proxy, or resort to history and use realized volatility as a proxy. We propose the use of the options market implied volatilities to estimate the current instantaneous forward volatilities. For risk management, a stochastic volatility model is usually needed. The model can be in the form of a stochastic differential equation, or in the form of a time series GARCH style model (see Tawfik (2025)[16] for an example). The volatilities estimated from the options market can represent the starting values of the stochastic model. The parameters of the model can then be estimated from the historical data. Even then, one has a choice of either, resorting to the history of swaption implied volatilities, or the history of realized swap volatilities.

9.3 Instantaneous Forward Volatility Model

9.4 Options on Different Tenors of Same Curve

Under the SABR model, the stochastic dependence across options written on different tenors of the same curve is made of two components. First is the correlation structure among the forwards themselves. Second is the correlation structure among the SABR volatility.

If F_t^i and σ_t^i are the price and the SABR volatility of the forward contract associated with the i's tenor at time t, then the following represents the associated Wiener increments

$$dW_t^i = \frac{dF_t^i}{\sigma_t^i\,(F_t^i)^\beta} \tag{9.8a}$$

$$dZ_t^i = \frac{dZ_t^i}{\nu^i\,\sigma_t^i} \tag{9.8b}$$

$$[dW_t^i,\, dZ_t^i] = \rho^i\,dt\,, \tag{9.8c}$$

where β^i and ν^i are the tenor specific SABR parameters. A time series for dW_t^i and dZ_t^i for each of the N_T tenors can be constructed based on the forward prices and the corresponding shortest expiry option prices. Given a forward price, SABR model parameters are obtained by fitting the model to the prices of options of same expiry and various strikes. Having the individual tenor-specific time series, of size N_t each, we then construct the two associated correlation matrices of the space of all tenors: the forward price correlation matrix C_f and the volatility correlation matrix C_σ. The correlation matrices $C^{N_t\times N_T}$ can then be diagonalized, using the singular value decomposition (SVD), into the form $U^{N_t\times N_t}\,\Sigma^{N_t\times N_T}\,V^{N_T\times N_T}$, where Σ is a diagonal matrix whose elements are in a non-decreasing order. It is typical for the correlation structure of the whole space to be explained be no more than the four largest values of Σ.

To include the stochastic dependence among options written on the same curve we modify Equations (1.1) to read

$$dF_t^i = (F_t^j)^\beta \sigma_t dW_t^j \qquad \beta\in[0,1],\ F_0^i>0 \tag{9.9a}$$

$$d\sigma_t^i = \nu\sigma_t^i dZ_t \qquad \nu\geq 0,\ \sigma_0^i>0 \tag{9.9b}$$

$$[dZ_t^i, dW_t^i] = \rho^i dt \qquad \rho\in[-1,1] \tag{9.9c}$$

$$dZ_t^i = (1-\sum_{j=1}^{N_\sigma}\gamma_j^2)^{\frac{1}{2}} d\tilde{Z}_t^i + \sum_{j=1}^{N_\sigma}\gamma_{ij} d\tilde{Z}_t^j \tag{9.9d}$$

$$dW_t^i = (1-\sum_{j=1}^{N_f}\lambda_{ij}^2)^{\frac{1}{2}} d\tilde{W}_t^i + \sum_{j=1}^{N_f}\lambda_{ij} d\tilde{W}_t^j \tag{9.9e}$$

where N_f and N_σ are the number of the largest eigenvalues λ_{ij} sufficient to explain the total variances of the set of forwards and their option volatilities, respectively.

9.5 Options on Same Underlying

When given a portfolio of options written on the same underlying, but with different strike prices and expiry dates, the time dependency of the SABR parameters is implicitly hidden in the prices of these options. In this section, we will develop an approach for extracting the implied risk-neutral time dependency information from an option portfolio.

Let Π be a portfolio of a collection of at-the-money options. Each option $C_i\in\Pi$ is associated with a tuple of SABR parameters $(\sigma,\nu,\rho,\beta,\tau)_i$. The SABR parameters of C_i are identified by examining

all available option prices of same expiry on the same underlying. For SABR parameter identification techniques, see

10 Implied Volatility Surface

Let $\mathbb{D}$ be defined as

$$\mathbb{D} = (0, \infty) \times [0, 1] \times [-1, 1] \times [0, \infty) \qquad x \in \mathbb{D} = (\sigma, \beta, \rho, \nu)\,, \tag{10.1}$$

where σ, β, ρ, ν are the SABR parameters associated with an option on a forward contract of tenor t and which expires at time τ. Also, let the functions $\hat{S}, \hat{T}$ be defined according to

$$\begin{aligned} &\hat{S} : I_1 \times I_2 \longrightarrow \mathbb{D} \\ &\hat{S} : (i, j) \mapsto (\sigma_{ij}, \beta_{i,j}, \rho_{i,j}, \nu_{i,j}) \\ &\hat{T} : I_1 \times I_2 \longrightarrow \mathbb{R}^+ \times \mathbb{R}^+ \\ &\hat{T} : (i, j) \mapsto (t_i, \tau_j) \end{aligned} \tag{10.2}$$

where, I_1 and I_2 are two finite ordered index sets. Now, we define the implied volatility surface as the function $S_h(t, \tau; \hat{T}^{-1}, \hat{S}) : \mathbb{R}^+ \times \mathbb{R}^+ \to \mathbb{D}$ for all $(t, \tau) \in \mathbb{R}^+ \times \mathbb{R}^+$, where h is the valuation date, which can be in the future. In other words, the volatility surface is a function that maps any tenor-expiry combination into a set of SABR parameters by interpolating the set of parameters given for a select finite set of options. Typically, the set of given parameters forms a grid pattern, *i.e.* $\hat{S}$ is surjective.

In order for the volatility surface, S_h, to be useful in proving a price for an option, it has to be accompanied by a price $F_h(t)$ for the underlying forward contract of tenor t. In case of the SABR model, F_h and S_h are simulated together due to their coupled processes.

11 Control Variable

It is not uncommon in a risk management scenario for the set $I_1 \times I_2$ to have a size in the hundreds. Estimating portfolio risk requires the simulation of the volatility surface over hundreds of forward horizons. It is not hard to see how the computational complexity grows very rapidly. As a result, each simulation of the volatility surface has to be limited to a number of paths that is far less than that which needed for accurate simulation of an option price. To alleviate that deficiency, we resort to a technique know as the *control variable* technique as a means of variance reduction and, consequently, accuracy improvement.

Our control variable approach uses the prices of four control options struck at different strikes, for every node on the implied volatility grid and for every simulation horizon, to adjust the distribution of the forward price F_h of the simulated sample paths. The philosophy behind this stems from the fact

that the true forward price distribution can be ascertained from the option prices of options struck at different strikes. Given a forward price $F_0(t)$ and an implied volatility set $S_0(t,\tau)$, one can accurately price options of various strikes that share the tenor t and the expiry h. This is usually done using a semi-analytic technique, such as those discussed in Section 6, additional Monte Carlo simulation or a partial differential equation (PDE) solver such as the finite difference method (FDM). Once the option prices are obtained, we compare their values with those predictable by our simulation. The discrepancy, is then, corrected by proportionately adjusting the values of F_h of the simulation paths.

12 The Algorithm

13 Negative Rates

14 Results

15 PCA Complexity Reduction

16 Bayesian Volatility

Assuming a standard filtered probability space $(\Omega, \mathcal{F}, \mathcal{F}_t, P)$, let the processes $F_t(F_0, \sigma_t, I_t),\ \sigma_t(\sigma_0)$ and $I_t(\sigma_t)$ be $\mathcal{F}_t$-measurable according to the SABR dynamics, stated in the preceding sections. In this section, we are concerned with finding the best estimate for σ_t, where F_t has been generated from an exogenous process. Using Bayesian statistics, we can write

$$P(\sigma_t = s|F_t = f) = \frac{P(F_t = x|I_t = a, \sigma_t = s)P(I_t = a|\sigma_t = s)P(\sigma_t = s)}{P(F_t = f)}\,. \tag{16.1}$$

Or more precisely,

$$P(\sigma_t = s|F_t = f) = \frac{P(F_t = f|I_t \in da, \sigma_t \in ds)P(I_t = a|\sigma_t \in ds)P(\sigma_t = s)}{P(F_t = f)} \tag{16.2a}$$

$$P(F_t = f) = \int_0^\infty \int_0^\infty P(F_t = f|I_t \in da, \sigma_t \in ds)\, P(I_t = a|\sigma_t \in ds)\, P(\sigma_t = s)\, da\, ds\,. \tag{16.2b}$$

Since we have already shown, in the early part of this paper, how to compute the distributions $P(F_t = f|I_t \in da, \sigma_t \in ds),\ P(I_t = a|\sigma_t \in ds)$ and $P(\sigma_t = s)$, we can now, at least numerically, compute the distribution $P(\sigma_t = s|F_t = f)$ and its CDF $Q_\sigma(s|F_t = f)$ defined as,

$$
\begin{aligned}
Q_\sigma(s|F_t = f) &= P(\sigma_t \leq s|F_t = f) \\
&= \int_0^s P(\sigma_t = s|F_t = f)\, ds\,.
\end{aligned}
\tag{16.3}
$$

To sample the distribution $Q_\sigma(.)$, we need to compute $Q_\sigma^{-1}(U)$, where $U \sim Uniform(0,1)$ and Q_σ^{-1} is the inverse of $Q_\sigma(.)$

References

[1] Abate, Joseph and Whitt, Ward, 2006, "A Unified Framework for Numerically Inverting Laplace Transforms". doi:10.1287/ijoc.1050.0137

[2] Choi, Jaehyuk; Hu, Lilian and Kwok, Yue Kuen, 2025, "Efficient and Accurate Simulation of the Stochastic-alpha-beta-rho Model," arXiv:2408.01898v2

[3] Choi, Jaehyuk; Kwak, Minsuk; Tee, Chyng Wen; and Wang, Yumeng, 2022, "A Black-Scholes user's guide to the Bachelier model," *Journal of Futures Markets* 42(5):959-980. https://ink.library.smu.edu.sg/lkcsb_research/6869

[4] Cai, Ning; Song, Yingda and Chen, Nan, 2017, "Exact Simulation of the SABR Model," *Operations Research* 65(4):931-951. doi:10.1287/opre.2017.1617

[5] Choi, Jaehyuk and Wu, Lixin, 2020, "The Equivalent Constant-elasticity-of-variance (CEV) Volatility of the Stochastic-Alpha-Beta-Rho (SABR) Model," arXiv:1911.13123v3

[6] Hagan, Patrick; Kumar, Deep; Lesnieweski, Andrew S. and Woodward, Diana, 2002, "Managing Smile Risk," *Wilmott* 1(2002):84-108.

[7] Hagan, Patrick; Kumar, Deep; Lesnieweski, Andrew S. and Woodward, Diana, 2014, "Arbitrage-Free SABR," *Wilmott* (2014):60-75. doi:10.1002/wilm.10290

[8] Heath, D., Jarrow, R. and Morton, A. (1992), "Bond Pricing and the Term Structure of Interest Rates: A New Methodology for Contingent Claims Valuation," *Econometrica* 60(1):77-105. doi:10.2307/2951677

[9] Islah, Othmane, 2009, "Solving SABR in Exact Form and Unifying It with LIBOR Market Model," SSRN: http://papers.ssrn.com/sol3/papers.cfm?astract-id=1489428

[10] Kang, C., 2014, "Simulation of the Shifted Poisson Distribution with an Application to the CEV Model," Management Science and Financial Engineering, 20:27-32. doi:10.7737/MSFE.2014.20.1.027

[11] Cheng, Luo, 2016, "On the Calibration of the SABR Model and its Extensions," Thesis submitted as part of the requirements for the award of the MSc in Mathematics and Finance, Imperial College London.

[12] Makarov, R.N. and Glew, D., 2010, "Exact Simulation of Bessel Diffusions," *Monte Carlo Methods and Applications* 16:312-347. doi:10.1515/mcma.2010.010. arXiv:0910.4177

[13] Matsumoto, Hiroyuki and Yor, Marc, 2005, "Exponential functionals of Brownian motion, I: Probability laws at fixed time," *Probability Surveys* 2:312-347. doi:10.1214/154957805100000159

[14] Paulot, Louis, 2015, "Asymptotic Implied Volatility at the Second Order with Application to the SABR Model," In: Friz, P., Gatheral, J., Gulisashvili, A., Jacquier, A., Teichmann, J. (eds) *Large Deviations and Asymptotic Methods in Finance*. Springer Proceedings in Mathematics & Statistics, (110) Springer, Cham. https://doi.org/10.1007/978-3-319-11605-1_2 arXiv:0906.0658v3

[15] van der Stoep, Anthonie and Grzelak, Lech Aleksander and Oosterlee, Cornelis W., 2015, "The Time-Dependent FX-SABR Model: Efficient Calibration Based on Effective Parameters," *International Journal of Theoretical and Applied Finance*, 18(6), SSRN: https://ssrn.com/abstract=2503891 doi:10.2139/ssrn.2503891

[16] Tawfik, Maged, 2025, "Monte Carlo Simulation of SOFR Yield Curve," *Working Paper*.