# SUMMARY OF PRODUCT CHARACTERISTICS

## 1. NAME OF THE MEDICINAL PRODUCT

Kestamol 500 mg film-coated tablets

## 2. QUALITATIVE AND QUANTITATIVE COMPOSITION

Each film-coated tablet contains 500 mg paracetamol.

The finished product complies with the British Pharmacopoeia (BP) monograph for Paracetamol Tablets.

For the full list of excipients, see section 6.1.

## 3. PHARMACEUTICAL FORM

Film-coated tablet.

White to off-white, capsule-shaped film-coated tablets, 17.5 mm × 7.3 mm, with a score line on one side and debossed with “K500” on the other. The score line is only to make swallowing easier and not to divide into equal doses.

## 4. CLINICAL PARTICULARS

### 4.1 Therapeutic indications

Kestamol 500 is indicated for the treatment of mild to moderate pain, including headache, toothache, period pain and muscular aches, and for the reduction of fever.

### 4.2 Posology and method of administration

**Posology**

| Age | Dose | Maximum in 24 hours |
|---|---|---|
| Adults and children 12 years and over | 1 to 2 tablets every 4 to 6 hours as required | 8 tablets (4 g) |
| Children 6 to 11 years | ½ to 1 tablet every 4 to 6 hours as required | 4 tablets (2 g) |
| Children under 6 years | Not recommended | |

Leave at least 4 hours between doses. Do not take for more than 3 days without consulting a doctor.

**Method of administration**

For oral use. Swallow the tablets whole with a glass of water.

### 4.3 Contraindications

Hypersensitivity to paracetamol or to any of the excipients listed in section 6.1.

### 4.4 Special warnings and precautions for use

Use with caution in patients with hepatic or renal impairment, chronic alcohol use, or glutathione depletion. Patients should be advised not to take other paracetamol-containing products at the same time.

### 4.5 Interaction with other medicinal products

The anticoagulant effect of warfarin may be enhanced by prolonged regular use of paracetamol. The rate of absorption of paracetamol may be increased by metoclopramide and reduced by colestyramine.

SPECIMEN – fictional test document. The company, product and all numbers are invented. Not medical advice.

### 4.6 Fertility, pregnancy and lactation

Paracetamol can be used during pregnancy and breast-feeding if clinically needed, at the lowest effective dose for the shortest possible time.

### 4.7 Effects on ability to drive and use machines

None known.

### 4.8 Undesirable effects

Adverse effects are rare at therapeutic doses. Very rare: hypersensitivity reactions including skin rash, blood dyscrasias, and serious skin reactions.

### 4.9 Overdose

Overdose may cause liver damage, which may not be apparent for 24 to 48 hours. Immediate medical attention is needed even if there are no symptoms. Treatment with acetylcysteine should be considered.

## 5. PHARMACOLOGICAL PROPERTIES

### 5.1 Pharmacodynamic properties

Pharmacotherapeutic group: other analgesics and antipyretics, anilides. ATC code: N02BE01.

Paracetamol has analgesic and antipyretic actions, thought to be mediated mainly through inhibition of prostaglandin synthesis in the central nervous system.

### 5.2 Pharmacokinetic properties

Paracetamol is rapidly and almost completely absorbed from the gastrointestinal tract, with peak plasma concentrations 30 to 60 minutes after an oral dose. It is metabolised mainly in the liver and excreted in the urine. The plasma half-life is 1 to 3 hours.

## 6. PHARMACEUTICAL PARTICULARS

### 6.1 List of excipients

*Tablet core:* maize starch, pregelatinised starch, povidone K30, stearic acid, magnesium stearate.

*Film coat:* hypromellose, titanium dioxide (E171), macrogol 400.

### 6.2 Incompatibilities

Not applicable.

### 6.3 Shelf life

3 years.

### 6.4 Special precautions for storage

Do not store above 30 °C. Store in the original package in order to protect from light and moisture.

### 6.5 Nature and contents of container

Clear PVC/aluminium foil blister strips of 10 tablets, packed in a printed cardboard carton with a package leaflet.

Pack sizes: 10 tablets (1 × 10) and 100 tablets (10 × 10). Not all pack sizes may be marketed.

### 6.6 Special precautions for disposal

No special requirements.

SPECIMEN – fictional test document. The company, product and all numbers are invented. Not medical advice.

## 7. MARKETING AUTHORISATION HOLDER

Kestrel Pharmaceuticals Pvt. Ltd.
4th Floor, Harbour View Tower, Nariman Point, Mumbai 400021, Maharashtra, India

## 8. MARKETING AUTHORISATION NUMBERS

| Country | Number | Date of first authorisation |
|---|---|---|
| India | NMLO/MA/2019/04417 | 12 March 2019 |
| Sri Lanka | SL/REG/2022/0815 | 4 August 2022 |

## 9. DATE OF REVISION OF THE TEXT

January 2025

SPECIMEN – fictional test document. The company, product and all numbers are invented. Not medical advice.