

Table 2 Baseline characteristics of the verified eligible cohorts

Footnote: baseline is the nearest valid preoperative examination. Show standardized differences in a supplementary column if useful; do not use baseline p-values to choose model predictors. Sex/laterality percentages use known documented categories and explicitly state the denominator.

Characteristic	TG-PRK + ACXL	ACXL alone	Missing / note
Eyes / unique patients	[n / N]	[n / N]	Report overall unique patients separately.
Age at index treatment, years	[n; mean(SD); median(IQR)]	[n; mean(SD); median(IQR)]	[missing n per cohort]
Recorded sex, categories	[n/N (%)]	[n/N (%)]	[unknown n]
OD / OS	[n/N (%)]	[n/N (%)]	[unknown n]
Baseline CDVA, logMAR	[n; mean(SD); median(IQR)]	[n; mean(SD); median(IQR)]	[missing n]
Baseline UDVA, logMAR	[n; mean(SD); median(IQR)]	[n; mean(SD); median(IQR)]	[missing n]
Sphere / cylinder / spherical equivalent, D	[separate rows in results]	[separate rows in results]	[missing n per measure]
K1 / K2 / Kmax, D	[separate rows in results]	[separate rows in results]	[missing n per measure]
TCT, micrometres	[n; mean(SD); median(IQR)]	[n; mean(SD); median(IQR)]	[missing n]
Amsler–Krumeich stage, if verified	[n/N (%) per stage]	[n/N (%) per stage]	[unknown n]
Baseline-to-surgery interval, days	[n; median(IQR)]	[n; median(IQR)]	[missing date n]