

2 Figure specifications

Figure	Specification	Readiness / interpretation
1. Cohort flow	Source records → verified cohorts → exclusions/pending → endpoint availability → prediction sample; eyes and patients at each stage.	Counts must reconcile; planned sample is a note, not a result.
2. Missingness map	Variables × record rows, with missing/observed/unverified distinctions; companion per-cohort percentage chart.	Can be produced from the current audit without claiming model readiness; no identifiers displayed.
3. Baseline-to-one-year CDVA	Paired eye trajectories or distribution plots by treatment; show paired n and actual window.	Only dated verified outcomes; do not connect unavailable observations.
4. Observed versus predicted CDVA	Held-out predictions versus observed logMAR; identity line, treatment colours and population n.	Requires valid grouped evaluation. Each eye's mean held-out prediction is a display summary, not extra data.
5. Calibration and residuals	Held-out calibration fit with identity line; residual-versus-prediction panel; show uncertainty and sparse-data limitations.	Continuous-outcome calibration; no arbitrary risk groups.
6. Predictor associations	Original-unit/standardized ridge coefficient plot with resampling variability and units.	Predictive associations; correlated features and shrinkage affect magnitude.
7. Error comparison	MAE/RMSE and paired ridge-minus-benchmark differences with approximate patient-cluster CIs.	No plot if model was not run; no clinical superiority or deployment claim.