

(3 pts) 1. Explain co-option of genes. Provide an example.

2. The longnose butterflyfish (right, on screen) has an unusually long mouth compared to other species of butterflyfishes (left). You plan to study the genetics of jaw length in the butterflyfishes.

(2 pts) (a) Identify one toolkit gene that you would be wise to study first.

(5 pts) (b) Explain the existing evidence that caused you to identify this toolkit gene.

(5 pts) 3. Describe one type of fossil or genetic evidence that explains why evolution across the Ediacaran-Cambrian boundary was not “explosive.”


(2 pts E.C.) 4. Name two of the maternal effect genes that control the earliest development of the *Drosophila* egg, before *Hox* genes are used.

(5 pts) 5. Why was photosynthesis by cyanobacteria important to the evolution of increased body size for single-celled eukaryotes?

(10 pts) 6. The snapping shrimp *Synalpheus brooksi* lives in sponges on coral reefs. Duffy (1996) showed that shrimp occupying nearby but different species of sponges rarely come into contact with each other because they don't usually leave their sponges. In the lab, a male shrimp from one sponge species is extremely aggressive towards females from a different sponge species, suggesting reproductive incompatibility. Analyses showed that shrimp from the same sponge species were genetically similar but shrimp from different sponge species were genetically distinct. Duffy argued the shrimp in different sponges are different species. If so, explain which prezygotic mechanism or mechanisms reproductively isolate shrimps in different sponge species.

- (5 pts) 7. Are dinosaurs extinct? Explain why or why not. Use the concepts of monophyly (or not monophyletic, if necessary) to justify your explanation.
- (5 pts) 8. Whales are most closely related to hippopotamuses but lack rear limbs. Explain how a change to *Hox* gene expression might cause the loss of rear limbs.
- (5 pts) 9. Explain ecological release and how it might have contributed to the Cambrian radiation.

The phylogeny below shows the relationships of gobies (fishes) in the genus *Elacatinus*. Each species has a stripe on that side. The color of the stripe for each species is in parentheses. The vertical line indicates the behavior of each species (sponge-dwelling, hovering, cleaning). Every species to the left of the vertical line has that behavior. Based on this information, answer the next two questions.



(5 pts) 10. Circle the *largest* monophyletic group you can identify based on color *or* behavior. Be clear which species are included / excluded. Explain why that group is monophyletic. Do not circle the entire tree. That would be incorrect.

(10 pts) 11. Does color or behavior better explain most differences among closely related species? Explain why, providing specific examples.

(3 pts) 12. Tell how heterozygosity changes in a population with non-random mating.

(12 pts) 13. The purple-throated carib is a species of hummingbird that lives on the Caribbean island of Dominica. Female caribs have longer, more curved bills than male caribs, which have shorter, straighter bills. Female hummingbirds get their nectar only from the red flowers of a plant called *Heliconia bihai*. Male hummingbirds get their nectar only from the yellow flowers of the same plant species. An individual plant has either red or yellow flowers, but not both. The flower shape of each color morph corresponds to the bill shape of each sex. Red flowers have a longer, more curved flower corolla, while yellow flowers have a shorter, straighter corolla. The hummingbirds function as pollinators for *H. bihai*. No other pollinators (including wind) for this plant are known. Based on this information only, can reproductive isolation evolve for (1) the hummingbirds and (2) the color morphs of *H. bihai*. If so, say how. Justify your answer.