

Problem 1

Theorem 1.1. Any tree T has at least $\Delta(T)$ leaves.

Lemma 1.1.1. Any connected subgraph of a tree is a tree as well.

Proof. If a graph $G = (V, E)$ is acyclic then E does not contain any cyclic subset and hence there is no cyclic subgraph of G . From these considerations, any connected subgraph of G is acyclic and therefore a tree. \square

Lemma 1.1.2. Any tree T with $\Delta(T) + 1$ vertices has $\Delta(T)$ leaves ($\Delta(T) \geq 1$).

Proof. Let $v_0 \in V_T$ denote the vertex of $T = (V_T, E_T)$ with $d(v_0) = \Delta(T)$.

For any $v, w \in V_T \setminus \{v_0\}$ ($v \neq w$):

- $d(v) \geq 1$: v_0 is adjacent to v since $|V_T| = \Delta(T) + 1 \Rightarrow d(v) \geq 1$
- $d(v) \leq 1$: Any edge $\{v, w\} \in E_T$ would create a cycle (v_0, v, w, v_0) which would render T invalid as a tree.

$\Rightarrow \forall v \in V_T \setminus \{v_0\} : d(v) = 1 \Leftrightarrow v$ is a leaf. Notably, $|V_T \setminus \{v_0\}| = \Delta(T)$. \square

Lemma 1.1.3. For any tree $T = (V_T, E_T)$ with $|V_T| > \Delta(T) + 1 \geq 3$, there exists a partition (S, S') of T with $\Delta(S) + \Delta(S') = \Delta(T)$.

Proof. Let $v_0 \in V_T$ denote a vertex with $d(v_0) = \Delta(T)$. Since $|V_T| > \Delta(T) + 1$, v_0 has at least one non-leaf adjacent vertex $v_1 \in V_T$. Now, let (S_0, S_1) denote the partition at the edge $\{v_0, v_1\}$ whereby S_0 contains v_0 and (S_1, v_1) respectively.

As seen in Lemma 1.1, S_0 and S_1 are trees.

- case $d(v_1) = 2$: In this case, a cut at the edge $\{v_0, v_1\}$ would make

\square

Proof by induction. Let $T = (V_T, E_T)$ be a tree.

Basis: $\Delta(T) = 1$

Inherently, $V_T = \{v_1, \dots, v_m\}$ ($v_i = v_j \Rightarrow i = j$) and $E_T = \{\{v_1, v_2\}, \dots, \{v_{m-1}, v_m\}\}$ ($m, i, j \in \mathbb{N}$). Any modification would violate our preconditions. v_1, v_m are the only vertices with degree 1. Therefore, the number of leaves is $2 \geq \Delta(T)$.

Step: For some $n \in \mathbb{N}$: Any $T = (V_T, E_T)$ with $\Delta(T) = n$ has at least n leaves.

Let $T' = (V_{T'}, E_{T'})$ be a tree with $\Delta(T') > n$. \square

Problem 2

Theorem 2.1. *If any removal of an edge increases the number of connected components of a graph G then G is acyclic.*

Proof. Let $S = (V_S, E_S)$ be a connected component of G . Removing any edge $e \in E_S$ increases the number of connected components and thereby disconnects S .

If S had contained a cycle C then there would not have been any unique paths in C . Hence, no edge removal would have an effect on the connectivity of S (removing an edge still leaves a path between any two vertices). Hereby, S had to be acyclic.

If any connected component of G was acyclic then G would be acyclic, too. \square

Theorem 2.2. *If adding any edge introduces a cycle in an acyclic graph $G = (V, E)$ then any two vertices in G are joined by a unique path.*

Proof. If adding an edge $\{v_0, v_1\}$ joining two non-adjacent vertices $v_0, v_1 \in V$ introduces a cycle (v_0, \dots, v_1, v_0) then there had to be *at least* one path from v_0 to v_1 .

Furthermore, if there was *more than one* path joining v_0 and v_1 then there would have already been a cycle (but G is acyclic). \Rightarrow Any vertex had to be joined by a unique path. \square

Theorem 2.3. *If any two vertices in a graph are joined by a unique path then any removal of an edge increases the number of connected components.*

Proof. Let $G = (V, E)$ be a graph in which all vertices are joined by a unique path. Let $e = \{v_0, v_1\} \in E$ be an edge. Thus, the unique path from v_0 to v_1 runs over (and is exactly) e . From these considerations, removing e would make v_1 inaccessible from v_0 and would thereby increase the number of connected components. \square

Problem 3

Theorem 3.1. *Either a graph or its complement is connected.*

Proof. If G was connected, we would already been done.

Let $G = (V, E)$ be disconnected. Moreover, let $S = (V_S, E_S)$ and $T = (V_T, E_T)$ be two connected components of G ($S \neq T$).

Now let $u, v \in V_S \cup V_T$ be vertices in these components.

- **Case 1:** $u \in V_S$ and $v \in V_T$.

Then G does not contain the edge $e = \{u, v\}$. Otherwise, S and T were interconnected. From these considerations, the graph's complement \bar{G} does contain e .

- **Case 2:** $u, v \in V_S$:

Considering that V_T is not empty, then there is a vertex $w \in V_T$ such that the edges $\{u, w\}$ and $\{v, w\}$ exist in \bar{G} (see Case 1). From these considerations, \bar{G} also contains the path (u, w, v) .

Therefore, all vertices in a *disconnected graph* G are connected in \bar{G} and hereby \bar{G} is connected.

□

Problem 4

Theorem 4.1. *If u and v are the only vertices of odd degree in a graph then there is a u - v -path.*

Proof.

□