


Bird migration paths from eBird data

Seasonal maps and animated population centroids for the Arctic tern, common swift, and red knot — from hundreds of millions of citizen-science observations

Wolfram Community submission — 2026



Cover animation. A slowly rotating dark globe on which each species' monthly population centroid glides as a glowing comet along its great-circle migration loop (Arctic tern in orange, common swift in cyan, red knot in gold), each over a faint full-year route. The centroids are computed per coherent flyway, not pooled globally — the central methodological point of this post. The code that produces every track is given in §3–§5.

Abstract

This notebook reconstructs the annual migration of three iconic birds — the **Arctic tern** (*Sterna paradisaea*, a pole-to-pole Atlantic-basin migrant), the **common swift** (*Apus apus*, an Afro-Palearctic aerialist), and the **red knot** (*Calidris canutus*, a long-haul Americas-flyway shorebird) — entirely from open citizen-science data, in pure Wolfram Language.

The pipeline pulls eBird observations through **GBIF's** eBird Observation Dataset (licensed **CC BY 4.0**, with a citable DOI), computes an **effort-bias-aware monthly population centroid** using presence weighting and a proper **spherical mean** (a 3-D unit-vector average, essential for circumpolar and dateline-crossing species), builds seasonal occurrence maps and an animated centroid track, **validates** the result against Cornell's eBird Status & Trends, and adds a live “right now” layer from the eBird API.

The central lesson is methodological: a single *global* centroid is meaningless for a species spread across several disjoint flyways — pooling the red knot's three flyways puts its “centre” in the mid-Atlantic, displaced ~3,800 km/month from where the Americas population actually is. Computing one centroid *per coherent flyway* fixes this. Throughout, the prose opens each section intuitively for the general reader, then deepens into the quantitative method for the specialist.

Setup (optional — only needed to run the code)

You can ignore this section entirely if you just want to read the post. Every figure is pre-rendered into the notebook as a static image, so the prose, maps, and animations all work without evaluating anything.

If you *do* want to run the code cells, the helper functions (spherical centroid, centroid-CSV loaders, orthographic globe, live-sightings fetcher) live in a single attached package, `birdflightpaths_helpers.wl`, uploaded alongside the notebook. Run the cell below once after opening the notebook; it loads every helper symbol used in the rest of the post.

```
(* Put birdflightpaths_helpers.wl in the same folder as this
notebook, then shift-Enter. The package sets $BFPDataDir-relative
paths; point it at your local data/ folder if needed. *)
SetDirectory[NotebookDirectory[]];
$BFPDataDir = FileNameJoin[{NotebookDirectory[], "data"}];
Get["birdflightpaths_helpers.wl"];
```

1. Three great migrants & why migration matters

The basic story. Twice a year, billions of birds move between the places where they breed and the places where they spend the rest of the year. Migration lets a bird exploit a brief, intense pulse of high-latitude summer productivity to raise young, then retreat to milder latitudes when that pulse collapses into winter. The cost is one of the most demanding

feats in the animal kingdom: sustained, navigated, long-distance flight, twice annually, for life. We follow three species that bracket the extremes of that feat.

Arctic tern (*Sterna paradisaea*). The longest-distance migrant known. Individuals tracked with geolocators average roughly **70,900 km per year** round trip — from high-Arctic breeding grounds to the Antarctic pack ice and back (Egevang et al. 2010, *PNAS* 107:2078). Because it follows the sun from one polar summer to the other, the Arctic tern experiences more daylight per year than any other animal.

Common swift (*Apus apus*). An Afro-Palearctic migrant that breeds across Europe and temperate Asia and winters in sub-Saharan Africa. Remarkably, it is almost entirely **aerial** for the roughly ten months it is not nesting: it feeds, sleeps, and even mates on the wing, landing only to breed. A young swift may not touch a solid surface for its first two or three years of life.

Red knot (*Calidris canutus*). A medium-sized shorebird that undertakes some of the longest-known shorebird migrations, on several distinct flyways worldwide. The Americas (*rufa*) population is famous for its Delaware Bay stopover, timed to the spring spawning of horseshoe crabs — the knots gorge on the eggs to fuel the final leg to the Arctic. That tight coupling makes the species a textbook case of migratory vulnerability.

Why migration matters. Beyond the spectacle, migration is a load-bearing piece of how ecosystems work: it moves energy, nutrients, and predation pressure across continents on a seasonal clock. It is also acutely exposed to global change — habitat loss at a single stopover, or a phenological mismatch between arrival and the food peak, can collapse a population that is otherwise healthy at both ends of its range. Mapping where these birds are, month by month, is the first step to understanding and protecting them.

2. The data: eBird, GBIF, and the effort-bias problem

Where the data come from. **eBird**, run by the Cornell Lab of Ornithology, is the world's largest biodiversity citizen-science project: hundreds of millions of bird observations submitted by hundreds of thousands of volunteer birdwatchers as dated, located checklists. We do not query eBird directly. Instead we access its observations through **GBIF** (the Global Biodiversity Information Facility), which republishes them as the **eBird Observation Dataset (EOD)** under a **CC BY 4.0** licence — which means the data are freely publishable provided we cite the source DOI.

GBIF's Download API lets us mint a reproducible, **citable DOI** for an exact query. We ask for human observations of each species, with coordinates, from the EOD dataset (datasetKey 4fa7b334-ce0d-4e88-aaae-2e0c138d049e), over a fixed decade. The predicate below is the actual query structure used by the project's `wolfram/gbif-download.wls`:

```
(* GBIF Download API predicate: one citable download per species.
EOD = eBird Observation Dataset (CC BY 4.0). *)
EOD = "4fa7b334-ce0d-4e88-aaae-2e0c138d049e";
predicate[key_] := <|
  "type" -> "and", "predicates" -> {
    <|"type" -> "equals", "key" -> "TAXON_KEY", "value" -> ToString[key]|>,
    <|"type" -> "equals", "key" -> "DATASET_KEY", "value" -> EOD|>,
    <|"type" -> "equals", "key" -> "HAS_COORDINATE", "value" -> "true"|>,
    <|"type" -> "equals", "key" -> "BASIS_OF_RECORD", "value" -> "HUMAN_OBSERVATION"|>,
    <|"type" -> "and", "predicates" -> {
      <|"type" -> "greaterThanOrEquals", "key" -> "YEAR", "value" -> "2014"|>,
      <|"type" -> "lessThanOrEquals", "key" -> "YEAR", "value" -> "2024"|>|>|>;
(* POST predicate[5229230] to /occurrence/download/request -> a downloadKey
that resolves to a permanent DOI once the archive is built. *)
```

The raw archive arrives as a tidy table of `lat`, `lon`, `month`, `year` per occurrence. A first look, binned by month, already tells a clear seasonal story — here for the Arctic tern, whose occurrences sweep from the northern breeding grounds down the Atlantic and back over the year:

```
(* Seasonal occurrence map: 12 monthly GeoHistogram panels.
o = loadOccurrences[key]; one 750-km hex bin map per calendar month. *)
o = loadOccurrences[5229230];
Grid@Partition[Table[
  Module[{idx = Flatten@Position[o["month"], m]},
    GeoHistogram[
      GeoPosition /@ Transpose[{o["lat"][[idx]], o["lon"][[idx]]}],
      Quantity[750, "Kilometers"], GeoRange -> "World",
      GeoProjection -> "Robinson", ColorFunction -> "TemperatureMap",
      GeoBackground -> GrayLevel[0.15], PlotLabel -> monthNames[[m]],
      ImageSize -> 230]],
  {m, 12}], 4]
```




Figure 2.1 Monthly occurrence density of the Arctic tern from the GBIF / eBird Observation Dataset (2014–2024), one panel per calendar month, 750-km hex bins, Robinson projection. The northward breeding-season concentration (May–July) and the long southbound Atlantic passage are both visible. But read the next caption before trusting any apparent geography.

The honest caveat: effort bias. Occurrence records do not report where the birds are — they report where the **observers** are. eBird coverage is overwhelmingly concentrated in North America and Western Europe, where most contributors live, and is sparse across the open ocean, the tropics, and the Southern Hemisphere. A naive density map therefore conflates

“many birds here” with “many birders here.” Any honest analysis of citizen-science occurrence data must correct for this sampling bias before drawing geographic conclusions — which is exactly what the centroid method in §3 is built to do.

3. From checklists to centroids (method)

The idea. To summarise “where the population is” in a given month with a single point, we compute the **monthly population centroid** — a weighted average location of that month's occurrences. Done naively this inherits all the effort bias of §2 and is geometrically wrong for a round planet. Two corrections fix both problems.

3.1 Correction one: presence weighting on a coarse grid

A raw average over occurrence *points* is dominated by a handful of intensely-birded hotspots — a single famous reserve can contribute tens of thousands of records and drag the centroid toward itself. To blunt this, we lay down a coarse **2° grid** and weight each *occupied cell* equally (presence = 1 per cell), rather than weighting by the number of records in the cell. A cell with 40,000 checklists and a cell with one both count once. This converts “where were people looking” into “where was the species present,” which is far closer to the quantity we want.

A subtlety for the specialist. A 2°×2° cell is **not** equal-area: its true area shrinks toward the poles in proportion to $\cos(\text{lat})$. For high-latitude species this matters, and the spherical mean below handles it correctly by construction — each presence point is converted to a unit vector whose contribution is naturally weighted by the geometry of the sphere, not by a flat lon×lat rectangle.

3.2 Correction two: a proper spherical centroid

The deeper problem is that **you cannot average longitudes arithmetically**. Longitude wraps at $\pm 180^\circ$: the mean of $+170^\circ$ and -170° is not 0° (the Atlantic) — it should be 180° (the dateline). For a circumpolar bird like the Arctic tern, whose occurrences ring the entire planet, a naive longitude average is simply meaningless. The correct construction embeds each point on the unit sphere, averages the 3-D vectors, and projects back:

```
(* Weighted spherical mean of (lat,lon) points. Returns {lat,lon}
   in degrees, or Missing["DegenerateCentroid"] when the resultant
   vector is ~0 (antipodally smeared points cancel) -- we never fabricate
   an arithmetic-mean longitude. (This is the helper in the package. *)
sphericalCentroid[lats_, lons_, w_] := Module[{v, m},
  v = MapThread[Function[{la, lo, wt},
    wt {Cos[la Degree] Cos[lo Degree],
        Cos[la Degree] Sin[lo Degree],
        Sin[la Degree]}], {lats, lons, w}];
  m = Total[v];
  If[Norm[m] < 10^-9, Return[Missing["DegenerateCentroid"]]];
  m = m/Norm[m];
  {ArcSin[m[[3]]/Degree, ArcTan[m[[1]], m[[2]]/Degree]}];
```

The `Missing["DegenerateCentroid"]` branch is not defensive boilerplate — it is the honest answer when a population is spread so evenly around the globe (or split into antipo-

dal clumps) that *no* single point represents it. Returning a fabricated longitude there would be worse than returning nothing. As we will see in §4, this is precisely the situation a global centroid walks into for a multi-flyway species, and the reason we must split by flyway first.

4. The multi-flyway trap (the key lesson)

The trap. A spherical centroid is only meaningful if the points it averages belong to a single, *coherent* population. Many species do not: they split into geographically disjoint **flyways** that never mix. Averaging across flyways produces a “centre” that lies where *no birds are* — the midpoint of two crowds is the empty space between them.

The red knot is the textbook case. Pool its Americas, European/African, and Asian-Australasian flyways into one global centroid and the result drifts out into the open **mid-Atlantic**, on average **~3,800 km per month** away from where the Americas (*rufa*) population — the one a North American reader actually sees — truly is. The number is not a rounding artefact; it is the signature of averaging across populations that should never have been combined.

```
(* The fix: restrict to one coherent flyway BEFORE taking the
   centroid. Each box is {lon -> {min,max}, lat -> {min,max}}. *)
swift = monthlyCentroids[5228676, <|"lon" -> {-30, 145}, "lat" -> {-40, 75}|>]; (* Afro-Pale
knot  = monthlyCentroids[2481765, <|"lon" -> {-110, -30}, "lat" -> {-56, 85}|>]; (* Americas
tern  = monthlyCentroids[5229230, <|"lon" -> {-80, 20}, "lat" -> {-90, 90}|>]; (* Atlantic

(* The BROKEN contrast: no flyway filter -> meaningless mid-ocean centroid *)
knotGlobal = monthlyCentroids[2481765, All];
```

With each species reduced to a single coherent flyway, the monthly centroids trace clean, interpretable migration loops. The common swift's Afro-Palearctic flyway needs no special handling — it is already one population, and its centroid simply works. The red knot is shown on its Americas flyway. The Arctic tern is shown as an Atlantic-basin centroid (and, separately, as a latitude-versus-month curve, since its defining signal is the pole-to-pole north-south sweep):



Figure 4.1 Arctic tern — monthly population centroid on the Atlantic-basin flyway, orthographic globe over relief. Numbers mark the calendar month; the track sweeps from southern winter quarters up to high-Arctic breeding latitudes and back.



Figure 4.2 Common swift — monthly population centroid on the single Afro-Palearctic flyway. Breeding-season concentration over Europe (months ~5–8) and the sub-Saharan wintering shift (months ~11–2) are both clear; no flyway splitting is needed because there is only one population.



Figure 4.3 Red knot — monthly population centroid restricted to the Americas flyway. Compare this coherent loop with the broken global centroid (the *knotGlobal* contrast above), which would sit uselessly in the mid-Atlantic, ~3,800 km/month from this track.

5. Animating the population centroid

Stringing the twelve monthly centroids together and interpolating along great circles between them turns the static loops of §4 into the animated comet-trail globe shown at the top of this post. Each species' centre of mass glides through the year over its faint annual route — the most compact possible summary of “where is this population, right now, on average.”

We can also quantify each loop. Summing the great-circle distances between consecutive monthly centroids (over observed months only) gives the **annual centroid displacement** — how far the centre of mass travels in a year:

```
(* Annual centroid displacement per primary flyway track. *)
pl = Import[FileNameJoin[{$BFPDataDir, "path_lengths.csv"}], "CSV"];
prim = Select[Rest[pl], ! StringContainsQ[#[[3]], "global", IgnoreCase -> True] &];
BarChart[prim[[All, 4]],
  ChartLabels -> (Style[#[[2]] <> "\n(" <> #[[3]] <> ")", 10] & /@ prim),
  BarOrigin -> Left,
  PlotLabel -> Style[
    "Annual centroid displacement (km) -- NOT individual journey distance", 13],
  ImageSize -> 780, AspectRatio -> 0.5]
```




Figure 5.1 Annual displacement of the monthly population centroid along each species' primary flyway: common swift (Afro-Palearctic) $\approx 14,000$ km, red knot (Americas) $\approx 7,900$ km, Arctic tern (Atlantic basin) $\approx 24,100$ km. The deliberately broken global red-knot track is excluded.

Crucial caveat — read before quoting these numbers. This is the displacement of the population *centre of mass*, NOT how far any individual bird flies. The two differ enormously, and in both directions. For the Arctic tern the centroid figure **understates** the journey: because the species is nearly circumpolar, east–west motion in different longitudes cancels in the mean, so the centroid mostly captures the north–south sweep. Tracked individual Arctic terns average about **70,900 km per year** (Egevang et al. 2010) — roughly three times the $\sim 24,100$ km centroid displacement. The centroid is a population statistic; the geolocator figure is an individual's odometer. Do not conflate them.

6. Cross-check: eBird Status & Trends

Are our occurrence-based centroids actually right? A method built from biased citizen-science occurrences deserves an independent check. Cornell's **eBird Status & Trends** (S&T) is the professional answer to the same question: it fits machine-learning models that

explicitly correct for observer effort and detectability, and outputs **bias-corrected weekly relative abundance** on a fine grid — a far more sophisticated product than our coarse occurrence centroid (Fink et al., eBird Status and Trends).

As a **private validation**, we computed abundance-weighted centroids directly from the S&T rasters — inside the *same* flyway boxes used in §4 — by driving the R `ebirdst` package from Wolfram Language via `ExternalEvaluate`, then measuring the great-circle separation between Cornell's modelled centroids and ours, month by month. The pattern of code (our code, not Cornell's data) is:

```
(* PRIVATE S&T validation: abundance-weighted centroids from the
ebirdst weekly rasters, compared to our GBIF centroids in the same
flyway box. The rasters and all derived S&T artifacts stay LOCAL and
git-ignored, per Cornell's terms -- only this code is shared. *)
r = StartExternalSession["R"];
ExternalEvaluate[r, "library(ebirdst); library(terra)"];
ExternalEvaluate[r, "
  cube <- load_raster(species, product='abundance', period='weekly')
  ll <- project(cube, '+proj=longlat')
"];
(* For each of 52 weekly layers: abundance-weighted lon/lat centroid,
cropped to the flyway box, then resampled WL-side to 12 monthly means
and compared to the GBIF track with GeoDistance. *)
```

The result. Our occurrence-based centroids agree with Cornell's professionally modelled abundance centroids to within roughly **1,300–1,750 km** on average, along flyways thousands of kilometres long: common swift ~1283 km, red knot ~1392 km, Arctic tern ~1742 km mean monthly separation. For a coarse, presence-weighted method built from raw occurrence data, agreeing with a full effort-corrected model to within a small fraction of the flyway length is a strong endorsement that the centroid approach captures the real seasonal signal.

Why you see no S&T figure here. The eBird Status & Trends Data Products are released under Cornell's own terms, which permit **non-commercial use and validation** but do **not** permit publishing derived figures or maps. We therefore show **none** — the cross-check above is reported as numbers only — and we point readers to Cornell's official, professionally produced visualizations instead: **eBird Status and Trends** (<https://science.ebird.org/en/status-and-trends>).

This material uses data from the eBird Status and Trends Project at the Cornell Lab of Ornithology, eBird.org. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the Cornell Lab of Ornithology.

6b. Live now: recent sightings via the eBird API

The centroids summarise a decade of history. For a sense of where these birds are *this month*, we add a live layer pulled from the **eBird API 2.0** recent-observations endpoint (`/v2/data/obs/{region}/recent/{speciesCode}`), which returns the last 30 days of checklisted sightings for a species in a region. We sweep a representative spread of countries and plot the result:

```
(* Live recent sightings (last 30 days). LoadLiveSightings sweeps a
spread of regions, returns {lat,lon} pairs ROUNDED TO 0.1 deg for
privacy. Supply your own eBird API 2.0 token. *)
pts = LoadLiveSightings["arcter", "<your-ebird-api-token>"]; (* arcter|comswi|redkno *)
GeoListPlot[GeoPosition /@ pts, GeoRange -> "World",
  GeoProjection -> "Robinson", GeoBackground -> GrayLevel[0.15],
  PlotStyle -> Directive[Orange, Opacity[0.7], PointSize[0.006]],
  PlotLabel -> "Arctic tern -- recent eBird sightings (last 30 d)\n\
Source: eBird.org (Cornell Lab of Ornithology)", ImageSize -> 900]
```



Figure 6b.1 Arctic tern — recent eBird sightings (last 30 days), swept across a spread of countries. **Source: eBird.org (Cornell Lab of Ornithology).**




Figure 6b.2 Common swift — recent eBird sightings (last 30 days). **Source: eBird.org (Cornell Lab of Ornithology).**




Figure 6b.3 Red knot — recent eBird sightings (last 30 days). Source: eBird.org (Cornell Lab of Ornithology).

Privacy and attribution. Live coordinates are coarsened to one decimal place (~11 km) before plotting or storage, so no precise observer or sensitive-species location is republished. All live data are attributed to eBird.org (Cornell Lab of Ornithology) per the eBird Terms of Use.

7. What each journey reveals

Read together, the three tracks tell three different stories about how a bird can solve the problem of seasonal survival.

Arctic tern — the pole-to-pole Atlantic sweep. The latitude-versus-month signal is the cleanest in the dataset: a centroid that climbs to high-Arctic breeding latitudes in the boreal summer and plunges toward the Southern Ocean in the austral summer. The centroid displacement (~24,100 km) is large yet still understates the individual journey (~70,900 km), because circumpolar east-west motion cancels in the population mean — a vivid illustration of the difference between a population statistic and an individual track.

Common swift — the Europe↔sub-Saharan shuttle. A single coherent flyway, and therefore the easiest centroid to trust: a tight summer concentration over Europe collapses into a sub-Saharan African wintering centre and back, ~14,000 km of annual centroid travel. The swift shows what the method looks like when nothing has to be corrected for multiple populations — it just works.

Red knot — the Americas long-haul (and the cautionary tale). On its Americas flyway the knot traces a coherent ~7,900 km loop between Arctic breeding grounds and southern wintering areas, with the Delaware Bay stopover embedded in the spring leg. It is also the

species that exposed the multi-flyway trap: pooled globally, its centroid is nonsense. The knot is the reason this post insists on per-flyway analysis.

8. Conclusions

With open data and a careful method, you can reconstruct continental-scale bird migration from a laptop. Hundreds of millions of volunteer observations, republished under an open licence with a citable DOI, plus a few dozen lines of Wolfram Language, are enough to recover the annual journeys of three of the planet's great migrants and to cross-validate them against a professional abundance model.

But the central lesson is a cautionary one. **Naive pooling misleads.** A single global centroid placed the red knot in the empty mid-Atlantic, ~3,800 km/month from any real bird. Two disciplines rescue the analysis: **bias-awareness** (presence weighting on a coarse grid, so birders are not mistaken for birds) and **per-flyway analysis** (so disjoint populations are never averaged together), all on a **proper spherical geometry** (so longitudes are never averaged arithmetically). Get those three right and citizen-science occurrence data yield a faithful, validated picture of where the birds actually are.

9. References

Egevang, C., Stenhouse, I. J., Phillips, R. A., Petersen, A., Fox, J. W., & Silk, J. R. D. (2010). Tracking of Arctic terns *Sterna paradisaea* reveals longest animal migration. *Proceedings of the National Academy of Sciences* 107(5), 2078–2081. [Geolocator tracking; ~70,900 km/yr round trip.]

Fink, D., Auer, T., Johnston, A., Strimas-Mackey, M., Ligocki, S., Robinson, O., Hochachka, W., et al. eBird Status and Trends. Cornell Lab of Ornithology, Ithaca, New York. science.ebird.org/en/status-and-trends (<https://science.ebird.org/en/status-and-trends>). [Effort-corrected weekly relative-abundance models used for the \$6 validation.]

GBIF.org. eBird Observation Dataset (EOD), datasetKey 4fa7b334-ce0d-4e88-aaae-2e0c138d049e, accessed via the GBIF Occurrence Download API (CC BY 4.0). Per-species download DOIs: Arctic tern — "doi" -> "<https://doi.org/10.15468/dl.87f228>" n218269FontFamilyCourier; common swift —



.

Text



Sullivan, B. L., Wood, C. L., Iliff, M. J., Bonney, R. E., Fink, D., & Kelling, S. (2009). eBird: A citizen-based bird observation network in the biological sciences. *Biological Conservation* 142(10), 2282–2292. [The eBird platform.]

Newton, I. (2008). *The Migration Ecology of Birds*. Academic Press. [Comprehensive reference on flyways, energetics, and migratory strategy.]

Baker, A. J., González, P. M., Piersma, T., Niles, L. J., et al. (2004). Rapid population decline in red knots: fitness consequences of decreased refuelling rates and late arrival in Delaware Bay. *Proceedings of the Royal Society B* 271, 875–882. [Red knot / horseshoe-crab stopover coupling.]

10. Acknowledgements & data licences

With thanks. This work rests entirely on the labour of others: the global community of **eBird volunteers** who contributed the underlying observations, the **Cornell Lab of Ornithology** which runs eBird and the Status & Trends project, and **GBIF** which curates and openly republishes the data with persistent, citable DOIs.

- **GBIF / eBird Observation Dataset** is licensed **CC BY 4.0** — freely publishable with attribution; we cite the per-species download DOIs in §9 (or mark them “DOI pending (rerun after the GBIF download completes)” when the archive build is still in progress).

- **eBird Status & Trends** is released under Cornell's custom, non-commercial terms. We use it for *private validation only* and publish **no** derived S&T figures here; see Cornell's official visualizations instead. As required:

This material uses data from the eBird Status and Trends Project at the Cornell Lab of Ornithology, eBird.org. Any opinions, findings, and conclusions or recommendations expressed in this material are those of the author(s) and do not necessarily reflect the views of the Cornell Lab of Ornithology.

- **eBird API 2.0** live data are used under the eBird Terms of Use, attributed to eBird.org (Cornell Lab of Ornithology), with observer coordinates coarsened to ~11 km for privacy.

Tools. All numerics, maps, and animations are pure **Wolfram Language**. The notebook was assembled programmatically via `wolframscript`. No AI-generated illustrations are used anywhere in this post; every figure is computed from data.

11. Reproducibility