

Reasoning Behavior Analysis Report

EXPERIMENTAL SETUP

=====

Data Folder: data/AIME2025__R10/deepseek/p2
Total Items: 300
Total Strategies: 9

OVERALL ACCURACY


=====

Total Problems: 300
Correct: 258
Incorrect: 42
Overall Accuracy: 86.00%

THINKING STRATEGY DISTRIBUTION

- =====
- A: Planning and Execution
Distribution: 82.7%
Accuracy: 87.50% (217/248)
 - B: Problem Decomposition
Distribution: 98.0%
Accuracy: 87.41% (257/294)
 - C: Formulate Hypotheses and Test Them
Distribution: 63.3%
Accuracy: 81.58% (155/190)
 - D: Guess the Answer First, Then Verify
Distribution: 5.7%
Accuracy: 23.53% (4/17)
 - E: Option Judgment and Elimination
Distribution: 31.7%
Accuracy: 84.21% (80/95)
 - F: Reverse Thinking
Distribution: 8.7%
Accuracy: 73.08% (19/26)
 - G: Divergent and Convergent Thinking
Distribution: 44.0%
Accuracy: 74.24% (98/132)
 - H: Counterexample Testing
Distribution: 21.0%
Accuracy: 73.02% (46/63)
 - I: Association Method
Distribution: 14.7%
Accuracy: 63.64% (28/44)

Thinking Strategy Distribution and Accuracy



Example for Strategy A: Planning and Execution

PROBLEM

N/A

GROUND TRUTH ANSWER

81

MODEL RESPONSE

Step 1. Understand the Problem

We want to fill a (3×9) grid with the numbers 1 through 9 such that:

- Each **row** is a permutation of $\{1, 2, \dots, 9\}$.
- Each of the three (3×3) **blocks** (columns 1–3, 4–6, 7–9) contains all numbers 1 through 9 exactly once.

This is equivalent to filling the first three rows of a Sudoku puzzle.

Step 2. A Step-by-Step Counting Plan

We break the grid into three blocks:

- **Block A**: columns 1–3
- **Block B**: columns 4–6
- **Block C**: columns 7–9

We proceed as follows:

1. Fill **Block A** (a (3×3) array) with numbers 1–9, each exactly once.
2. Fill **Block B** such that:
 - It is a permutation of 1–9.
 - For each row (i) , the numbers in Block B do **not** include those already in Block A.

CHAIN OF THOUGHT (First 1500 chars)

We are given: "The 27 cells of a 3×9 grid are filled in using the numbers 1 through 9 so that each row contains 9 different numbers, and each of the three 3×3 blocks heavily outlined in the example below contains 9 different numbers, as in the first three rows of a Sudoku puzzle.

Example grid:

Row1: 4,2,8,9,6,3,1,7,5

Row2: 3,7,9,5,2,1,6,8,4

Row3: 5,6,1,8,4,7,9,2,3

The number of different ways to fill such a grid can be written as $p^a \cdot q^b \cdot r^c \cdot s^d$, where p, q, r, s and a, b, c, d are distinct prime numbers and a, b, c, d are positive integers. Find $p \cdot a + q \cdot b + r \cdot c + s \cdot d$."

Interpretation: We have a grid of size 3 rows x 9 columns. Conditions:

- Each row is a permutation of numbers 1, ..., 9.
- The grid is partitioned into three blocks: columns 1-3, columns 4-6, columns 7-9. In each block (3x3 square), all numbers 1, ..., 9 appear exactly once.

We want: number of ways to fill such grid. But careful: The grid is not necessarily a Sudoku puzzle's first three rows? Actually it says: "as in the first three rows of a Sudoku puzzle." That means that if we take a Sudoku puzzle's first three rows, they satisfy: each row is a permutation, and each of the three 3x3 blocks (columns 1-3, 4-6, 7-9) are permutations. But there is no condition from columns? Actually in a Sudoku, columns also have all numbers, but here we are only filling the first three rows. So it's exactly: We fill a 3x9 grid such that each row is a permutat...

Example for Strategy B: Problem Decomposition

PROBLEM

=====

N/A

GROUND TRUTH ANSWER

=====

113

MODEL RESPONSE

=====

The problem involves finding the number of ways to draw 12 segments of equal length on a regular 24-gon such that each vertex is an endpoint of exactly one segment. This requires pairing the vertices into 12 pairs, where each pair is connected by a chord of the same length. The chord length corresponds to the step size k between vertices, where $(1 \leq k \leq 12)$ since for $(k > 12)$, the chord length is equivalent to $(24-k)$.

A perfect matching using only chords of length k is possible only if $(\gcd(k, 24))$ is such that $(24 / \gcd(k, 24))$ is even. This excludes $(k = 8)$ because $(\gcd(8, 24) = 8)$ and $(24/8 = 3)$ is odd, making a perfect impossible.

For each valid k , the number of perfect matchings is calculated based on $(\gcd(k, 24))$...

CHAIN OF THOUGHT (First 1500 chars)

=====

First, I need to understand the problem. There's a regular 24-gon, so it has 24 vertices. I have to draw 12 segments, each of equal length, and each vertex must be an endpoint of exactly one segment. That means I'm pairing up the vertices into 12 pairs, and each pair is connected by a segment of equal length. Since the polygon is regular, the segments must be chords of equal length, but since all segments are equal, I need to find how many ways to pair the vertices such that the distance between each pair is the same.

The segments are of equal lengths, so in a regular polygon, chords of equal length correspond to vertices that are a certain number of steps apart. For example, in a regular n -gon, chords between vertices that are k steps apart have the same length for fixed k . But here, $n=24$, and I need to pair all vertices with segments of the same length, so all pairs must have the same chord length, meaning that for each pair, the vertices are separated by the same number of steps around the polygon.

Let me denote the vertices as points on a circle, equally spaced. The distance between two vertices depends on the chord length, which is determined by the central angle. The central angle between adjacent vertices is $360^\circ/24 = 15^\circ$. So, if two vertices are k steps apart, the central angle is $k \cdot 15^\circ$, and the chord length is $2 \cdot \sin((k \cdot 15^\circ)/2)$ times the radius, but since we care about equal length, we need the same k for all pairs.

But k can be from 1 to 12, since for $k > 12$, it wou...

Example for Strategy C: Formulate Hypotheses and Test Them

PROBLEM

N/A

GROUND TRUTH ANSWER

240

MODEL RESPONSE

The function $f(x) = \frac{(x - 18)(x - 72)(x - 98)(x - k)}{x}$ achieves its minimum value at exactly two positive real numbers x for three values of k . These values are found by ensuring that the two local minima of $f(x)$ have the same value, which leads to the equation $(s^2 - 376s + 25228)^2 = 1016064(s - 94)$, where $(s = a + b)$ and (a) and (b) are the (x) -coordinates of the local minima.

Solving this equation yields $(s = 98)$, $(s = 110)$, and $(s = 194)$. Using $(k = 2s - 188)$, the corresponding values of (k) are (8) , (32) , and (200) . The sum of these values is $(8 + 32 + 200 = 240)$.

Thus, the sum of the three values of (k) is 240.

$\boxed{240}$

CHAIN OF THOUGHT (First 1500 chars)

I need to find the sum of three positive real numbers k such that the function $f(x) = [(x - 18)(x - 72)(x - 98)(x - k)] / x$ has its minimum at exactly two positive real numbers x . Since x is positive, we can work with this function.

First, I should simplify $f(x)$. Since x is positive, we can write $f(x)$ as:

$$f(x) = (x - 18)(x - 72)(x - 98)(x - k) / x$$

But that might be messy. I could expand the numerator and then divide by x , but that seems complicated. Notice that $f(x)$ can be written as:

$$f(x) = [(x - 18)(x - 72)(x - 98)(x - k)] / x$$

I recall that for functions like this, the minimum might occur where the derivative is zero, and we need exactly two points where the minimum is achieved, meaning that there are two points with the same minimum value, and that value is the global minimum.

Since it's a minimum over positive reals, and the function might go to infinity as x approaches 0 or infinity, so there should be a minimum somewhere.

I should consider the derivative of $f(x)$. Let me set $g(x) = \ln f(x)$ or something, but since we have multiple factors, perhaps it's better to handle $f(x)$ directly.

Let me write $f(x)$ as:

$$f(x) = (x - 18)(x - 72)(x - 98)(x - k)/x$$

But to find the critical points, I need $f'(x) = 0$.

Computing $f'(x)$ might be messy because of the product. I could write $f(x)$ as a product and use logarithmic differentiation, but since we need the minimum, we need where $f'(x) = 0$.

$$\text{Set } u(x) = (x - 18)(x - 72)(x - 98)(x - k)$$

$$\text{Then } f(x) = u(x)/x$$

$$\text{So } f'(x) = [u'(x)x - \dots]$$

Example for Strategy D: Guess the Answer First, Then Verify

PROBLEM

=====

N/A

GROUND TRUTH ANSWER

=====

610

MODEL RESPONSE

=====

The greedy algorithm for making change with 1-cent, 10-cent, and 25-cent coins succeeds when it uses the minimum number of coins for a given value N . The algorithm fails only when $(N \bmod 25)$ is in the set $\{5, 6, 7, 8, 9, 15, 16, 17, 18, 19\}$, but this condition applies only for $(N \geq 25)$. For $(N < 25)$, the greedy algorithm always succeeds because only 1-cent and 10-cent coins are used, and the algorithm is optimal for these coins.

For N from 1 to 24, all values succeed, giving 24 values.

For N from 25 to 1000, there are 976 values. The number of values where the greedy algorithm fails is determined by the remainders modulo 25. The remainders that cause failure are 5, 6, 7, 8, 9, 15, 16, 17, 18, and 19. Each of these remainders (from 1 to 24) ap...

CHAIN OF THOUGHT (First 1500 chars)

=====

I need to find how many values of N from 1 to 1000 have the property that the greedy algorithm for making change with 1, 10, and 25 cent coins uses the minimum number of coins. The greedy algorithm always takes the largest coin possible without exceeding N , and we need to ensure that no other combination uses fewer coins.

First, I should figure out when the greedy algorithm fails. From the example, for $N=42$, it fails because we can use fewer coins: 4 dimes and 2 pennies instead of 1 quarter, 1 dime, and 7 pennies. The greedy used 9 coins, but minimal is 6 coins.

The minimal number of coins for any N can be found using dynamic programming, but since N is only up to 1000, I could compute the minimal coins for each N and compare it to the greedy approach. But that might be tedious, and there might be a smarter way.

I recall that for coin systems, the greedy algorithm is optimal only if the coin system is canonical, which means that the greedy algorithm always gives the minimal number of coins. But with 1, 10, 25, it is not canonical because of examples like 30 cents: greedy would be 25 and then 5 pennies, so 6 coins, but 3 dimes would be 3 coins, so better. So for $N=30$, greedy fails.

But we need to find for which N it succeeds.

Since the coins are 1, 10, 25, the minimal coins can be found by considering how many quarters, dimes, and pennies to use. Let Q be the number of quarters, D dimes, P pennies, so $25Q + 10D + P = N$, and we want to minimize total coins $C = Q + D + P$.

...

Example for Strategy E: Option Judgment and Elimination

PROBLEM

N/A

GROUND TRUTH ANSWER

70

MODEL RESPONSE

The problem requires finding all integer bases $(b > 9)$ such that (17_b) divides (97_b) .

Convert (17_b) and (97_b) to base 10:

- $(17_b = 1 \cdot b + 7 = b + 7)$
- $(97_b = 9 \cdot b + 7 = 9b + 7)$

Thus, $(b + 7)$ must divide $(9b + 7)$. Set $(d = b + 7)$, so $(b = d - 7)$. Substitute:

$$9b + 7 = 9(d - 7) + 7 = 9d - 63 + 7 = 9d - 56$$

Since (d) divides $(9d - 56)$, and (d) divides $(9d)$, it follows that (d) divides (56) . The divisors of 56 are 1, 2, 4, 7, 8, 14, 28, 56.

Given $(b > 9)$, then $(d = b + 7 > 16)$. Thus, possible values for (d) are 28 and 56.

- If $(d = 28)$, then $(b = 28 - 7 = 21)$
- If $(d = 56)$, then $(b = 56 - 7 = 49)$

Verify both bases:

- For $(b = 21)$: $(17_{21} = 21 + 7 = 28)$, $(97_{21} = 21 \cdot 9 + 7 = 196)$

CHAIN OF THOUGHT (First 1500 chars)

First, I need to find all integer bases $b > 9$ such that 17_b divides 97_b . Since $b > 9$, both numbers are written in base b , so I need to convert them to base 10 to work with them.

17_b in base 10 is $1 \cdot b + 7 \cdot 1 = b + 7$.

Similarly, 97_b in base 10 is $9 \cdot b + 7 \cdot 1 = 9b + 7$.

So, I need $b + 7$ to divide $9b + 7$. That is, $(b + 7) \mid (9b + 7)$.

In terms of division, I can write this as:

$$9b + 7 \equiv 0 \pmod{b + 7}$$

I need to find values of $b > 9$ where this holds.

I can manipulate the expression. Let me set $d = b + 7$, so $b = d - 7$.

Then $9b + 7 = 9(d - 7) + 7 = 9d - 63 + 7 = 9d - 56$.

So, I need d to divide $9d - 56$, which means d divides 56.

Because if d divides $9d$, and d divides $9d - 56$, then d must divide 56.

So, $d \mid 56$.

Since $d = b + 7$ and $b > 9$, so $b > 9$ implies $d > 9 + 7 = 16$.

So $d > 16$.

Now, d divides 56. The divisors of 56 are 1, 2, 4, 7, 8, 14, 28, 56.

Since $d > 16$, possible d are 28 and 56.

So, $d = 28$ or $d = 56$.

Then, since $d = b + 7$, so $b = d - 7$.

If $d = 28$, $b = 28 - 7 = 21$.

If $d = 56$, $b = 56 - 7 = 49$.

Now, b must be greater than 9, which both 21 and 49 are, so they should be valid.

I need to confirm that 17_b indeed divides 97_b for these bases.

First, $b=21$.

$$17_{21} = 1 \cdot 21 + 7 = 28$$

$$97_{21} = 9 \cdot 21 + 7 = 189 + 7 = 196$$

Now, 28 divides 196? $196 / 28 = 7$, yes, since $28 \cdot 7 = 196$.

Second, $b=49$.

$$17_{49} = 1 \cdot 49 + 7 = 56$$

$$97_{49} = 9 \cdot 49 + 7 = 441 + 7 = 448$$

Now, 56 divides 448? $448 / 56 = 8$, yes, since $56 \cdot 8 = 448$.

So both work.

The problem asks for the sum of all integer bas...

Example for Strategy F: Reverse Thinking

PROBLEM

=====

N/A

GROUND TRUTH ANSWER

=====

240

MODEL RESPONSE

=====

The function $f(x) = \frac{(x - 18)(x - 72)(x - 98)(x - k)}{x}$ achieves its minimum value at exactly two positive real numbers x for exactly three positive values of k . These values are $k = 8$, $k = 32$, and $k = 200$.

The sum of these values is $8 + 32 + 200 = 240$.

This result is derived from the condition that the line $y = Cx$ is tangent to the polynomial $u(x) = (x-18)(x-72)(x-98)(x-k)$ at two points, ensuring that $f(x) \geq C$ for all $x > 0$ with equality only at those two points. The values of k satisfy the equation obtained from comparing coefficients, leading to the solutions $k = 8$, $k = 32$, and $k = 200$.

$\boxed{240}$

CHAIN OF THOUGHT (First 1500 chars)

=====

I need to find the sum of three positive real numbers k such that the function $f(x) = [(x - 18)(x - 72)(x - 98)(x - k)] / x$ has its minimum at exactly two positive x values. First, since x is positive, I can work with this function.

I recall that for a function to have a minimum at a point, the derivative should be zero, and the second derivative should be positive, or we need to check the behavior around that point. But here, it says the minimum value is achieved at exactly two points, which might mean that there are two points where the function attains the same minimum value, and that value is the global minimum or something.

I should find the derivative of $f(x)$ and set it to zero to find critical points. But since k is a parameter, I need to find k such that there are exactly two critical points where the function has a minimum, and perhaps these two points are both minima with the same value.

The function is a rational function, and since $x > 0$, I can consider $f(x)$ as a function defined for $x > 0$.

Let me write $f(x)$ as:

$$f(x) = [(x - 18)(x - 72)(x - 98)(x - k)] / x$$

I can expand the numerator, but that might be messy. Since we have factors, perhaps I can think in terms of logarithms or something to simplify the derivative.

I remember that for functions with products, taking the logarithm can help with differentiation. But here, since we have division by x , I can write:

$$f(x) = (x - 18)(x - 72)(x - 98)(x - k) / x$$

Let me denote $g(x) = \ln f(x) = \ln[(x - 18)(x - 72)(x - 98)(x - k)] - \ln x$.

Example for Strategy G: Divergent and Convergent Thinking

PROBLEM

N/A

GROUND TRUTH ANSWER

510

MODEL RESPONSE

Step 1. Rewrite the Inequalities

We consider points $((x, y, z))$ in the plane:

$$\begin{aligned} & \\ x + y + z &= 75 \\ & \end{aligned}$$

that satisfy:

$$\begin{aligned} & \\ x - yz &< y - zx < z - xy \\ & \end{aligned}$$

This implies two inequalities:

1. $(x - yz < y - zx)$
2. $(y - zx < z - xy)$

Rewriting:

- From (1):
$$\begin{aligned} & \\ x - yz - y + zx &< 0 \implies (1 + z)(x - y) < 0 \\ & \end{aligned}$$
- From (2):
$$\begin{aligned} & \\ y - zx - z + xy &< 0 \implies (1 + x)(y - z) < 0 \\ & \end{aligned}$$

From the chain $(x - yz < y - zx < z - xy)$, it follows that:

$$\begin{aligned} & \\ x - yz &< z - xy \implies (1 + y)(x - z) < 0 \\ & \end{aligned}$$

Thus, the three inequalities become:

- (A) $(1 + z)(x - y) < 0$
- (B) $(1 + x)(y - z) < 0$
- (C) $(1 + y)(x - z) < 0$

Step 2. Determine the Sign Patterns

Each inequality implies that the factors have opposite signs. There ar...

CHAIN OF THOUGHT (First 1500 chars)

We are given: "The set of points in 3-dimensional coordinate space that lie in the plane $x+y+z=75$ whose coordinates satisfy the inequalities $x-yz < y-zx < z-xy$ forms three disjoint convex regions. Exactly one of those regions has finite area. The area of this finite region can be expressed in the form $a\sqrt{b}$, where a and b are positive integers and b is not divisible by the square of any prime. Find $a+b$."

Interpretation: Points (x,y,z) in $\mathbb{R}^3$, but they lie on the plane: $x+y+z = 75$. And they satisfy:

- (1) $x - yz < y - zx$
- (2) $y - zx < z - xy$

Or equivalently:

$$x - yz < y - zx \implies x - yz - (y - zx) < 0 \implies x - yz - y + zx < 0 \implies x(1+z) - y(1+z) < 0? \text{ Actually careful:}$$

$$x - yz < y - zx$$

$$\implies x - yz - y + zx < 0$$

$$\implies x(1+z) - y(1+z) < 0? \text{ Actually careful: } x - yz + zx - y = x(1+z) - y(1+z) = (1+z)(x-y) < 0.$$

$$\text{So inequality (1) becomes: } (1+z)(x-y) < 0.$$

$$\text{Similarly, (2): } y - zx < z - xy$$

$$\implies y - zx - z + xy < 0$$

$$\implies y(1+x) - z(1+x) = (1+x)(y-z) < 0.$$

And also there is an implied third inequality? Actually the condition is: $x-yz < y-zx < z-xy$. That gives two inequalities: (1) and (2). But there is also implicitly that the ordering is strict. But we don't have an inequality like: $x-yz < z-xy$? That would be automatically if the chain holds? Actually careful: The condition is: $x-yz < y-zx$ and $y-zx < z-xy$. That does not necessarily imply $x-yz < z-xy$? Actually if $a < b$ and $b < c$ then $a < c$. So it implies $(1+z)(x-y) < 0$ and $(1+x)(y-z) < 0$, and then also $(1+y)(z-x) < 0$? Becaus...

Example for Strategy H: Counterexample Testing

PROBLEM

N/A

GROUND TRUTH ANSWER

16

MODEL RESPONSE

The valid distributions of flavors that satisfy the conditions are (5,3,1), (6,2,1), and (4,3,2), where the numbers represent the counts of chocolate, vanilla, and strawberry, respectively.

For each distribution, the number of ways to assign flavors to the distinct players is calculated using multinomial coefficients:

- For (5,3,1): $\frac{9!}{5!3!1!} = 504$
- For (6,2,1): $\frac{9!}{6!2!1!} = 252$
- For (4,3,2): $\frac{9!}{4!3!2!} = 1260$

The total number of assignments (N) is the sum:

$$504 + 252 + 1260 = 2016$$

The remainder when (N) is divided by 1000 is:

$$2016 \bmod 1000 = 16$$

Thus, the remainder is 16.

$\boxed{16}$

CHAIN OF THOUGHT (First 1500 chars)

I need to find the number of ways to assign flavors to the nine players such that each flavor is used at least once, and the number of chocolate is greater than vanilla, which is greater than strawberry. Then I need to find that number modulo 1000.

First, since there are three flavors and nine players, and each flavor has at least one, I need to consider the possible distributions of the flavors. Let C be the number of chocolate, V be vanilla, and S be strawberry. We have $C > V > S$, and $C + V + S = 9$, with each at least 1.

Since $C > V > S$, and all are integers, S must be at least 1, V at least 2, and C at least 3, but since $C > V$, and $V > S$, it might be tighter. Let me list all possible triplets (C,V,S) such that $C + V + S = 9$, $C > V > S$, and $S \geq 1$, $V \geq 2$, $C \geq 3$.

I'll start with S=1. Then $V > S$, so $V \geq 2$, and $C > V$, so $C \geq 3$. $C + V + 1 = 9$, so $C + V = 8$. Since $C > V$, possible pairs (C,V) with $C + V = 8$ and $C > V$: C=5,V=3; C=6,V=2 but V=2, S=1, so $V > S$, ok? C=6,V=2, but V=2 and S=1, so $V > S$, yes. C=7,V=1 but V=1 not greater than S=1, since V must be greater than S, so V cannot be 1 if S=1. Similarly, C=4,V=4 but C not greater than V. So for S=1, $C + V = 8$, $C > V$, so possible: C=5,V=3; C=6,V=2; C=7,V=1 but V=1 not greater than S=1, so invalid. So only C=5,V=3 or C=6,V=2.

C=6,V=2,S=1: here $V=2 > S=1$, yes. And $C=6 > V=2$, yes.

C=7,V=1,S=1: but $V=1$ not greater than $S=1$, so no.

So for S=1, we have (5,3,1) and (6,2,1).

Now S=2. Then $V > S$, so $V \geq 3$, and $C > V$, so $C \geq 4$. $C + V + 2 = 9$, so $C + V = 7$. $C > V$, so possible:...

Example for Strategy I: Association Method

PROBLEM

N/A

GROUND TRUTH ANSWER

81

MODEL RESPONSE

Step 1. Understand the Problem

We have a (3×9) grid. The numbers $(1, 2, \dots, 9)$ are to be placed so that:

- Each **row** contains all 9 numbers (i.e., each row is a permutation of $\{1, \dots, 9\}$).
- Each of the three (3×3) **blocks** (columns 1–3, 4–6, and 7–9) contains all 9 numbers.

There is **no condition** on the columns (other than within the blocks).

Step 2. Break the Grid into Blocks

Label the rows as 1, 2, 3 and the blocks as:

- **Block 1**: columns 1–3
- **Block 2**: columns 4–6
- **Block 3**: columns 7–9

Each number appears **exactly once** in each row and **exactly once** in each block.

Step 3. Count the Number of Ways to Fill the Grid

We proceed by:

1. Filling **Block 1**
2. Filling **Block 2**
3. Fi...

CHAIN OF THOUGHT (First 1500 chars)

We are given: "The 27 cells of a 3×9 grid are filled in using the numbers 1 through 9 so that each row contains 9 different numbers, and each of the three 3×3 blocks heavily outlined in the example below contains 9 different numbers, as in the first three rows of a Sudoku puzzle.

	4		2		8		9		6		3		1		7		5	
	3		7		9		5		2		1		6		8		4	
	5		6		1		8		4		7		9		2		3	

The number of different ways to fill such a grid can be written as $p^a \cdot q^b \cdot r^c \cdot s^d$, where p, q, r, s and ss are distinct prime numbers and a, b, c, d and d are positive integers. Find $p \cdot a + q \cdot b + r \cdot c + s \cdot d$."

Interpretation: We have a grid that is 3 rows and 9 columns. It is partitioned into three blocks: columns 1-3, columns 4-6, columns 7-9. Condition: Each row is a permutation of numbers 1,...,9. And each block (3x3 square) contains numbers 1,...,9 exactly once.

We want: number of ways to fill such a grid. But careful: The grid is filled with numbers 1,...,9. But note: The numbers are not necessarily placed in a pattern that is the first three rows of a Sudoku? Actually it says: "as in the first three rows of a Sudoku puzzle." That means that if we take a full Sudoku grid, the first three rows satisfy that each row is a permutation of 1,...,9 and each of the three 3x3 blocks (columns 1-3, 4-6, 7-9) are also permutations. But wait: There is also condition that columns? Actually no: In Sudoku, columns also ha...