AMT-130: a historical prior from FDA advisory committee votes

5 November 2026

Summary

uniQure's AMT-130 is a gene therapy for Huntington's disease. The company has said that the U.S. Food and Drug Administration (FDA) accepted its biologics license application for accelerated approval, granted priority review, and set a target decision date of 1 May 2027, and that an advisory committee meeting is planned and not yet dated. Those statements come from uniQure releases on 2 and 4 November 2026. They are company statements until an FDA letter is in hand. The orphan-drug designation is an FDA record: designation 525216, dated 27 September 2017. Orphan designation is for a disease that affects fewer than 200,000 people in the United States. The wording of the committee's question is unset, so the rates below pool approval questions and effectiveness questions.

The meeting table holds 156 rows. The base count is 150, after dropping 4 ties, 1 meeting superseded by a later vote on the same product, and the Lynparza meeting that was favourable only inside a genetic subgroup. Of those 150, 116 asked about approval, licensure, or whether benefits outweigh risks, and 25 asked about effectiveness. Those 141 are the meetings in this calculation. Nine asked about a label, a delay, applicability to U.S. patients, or a surrogate marker. They are listed in the appendix and stay out of the rates. One of the 141, deramiocel on 29 July 2026, has no final action yet. It stays in the vote count and drops out of the approval count, which uses 140 meetings.

On the 141, 92 votes were favourable. The rate is 0.65, and the Jeffreys interval, a range from the counts alone, runs from 0.57 to 0.73. On the 140, 95 products were later approved. The rate is 0.68, and the interval runs from 0.60 to 0.75. A favourable vote means more yes votes than no votes. Given a favourable vote, 86 of 92 products were approved. Given an unfavourable vote, 9 of 48 were approved. Of the 141 votes, 125 come from FDA minutes or transcripts and 16 from press reports. Of the 140 outcomes, 57 come from an FDA document, 44 from openFDA, the agency's public file of drug-application records, 33 from press reports, and 6 from a Securities and Exchange Commission filing or another company document.

Six facts known before the meeting were tested one at a time, and an opioid column was added as a seventh. A meeting was dropped only when that one fact was unrecorded. The test is Fisher's exact test on the two-by-two table. The p-values are two-sided and are not adjusted for the seven factors. For the vote, three factors come in under 0.05. An orphan product was favourable in 40 of 51 meetings, against 50 of 88 ($p = 0.011$). An opioid was favourable in 5 of 15, against 87 of 126 ($p = 0.009$). Priority review was favourable in 57 of 71, against 34 of 67 with standard review ($p = 0.0003$), on the 138 meetings with a recorded review type. Cell and gene therapy was favourable in 10 of 12, against 82 of 129 ($p = 0.22$). The biologics center, a neurology committee, and an initial application stay above 0.05.

For eventual approval, opioid and priority review come in under 0.05. Opioid is 5 of 15 against 90 of 125 ($p = 0.006$). Priority review is 56 of 70 against 39 of 67 ($p = 0.009$). Orphan status is 39 of 50 against 54 of 88 ($p = 0.059$).

Three review types in the whole table are still unverified. Filling those three as standard, and then as priority, gives the same direction on both outcomes. Priority review therefore enters the joint model. The model the vote screen selects is orphan status, opioid, and priority review. Read for AMT-130, which is an orphan product with

priority review and is not an opioid, it gives 0.83 for a favourable vote and 0.81 for eventual approval, from cells of 31 of 38 and 30 of 37. A four-factor model adds cell and gene therapy. It gives 0.84 and 0.87, from cells of 8 of 9 and 8 of 8. The fourth factor does not earn its place.




Figure 1. Penalized log-odds for a favourable vote and for later FDA approval. A circle is the factor on its own, using every meeting that records it. A square is the same factor inside the model that contains orphan status, opioid, and priority review. The line is a bootstrap 95 percent interval. A positive value means more of the outcome. The vote panel uses 141 meetings. The approval panel uses 140. Orphan status is missing for 2 of them. Priority review is missing for 3.

After the vote is known, a model with the vote alone puts eventual approval at 0.90 after a favourable vote and at 0.26 after an unfavourable vote.

Introduction

An FDA advisory committee votes. The agency decides. AMT-130 has not had that meeting. The date, the committee, and the wording of the question are unset. A question about effectiveness is a plausible pivotal question, and so is a question about approval or benefit against risk. The calculation uses both. The answer is a base rate from past cases. It does not fold in the Huntington's data package.

The order is the same as before. Which factors are associated with the vote, and with later approval, when each factor keeps every meeting that records it? What happens to the size and the test when the factors that clear the screen are put in one model? The meeting file and the model output sit beside this report.

Method

Each row is one advisory committee meeting and one product, from 4 November 2016 through 2026, where the committee took a vote on that product. The vote recorded is the pivotal vote. The base count of 150 drops a superseded meeting, a tie, and the Lynparza subgroup vote. An outcome that is still pending is left out of the approval likelihood and kept in the vote count. Deramiocel is that pending row. Its effectiveness vote was 3 yes and 9 no.

The nine meetings left out of the 141 asked a narrower question: a dose in the label, whether to wait for another trial, whether a foreign trial applies to U.S. patients, or whether a surrogate marker is reasonably likely to predict benefit. Those are real meetings. They are a different question from whether the product works or should be

approved. The final action counts a later approval as approved. The one-year action, approval within 365 days, is a sensitivity in the appendix.

Seven factors are coded yes or no. Orphan status means an FDA orphan designation for the indication reviewed. Two cells stay marked "review" and drop out of models that use this factor: Darzalex Faspro for high-risk smoldering myeloma, and Tudriqev. Priority review means the original application's designation. A supplement or a later cycle takes that original designation when a source names it. Where no source names it, the designation is calculated from the initial submission to the original goal date, and the basis says so. A missing day is taken as the 15th. Three rows stay blank, because the dates still do not pick one anniversary: nicotine mouth spray, Tookad, and AR19. A blank drops out of models that use this factor and stays in models that do not. Opioid means an active moiety is an opioid agonist. Every row is yes or no. The fifteen opioid meetings inside the 141 are the ones that are not a tie. Cell and gene therapy means the Cellular, Tissue, and Gene Therapies Advisory Committee, or Kymriah, Ryoncil, Carvykti, or Abecma. The neurology factor means the Peripheral and Central Nervous System Drugs Advisory Committee or the Psychopharmacologic Drugs Advisory Committee. Center means the Center for Biologics Evaluation and Research. Initial application means a first license, as against a supplement. AMT-130 is an initial biologics application at the biologics center, a cell and gene therapy, with orphan status and, on the company's account, priority review. It is not an opioid. The committee is unset. The models below that omit the neurology factor do not move when that assignment is made.

The correlation is the phi coefficient, the ordinary correlation between two yes-or-no columns. The one-at-a-time test is Fisher's exact test. Where the unpenalized logistic fit stays finite, the appendix also gives the likelihood-ratio p-value. The effect size throughout is the penalized coefficient. The penalty matches a normal prior with standard deviation 2.5 on the intercept and standard deviation 1 on each factor. The same penalty is used alone and in a combined model.

The joint model is every factor that clears 0.05 for the vote, after a sensitivity on priority review. That sensitivity fills the blank review types once as standard and once as priority. Priority review enters only when those two fills agree and the recorded-review sample does not remove another factor's association. With three blanks left, the two fills agree, so the joint model is orphan status, opioid, and priority review. Orphan status plus opioid, the same pair with cell and gene therapy added, and a four-factor model that adds cell and gene therapy to the joint model, are reported beside it. On this set the unpenalized fits stay finite.

The bootstrap resamples the meetings 400 times with a fixed seed. The stratum interval is a Jeffreys interval on the meetings that match the factor pattern. When that cell is small, its interval is the description of the cell.

Results

One factor at a time

"Yes" is the count when the factor is present. "No" is the count when it is absent. The last column is the Fisher p-value.

Factor	Meetings	Yes	No	Correlation	Log-odds	p
Favourable vote						
Opioid	141	5/15	87/126	-0.23	-1.13	0.009
Orphan status	139	40/51	50/88	0.22	0.88	0.011
Priority review	138	57/71	34/67	0.31	1.20	0.0003
Biologics center	141	15/18	77/123	0.15	0.78	0.11

Factor	Meetings	Yes	No	Correlation	Log-odds	p
Cell and gene therapy	141	10/12	82/129	0.12	0.67	0.22
Initial application	141	71/110	21/31	-0.03	-0.11	0.83
Neurology committees	141	11/19	81/122	-0.06	-0.29	0.61
Eventual approval						
Opioid	140	5/15	90/125	-0.26	-1.23	0.006
Biologics center	140	15/17	80/123	0.16	0.92	0.093
Cell and gene therapy	140	10/11	85/129	0.14	0.89	0.10
Orphan status	138	39/50	54/88	0.17	0.69	0.059
Priority review	137	56/70	39/67	0.24	0.92	0.009
Initial application	140	73/109	22/31	-0.04	-0.14	0.83
Neurology committees	140	12/19	83/121	-0.04	-0.19	0.61

Opioid, orphan status, and priority review clear 0.05 for the vote. Opioid and priority review also clear it for approval. The fifteen opioid meetings are the opioid rows that are not a tie. Five of them were favourable, and five were later approved. Those are not the same five: one favourable vote was followed by a complete response, and one unfavourable vote was followed by approval. Priority review's 138 leave out 3 votes with no recorded review type. On the 140 with a final action, those 3 include 1 favourable vote and no approvals. The R-squared is 0.07 for opioid on the vote and 0.08 on approval, 0.07 for orphan status on the vote, and 0.13 for priority review on the vote. Cell and gene therapy's vote interval in Figure 1 crosses zero. The two unfavourable votes in that group are NurOwn, 1 yes and 17 no, with the application later withdrawn, and deramiocel, 3 yes and 9 no, still pending. Both were effectiveness questions. NurOwn's review type is standard on the calculated clock.

For approval, the Fisher test and the likelihood-ratio test disagree at two rows that the Fisher test does not carry forward. Orphan status has Fisher p = 0.059 and likelihood-ratio p = 0.041. The biologics center has Fisher p = 0.093 and likelihood-ratio p = 0.038. Opioid's likelihood-ratio p-value on approval is 0.004. Priority review's likelihood-ratio p-value on approval is 0.005.

Why priority review is in the joint model

The recorded-review comparison for the vote is 57 of 71 against 34 of 67 (p = 0.0003, log-odds 1.20). Coding the three blanks as standard review gives 57 of 71 against 35 of 70 (p = 0.0002, log-odds 1.23). Coding them as priority review gives 58 of 74 against 34 of 67 (p = 0.0007, log-odds 1.11). For approval the same two fills are p = 0.004 (log-odds 1.01) and p = 0.029 (log-odds 0.76). The direction does not change. The three meetings with no recorded review type had one favourable vote and no approvals. Restricting to the recorded-review meetings leaves the vote associations for orphan status and for opioid in place. Priority review enters the joint model.

Orphan status and opioid together

This is the pair from the earlier screen, kept so the priority term can be compared with it. It uses the meetings that record orphan status. The fit uses 139 vote meetings and 138 approval meetings. On those meetings, before the factors are combined, orphan status is 40 of 51 against 50 of 88 (p = 0.011, log-odds 0.88) and opioid is 5 of 15 against 85 of 124 for the vote (p = 0.010, log-odds -1.11).

Factor	Vote, alone	Vote, combined	Approval, alone	Approval, combined
Orphan status	0.88	0.75	0.69	0.53
Opioid	-1.11	-0.90	-1.21	-1.06

The vote drop-one p-values are 0.045 for orphan status and 0.044 for opioid. The criterion is 177.5 for orphan status alone and 175.5 with opioid, so the second factor earns its place on the vote. The joint bootstrap intervals run from 0.08 to 1.52 and from -1.69 to -0.13. For approval, opioid's drop-one p-value is 0.016 and orphan's is 0.18. The approval bootstrap interval for orphan status crosses zero. Figure 1 now puts the squares on the three-factor screen, which adds priority review.

Orphan status, opioid, and cell and gene therapy

This is the two-factor model with cell and gene therapy added. It uses the same 139 vote meetings and 138 approval meetings, because opioid and cell and gene therapy are filled on every row and the only rows dropped are the two orphan cells marked "review." Tudriqev is one of those two, and it is a cell and gene therapy, so the class count inside this fit is 9 of 11 for the vote and 9 of 10 for approval. The one-at-a-time count of 10 of 12 keeps Tudriqev. On these meetings, before the factors are combined, the vote log-odds are 0.88 for orphan status, -1.11 for opioid, and 0.61 for cell and gene therapy (9 of 11 against 81 of 128, p = 0.33).

Factor	Vote, alone	Vote, combined	Approval, alone	Approval, combined
Orphan status	0.88	0.71	0.69	0.45
Opioid	-1.11	-0.89	-1.21	-1.06
Cell and gene therapy	0.61	0.21	0.83	0.57

The vote drop-one p-values are 0.081 for orphan status, 0.044 for opioid, and 0.75 for cell and gene therapy. The criterion is 175.5 for orphan status and opioid, and 177.4 with the third factor, so the third factor does not earn its place on the vote. The bootstrap intervals are 0.02 to 1.50, -1.68 to -0.12, and -0.66 to 1.09. For approval the drop-one p-values are 0.38, 0.016, and 0.27, and the criterion moves from 170.2 to 171.0. The approval bootstrap intervals are -0.22 to 1.21, -1.87 to -0.27, and -0.34 to 1.26. The cell-and-gene-therapy interval crosses zero on both outcomes. Among orphan products, the class is 9 of 11 against 31 of 40 for the vote (p = 1.00) and 9 of 10 against 30 of 40 for approval (p = 0.42).

Priority review, orphan status, opioid, and cell and gene therapy

This is the screen-selected model with cell and gene therapy added. It uses 136 vote meetings and 135 approval meetings. The two orphan cells marked "review" drop out, and so do the three meetings with no review type. On these meetings, before the factors are combined, the vote log-odds are 0.85 for orphan status, 1.21 for priority review, -1.14 for opioid, and 0.59 for cell and gene therapy (9 of 11 against 80 of 125, p = 0.33).

Factor	Vote, alone	Vote, combined	Approval, alone	Approval, combined
Orphan status	0.85	0.39	0.61	0.13
Priority review	1.21	1.03	0.93	0.77
Opioid	-1.14	-0.86	-1.28	-1.09
Cell and gene therapy	0.59	0.13	0.79	0.52

The vote drop-one p-values are 0.46 for orphan status, 0.004 for priority review, 0.058 for opioid, and 0.86 for cell and gene therapy. The criterion is 164.2 without the fourth factor and 166.1 with it, so cell and gene therapy does not earn its place. The bootstrap intervals are -0.26 to 1.13, 0.42 to 1.76, -1.67 to -0.08, and -0.75 to 1.00. For approval the drop-one p-values are 0.94, 0.034, 0.013, and 0.30, and the criterion moves from 160.7 to 161.6. The approval bootstrap intervals are -0.56 to 0.85, 0.06 to 1.46, -1.91 to -0.34, and -0.25 to 1.14. Orphan status and cell and gene therapy cross zero on both outcomes once priority review is in the model.

Orphan status and cell and gene therapy together

This pair leaves opioid out. On the 139 vote meetings the coefficients are 0.84 for orphan status and 0.22 for cell and gene therapy. The drop-one p-values are 0.021 and 0.75. The criterion is 177.5 for orphan status alone and 179.4 with the second factor. On the 138 approval meetings the coefficients are 0.61 and 0.58, and the drop-one p-values are 0.13 and 0.27.

What the fits give for AMT-130

Model	Outcome	Probability	Bootstrap 95%	Matching meetings	Interval for that cell
Orphan status	Favourable vote	0.77	0.65-0.87	40/51	0.66-0.88
Orphan status	Eventual approval	0.77	0.66-0.86	39/50	0.65-0.88
Orphan, not an opioid	Favourable vote	0.77	0.65-0.87	40/51	0.66-0.88
Orphan, not an opioid	Eventual approval	0.77	0.66-0.87	39/50	0.65-0.88
Orphan, not an opioid, cell and gene therapy	Favourable vote	0.80	0.62-0.92	9/11	0.53-0.96
Orphan, not an opioid, cell and gene therapy	Eventual approval	0.84	0.69-0.92	9/10	0.62-0.99
Orphan, priority review, not an opioid	Favourable vote	0.83	0.73-0.91	31/38	0.67-0.91
Orphan, priority review, not an opioid	Eventual approval	0.81	0.70-0.90	30/37	0.66-0.91
Those three, plus cell and gene therapy	Favourable vote	0.84	0.67-0.94	8/9	0.59-0.99
Those three, plus cell and gene therapy	Eventual approval	0.87	0.76-0.94	8/8	0.74-1.00
Orphan and cell and gene therapy	Favourable vote	0.80	0.62-0.91	9/11	0.53-0.96
Orphan and cell and gene therapy	Eventual approval	0.84	0.69-0.92	9/10	0.62-0.99

AMT-130 is not an opioid, and on the company's account it has priority review. The screen-selected model is read at orphan status yes, priority review yes, and opioid no. It gives 0.83 and 0.81. The four-factor model also sets cell and gene therapy to yes. It gives 0.84 and 0.87. The matching cells are 8 of 9 and 8 of 8. NurOwn is standard review on the calculated clock, so it is not in those cells. Deramiocel is the unfavourable vote in the 8 of 9, and it has no final action, which is why the approval cell is 8 of 8. There is no completed meeting in that cell with an unfavourable vote. Tudriqev stays out because its orphan label is still "review." All ten completed cell-and-gene-therapy votes in the orphan cell come from FDA minutes or transcripts. Two of the ten outcomes come from an FDA document, four from openFDA, and four from press reports.

Once the vote is known, the pre-meeting labels add little. The vote-only model has an R-squared of 0.64. Eventual approval is 0.90 after a favourable vote and 0.26 after an unfavourable vote. The bootstrap intervals run from 0.85 to 0.94 and from 0.17 to 0.36.

Discussion

Adding the 25 effectiveness questions changes the cell-and-gene-therapy count from a run of favourable votes to 10 of 12. NurOwn and deramiocel are the two unfavourable votes, and both were effectiveness questions. That is why a model fit on approval questions only made the class look separate from everything else, and why the same factor has a Fisher p-value of 0.22 once those questions are included. Ryoncil, also an effectiveness question, was favourable and was approved in 2024 after a complete response in 2020. Tudriqev was favourable and approved, and it stays out of the orphan models because the designation text does not settle the indication.

Opioid is the new association, and it runs against approval. Five of fifteen opioid meetings were favourable, and five of fifteen were approved. The products are oxycodone, hydrocodone, buprenorphine, tramadol, sufentanil, oliceridine, and oxycodegol, including buprenorphine for opioid use disorder and for depression. The column is filled for every meeting, so the estimate does not depend on dropping a blank.

Priority review clears 0.05 for the vote and for approval on the meetings with a recorded original designation, 138 and 137 of them. The sensitivity agrees. Three cells in the whole table remain unverified, and all three are inside the 141. One of those votes was favourable, and none of the three products was approved. The flag is the original application's designation, including where the committee reviewed a supplement or a resubmission. Twenty-one of the standard values are calculated from the submission date and the goal date, and the basis says so.

The biologics center is the other near result for approval, at Fisher p = 0.093. In a model with every factor except priority review, the opioid vote coefficient is -0.86 and the biologics-center coefficient is 0.50. The cell-and-gene-therapy vote coefficient is -0.08. With priority review added, and with the three blanks dropped, the priority vote coefficient is 1.04 and the opioid coefficient is -0.85. The neurology committees lean slightly the other way on this wider set (11 of 19 favourable). No cell-and-gene-therapy meeting in the table was held by either of those committees.

The vote remains the strong association. An R-squared of 0.18 for orphan status, opioid, and priority review together, and 0.64 for the vote itself, is the scale of this table. The final action and the one-year action disagree on 14 of the 81 meetings that have both labels. On those 81, 54 products were approved within a year. The one-year screen does not clear 0.05 for any factor. The FDA calls a committee when the decision is uncertain enough to want one. The 0.65 and 0.68 figures are rates among products that reached a vote. The table has no column for the strength of the evidence, for an external control, or for accelerated approval. Sixteen of the 141 votes and 39 of the 140 outcomes rest on press or company documents.

Conclusions

Of 150 pivotal meetings, 141 asked an approval, benefit-risk, or effectiveness question. The chance of a favourable vote on those 141 is 0.65 (92 of 141). The chance of eventual approval on the 140 with a final action is 0.68 (95 of 140). Given a favourable vote, 86 of 92 were approved. Given an unfavourable vote, 9 of 48 were approved. The vote-only model puts those conditional chances at 0.90 and 0.26.

Taken one at a time, opioid, orphan status, and priority review clear a Fisher test at 0.05 for the vote: 5 of 15 against 87 of 126 (p = 0.009), 40 of 51 against 50 of 88 (p = 0.011), and 57 of 71 against 34 of 67 (p = 0.0003). Cell and gene therapy is 10 of 12 (p = 0.22). For approval, opioid is 5 of 15 against 90 of 125 (p = 0.006) and priority review is 56 of 70 against 39 of 67 (p = 0.009). Orphan status is 39 of 50 against 54 of 88 (p = 0.059).

The sensitivity on the three blank review types agrees with itself, so priority review is in the joint model. In the model with orphan status, opioid, and priority review, the vote coefficients are 0.41, -0.86, and 1.03. The drop-one p-values are 0.40, 0.058, and 0.004. Read as an orphan product with priority review that is not an

opioid, that model gives 0.83 and 0.81, from cells of 31 of 38 and 30 of 37. Adding cell and gene therapy moves the vote coefficients to 0.39, 1.03, -0.86, and 0.13. The fourth factor's drop-one p-value is 0.86, and the criterion rises from 164.2 to 166.1. Read as an orphan gene therapy with priority review that is not an opioid, the four-factor model gives 0.84 and 0.87, from cells of 8 of 9 and 8 of 8.

Orphan status is taken from FDA designation 525216. Priority review for AMT-130 is a company statement. The clinical evidence is outside this calculation. Nine meetings with a narrower question are outside the 141.

Suggested improvements

Review type is still unverified for 3 of the 141 votes: nicotine mouth spray, Tookad, and AR19. The flag is the original application's review type. A calculated value is tagged in the basis, including where a missing day was taken as the 15th or where the complete-response date was used because it landed on the anniversary. The page checked for each remaining blank is named in the factor log.

Darzalex Faspro for high-risk smoldering myeloma, and Tudriqev, should be set to yes or no only if a source settles the indication. Deramiocel's final action is open until the agency decides. The one-year label is thin on this set and was left unverified where no source dated the action.

A sourced column for accelerated approval, for a resubmission, and for an external control would address the largest hole in the factor list. The residual of about 19 meetings against the Daval count for 2016 to 2021 is a difference in published totals. That paper does not list the meetings.

Appendix. The second test

The likelihood-ratio p-value is from the unpenalized logistic fit on the same meetings. R-squared is Nagelkerke's.

Factor	Meetings	R-squared	Likelihood-ratio p
Favourable vote			
Opioid	141	0.07	0.008
Orphan status	139	0.07	0.009
Priority review	138	0.13	0.0002
Biologics center	141	0.03	0.069
Cell and gene therapy	141	0.02	0.15
Neurology committees	141	0.00	0.47
Initial application	141	0.00	0.74
Eventual approval			
Opioid	140	0.08	0.004
Orphan status	138	0.04	0.041
Biologics center	140	0.04	0.038
Cell and gene therapy	140	0.04	0.060
Priority review	137	0.08	0.005
Neurology committees	140	0.00	0.64
Initial application	140	0.00	0.67

These rows are not ranked against each other by a criterion. Priority review's row sums fewer meetings.

Appendix. More than two factors

The model without priority review drops the two orphan cells marked "review." The unpenalized fit is stable. The model with priority review also drops the three blank review types. Its unpenalized fit is stable, and the coefficients below are the penalized ones.

Factor	Vote, without priority	Approval, without priority	Vote, all seven	Approval, all seven
Orphan status	0.69	0.43	0.35	0.09
Opioid	-0.86	-1.02	-0.85	-1.10
Cell and gene therapy	-0.08	0.27	-0.03	0.32
Priority review			1.04	0.77
Biologics center	0.50	0.58	0.25	0.39
Neurology committees	-0.10	0.00	-0.23	-0.09
Initial application	-0.05	-0.03	0.06	0.13

On the recorded-review sample the priority vote coefficient stays near 1.0 and the orphan coefficient falls to 0.35. A year term, added to the screen-selected model and evaluated as a line centered at 2020, moves the criterion from 160.4 to 158.3 for the vote and from 160.7 to 161.0 for approval. It is not added.

Appendix. Approval within one year

Eighty-one of the 140 meetings with a final action also have a one-year label. Fifty-four of the 81 were approved within a year. The two labels disagree 14 times. Priority review is the one factor whose Fisher test clears 0.05 on this outcome. The p column is that test.

Factor	Meetings	Yes	No	Correlation	Log-odds	p
Orphan status	79	22/30	31/49	0.10	0.38	0.46
Priority review	80	30/37	24/43	0.27	0.97	0.019
Initial application	81	52/75	2/6	0.20	0.85	0.092
Neurology committees	81	10/13	44/68	0.10	0.41	0.53
Opioid	81	3/6	51/75	-0.10	-0.43	0.39
Cell and gene therapy	81	2/4	52/77	-0.08	-0.35	0.60
Biologics center	81	2/4	52/77	-0.08	-0.35	0.60

The orphan-only probability on this outcome is 0.72 (bootstrap 0.57 to 0.85), from 22 of 30. The screen-selected model, read at orphan status yes, priority review yes, and opioid no, gives 0.79.

Appendix. Approval given the vote

The model with the vote and no pre-meeting factor is stable, uses the 140 meetings with a final action, and has an R-squared of 0.64.

Vote	Probability	Bootstrap 95%	Meetings	Interval for that cell
Favourable	0.90	0.85-0.94	86/92	0.87-0.97
Unfavourable	0.26	0.17-0.36	9/48	0.10-0.31

Adding orphan status, opioid, and priority review puts the favourable-vote probability at 0.91, on a cell of 30 of 31, and the unfavourable probability at 0.29, on a cell of 0 of 6. Adding cell and gene therapy puts the favourable-vote probability at 0.93, on a cell of 8 of 8. The unfavourable cell for that pattern is empty: NurOwn, the unfavourable cell-and-gene-therapy vote with a final action, is standard review on the calculated clock, and deramiocel has no final action.

Appendix. Cell and gene therapy meetings

These rows are in the 141. Tudriqev is in the vote and approval counts and drops out of any model that uses orphan status. Deramiocel is in the vote count only.

Date	Committee	Product	Question	Yes-no	Vote	Final action	Orphan
12 Jul 2017	ODAC	Kymriah	Approval	10-0	Favourable	Approved	Yes
12 Oct 2017	CTGTAC	Luxturna	Approval	16-0	Favourable	Approved	Yes
13 Aug 2020	ODAC	Ryoncil	Effectiveness	8-2	Favourable	Approved in 2024	Yes
15 Apr 2021	CTGTAC	Lantidra	Approval	12-4	Favourable	Approved	Yes
9 Jun 2022	CTGTAC	Skysona	Approval	15-0	Favourable	Approved	Yes
10 Jun 2022	CTGTAC	Zynteglo	Approval	13-0	Favourable	Approved	Yes
12 May 2023	CTGTAC	Elevidys	Approval	8-6	Favourable	Approved	Yes
27 Sep 2023	CTGTAC	NurOwn	Effectiveness	1-17	Unfavourable	Withdrawn	Yes
15 Mar 2024	ODAC	Carvykti	Approval	11-0	Favourable	Approved	Yes
15 Mar 2024	ODAC	Abecma	Approval	8-3	Favourable	Approved	Yes
29 Jul 2026	CTGTAC	Deramiocel	Effectiveness	3-9	Unfavourable	Pending	Yes
30 Jul 2026	CTGTAC	Tudriqev	Effectiveness	10-3	Favourable	Approved	Review

ODAC is the Oncologic Drugs Advisory Committee. CTGTAC is the Cellular, Tissue, and Gene Therapies Advisory Committee. The ten completed orphan rows were favourable in 9 and approved in 9. NurOwn is the failure in both columns.

Appendix. The nine meetings left out

Each pivotal question was about a label, a delay, applicability, or a surrogate, so the meeting is outside the 141.

Date	Product	What the question asked
8 Mar 2018	Xeljanz	Whether a 10 mg maintenance dose belonged in the label
7 Aug 2018	Arikayce	Coded restriction/other; the stored text is the agenda
26 Feb 2019	Xpovio	Whether approval should wait for a randomized trial
9 Feb 2021	Keytruda	Whether the decision should be deferred for more data
24 Jun 2021	Retifanlimab	Coded restriction/other
10 Feb 2022	Sintilimab	Whether another trial in U.S. patients should be required first
22 Mar 2023	Qalsody	Whether a reduction in a blood marker was reasonably likely to predict benefit
5 Oct 2023	Lumakras	Whether progression-free survival could support the application
20 May 2025	Columvi	Whether a foreign trial's population applied to U.S. patients

Qalsody is a treatment for a neurologic disease, reviewed by a neurology committee. Its question was about a surrogate marker. It is part of the nine, and it is not in the 0.65 and 0.68 rates.

Appendix. Where the meetings come from

The meeting list began as FDA advisory committee notices in the Federal Register. Votes were taken from FDA minutes, voting results, and transcripts, with press and company filings where the minutes could not be recovered. A published count of meetings from 2016 to 2021, Daval and colleagues in JAMA Health Forum in 2023, was the completeness check. That paper publishes committee-year totals. About 19 of those counts remain unmatched after the start date of this table, pharmacy-compounding meetings, and joint meetings filed under another committee. One row was added from that search: solithromycin, on 4 November 2016, a benefit-risk vote of 7 yes and 6 no.

The whole table has 156 rows. Votes come from FDA minutes or transcripts for 137 and from press reports for 19. Outcomes come from an FDA document or openFDA for 112, from press reports for 38, and from a company filing for 6.

Appendix. Files

The meeting table is `adcom_full_pivotal.csv`. The four-cell counts are `two_by_two.csv`. The factor lookups are `factor_verification.csv`. This screen is `amt130_one_at_a_time.csv` and `amt130_factor_compare.json`. The figure is `charts/amt130_factor_compare.png`. The fit is `scripts/amt130_factor_compare.py`, run from this folder. Corrections to the meeting table are made in `scripts/cur.py` and applied when the table is rebuilt.