

IPML

IMPRECISE
PROBABILISTIC
MACHINE LEARNING

Lecture 9: Conformal Prediction and Calibration

Krikamol Muandet
16 January 2026



Outline

1. Conformal Prediction
2. Probabilistic Calibration
3. Applications

Conformal Prediction

Conformal Prediction

- Conformal prediction is a *frequentist* approach to **distribution-free** uncertainty quantification that is:
 1. Agnostic to the model
 2. Agnostic to data distribution
 3. Valid in finite sample



Conformal Coverage Guarantee

- Given a predictive model $\hat{f}: \mathcal{X} \rightarrow \Delta(\mathcal{Y})$ and the test point $(X_{\text{test}}, Y_{\text{test}})$, we seek to construct a **prediction set** $\mathcal{C}(X_{\text{test}}) \subset \mathcal{Y}$ that is **valid**, meaning

$$P(Y_{\text{test}} \in \mathcal{C}(X_{\text{test}})) \geq 1 - \alpha$$

where $\alpha \in [0,1]$ is a user-chosen error rate.

- Marginal coverage:** *The probability that the prediction set contains the correct label is almost exactly $1 - \alpha$.*

Conformal Coverage Guarantee



{
fox
squirrel
0.99
}



{
fox
squirrel, gray, bucket, rain
0.82, fox, 0.03, barrel
0.02, 0.02
}



{
marmot, fox
0.30, squirrel, 0.22, mink, weasel, beaver, polecat
0.18, 0.16, 0.03, 0.01
}

We want the prediction set to contains the correct label **with high probability**.
The prediction set $\mathcal{C}(X_{\text{test}})$ captures the model's uncertainty on the data X_{text} .

Split Conformal Prediction



Given a **calibration set** $(X_1, Y_1), \dots, (X_n, Y_n)$:

1. Identify a heuristic uncertainty using the pre-trained model
2. Define the **score function** $s(x, y) \in \mathbb{R}$
3. Compute \hat{q} as the $\frac{[(n+1)(1-\alpha)]}{n}$ quantile of the calibration scores

$$s_1, s_2, \dots, s_n := s(X_1, Y_1), s(X_2, Y_2), \dots, s(X_n, Y_n)$$

4. Form the prediction sets of the new example X_{test} as

$$\mathcal{C}(X_{\text{test}}) = \{y : s(X_{\text{test}}, y) \leq \hat{q}\}$$



Coverage Property

Suppose $(X_1, Y_1), \dots, (X_n, Y_n)$ and $(X_{\text{test}}, Y_{\text{test}})$ are i.i.d and we define \hat{q} as

$$\hat{q} = \inf\left\{q : \frac{|\{i : s(X_i, Y_i) \leq q\}|}{n} \geq \frac{\lceil (n+1)(1-\alpha) \rceil}{n}\right\}$$

The prediction set $C(X) = \{y : s(X, y) \leq \hat{q}\}$ has a **valid marginal coverage**.

- WOLG, assume that the calibration scores are sorted: $s_1 < s_2 < \dots < s_n$.
 1. If $\alpha < 1/(n+1)$, $\hat{q} = \infty$. Then, $C(X) = \mathcal{Y}$. **Valid!**
 2. If $\alpha \geq 1/(n+1)$, $\hat{q} = s_{\lceil (n+1)(1-\alpha) \rceil}$.
- Note that $\{Y_{\text{test}} \in C(X_{\text{test}})\} = \{s_{\text{test}} \leq \hat{q}\} = \{s_{\text{test}} \leq s_{\lceil (n+1)(1-\alpha) \rceil}\}$
- By exchangeability, $P(s_{\text{test}} \leq s_k) = k/(n+1)$.
- $P(s_{\text{test}} \leq s_{\lceil (n+1)(1-\alpha) \rceil}) = \lceil (n+1)(1-\alpha) \rceil / (n+1) \geq 1 - \alpha$.

Choice of Score Function

The **score function** $s(x, y) \in \mathbb{R}$ encodes *disagreement* between x and y . The larger the score $s(x, y)$, the worse the agreement between x and y .



While the prediction set may provide a valid marginal coverage, the set can be useless if the score function is uninformative.

Adaptive Prediction Sets


- Assume a predictive model $\hat{f}: \mathcal{X} \rightarrow \Delta(\mathcal{Y})$ where $\mathcal{Y} = \{1, \dots, K\}$.
- Let $\pi_1(x), \dots, \pi_K(x)$ be the permutation of $\{1, \dots, K\}$ that sorts $\hat{f}(x)$ from most likely to least likely.
- Then, we define a score function as

$$s(x, y) = \sum_{j=1}^k \hat{f}(x)_{\pi_j(x)}, \quad \text{with } y = \pi_k(x)$$

- Calculate the quantile $\hat{q} = \text{quantile}\left(s_1, \dots, s_n; \frac{\lceil (n+1)(1-\alpha) \rceil}{n}\right)$ and form the set

$$C(x) = \{\pi_1(x), \dots, \pi_k(x)\}, \quad k = \sup \left\{ k' : \sum_{j=1}^{k'} \hat{f}(x)_{\pi_j(x)} < \hat{q} \right\} + 1$$

Adaptive Prediction Sets




Conformalized Quantile Regression

- **Quantile regression:** learn the γ quantile t_γ of $Y_{\text{test}} \mid X_{\text{test}} = x$
- The quantile $[\hat{t}_{0.05}(x), \hat{t}_{0.95}(x)]$ has approximately 90% coverage.

$$s(x, y) = \max\{\hat{t}_{\alpha/2}(x) - y, y - \hat{t}_{1-\alpha/2}(x)\}$$

$$C(x) = [\hat{t}_{\alpha/2}(x) - \hat{q}, \hat{t}_{1-\alpha/2}(x) + \hat{q}]$$

$$\hat{q} = \text{Quantile}\left(s_1, \dots, s_n; \frac{[(n+1)(1-\alpha)]}{n}\right)$$



Conformalized Uncertainty Estimates

Given some **point prediction** $\hat{f}(x)$ and some **uncertainty scalar** $u(x)$, we can provide a conformal guarantee:

- **Standard deviation** $\hat{\sigma}(x)$: We can assume that

$$Y_{\text{test}} \mid X_{\text{test}} = x \sim N(\mu(x), \sigma(x))$$

with models $\hat{f}(x)$ and $\hat{\sigma}(x)$. Conformal prediction gives $\hat{f}(x) \pm \hat{q}\hat{\sigma}(x)$.

- **Magnitude of the residual** $\hat{r}(x)$: After fitting a model \hat{f} , we fit a second model \hat{r} that predicts $|y - \hat{f}(x)|$.

$$s(x, y) = \frac{|y - \hat{f}(x)|}{u(x)}, \quad C(x) = [\hat{f}(x) - u(x)\hat{q}, \hat{f}(x) + u(x)\hat{q}]$$

Example

Given a trained probabilistic classifier and a calibration set, compute a prediction set $\mathcal{C}(x) \subseteq \{A, B, C\}$ for a new input x_{test} , using split conformal in the multiclass classification setting.

- For a new x_{test} , the classifier outputs:
 - $\hat{p}(A \mid x_{\text{test}}) = 0.50$
 - $\hat{p}(B \mid x_{\text{test}}) = 0.35$
 - $\hat{p}(C \mid x_{\text{test}}) = 0.15$
- We want 90% marginal coverage, so $\alpha = 0.1$.

i	True y_i	$\hat{p}(y_i x_i)$
1	A	0.80
2	B	0.55
3	C	0.60
4	A	0.40
5	B	0.70
6	C	0.30
7	A	0.90
8	B	0.20
9	C	0.45
10	A	0.65

Example

- Define the non-conformity score as

$$s(x, y) = 1 - \hat{p}(y | x)$$

- Let $n = 10$. Then, compute the index

$$k = \lceil (n + 1)(1 - \alpha) \rceil = 10$$

- Sort s_1, \dots, s_{10} increasingly and set the threshold $\tau = s_{(k)}$ (the k -th smallest) [$\tau = 0.8$]
- For each label $y \in \{A, B, C\}$, compute $s(x_{\text{test}}, y)$ and include y in the set if $s(x_{\text{test}}, y) \leq \tau$.

i	True y_i	$\hat{p}(y_i x_i)$	s_i
1	A	0.80	0.20
2	B	0.55	0.45
3	C	0.60	0.40
4	A	0.40	0.60
5	B	0.70	0.30
6	C	0.30	0.70
7	A	0.90	0.10
8	B	0.20	0.80
9	C	0.45	0.55
10	A	0.65	0.35

Sol: $s(x_{\text{test}}, A) = 0.5, s(x_{\text{test}}, B) = 0.65, s(x_{\text{test}}, C) = 0.85 \Rightarrow \mathcal{C}, (x_{\text{test}}) = \{A, B\}$

Adaptivity

The procedure should return **large sets** for harder inputs and **smaller sets** for easier inputs.



{
fox
squirrel
0.99
}



{
fox, gray, bucket, rain
squirrel, fox, barrel
0.82, 0.03, 0.02, 0.02
}



{
marmot, fox, mink, weasel, beaver, polecat
0.30, squirrel, 0.22, 0.18, 0.16, 0.03, 0.01
}


Conditional Coverage

- The **conditional coverage** property:

$$P(Y_{\text{test}} \in C(X_{\text{test}}) \mid X_{\text{test}}) \geq 1 - \alpha$$

- **Interpretation:** *for every value of the input X_{test} , we seek to return prediction sets with $1 - \alpha$ coverage.*
 - If we have two groups, A and B, comprising 90% and 10% of the population, conditional coverage requires that the prediction sets cover Y with at least 90% probability within each group.
- This is a stronger property than the *marginal coverage* property and is **impossible** to achieve in general.

Conditional Coverage



Extensions

- Group-balanced conformal prediction:

$$P(Y_{\text{test}} \in C(X_{\text{test}}) \mid X_{\text{test},1} = g) \geq 1 - \alpha, \quad \forall g \in \{1, \dots, G\}$$

- Class-conditional conformal prediction:

$$P(Y_{\text{test}} \in C(X_{\text{test}}) \mid Y_{\text{test}} = y) \geq 1 - \alpha, \quad \forall y \in \{1, \dots, K\}$$

- Conformal risk control:

$$\mathbb{E}[\ell(C(X_{\text{test}}), Y_{\text{test}})] \leq \alpha, \quad \forall y \in \{1, \dots, K\}$$

- Conformal prediction under distribution shift


Probabilistic Calibration

Probabilistic Calibration

- Calibration ensures **honesty** in probability estimates overall.
- A probabilistic predictor \hat{f} is *calibrated* if predicted probabilities match empirical frequencies:

$$P[Y = 1 \mid \hat{f}(x) = \rho] = \rho, \quad \forall \rho \in [0,1]$$

- **Interpretation:** Among all instances predicted with probability ρ , about ρ fraction of them should be positive.
- Violations indicate **systematic misrepresentation** of uncertainty.




Multi-Calibration

- A predictor is *multi-calibrated* if it is calibrated **simultaneously** across many subpopulations:

$$P[Y = 1 \mid \hat{f}(x) = \rho, G = g] = \rho, \quad \forall \rho \in [0,1], \forall g \in \{1, \dots, K\}$$



- Calibration holds not only overall, but for *every relevant group*.



Applications

Multilabel Classification


The model's output is thresholded to get the subset of K classes, $C_\lambda(x) = \{y : \hat{f}(x) \geq \lambda\}$. The conformal risk control is used to pick the threshold λ certifying a low false negative rate (FNR).



Red = false negatives, Blue = false positives, Black = true positives

Tumor Segmentation



The model's output is thresholded to get the predicted binary mask, $C_\lambda(x) = \{(i, j) : \hat{f}(x)_{(i, j)} \geq \lambda\}$. The conformal risk control is used to pick the threshold λ certifying a low false negative rate (FNR).



Red = false negatives, Blue = false positives, Black = true positives

Weather Prediction with Time-Series

We seek to predict the temperature of different locations on Earth given covariates such as the latitude, longitude, altitude, atmospheric pressure, and so on.



Exchangeability is violated!

Recommended Reading

- Anastasios N. Angelopoulos and Stephen Bates. [**A Gentle Introduction to Conformal Prediction and Distribution-Free Uncertainty Quantification.**](#) arXiv:2107.07511 (2022).
- Glenn Shafer, Vladimir Vovk. [**A Tutorial on Conformal Prediction.**](#) Journal of Machine Learning Research, 9(12):371–421, 2008.

Exercise I

You are given a trained probabilistic classifier over 4 classes $\{A, B, C, D\}$ and a calibration set of $n = 12$ examples. Your goal is to compute a split conformal prediction set $\mathcal{C}(x_{\text{test}}) \subseteq \{A, B, C, D\}$ for a new input x_{test} , targeting 80% marginal coverage.

For the new input x_{test} , the classifier outputs:

- $\hat{p}(A \mid x_{\text{test}}) = 0.34$
- $\hat{p}(B \mid x_{\text{test}}) = 0.27$
- $\hat{p}(C \mid x_{\text{test}}) = 0.22$
- $\hat{p}(D \mid x_{\text{test}}) = 0.17$

i	True y_i	$\hat{p}(y_i x_i)$
1	A	0.88
2	B	0.62
3	C	0.51
4	D	0.44
5	A	0.73
6	B	0.39
7	C	0.81
8	D	0.57
9	A	0.29
10	B	0.76
11	C	0.35
12	D	0.68

Exercise II

You are given a trained probabilistic classifier over 4 classes $\{A, B, C, D\}$ and a calibration set of $n = 10$ examples. Your goal is to compute a split conformal prediction set $\mathcal{C}(x_{\text{test}}) \subseteq \{A, B, C, D\}$ for a new input x_{test} , targeting 80% marginal coverage.

For the new input x_{test} , the classifier outputs:

- $\hat{p}(A \mid x_{\text{test}}) = 0.41$
- $\hat{p}(B \mid x_{\text{test}}) = 0.33$
- $\hat{p}(C \mid x_{\text{test}}) = 0.16$
- $\hat{p}(D \mid x_{\text{test}}) = 0.10$

i	True y_i	$\hat{p}(y_i x_i)$
1	A	0.91
2	B	0.78
3	C	0.66
4	A	0.59
5	D	0.84
6	B	0.72
7	C	0.63
8	D	0.55
9	A	0.47
10	B	0.40

Exercise III

You are given a trained probabilistic classifier over 4 classes $\{A, B, C, D\}$ and a calibration set of $n = 10$ examples. Your goal is to compute a split conformal prediction set $\mathcal{C}(x_{\text{test}}) \subseteq \{A, B, C, D\}$ for a new input x_{test} , targeting 80% marginal coverage.

For the new input x_{test} , the classifier outputs:

- $\hat{p}(A \mid x_{\text{test}}) = 0.38$
- $\hat{p}(B \mid x_{\text{test}}) = 0.26$
- $\hat{p}(C \mid x_{\text{test}}) = 0.21$
- $\hat{p}(D \mid x_{\text{test}}) = 0.15$

i	True y_i	$\hat{p}(y_i x_i)$
1	A	0.93
2	B	0.81
3	C	0.74
4	D	0.70
5	A	0.65
6	B	0.61
7	C	0.58
8	D	0.54
9	A	0.49
10	B	0.42