Three Triangles Integer Factorization Algorithm

"Discovery consists of seeing what everybody has seen and thinking what nobody has thought."

— Albert Szent-Györgyi

Abstract:

Factor the composite number C = (a+1)(a+b) by finding the solution to C = T(a) + T(a+b) - T(b-1) where T(n) is the nth triangular number.

Definitions:

- a, b := natural numbers
- T(n) := n(n+1)/2
- S(a, b) := T(a)+T(a+b)-T(b-1)
- make_even(n) := if n is odd return n-1 else return n

Theorem:

For every composite number C = (a+1)(a+b) there exist three triangular numbers such that C = T(a) + T(a+b) - T(b-1).

Proof:

```
1. (a+1)(a+b)=T(a)+T(a+b)-T(b-1)
2. a^2+ab+a+b=a(a+1)/2+(a+b)(a+b+1)/2-(b-1)b/2
3. 2a^2+2ab+2a+2b=a(a+1)+(a+b)(a+b+1)-(b-1)b
4. 2a^2+2ab+2a+2b=a^2+a+a(a+b+1)+b(a+b+1)-b^2+b
5. 2a^2+2ab+2a+2b=a^2+a+a^2+ab+a+ba+b^2+b-b^2+b
```

6. $2a^2+2ab+2a+2b=2a^2+2ab+2a+2b$

7. 0=0

Theorem:

For natural numbers a, b it holds that S(a, b) > S(a-1, b+1).

Proof:

```
1. S(a, b) > S(a-1, b+1)
2. (a+1)(a+b) > (a+1-1)(a-1+b+1)
3. a(a+b)+a+b > a(a+b)
4. a+b > 0
```

Theorem:

For natural number a, b it holds that S(a, b) < S(a, b + max(1, ceil((C-S(a, b))/(a+1)))).

Proof:

```
1. S(a, b) < S(a, b+max(1, ceil((C-S(a, b))/(a+1))))
2. (a+1)(a+b) < (a+1)(a+b+max(1, ceil((C-S(a, b))/(a+1))))
3. a+b < a+b+max(1, ceil((C-S(a, b))/(a+1)))
4. 0 < max(1, ceil((C - S(a, b))/(a+1)))
5. 0 < 1
```

Algorithm:

```
input: natural number C
output: found factors, or 1 and C if C is a prime number
   1. let a = make_even( floor( sqrt( C ) ) )
```

```
2. let b = 1
3. if S(a, b) > C then a=a-1, b=b+1
```

- 4. if S(a, b) < C then b=b+max(1, ceil((C-S(a, b))/(a+1)))
- 5. if S(a, b) == C then exit: found factors (a+1) and (a+b)

```
6. if a == 0 then exit: C is a prime number
```

7. goto step 3.

Examples

Factor 51

```
1. S(6, 1) = 49 \Rightarrow b = b + max(1, ceil(2/7))

2. S(6, 2) = 56 \Rightarrow a = a - 2, b = b + 2

3. S(4, 4) = 40 \Rightarrow b = b + max(1, ceil(11/5))

4. S(4, 7) = 55 \Rightarrow a = a - 2, b = b + 2

5. S(2, 9) = 33 \Rightarrow b = b + max(1, ceil(18/3))

6. S(2, 15) = 51 \Rightarrow 51 = 3 * 17
```

Factor 23

```
1. S(2, 1) = 9 => b = b + max(1, ceil(14 / 3))
2. S(2, 6) = 24 => a = a - 2, b = b + 2
3. a = 0 => 23 = 1 * 23
```

factor 221

```
1. S(12, 1) = 169 => b = b + max(1, ceil(52 / 13))
2. S(12, 5) = 221 => 221 = 13 * 17
```

Conclusion

To my knowledge this is not based on any existing solutions. I do not claim it to be efficient or useful, I'm only concerned with its correctness and completeness. One potential benefit I see is the reduced magnitude of dividends used in divisions. Hope it inspires some ideas.