

UnitedHealthcare® Commercial and Individual Exchange *Medical Policy*

Preventive Care Services

Policy Number: MP.016.56 Effective Date: October 1, 2025

Instructions for Use

Table of Contents	Page
Application	
Coverage Rationale	
Frequently Asked Questions	4
Definitions	
Applicable Codes	5
Benefit Considerations	
Clinical Evidence	
References	
Policy History/Revision Information	
Instructions for Use	

Related Commercial/Individual Exchange Policies

- Breast Imaging for Screening and Diagnosing Cancer
- Cardiovascular Disease Risk Tests
- Consultation Services
- Cytological Examination of Breast Fluids for Cancer Screening or Diagnosis
- Gastrointestinal Colonoscopy Procedure Guidelines
- Genetic Testing for Hereditary Cancer
- Long-Acting Injectable Antiretroviral Agents for HIV (for Commercial Only)
- Long-Acting Injectable Antiretroviral Agents for HIV (for Individual Exchange Only)
- Magnetic Resonance Imaging (MRI) and Computed Tomography (CT) Scan – Site of Service
- Ocular Photoscreening
- Outpatient Surgical Procedures Site of Service
- Preventive Medicine and Screening
- Preventive Vaccines (Immunizations) (for Commercial Only)
- Preventive Vaccines (Immunizations) (for Individual Exchange Only)
- Screening Colonoscopy Procedures Site of Service

Application

UnitedHealthcare Commercial

This Medical Policy applies to UnitedHealthcare Commercial benefit plans.

UnitedHealthcare Individual Exchange

This Medical Policy applies to Individual Exchange benefit plans.

Coverage Rationale

Indications for Coverage Introduction

UnitedHealthcare covers certain medical services under the preventive care services benefit. The federal Patient Protection and Affordable Care Act (PPACA) requires non-grandfathered health plans to cover certain "recommended preventive services" as identified by PPACA under the preventive care services benefit, without cost sharing to members when provided by network providers. This includes:

• Evidence-based items or services that have in effect a rating of "A" or "B" in the current recommendations of the United States Preventive Services Task Force.

Preventive Care Services

Page 1 of 46
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Page 1 of 46
Effective 10/01/2025

- Immunizations for routine use in children, adolescents and adults that have in effect a recommendation from the *Advisory Committee on Immunization Practices of the Centers for Disease Control and Prevention*.
- With respect to infants, children and adolescents, evidence-informed preventive care and screenings provided for in the comprehensive guidelines supported by the *Health Resources and Services Administration*.
- With respect to women, such additional preventive care and screenings as provided for in comprehensive guidelines supported by the *Health Resources and Services Administration*.

Member Cost-Sharing Non-Grandfathered Plans

- Non-grandfathered plans provide coverage for preventive care services with no member cost sharing (i.e., covered at 100% of Allowed Amounts without deductible, coinsurance, or copayment) when services are obtained from a Network provider.
- Under PPACA, services obtained from an out-of-network provider are not required to be covered under a plan's preventive benefit and may be subject to member cost sharing. Refer to the member specific benefit plan document for out-of-network benefit information, if any.

Grandfathered Plans

- Plans that maintain grandfathered status under PPACA are not required by law to provide coverage for these preventive services without member cost sharing; although a grandfathered plan may choose to voluntarily amend its plan document to include these preventive benefits.
- Except where there are state mandates, a grandfathered plan might include member cost sharing, or exclude some of the preventive care services identified under PPACA.
- Refer to the member specific benefit plan document for details on how benefits are covered under a grandfathered plan.

Preventive vs. Diagnostic Services

Certain services can be done for preventive or diagnostic reasons. When a service is performed for the purpose of preventive screening and is appropriately reported, it will be considered under the preventive care services benefit. This includes services directly related to the performance of a covered preventive care service (refer to the <u>Frequently Asked Questions</u> section for additional information.)

Preventive services are those performed on a person who:

- has not had the preventive screening done before and does not have symptoms or other abnormal studies suggesting abnormalities*; or
- has had screening done within the recommended interval with the findings considered normal; or
- has had diagnostic services results that were normal after which the physician recommendation would be for future preventive screening studies using the preventive services intervals.

*In the case of a colonoscopy done as a follow-up to a positive stool-based screening (e.g., FIT, FOBT, and fecal DNA) or direct visualization screening (e.g., sigmoidoscopy or CT colonography), refer to FAQ#4 below.

When a service is done for diagnostic purposes, it will be considered under the applicable non-preventive medical benefit. Diagnostic services are done on a person who:

- had abnormalities found on previous preventive or diagnostic studies that require further diagnostic studies; or
- had abnormalities found on previous preventive or diagnostic studies that would recommend a repeat of the same studies within shortened time intervals from the recommended preventive screening time intervals; or
- had a symptom(s) that required further diagnosis; or
- does not fall within the applicable population for a recommendation or guideline.

Covered Breastfeeding Equipment

Personal-use electric breast pump:

- The purchase of a personal-use electric breast pump (HCPCS code E0603).
 - This benefit is limited to one pump per birth. In the case of a birth resulting in multiple infants, only one breast pump is covered.
 - A breast pump purchase includes the necessary supplies for the pump to operate.
- Replacement breast pump supplies necessary for the personal-use electric breast pump to operate. This includes: standard power adaptor, tubing adaptors, tubing, locking rings, bottles specific to breast pump operation, caps for bottles that are specific to the breast pump, valves, filters, and breast shield and/or splash protector for use with the breast pump.

Breastmilk storage bags (HCPCS code A4287).

Colonoscopies

Colonoscopy – Preventive Care Services Benefit (Without Member Cost-Sharing)

Member cost-sharing for colonoscopy is waived when all of the following apply:

- The patient's age is 45-75 years (ends on 76th birthday) as recommended by the USPSTF; and
- The provider is participating in the network; and
- When billed in accordance with the coding in the Colorectal Cancer Screening row listed in this policy.

Colonoscopy may require a site of service review. Refer to the Medical Policies titled <u>Screening Colonoscopy Procedures</u> <u>— Site of Service</u> and <u>Outpatient Surgical Procedures — Site of Service</u>.

Colonoscopy – Medical Benefit (With Member Cost-Sharing)

Member cost-sharing may apply when a colonoscopy is done in any one of the following scenarios:

- The patient's age is outside of the age recommendation of the USPSTF (age 45-75 years); or
- The provider is non-network; or
- Colonoscopy performed with a shortened time interval outside of the USPSTF recommendations; or
- Colonoscopy performed for diagnostic purposes; or
- Colonoscopy performed for surveillance purposes (e.g., a follow-up colonoscopy performed after identification or removal of a polyp or cancer on a previous colonoscopy); or
- Colonoscopy performed for therapeutic/treatment purposes.

The above colonoscopies may require advanced notification and/or site of service review. Refer to the <u>Gastrointestinal</u> <u>Colonoscopy Procedure Guidelines</u> and the Medical Policy titled <u>Outpatient Surgical Procedures – Site of Service</u>.

Coverage Limitations and Exclusions

- Services not covered under the preventive care benefit may be covered under another portion of the medical benefit plan.
- The coverage outlined in this policy does not address certain outpatient prescription medications, tobacco cessation drugs and/or over the counter items, as required by PPACA. These preventive benefits are administered by the member's pharmacy plan administrator. For details on coverage, refer to the member-specific pharmacy plan administrator.
- A vaccine (immunization) is not covered if it does not meet company vaccine policy requirements for FDA labeling and
 if it does not have explicit ACIP recommendations for routine use published in the Morbidity and Mortality Weekly
 Report (MMWR) of the Centers for Disease Control and Prevention (CDC) and is not listed on the applicable
 immunization schedule of ACIP. (Refer to the Preventive Care Services: Vaccine Codes.)
- Examinations, screenings, testing, or vaccines (immunizations) are not covered when:
 - o required solely for the purposes of career or employment, school or education, sports or camp, travel [including travel vaccines (immunizations)], insurance, marriage, or adoption; or
 - o related to judicial or administrative proceedings or orders; or
 - o conducted for purposes of medical research; or
 - o required to obtain or maintain a license of any type.
- Services that are investigational, experimental, unproven, or not medically necessary are not covered.
- Breastfeeding equipment and supplies not listed above. This includes, but is not limited to:
 - Manual breast pumps and all related equipment and supplies.
 - Hospital-grade breast pumps and all related equipment and supplies.
 - Equipment and supplies not listed in the <u>Covered Breastfeeding Equipment</u> section above, including but not limited to:
 - Batteries, battery-powered adaptors, and battery packs.
 - Electrical power adapters for travel.
 - Bottles which are not specific to breast pump operation. This includes the associated bottle nipples, caps, and lids.
 - Travel bags, and other similar travel or carrying accessories.
 - Breast pump cleaning supplies including soap, sprays, wipes, steam cleaning bags and other similar products.
 - Baby weight scales.
 - Garments or other products that allow hands-free pump operation.

- Breast milk storage accessories such as ice-packs, labels, labeling lids, and other similar products. The breastmilk storage accessories exclusion does not apply to breastmilk storage bags (HCPCS code A4287).
- Nursing bras, bra pads, breast shells, nipple shields, and other similar products.
- Creams, ointments, and other products that relieve breastfeeding related symptoms or conditions of the breasts or nipples.

Note: Refer to the <u>Indications for Coverage</u> section above for covered breastfeeding equipment.

Frequently Asked Questions (FAQ)

1	Q:	If a polyp is encountered during a preventive screening colonoscopy, are future colonoscopies considered under the preventive care services benefit?			
	A:	No. If a polyp is removed during a preventive screening colonoscopy, future colonoscopies would normally be considered to be diagnostic because the time intervals between future colonoscopies would be shortened.			
2	Q:	Are the related therapeutic services for a preventive colonoscopy covered under the preventive care benefit?			
	A:	Yes, related services integral to a colonoscopy are covered under the preventive care services benefit including: pre-operative examination, the associated facility, anesthesia, polyp removal (if necessary), pathologist and physician fees. However, the preventive benefit does not include a post-operative examination.			
3	Q:	Do any preventive care services require prior-authorization?			
	A:	Certain services require prior-authorization on most benefit plans. This includes, but may not be limited to: BRCA lab screening, computed tomographic colonography (virtual colonoscopy), and screening for lung cancer with low-dose computed tomography.			
4	If a member in the age range of 45-75 years has a positive stool-based colorectal cancer screening test (e.g., FIT, FOBT, and fecal DNA) or direct visualization screening test (e.g., sigmoidoscopy or CT colonography), and has a follow up colonoscopy, is the colonoscopy included in the preventive care services benefit?				
	A:	Yes, in this situation, the colonoscopy would be considered under the preventive care services benefit when billed in accordance with the coding in the Colorectal Cancer Screening row listed in this policy.			
5	Q:	need to be the primary diagnosis on the claim?			
	A:	In general, most preventive services do not require the preventive diagnosis code to be in the primary position. However, certain preventive services do require the diagnosis code to be in the primary position, which include: (1) Chemoprevention of Breast Cancer (Counseling), (2) Genetic Counseling and Evaluation for BRCA Testing, and (3) Prevention of Human Immunodeficiency Virus (HIV) Infection.			
6	Q:	If a woman has an abnormal finding on a preventive screening mammography and the follow up mammogram was found to be normal, will UnitedHealthcare cover her future mammograms under the preventive care services benefit?			
	A:	Yes, if the member was returned to normal mammography screening protocol, her future mammography screenings would be considered under the preventive care services benefit.			
7	Q:	If a member had elevated cholesterol on a prior preventive screening, are future cholesterol tests considered under the preventive care services benefit?			
	A:	Once the diagnosis has been made, further testing is considered diagnostic rather than preventive. This is true whether or not the member is receiving pharmacotherapy.			
8	Q:	Are the related services for a woman's outpatient sterilization or other contraceptive procedure covered under the preventive care benefit?			
	A:	Related services for a woman's outpatient sterilization or other contraceptive procedure are covered under the preventive care services benefit. This includes associated implantable devices, facility fee, anesthesia, and surgeon/physician fees. If a woman is admitted to an inpatient facility for another reason (e.g., maternity/delivery), and has a sterilization or other contraceptive procedure performed during that admission, the sterilization or other contraceptive procedure fees mare covered under			

		the preventive care services benefit. This includes associated sterilization/ contraception surgical fees, device fees, anesthesia, pathology, and physician fees. However, the facility fees are not covered under the preventive care benefit since the sterilization or other contraceptive procedure is incidental to, and is not the primary reason, for the inpatient admission.
9	Q:	Are blood draws/venipunctures included in the preventive care benefit?
	A:	Yes, blood draws/venipunctures are considered under the preventive benefit if billed for a covered preventive lab service that requires a blood draw.
10	Q:	Is a newly-combined vaccine (a vaccine with several individual vaccines combined into one) covered under preventive care benefits?
	A:	A new vaccine that is pending ACIP recommendations, but is a combination of previously approved individual components, may be eligible under the preventive care benefit.
11	Q:	Are preventive care services affected by other policies?
	A:	Yes, including for example, the Reimbursement Policy titled <u>Preventive Medicine and Screening</u> <u>Policy</u> describes situations which may affect reimbursement of preventive care services.
12	Q:	Are travel vaccines covered under preventive care benefits?
	A:	Benefits for preventive care services include vaccines for routine use in children, adolescents and adults that have in effect a recommendation from ACIP with respect to the individual involved. Vaccines that are specific to travel (e.g., typhoid, yellow fever, cholera, plague, and Japanese encephalitis virus) are excluded from the preventive care services benefit.
13	Q:	Does the preventive care services benefit include prescription or over the counter (OTC) items?
	A:	Refer to the plan's pharmacy benefit plan administrator for details on prescription medications and OTCs available under the plan's preventive benefit.

Definitions

The following definitions may not apply to all plans. Refer to the member specific benefit plan document for applicable definitions.

Modifier 33: Preventive service; when the primary purpose of the service is the delivery of an evidence based service in accordance with a US Preventive Services Task Force A or B rating in effect and other preventive services identified in preventive services mandates (legislative or regulatory), the service may be identified by adding 33 to the procedure. For separately reported services specifically identified as preventive, the modifier should not be used.

Note: UnitedHealthcare considers the procedures and diagnostic codes and Preventive Benefit Instructions listed in the table below in determining whether preventive care benefits apply. While Modifier 33 may be reported, it is not used in making preventive care benefit determinations.

Acronyms

Throughout this document the following acronyms are used:

- USPSTF: United States Preventive Services Task Force
- PPACA: Patient Protection and Affordable Care Act of 2010
- ACIP: Advisory Committee on Immunization Practices
- HRSA: Health Resources and Services Administration

Applicable Codes

The following list(s) of procedure and/or diagnosis codes is provided for reference purposes only and may not be all inclusive. Listing of a code in this policy does not imply that the service described by the code is a covered or non-covered health service. Benefit coverage for health services is determined by the member specific benefit plan document and applicable laws that may require coverage for a specific service. The inclusion of a code does not imply any right to reimbursement or guarantee claim payment. Other Policies and Guidelines may apply. CPT® is a registered trademark of the American Medical Association.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

rating was released,	not when	the
benefit is effective.		

Service

A date in this column is when the listed

Abdominal Aortic Aneurysm Screening

USPSTF Rating (Dec. 2019): B

The USPSTF recommends 1-time screening for abdominal aortic aneurysm (AAA) with ultrasonography in men aged 65-75 years who have ever smoked.

Code(s)

Procedure Code(s):

Ultrasound Screening Study for Abdominal Aortic Aneurysm: 76706

Diagnosis Code(s):

F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Age 65 through 75 (ends on 76th birthday).

Requires at least one of the diagnosis codes listed in this row.

Bacteriuria Screening

USPSTF Rating (Sept. 2019): A

The USPSTF recommends screening for asymptomatic bacteriuria using urine culture in pregnant persons.

Procedure Code(s):

81007, 87086, 87088

Diagnosis Code(s):

Pregnancy Diagnosis Codes

Requires a Pregnancy Diagnosis Code.

Chlamydia Infection Screening

USPSTF Rating (Sept. 2021): B

The USPSTF recommends screening for chlamydia in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection.

Notes:

- This recommendation applies to asymptomatic, sexually active adolescents and adults, including pregnant persons.
- Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.

Procedure Code(s):

Chlamydia Infection Screening: 86631, 86632, 87110, 87270, 87320, 87490, 87491, 87492, 87801, 87810

Blood Draw:

36415, 36416

Blood draw codes only apply to lab codes 86631 or 86632

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

Other: Z11.3, Z11.4, Z11.8, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51,

Z72.52, Z72.53

Chlamydia Infection Screening: Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis codes

listed in this row.

Blood Draw:

Required to be billed with 86631 or 86632 and

- One of the Screening diagnosis codes listed in this row or
- With a Pregnancy Diagnosis Code.

Gonorrhea Screening

USPSTF Rating (Sept. 2021): B

The USPSTF recommends screening for gonorrhea in all sexually active women 24 years or younger and in women 25 years or older who are at increased risk for infection.

Notes:

This recommendation applies to asymptomatic, sexually active

Procedure Code(s):

87590, 87591, 87592, 87801, 87850

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129 Requires either a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row.

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

Certain codes may not be payable in all cir	rcumstances due to other policies or				
Certain codes may not be payable in all circumstances due to other policies or guidelines.					
For preventive care medications, refer to the pharmacy plan administrator.					
adolescents and adults Oth	de(s) er: Z11.3, Z11.4, Z11.9, Z20.2, 0.6, Z29.81, Z72.51, Z72.52, 2.53	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.			
Infection Screening Pregnant Women: USPSTF Rating (July 2019): A The USPSTF recommends screening for hepatitis B virus (HBV) infection in pregnant women at their first prenatal visit. Adolescents and Adults at Increased Risk for Infection: USPSTF Rating (Dec. 2020): B The USPSTF recommends screening for hepatitis B virus (HBV) infection in persons at high risk for infection. Bright Futures (July 2022): Bright Futures recommends screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).	cedure Code(s): patitis B Virus Infection Screening: 104, 86706, 87340, 87341, 87467, 1499 od Draw: 115, 36416 gnosis Code(s): gnancy: gnancy Diagnosis Codes eening: 11t: Z00.00, Z00.01 1d: Z00.121, Z00.129 er: Z11.3, Z11.4, Z20.2, Z20.6, 1.59, Z29.81, Z57.8, Z72.51, 2.52, Z72.53	Hepatitis B Virus Infection Screening: Requires a Pregnancy Diagnosis Code or one of the Screening diagnosis codes listed in this row. Blood Draw: Requires one of the listed Hepatitis B Virus Infection Screening procedure codes listed in this row and • A Pregnancy Diagnosis Code or • One of the Screening diagnosis codes listed in this row.			

Hepatitis C Virus Infection Screening

USPSTF Rating (March 2020): B

The USPSTF recommends screening for hepatitis C virus infection in adults aged 18-79

Bright Futures (March 2021)

Bright Futures recommends screening all individuals ages 18 to 79 years at least once for hepatitis C virus infection (HCV).

Procedure Code(s):

Hepatitis C Virus Infection Screening: 86803, 86804, G0472

Blood Draw: 36415, 36416

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Hepatitis C Virus Infection Screening:

Does not have diagnosis code requirements for the preventive benefit to apply.

Blood Draw:

Requires one of the Hepatitis C Virus Infection Screening procedure codes listed in this row

HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults

Procedure Code(s):

HIV (Human Immunodeficiency Virus) Screening:

86689, 86701, 86702, 86703, 87389, 87390, 87391, 87534, 87535, 87536, No age limits.

HIV - Human Immunodeficiency Virus -Screening:

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

USPSTF Rating (June 2019): A

The USPSTF recommends that clinicians screen for HIV infection in:

- Adolescents and adults aged 15-65 years. Younger adolescents and older adults who are at increased risk of infection should also be screened.
- All pregnant persons, including those who present in labor or at delivery whose HIV status is unknown.

Note: Bright Futures recommends HIV screening lab work be conducted at least once between ages 15-21 years. Also recommended anytime between ages 11-14 years, when a risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.

Code(s)

87537, 87538, 87539, 87806, G0432, G0433, G0435, G0475, S3645

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129, Other: Z11.3, Z11.4, Z11.59, Z11.9, Z20.2, Z20.6, Z22.6, Z22.8, Z22.9,

Z29.81, Z72.51, Z72.52, Z72.53 Also see Expanded Women's Preventive Health section.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Requires a <u>Pregnancy Diagnosis Code</u> **or** one of the Screening diagnosis codes listed in this row.

Blood Draw:

Requires **both** of the following:

- One of the listed HIV Screening procedure codes listed in this row and
- One of the Screening diagnosis codes listed in this row or a Pregnancy Diagnosis Code.

RH Incompatibility Screening

USPSTF Rating (Feb. 2004): A

Rh (D) blood typing and antibody testing for all pregnant women during their first visit for pregnancy-related care.

USPSTF Rating (Feb. 2004): B

Repeated Rh (D) antibody testing for all unsensitized Rh (D)-negative women at 24-28 weeks' gestation, unless the biological father is known to be Rh (D)-negative.

Procedure Code(s):

RH Incompatibility Screening: 86850, 86901

Blood Draw: 36415, 36416

Diagnosis Code(s):

Pregnancy Diagnosis Codes

RH Incompatibility Screening:

Requires a Pregnancy Diagnosis Code.

Blood Draw:

Required to be billed with 86850 or 86901 **and** with a <u>Pregnancy Diagnosis</u> Code.

Syphilis Screening

Non-Pregnant Adolescents and Adults at Increased Risk:

USPSTF Rating (Sept. 2022): A

The USPSTF recommends screening for syphilis infection in persons who are at increased risk for infection (asymptomatic,

Procedure Code(s):

Syphilis Screening: 0064U, 0065U, 0210U, 86592, 86593, 86780

Blood Draw: 36415, 36416

Syphilis Screening:

Requires a <u>Pregnancy Diagnosis Code</u> **or** one of the Screening diagnosis code listed in this row.

Blood Draw:

Requires **both** of the following:

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

nonpregnant adolescents and adults who are at increased risk for syphilis infection).

Pregnant Women:

USPSTF Rating (Sept. 2018): A

The USPSTF recommends early screening for syphilis infection in all pregnant women.

Note: Bright Futures recommends sexually transmitted infection screening be conducted if risk assessment is positive between ages 11-21 years.

Code(s)

Diagnosis Code(s):

Pregnancy:

Pregnancy Diagnosis Codes

or

Screening:

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

Other: Z11.2, Z11.3, Z11.4, Z11.9, Z20.2, Z20.6, Z29.81, Z72.51,

Z72.52, Z72.53

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

- One of the listed Syphilis Screening procedure codes listed in this row and
- One of the Screening diagnosis codes listed in this row or a Pregnancy Diagnosis Code.

Genetic Counseling and Evaluation for BRCA Testing; and BRCA Lab Screening

USPSTF Rating (Aug. 2019): B

The USPSTF recommends that primary care clinicians assess women with a personal or family history of breast, ovarian, tubal, or peritoneal cancer or who have an ancestry associated with breast cancer susceptibility 1 and 2 (BRCA1/2) gene mutations with an appropriate brief familial risk assessment tool. Women with a positive result on the risk assessment tool should receive genetic counseling and, if indicated after counseling, genetic testing.

Refer to the Medical Policy titled Genetic Testing for Hereditary Cancer.

Genetic Counseling and Evaluation

Procedure Code(s):

Medical Genetics and Genetic Counseling Services:

96041, S0265

Evaluation and Management (Office Visits):

99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, 99385, 99386, 99387, 99395, 99396, 99397, G0463

Diagnosis Code(s):

Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43

BRCA Lab Screening Procedure Code(s):

81162, 81163, 81164, 81165, 81166, 81167, 81212, 81215, 81216, 81217

Blood Draw: 36415, 36416

Diagnosis Code(s):

Family History or Personal History of breast cancer and/or ovarian cancer: Z15.01, Z15.02, Z80.3, Z80.41, Z85.3, Z85.43

Genetic Counseling and Evaluation

*Medical Necessity plans require genetic counseling before BRCA Lab Screening.

Requires one of the Genetic Counseling and Evaluation diagnosis codes listed in this row in the primary position.

BRCA Lab Screening

*Prior authorization requirements apply to BRCA lab screening.

Applies to **age 18+** when billed with one of the BRCA Lab Screening diagnosis codes listed in this row.

Blood Draw:

Requires one of the BRCA Lab Screening procedure codes listed in this row **and** one of the BRCA Lab Screening diagnosis codes listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Screening for Pre-Diabetes and Type 2 Diabetes

USPSTF Rating (Aug. 2021): B

The USPSTF recommends screening for prediabetes and type 2 diabetes in adults aged 35 to 70 years who have overweight or obesity. Clinicians should offer or refer patients with prediabetes to effective preventive interventions.

Refer to Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors:
Behavioral Counseling Interventions for intensive behavioral counseling interventions.

For additional diabetes screening benefits, also see the *Expanded Women's Preventive Health* section for <u>Screening for Diabetes in Pregnancy</u> and <u>Screening for Diabetes After Pregnancy</u>.

Code(s)

Pre-Diabetes Preventive Interventions

Procedure Code(s):

Medical Nutrition Therapy or Counseling: 97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling: 99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0447, G0473, G9886

Diagnosis Code(s):

R73.03 (prediabetes)

Diabetes Screening

Procedure Code(s): Diabetes Screening:

82947, 82948, 82950, 82951, 82952, 83036

Blood Draw:

36415, 36416

Diagnosis Code(s):

Required Diagnosis Codes (requires at least one):

Z00.00, Z00.01, Z13.1

And one of the following additional diagnosis codes as follows:

Additional Diagnosis Codes (requires at least one):

Overweight:

E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29

Obesitv:

E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Body Mass Index 30.0 - 39.9:

Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37,

Z68.38, Z68.39

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Pre-Diabetes Preventive Interventions

Limited to age 35-70 years (ends on 71st birthday).

Requires diagnosis code R73.03.

Diabetes Screening

Limited to age 35-70 years (ends on 71st birthday).

Diabetes Screening:

Requires one of the Required Diagnosis Codes listed in this row **and** one of the listed Additional Diagnosis Codes in this row.

Blood Draw:

Requires all of the following:

- One of the listed Diabetes
 Screening procedure codes listed in this row and
- One of the listed Required Diagnosis Codes and
- One of the listed Additional Diagnosis Codes.

Preventive Benefit Does Not Apply:

If a Diabetes Diagnosis Code is present in any position, the preventive benefit does **not** apply; see the <u>Diabetes</u> <u>Diagnosis Code List</u>.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refe	For preventive care medications, refer to the pharmacy plan administrator.					
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.				
	Body Mass Index 40.0 and Over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 See the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy.					
Gestational Diabetes Screening USPSTF Rating (Aug. 2021): B The USPSTF recommends screening for gestational diabetes mellitus in asymptomatic pregnant persons at 24 weeks of gestation or after. For additional diabetes screening benefits, also see the Screening for Pre-Diabetes and Type 2 Diabetes row. Also see the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy and Screening for Diabetes After Pregnancy.	See the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy codes.	See the Expanded Women's Preventive Health section for Screening for Diabetes in Pregnancy preventive benefit instructions. Note: This benefit applies regardless of the gestational week.				
Screening Mammography	Procedure Code(s): 77063, 77067	No age limits. Does not have diagnosis code				
USPSTF Rating (2002): B The USPSTF recommends	Revenue Code: 0403	requirements for the preventive benefit to apply.				
screening mammography, with or without clinical breast examination (CBE), every 1-2 years for women aged 40 and older.	Diagnosis Code(s): Does not have diagnosis code requirements for the preventive benefit to apply.	Note : This benefit only applies to screening mammography.				
Also refer to the Medical Policy titled Breast Imaging for Screening and Diagnosing Cancer. Also see the Breast Cancer Screening for Average-Risk Women recommendation in the Expanded Women's Preventive Health section.						
Cervical Cancer	Human Papillomavirus DNA	Human Papillomavirus DNA				
Screening	Testing (HPV)	Testing (HPV)				
USPSTF Rating (Aug. 2018): A	Procedure Code(s): 0502U, 87624, 87625, 87626, G0476	Age 30 years and up.				

Preventive Care Services

United Healthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Age 21-29 years:

The USPSTF recommends screening for cervical cancer every 3 years with cervical cytology alone in women aged 21-29 years.

Age 30-65 years:

For women aged **30 to 65 years**, the USPSTF recommends:

- Screening every 3 years with cervical cytology alone,
- Every 5 years with high-risk human papillomavirus (hrHPV) testing alone, or
- Every 5 years with hrHPV testing in combination with cytology (co-testing).

Bright Futures, March 2014:

Adolescents should no longer be routinely screened for cervical dysplasia until age 21.

Also see <u>Screening for Cervical</u>
<u>Cancer</u> in the *Expanded Women's*Preventive Health section.

Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - Cholesterol Screening (Lipid Disorders Screening)

USPSTF Rating (August 2022): B

The USPSTF recommends that clinicians prescribe a statin for the primary prevention of CVD for adults aged 40 to 75 years who have 1 or more CVD risk factors (i.e. dyslipidemia, diabetes, hypertension, or smoking) and an estimated 10-year risk of a cardiovascular event of 10% or greater.

Notes:

 For statin medications benefits, refer to the pharmacy plan administrator.

Code(s)

Diagnosis Code(s):

Z00.00, Z00.01, Z01.411, Z01.419, Z11.51, Z12.4

Cervical Cytology (Pap Test) Code Group 1 Procedure Code(s):

G0101, G0123, G0124, G0141, G0143, G0144, G0145, G0147, G0148, Q0091, P3000, P3001

Code Group 1 Diagnosis Code(s):

Does not have diagnosis code requirements for preventive benefit to apply.

Code Group 2 Procedure Code(s):

88141, 88142, 88143, 88147, 88148, 88150, 88152, 88153, 88155, 88164, 88165, 88166, 88167, 88174, 88175

Code Group 2 Diagnosis Code(s):

Z00.00, Z00.01, Z01.411, Z01.419, Z12.4

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Requires one of the diagnosis codes listed in this row.

Cervical Cytology (Pap Test) Code Group 1:

Limited to age 21-65 years (ends on 66th birthday).

Does not have diagnosis code requirements for preventive benefits to apply.

Code Group 2:

Limited to age 21–65 years (ends on 66th birthday).

Requires one of the Code Group 2 diagnosis codes listed in this row.

Procedure Code(s):

Cholesterol Screening:

80061, 82465, 83718, 83719, 83721, 83722, 84478

Blood Draw: 36415, 36416

ASCVD Risk Assessment and Risk Management Services: G0537, G0538

Diagnosis Code(s):

Z00.00, Z00.01, Z13.220

Cholesterol Screening:

Ages 40-75 years (ends on 76th birthday).

Requires one of the diagnosis codes listed in this row for 80061, 82465, 83718, 83719, 83721, 83722, 84478.

Blood Draw:

Ages 40-75 years (ends on 76th birthday): Requires one of the listed Cholesterol Screening procedure codes **and** one of the Diagnosis Codes listed in this row.

ASCVD Risk Assessment and Management Services:

Ages 40-75 years (ends on 76th birthday). The diagnosis codes listed in this row are not required for G0537 and G0538.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

 See <u>Dyslipidemia Screening</u> (<u>Bright Futures</u>) for recommendations for children.

Code(s)

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Preventive Benefit Does Not Apply:

For all ages above, if any of the following lipid disorders diagnosis codes are present in any position, the preventive benefit does **not** apply: E71.30, E75.5, E78.00, E78.010, E78.011, E78.019, E78.2, E78.3, E78.41, E78.49, E78.5, E78.79, E78.81, E78.89, E88.2, E88.89

Colorectal Cancer Screening

USPSTF Rating (May 2021): B

The USPSTF recommends screening for colorectal cancer in adults aged 45 to 49 years.

USPSTF Rating (May 2021): A

The USPSTF recommends screening for colorectal cancer in all adults aged 50 to 75 years.

Also refer to the Medical Policies titled Outpatient Surgical
Procedures - Site of Service;
Screening Colonoscopy Procedures
- Site of Service; and Magnetic
Resonance Imaging (MRI) and
Computed Tomography (CT) Scan
- Site of Service.

Also see the <u>Frequently Asked</u> Questions section.

Colonoscopy Procedure Code(s):

Preventive Colonoscopy: G0105, G0121

Preventive Colonoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right): 44388*, 44389*, 44392*, 44394*, 45378*, 45380*, 45381*, 45384*, 45385*, 45388*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Note: Also see the <u>Colonoscopy Pre-</u> Op Consultation row below.

Colonoscopy

Age Limits: 45-75 years (ends on 76th birthday).

Codes G0105 and G0121 do not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to G0104, G0105, G0121, G0328 or S0285

Sigmoidoscopy

Procedure Code(s):

Preventive Sigmoidoscopy: G0104

Preventive Sigmoidoscopy When Billed with Certain Codes (see Preventive Benefit Instructions to the right):

45330*, 45331*, 45333*, 45338*, 45346*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0,

Sigmoidoscopy

Age Limits: 45-75 years (ends on 76th birthday).

Code G0104 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to codes G0104, G0105, G0121, G0328 or S0285

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79,

Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy)

Procedure Code(s):

Pathology: 88304, 88305

Anesthesia: 00811, 00812, 99152, 99153, 99156, 99157, G0500

Diagnosis Code(s):

Applies to the Pathology and Anesthesia codes listed above: Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT) Procedure Code(s):

Preventive: G0328

Preventive When Billed with Certain Codes (see Preventive Benefit Instructions to the right): 82270*, 82274*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above:

Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Fecal DNA

Procedure Code(s):

0464U, 81528

Diagnosis Code(s):

Code 81528 does not have diagnosis code requirements for preventive benefits to apply.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy)

Age Limits: 45-75 years (ends on 76th birthday).

Requires **both** of the following:

- One of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); and
- One of the procedure codes listed in the Colonoscopy row, or the Sigmoidoscopy row.

Note: Preventive benefits apply when the surgeon's claim is preventive.

Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT)

Age Limits: 45-75 years (ends on 76th birthday).

Code G0328 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when:

- Billed with one of the diagnosis codes listed in this row (Z00.00, Z00.01, Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79); or
- Billed in addition to G0104, G0105, G0121, G0328 or S0285.

Fecal DNA

Age Limits: 45-75 years (ends on 76th birthday).

Benefit is limited to once every 3 years.

Codes 0464U and 81528 do not have diagnosis code requirements for preventive benefits to apply.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Pre-Op Consultation Procedure Code(s):

Preventive:

S0285

Preventive when billed with one of the diagnosis codes listed in this row: 99202*, 99203*, 99204*, 99205*, 99211*, 99212*, 99213*, 99214*, 99215*, 99242*, 99243*, 99244*, 99245*, 99417*

Diagnosis Code(s):

Applies to Procedure Codes with asterisk(*) above: Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79

Note: For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Pre-Op Consultation

Age Limits: 45-75 years (ends on 76th birthday).

Code S0285 does not have diagnosis code requirements for preventive benefits to apply.

Codes with an asterisk(*) are preventive when billed with one of the diagnosis codes listed in this row (Z12.10, Z12.11, Z12.12, Z15.060, Z15.068, Z80.0, Z83.710, Z83.711, Z83.718, Z83.719, Z83.72, Z83.79).

Computed Tomographic Colonography (Virtual Colonoscopy)

Procedure Code(s):

74263

Diagnosis Code(s):

Does not have diagnosis code requirements for preventive benefit to apply.

Computed Tomographic Colonography (Virtual Colonoscopy)

Age Limits: 45-75 years (ends on 76th birthday).

Does not have diagnosis code requirements for preventive benefit to apply.

Prior authorization requirements may apply, depending on plan.

Wellness Examinations

(well-baby, well-child, well-adult)

USPSTF Rating: None

UnitedHealthcare supports AAP and AAFP age and frequency quidelines.

HRSA Requirements:

The Wellness Examinations codes in this row include the following HRSA requirements for women, where applicable:

Procedure Code(s):

Medicare Wellness Exams: G0402, G0438, G0439

STIs Behavioral Counseling: G0445

Annual Gynecological Exams: S0610, S0612, S0613

Pelvic Examination (add-on code): 99459

Does not have diagnosis code requirements for the preventive benefit to apply.

G0445 is limited to twice per year.

G0296 is limited to age 50 to 80 years (ends on 81st birthday).

Pelvic Examination add-on code 99459:

Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

- Breastfeeding support, counseling, and education
- Contraceptive methods and sterilizations (counseling and follow-up care)
- Screening and counseling for interpersonal domestic violence
- Screening for human immunodeficiency virus infection (HIV); education and risk assessment
- Counseling for sexually transmitted infections (STIs)
- Well-woman preventive visits
- Screening for urinary incontinence
- Obesity prevention in midlife women (counseling)

Code(s)

Preventive Medicine Services (Evaluation and Management): 99381, 99382, 99383, 99384, 99385, 99386, 99387 99391, 99392, 99393, 99394, 99395, 99396, 99397

Preventive Medicine, Individual Counseling:

99401, 99402, 99403, 99404

Preventive Medicine, Group Counseling: 99411, 99412

Newborn Care (evaluation and management):
99461

Counseling Visit (to discuss the need for Lung Cancer Screening (LDCT) using Low Dose CT Scan):

G0296

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Also see the <u>Expanded Women's</u> Preventive Health section.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's

sections above for additional instructions.

not be payable in all circumstances due to other policies or quidelines.

Newborn Screenings

All newborns

USPSTF Rating (March 2008): A

Hypothyroidism Screening: Screening for congenital hypothyroidism in newborns.

USPSTF Rating (March 2008): A

Phenylketonuria Screening: Screening for phenylketonuria (PKU) in newborns.

USPSTF Rating (Sept. 2007): A

Sickle Cell Screening: Screening for sickle cell disease in newborns.

Note: For Bright Futures hearing screening, see <u>Hearing Tests</u> (Bright Futures).

Procedure Code(s):

Hypothyroidism Screening: 84437, 84443

Phenylketonuria Screening: 84030, S3620

Sickle Cell Screening: 83020, 83021, 83030, 83033, 83051,

Blood Draw: 36415, 36416

S3850

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Newborn Screenings:

Age 0-90 days. Does not have diagnosis code requirements for the preventive benefit to apply.

Blood Draw:

Age 0-90 days, requires one of the listed Hypothyroidism Screening, Phenylketonuria Screening, or Sickle Cell Screening procedure codes.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Metabolic Screening	
benefit is effective.	
rating was released, not when the	

A date in this column is when the listed

Code(s)

Preventive Benefit Instructions

Metabolic Screening Panel:

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Age 0-90 days. Does not have diagnosis code requirements for the preventive

Panel (Newborns)

Service

Procedure Code(s):

Metabolic Screening Panel:

82017, 82136, 82261, 82775, 83020, 83498, 83516, 84030, 84437, 84443, S3620

Blood Draw: 36415, 36416

Blood Draw:

benefit to apply.

Age 0-90 days. Requires one of the listed Metabolic Screening Panel procedure codes listed in this row.

Osteoporosis Screening

USPSTF Rating (June 2018): B

Women 65 and older: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in women 65 vears and older.

Procedure Code(s): 76977, 77080, 77081, G0130

Diagnosis Code(s):

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive

> Requires one of the diagnosis codes listed in this row.

USPSTF Rating (June 2018): B

Postmenopausal women younger than 65 years at increased risk of osteoporosis: The USPSTF recommends screening for osteoporosis with bone measurement testing to prevent osteoporotic fractures in postmenopausal women younger than 65 years who are at increased risk of osteoporosis, as determined by a formal clinical risk assessment tool.

Z00.00, Z00.01, Z13.820, Z82.62

benefit to apply.

Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults

USPSTF Rating (Nov. 2018): B

The USPSTF recommends screening for unhealthy alcohol use in primary care settings in adults 18 years or older, including pregnant women, and providing persons

Procedure Code(s):

Alcohol or Drug Use Screening: 99408, 99409

Annual Alcohol Screening: G0442

Brief Counseling for Alcohol: G0443

Diagnosis Code(s):

Does not have diagnosis code requirements for preventive benefit to apply.

Does not have diagnosis code requirements for preventive benefits to apply.

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

Preventive Care Services Also see the Expanded Women's Preventive Health section. Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications, refer to the pharmacy plan administrator. Service **Preventive Benefit Instructions** A date in this column is when the listed Refer to Coverage Rationale, and FAQ's rating was released, not when the Code(s) sections above for additional instructions. benefit is effective. engaged in risky or hazardous drinking with brief behavioral counseling interventions to reduce unhealthy alcohol use. **Bright Futures (April 2017):** Bright Futures recommends alcohol or drug use assessments from age 11-21 years. Also see rows: Unhealthy Drug Use Screening (Adults); and Tobacco, Alcohol, or Drug Use Assessment (Bright Futures). **Unhealthy Drug Use Procedure Code(s):** Does not have diagnosis code requirements for preventive benefits to Screening (Adults) Alcohol or Drug Use Screening: apply. 99408, 99409 USPSTF Rating (June 2020): B **Diagnosis Code(s):** The USPSTF recommends screening by asking questions Does not have diagnosis code about unhealthy drug use in adults requirements for preventive benefit to age 18 years or older. Screening apply. should be implemented when services for accurate diagnosis, effective treatment, and appropriate care can be offered or referred. (Screening refers to asking questions about unhealthy drug use, not testing biological specimens.) **Bright Futures (April 2017):** Bright Futures recommends alcohol or drug use assessments from age 11-21 years. Also see rows: Screening and

High Blood Pressure in Adults – Screening

in Primary Care to Reduce
Unhealthy Alcohol Use in Adults;
and Tobacco, Alcohol, or Drug Use
Assessment (Bright Futures).

Behavioral Counseling Interventions

Blood Pressure Measurement in a Clinical Setting N/A

Blood Pressure Measurement in a Clinical Setting

This service is included in a preventive care wellness examination.

Page 18 of 46

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	m		

A date in this column is when the listed rating was released, not when the benefit is effective.

USPSTF Rating (April 2021): A

The USPSTF recommends screening for hypertension in adults 18 years or older with office blood pressure measurement.

The USPSTF recommends obtaining blood pressure measurements outside of the clinical setting for diagnostic confirmation before starting treatment.

Code(s)

Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting)

Procedure Code(s):

Ambulatory Blood Pressure Measurement: 93784, 93786, 93788 or 93790

Diagnosis Code(s):

Abnormal Blood-Pressure Reading Without Diagnosis of Hypertension: R03 0

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Ambulatory Blood Pressure Measurement (Outside of a Clinical Setting)

Age 18 years and older. Requires the diagnosis code listed in this row.

Breast Cancer: Medication Use to Reduce Risk

USPSTF Rating (Sept. 2019): B

The USPSTF recommends that clinicians offer to prescribe risk-reducing medications, such as tamoxifen, raloxifene, or aromatase inhibitors, to women who are at increased risk for breast cancer and at low risk for adverse medication effects.

Procedure Code(s):

Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99385, 99386, 99387, 99395, 99396, 99397,

Diagnosis Code(s):

99417, G0463

Z80.3, Z80.41, Z15.01, Z15.02

Requires one of the diagnosis codes listed in this row in the primary position.

Primary Care Interventions to Promote Breastfeeding

USPSTF Rating (Oct. 2016): B

The USPSTF recommends providing interventions during pregnancy and after birth to support breastfeeding.

N/A

Also see the Expanded Women's Preventive Health section

Included in primary care or OB/GYN office visits.

Depression in Adults (Screening)

USPSTF Rating (June 2023): B

The USPSTF recommends screening for depression in the adult population, including pregnant and postpartum persons, as well as older adults.

Bright Futures (February 2017):

Maternal Depression Screening: Routine screening for postpartum depression should be integrated

Procedure Code(s):

96127, 96161, G0136, G0444

Diagnosis Code(s):

Required for 96127 Only: Encounter for Screening for Depression: Z13.31, Z13.32 Requires one of the diagnosis codes listed in this row, for 96127.

The diagnosis codes listed in this row are **not** required, for G0136, G0444, and 96161.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Futures).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Tor preventive care medications, rele	r to the pharmacy plan administrator.	
Service A date in this column is when the listed rating was released, not when the benefit is effective. into well-child visits at 1, 2, 4, and 6	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
months of age. Also see the rows for <u>Screening for Anxiety</u> (HRSA); <u>Anxiety Disorders in Adults Screening</u> (USPSTF): Depression in Children and Adolescents (Screening) (USPSTF); <u>Perinatal Depression – Preventive Interventions</u> (<u>Counseling</u>) (USPSTF); and <u>Depression and Suicide Risk</u> <u>Screening</u> (Bright Futures).		
Depression in Children and Adolescents (Screening)	Procedure Code(s): 96127, 96161, G0136, G0444	Requires one of the diagnosis codes listed in this row, for 96127. The diagnosis codes listed in this row
USPSTF Rating (October 2022): B The USPSTF recommends screening for major depressive disorder (MDD) in adolescents aged 12-18 years.	Diagnosis Code(s): Required for 96127 Only: Encounter for Screening for Depression: Z13.31, Z13.32	are not required for G0136, G0444, and 96161.
Bright Futures (February 2017): Maternal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.		
Note : The Bright Futures Periodicity Schedule recommends depression screening begin at age 12-21 years.		
Also see the rows for Anxiety Disorders in Adults Screening (USPSTF): Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright		

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

64 or younger.

Service

ımn is when the listed rating was released, not when the benefit is effective.

Code(s)

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Preventive Benefit Instructions

Anxiety Disorders in Adults (Screening)

USPSTF Rating (June 2023): B The USPSTF recommends screening for anxiety in adults, including pregnant and postpartum persons. This applies to adults age

Also see the rows for Screening for Anxiety (HRSA); and Screening for Anxiety in Children and Adolescents (USPSTF).

Procedure Code(s):

Diagnosis Code(s):

Encounter for Screening Examination for Other Mental Health and Behavioral Disorders: Z13.39

Requires the diagnosis code listed in this row.

Screening for Anxiety in Children and Adolescents

USPSTF Rating (October 2022): B

The USPSTF recommends screening for anxiety in children and adolescents aged 8 to 18 years.

Also see the rows for Anxiety Disorders in Adults Screening (USPSTF): Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression - Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).

Procedure Code(s):

96127

Diagnosis Code(s):

Encounter for Screening Examination for Other Mental Health and Behavioral Disorders: 713 39

Requires the diagnosis code listed in this row.

Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors: Behavioral **Counseling Interventions**

USPSTF Rating (Nov. 2020): B

The USPSTF recommends offering or referring adults with cardiovascular disease risk factors to behavioral counseling interventions to promote a healthy diet and physical activity.

Procedure Code(s):

Medical Nutrition Therapy or Counseling:

97802, 97803, 97804, G0270, G0271

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886

ASCVD Risk Assessment and Risk Management Services: G0537, G0538

Diagnosis Code(s):

Screening: Z13.220

Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886

The diagnosis codes listed in this row are **not** required for G0446. G0447. G0473, G0537, and G0538.

G0446 is limited to once per year.

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

M		

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Nicotine Dependence, Tobacco Use, or Family History of IHD: F17.210, F17.211, F17.213, F17.218, F17.219, Z72.0, Z87.891, Z82.49

Overweight:

E66.3, Z68.25, Z68.26, Z68.27, Z68.28, Z68.29

Body Mass Index 30.0 – 39.9: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39

Body Mass Index 40.0 and Over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Impaired Fasting Glucose: R73.01

Metabolic Syndrome; Insulin Resistance Syndrome Type A; Other Insulin Resistance: E88.810, E88.811, E88.818, E88.819

Hyperlipidemia / Dyslipidemia: E78.00, E78.010, E78.011, E78.019, E78.1, E78.2, E78.3, E78.41, E78.49, E78.5

Obesity:

E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82, Z68.41, Z68.42, Z68.43, Z68.44, Z68.45

Essential Hypertension:

110

Resistant Hypertension:

I1A.0

Secondary Hypertension: 115.0, 115.1, 115.2, 115.8, 115.9, N26.2

Hypertension Complicating Pregnancy, Childbirth, and the Puerperium: O10.011, O10.012, O10.013,

O10.019, O10.02, O10.03, O10.111, O10.112, O10.113, O10.119, O10.12,

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

	m		

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

O10.13, O10.211, O10.212, O10.213, O10.219, O10.22, O10.23, O10.311, O10.312, O10.313, O10.319, O10.32, O10.33, O10.411, O10.412, O10.413, O10.419, O10.42, O10.43, O10.911, O10.912, O10.913, O10.919, O10.92, O10.93, O11.1, O11.2, O11.3, O11.4, O11.5, O11.9, O13.1, O13.2, O13.3, O13.4, O13.5, O13.9, O16.1, O16.2, O16.3, O16.4, O16.5, O16.9

Urgent/Emergency/Crisis Hypertension: 116.0, 116.1, 116.9

Diabetes:

Diabetes Diagnosis Code List

Atherosclerosis:

Atherosclerosis Diagnosis Code List

Coronary Atherosclerosis: 125.10, 125.110, 125.111, 125.112, 125.118, 125.119, 125.700, 125.701, 125.702, 125.708, 125.709, 125.710, 125.711, 125.712, 125.718, 125.719, 125.720, 125.721, 125.722, 125.728, 125.729, 125.730, 125.731, 125.732, 125.738, 125.739, 125.750, 125.751, 125.752, 125.758, 125.759, 125.760, 125.761, 125.762, 125.762, 125.768, 125.799, 125.790, 125.791, 125.792, 125.798,

Weight Loss to Prevent Obesity-Related Morbidity and Mortality in Adults: Behavioral Interventions

USPSTF Rating (Sept. 2018): B

The USPSTF recommends that clinicians offer or refer adults with a body mass index (BMI) of 30 or higher (calculated as weight in kilograms divided by height in meters squared) to intensive multicomponent behavioral interventions.

Procedure Code(s):

Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271

125.799, 125.810, 125.811, 125.812

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886

Diagnosis Code(s):

Body Mass Index 30.0-39.9: Z68.30, Z68.31, Z68.32, Z68.33, Z68.34, Z68.35, Z68.36, Z68.37, Z68.38, Z68.39

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886

G0446 is limited to once per year. The diagnosis codes listed in this row are **not** required for G0446, G0447, and G0473.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refe	r to the pharmacy plan administrator.	
Service A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
	Body Mass Index 40.0 and over: Z68.41, Z68.42, Z68.43, Z68.44, Z68.45 Obesity: E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9, E88.82	
High Body Mass Index in Children and Adolescents USPSTF Rating (June 2024): B The USPSTF recommends that clinicians provide or refer children and adolescents 6 years or older with a high body mass index (BMI) (≥ 95 th percentile for age and sex) to comprehensive, intensive behavioral interventions. See the Practice Considerations section of the published USPSTF recommendation or more information about behavioral interventions.	Procedure Code(s): Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271 Preventive Medicine Individual Counseling: 99401, 99402, 99403, 99404 Behavioral Counseling or Therapy: 0403T, G0446, G0447, G0473, G9886 Also see the codes in the Wellness Examinations row above. Diagnosis Code(s): Obesity: E66.01, E66.09, E66.1, E66.2, E66.811, E66.812, E66.813, E66.89, E66.9. E88.82 Pediatric BMI of 120% or more of the 95th percentile for age: Z68.54, Z68.55, Z68.56	Requires one of the diagnosis codes listed in this row for 0403T, 97802-97804, 99401-99404, G0270, G0271, and G9886 G0446 is limited to once per year. The diagnosis codes listed in this row are not required for G0446, G0447, and G0473.
Healthy Weight and Weight Gain During Pregnancy: Behavioral Counseling Interventions USPSTF Rating (May 2021): B The USPSTF recommends that clinicians offer pregnant persons effective behavioral counseling interventions aimed at promoting healthy weight gain and preventing excess gestational weight gain in pregnancy.	Procedure Code(s): Medical Nutrition Therapy: 97802, 97803, 97804, G0270, G0271 Preventive Medicine Individual Counseling: 99401, 99402, 99403, 99404 Behavioral Counseling or Therapy: G0447, G0473 Diagnosis Code(s): Pregnancy Diagnosis Codes	Requires one of the diagnosis codes listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Behavioral Counseling to Prevent Sexually Transmitted Infections

USPSTF Rating (Aug. 2020): B

The USPSTF recommends behavioral counseling for all sexually active adolescents and for adults who are at increased risk for sexually transmitted infections (STIs).

Interventions for Tobacco Smoking Cessation in Adults, including Pregnant Persons

USPSTF Rating (Jan. 2021): A Pregnant Persons (A): The USPSTF recommends that clinicians ask all pregnant persons about tobacco use, advise them to stop using tobacco, and provide behavioral interventions for cessation to pregnant persons who use tobacco.

Nonpregnant Adults (A): The USPSTF recommends that clinicians ask all adults about tobacco use, advise them to stop using tobacco, and provide behavioral interventions and US Food and Drug Administration (FDA)—approved pharmacotherapy for cessation to nonpregnant adults who use tobacco.

Note: Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.

Also see rows: <u>Unhealthy Drug Use</u>
<u>Screening (Adults)</u>; and <u>Tobacco</u>,
<u>Alcohol</u>, or <u>Drug Use Assessment</u>
(Bright Futures).

Code(s)

Procedure Code(s):

STIs Behavioral Counseling: G0445

Preventive Medicine Individual Counseling 99401, 99402, 99403, 99404

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Procedure Code(s):

Behavioral Interventions: 99406, 99407

Preventive Medicine, Individual Counseling: 99401, 99402, 99403, 99404

Also see the codes in the Wellness Examinations row above.

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Does not have diagnosis code requirements for the preventive benefit to apply.

G0445 is limited to twice per year.

Does not have diagnosis code requirements for the preventive benefit to apply.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Primary Care Interventions To Prevent Tobacco Use In Children And Adolescents

USPSTF Rating (April 2013): B

The USPSTF recommends that primary care clinicians provide interventions, including education or brief counseling, to prevent initiation of tobacco use among school-aged children and adolescents.

Bright Futures (April 2017):

Bright Futures recommends tobacco use assessments from age 11-21 years.

Also see rows: <u>Unhealthy Drug Use</u>
<u>Screening (Adults)</u>; and <u>Tobacco</u>,
<u>Alcohol</u>, or <u>Drug Use Assessment</u>
(Bright Futures).

Code(s)

Procedure Code(s):

Smoking and Tobacco Use Cessation Counseling Visit: 99406, 99407

Preventive Medicine, Individual Counseling: 99401, 99402, 99403, 99404
Also see the codes in the Wellness Examinations row above.

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Does not have diagnosis code requirements for the preventive benefit to apply.

Screening for Visual Impairment in Children

USPSTF Rating (Sept. 2017): B

The USPSTF recommends vision screening at least once in all children age 3 to 5 years to detect amblyopia or its risk factors.

Bright Futures:

Visual acuity screening is recommended for age 4 and 5 years as well as in cooperative 3 year olds.

Instrument-based screening recommended for age 12 and 24 months, in addition to the well visits at 3-5 years of age.

Behavioral Counseling to Prevent Skin Cancer

USPSTF Rating (March 2018): B

The USPSTF recommends counseling young adults, adolescents, children, and parents of young children about minimizing exposure to ultraviolet (UV) radiation for persons ages 6 months

Procedure Code(s):

Visual Acuity Screening (e.g., Snellen chart):

99173

Instrument-Based Screening: 99174, 99177

Diagnosis Code(s):

See the Preventive Benefit Instructions.

Visual Acuity Screening (99173):

Up to age 21 years (ends on 22nd birthday). Does not have diagnosis code requirements for preventive benefits to apply.

Instrument-Based Screening (99174 and 99177):

- Age 1 to 5 (ends on 6th birthday):
 Does not have diagnosis code requirements for preventive benefits to apply.
- Age 6 to 21 years (ends on 22nd birthday): Refer to the Medical Policy titled <u>Ocular Photoscreening</u> for allowable diagnoses.

N/A

This service is included in a preventive care wellness examination or focused E&M visit.

Preventive Care Services
UnitedHealthcare Commercial and Individual Exchange Medical Policy

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

For preventive care	medications, refe	r to the pharmacy plan administrator.	
Service A date in this column rating was released, r benefit is effective.		Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
to 24 years with fair reduce their risk of			
Falls Preventi Community-D Older Adults		N/A	This service is included in a preventive care wellness examination or focused E&M visit.
USPSTF Rating (January The USPSTF recommendation interventions to prevention to prevention or older who are at for falls.	nmends exercise vent falls in adults 65 years		
Screening for Partner Violen		N/A	This service is included in a preventive care wellness examination.
USPSTF Rating (O The USPSTF reconclinicians screen for violence in women of age and provide or who screen positive support services.	nmends that intimate partner of reproductive refer women		
Also see <u>Screening</u> <u>for Interpersonal an</u> <u>Violence</u> in the <i>Expo</i> <u>Preventive Health</u> s	d Domestic anded Women's		
Screening for Cancer with L		Procedure Code(s): 71271	Requires one of the diagnosis codes listed in this row.
Computed Tor		Diagnosis Code(s):	Limitations:
USPSTF Rating (M	nmends annual	F17.210, F17.211, F17.213, F17.218, F17.219, Z87.891	 Limited to one per year, and All of the following criteria: Age 50 to 80 years (ends on
dose computed tom	screening for lung cancer with low- dose computed tomography (LDCT) in adults aged 50 to 80 years who	Codes for Reporting Purposes: G9275, G9276	81 st birthday), and O At least 20 pack-years* of
have a 20 pack-yea history and currently quit within the past	y smoke or have	Note : Codes G9275 and G9276, are for reporting purposes only, if applicable. These codes are not	smoking history, and Either a current smoker or has quit within the past 15 years
Screening should be once a person has a 15 years or develop	not smoked for	separately reimbursable.	Note : Prior authorization requirements may apply, depending on plan.
problem that substate expectancy or the a willingness to have surgery.	ntially limits life bility or		*A pack-year is a way to measure the amount a person has smoked over a long period of time. It is calculated by multiplying the number of packs of cigarettes smoked per day by the number of years the person has

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy number of years the person has smoked. For example, 1 pack year is equal to smoking 1 pack per day for 1

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

A date in this column is when the listed
rating was released, not when the
benefit is effective.

Service

Code(s)

Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's

sections above for additional instructions.
year, or 2 packs per day for half a year,
and so on. Source: National Institutes of
Health, National Cancer Institute
Dictionary of Cancer Terms, pack year
definition web page.

https://www.cancer.gov/publications/dictionaries/cancer-terms/def/pack-year

Fluoride Application in Primary Care

USPSTF Rating (May 2014): B

Children From Birth Through Age 5 Years. The USPSTF recommends that primary care clinicians apply fluoride varnish to the primary teeth of all infants and children starting at the age of primary tooth eruption.

Bright Futures (July 2022):

Bright Futures adopted the May 2014 recommendation of the USPSTF and further recommends, once teeth are present, apply fluoride varnish to all children every 3 to 6 months in the primary care or dental office, based on caries risk.

Procedure Code(s):

Application of Topical Fluoride by Physician or Other Qualified Health Care Professional: 99188

Diagnosis Code(s):

Does not have diagnosis code requirements for the preventive benefit to apply.

Age 0-5years (ends on 6th birthday).

Does not have diagnosis code requirements for the preventive benefit to apply.

Latent Tuberculosis Infection in Adults: Screening

USPSTF Rating (May 2023): B

The USPSTF recommends screening for latent tuberculosis infection (LTBI) in populations at increased risk. This recommendation applies to asymptomatic adults 18 years or older at increased risk for tuberculosis (TB).

Procedure Code(s):

Screening: 86480, 86481, 86580

Follow-Up Visit to Check Results: 99211

Blood Draw: 36415, 36416

Diagnosis Code(s):

R76.11, R76.12, Z00.00, Z00.01, Z11.1, Z11.7, Z20.1

Note for age 18-21 years (ends on 22nd birthday): In addition to the codes in this row, the preventive benefit also applies to the diagnosis codes listed in the Bright Futures row: Tuberculosis (TB) Testing.

Screening:

Ages 18 years and up.

Requires one of the diagnosis codes listed in this row for CPT code 86480, 86481, and 86580.

Follow-Up Visit to Check Results (99211):

CPT code 99211 requires diagnosis code R76.11 or R76.12.

Blood Draw:

Ages 18 years and up.

Required to be billed with 86480 or 86481 **and** one of the diagnosis codes listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

Hypertensive Disorders of Pregnancy – Screening

USPSTF Rating (Sept. 2023): B

The USPSTF recommends screening for hypertensive disorders in pregnant persons with blood pressure measurements throughout pregnancy.

Perinatal Depression – Preventive Interventions (Counseling)

USPSTF Rating (Feb. 2019): B

The USPSTF recommends that clinicians provide or refer pregnant and postpartum persons who are at increased risk of perinatal depression to counseling interventions.

Note: This policy addresses coding for interventions from a network medical provider only. For perinatal depression preventive interventions with a mental health provider, refer to the plan's mental health plan benefit administrator.

Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); and Depression Screening (Bright Futures).

Prevention of Human Immunodeficiency Virus (HIV) Infection: Preexposure Prophylaxis

USPSTF Rating (Aug. 2023): A

The USPSTF recommends that clinicians prescribe preexposure prophylaxis using effective antiretroviral therapy to persons

Code(s)

Hypertensive disorders of pregnancy screening by blood pressure measurement is included in the code for a prenatal care office visit. See the following code groups in the Expanded Women's Preventive Health section:

- Prenatal Office Visits
- Prenatal Care Visits
- Global Obstetrical Codes

Code Group 1 Procedure Code(s):

Preventive Medicine Individual Counseling:

99401, 99402, 99403, 99404

Preventive Medicine, Group Counseling: 99411, 99412

Prenatal Care Visits: 59425, 59426

Preventive Medicine Services (Evaluation and Management): 99381, 99382, 99383, 99384, 99385, 99386, 99387, 99391, 99392, 99393, 99394, 99395, 99396, 99397

Code Group 2 Procedure Code(s):

Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, G0463

Code Group 2 Diagnosis Code(s):

A <u>Pregnancy Diagnosis Code</u>; **or** Z39.2 (encounter for routine postpartum follow-up); **or** Z13.32 (encounter for screening for maternal depression)

Procedure Code(s):

Kidney Function Testing (Creatinine): 82565, 82575

Pregnancy Testing: 81025, 84702, 84703

Office Visits:

99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417,

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

See the following code groups in the Expanded Women's Preventive Health section:

- Prenatal Office Visits
- Prenatal Care Visits
- Global Obstetrical Codes

Code Group 1:

Does not have diagnosis code requirements for the preventive benefit to apply.

Code Group 2:

Requires one of the Code Group 2 diagnosis codes listed in this row.

Requires one of the diagnosis codes listed in this row in the primary position.

Note: Prior authorization requirements may apply, depending on plan. Refer to the Medical Benefit Drug Policy titled Long-Acting Injectable Antiretroviral Agents for HIV (for Commercial Only) or Long-Acting Injectable Antiretroviral

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

who are at increased risk of HIV acquisition to decrease the risk of acquiring HIV.

Note: This benefit also includes:

- Kidney function testing (creatinine)
- Serologic testing for hepatitis B and C virus
- Testing for other STIs
- Pregnancy testing when appropriate
- Ongoing follow-up and monitoring including HIV testing every 3 months

Refer to the plan's pharmacy benefit plan administrator for details on prescription medications available under the plan's preventive benefit.

Code(s)

G0463

(also see codes in the Wellness Examinations section)

Antiretroviral Therapy Injection: 96372 (Administration) J0739 (Injection cabotegravir, 1mg) G0012 (Administration)

Counseling for PrEP to prevent HIV: G0011, G0013

Pharmacy Supplying Fee for HIV PrEP: Q0521

Diagnosis Code(s):

Z11.3, Z11.4, Z20.2, Z20.6, Z29.81, Z72.51, Z72.52, Z72.53

Also see the sections for:

- Behavioral Counseling to Prevent Sexually Transmitted Infections
- Chlamydia Infection Screening
- Gonorrhea Screening
- Hepatitis B Virus Infection Screening
- Hepatitis C Virus Infection Screening
- HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults
- Syphilis Screening

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Agents for HIV (for Individual Exchange Only).

Bright Futures

Anemia Screening in Children

(Bright Futures)

Procedure Code(s):

Anemia Screening in Children: 85014, 85018

Blood Draw: 36415, 36416

Diagnosis Code(s):

Z00.110, Z00.111, Z00.121, Z00.129, Z13.0

Anemia Screening in Children: Ages prenatal to 21 (ends on 22nd birthday). No frequency limit.

Requires one of the diagnosis codes listed in this row

Blood Draw:

Ages prenatal to 21 (ends on 22nd birthday).

Required to be billed with 85014 or 85018 **and** one of the diagnosis codes listed in this row.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

÷	Δ	r۱	/1	^	Δ

A date in this column is when the listed rating was released, not when the benefit is effective.

Code(s)

Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's

Hearing Tests

Bright Futures (April 2017):

Hearing Tests: Recommended at ages: Newborn; between 3-5 days to 2 months; 4 years; 5 years, 6 years; 8 years; 10 years; once between age 11-14 years; once between age 15-17 years; once between age 18-21 years; also recommended for those that have a positive risk assessment.

Risk Assessment: Recommended at ages: 4 mo, 6 mo, 9 mo, 12 mo, 15 mo, 18 mo, 24 mo, 30 mo, 3 years, 7 years, and 9 years.

Procedure Code(s):

Hearing Tests: 92551, 92552, 92553, 92558, 92587, 92588, 92650, 92651, V5008

Diagnosis Code(s):

Examination of Hearing: Z01.10 Routine Child: Z00.121, Z00.129 General Exam (for 18-21 years): Z00.00, Z00.01

Note: A risk assessment is included in the code for a wellness examination visit: see the codes in the Wellness Examinations row above.

sections above for additional instructions.

Ages 0-90 days: Does not have diagnosis code requirements for the preventive benefit to apply.

Ages 91 days to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row. Limit of once per year.

Screening for Visual Impairment in Children

(Bright Futures)

See row above for Screening for Visual Impairment in Children.

See row above Screening for Visual Impairment in Children.

Formal Developmental/ Autism Screening

Bright Futures:

- A formal, standardized developmental screen is recommended during the 9 month visit.
- A formal, standardized developmental screen is recommended during the 18 month visit, including a formal autism screen.
- A formal, standardized autism screen is recommended during the 24 month visit.
- A formal, standardized developmental screen is recommended during the 30 month visit.

Procedure Code(s):

96110

Diagnosis Code(s):

Z00.121, Z00.129, Z13.40, Z13.41, Z13.42, Z13.49

Ages prenatal to 2 years (ends on 3rd birthday).

No frequency limit.

Requires one of the diagnosis codes listed in this row.

Lead Screening

Bright Futures:

Screening Lab Work: Conduct risk assessment or screening, as appropriate, at the following intervals: 12 mo and 24 mo.

Risk Assessment, and Screening if positive: Recommended at 6 mo, 9

Procedure Code(s):

Lead Screening: 83655

Blood Draw: 36415, 36416

Diagnosis Code(s):

Z00.121, Z00.129, Z77.011

Lead Screening:

Ages 6 months through age 6 years (ends on 7th birthday). No frequency limit.

Requires one of the diagnosis codes listed in this row.

Blood Draw:

Ages 6 months through age 6 years (ends on 7th birthday).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

•	^	BAN	,i	_	^
7	e	r١	,,		н

A date in this column is when the listed rating was released, not when the benefit is effective.

mo. 12 mo. 18 mo. 24 mo. 3 years. 4 years, 5 years, and 6 years.

Tuberculosis (TB) **Testing**

Bright Futures

For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults.

Procedure Code(s):

Screening: 86580

Code(s)

Follow-Up Visit to Check Results: 99211

Diagnosis Code(s):

R76.11, R76.12, Z20.1, Z00.121, Z00.129, Z11.1, Z11.7

Note for age 18 years and older: In addition to these codes, the preventive benefit also applies to all codes listed in the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions. Required to be billed with 83655 and one of the diagnosis codes in this row.

Ages prenatal to 21 (ends on 22nd birthday).

Note: For age 18 years and older, also refer to the USPSTF recommendation above for Latent Tuberculosis Infection: Screening, Adults

No frequency limit.

CPT code 86580 requires one of the diagnosis codes listed in this row.

CPT code 99211 requires diagnosis code R76.11, R76.12, or Z11.1.

Dyslipidemia Screening

Bright Futures (April 2014):

Risk Assessment: Recommended at 24 mo, 4 years, 6 years, 8 years, 12 years, 13 years, 14 years, 15 years, 16 years.

Screening Lab Work: Conduct if risk assessment is positive, or, at the following intervals: once between age 9-11 years; once between age 17-21 years

Procedure Code(s):

Dyslipidemia Screening Lab Work: 80061, 82465, 83718, 83719, 83721, 83722, 84478

Blood Draw: 36415, 36416

Diagnosis Code(s):

Z00.121, Z00.129, Z13.220

Note: A risk assessment is included in the code for a wellness examination visit: see the Wellness Examinations row above.

Dyslipidemia Screening Lab Work: Ages 24 months to 21 years (ends on 22nd birthday). Requires one of the diagnosis codes listed in this row.

Blood Draw:

Ages 24 months to 21 years (ends on 22nd birthday).

Requires one of the listed Dyslipidemia Screening procedure codes listed in this row and one of the diagnosis codes listed in this row.

Tobacco, Alcohol, or Drug Use Assessment

Bright Futures (April 2017):

Bright Futures recommends tobacco, alcohol, or drug use assessment from age 11-21 years. See codes in the rows above:

- Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents
- Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults
- Unhealthy Drug Use Screening (Adults)

An assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.

See the rows above:

- Primary Care Interventions To Prevent Tobacco Use in Children and Adolescents
- Screening and Behavioral Counseling Interventions in Primary Care to Reduce Unhealthy Alcohol Use in Adults
- Unhealthy Drug Use Screening (Adults)

See the Wellness Examinations row above.

Behavioral/Social/ **Emotional Screening**

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

Preventive Care Services				
Also see the Expanded Women's Preventive Health section.				
	Certain codes may not be payable in all circumstances due to other policies or guidelines. For preventive care medications, refer to the pharmacy plan administrator.			
Service	to the pharmacy plan administrator.			
A date in this column is when the listed		Preventive Benefit Instructions		
rating was released, not when the	Codo(a)	Refer to Coverage Rationale, and FAQ's		
benefit is effective.	Code(s)	sections above for additional instructions.		
Bright Futures (July 2022): Bright Futures recommends behavioral/social/emotional screening annually from newborn to 21 years.				
Also see the rows for Screening for Anxiety (HRSA); Screening for Depression in Adults (USPSTF); Perinatal Depression – Preventive Interventions (Counseling) (USPSTF); and Depression and Suicide Risk Screening (Bright Futures).				
Depression and Suicide Risk Screening	See the codes in the <u>Depression in Children and Adolescents</u> (Screening) row above.	See the <u>Depression in Children and</u> <u>Adolescents (Screening)</u> row above.		
Bright Futures (July 2022): Bright Futures recommends screening adolescents age 12-21 years for depression and suicide risk, making every effort to preserve confidentiality of the adolescent.				
Bright Futures (February 2017): Matemal Depression Screening: Routine screening for postpartum depression should be integrated into well-child visits at 1, 2, 4, and 6 months of age.				
Also see the rows for Screening for Anxiety (HRSA); Depression in Children and Adolescents (Screening) (USPSTF); and Perinatal Depression – Preventive Interventions (Counseling).				
Sexually Transmitted	See the codes in the Chlamydia	See the Chlamydia Infection Screening		
Infections (STI)	Infection Screening and Gonorrhea Screening rows above.	and Gonorrhea Screening rows above.		
Bright Futures (April 2017):	Ocidentify tows above.			
Bright Futures recommends the				
following:				
STI Risk Assessment: Conduct risk assessment at each of the				
recommended visits between 11				
years – 21 years.				

STI Lab Work: Conduct if risk assessment is positive.

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service

A date in this column is when the listed rating was released, not when the benefit is effective.

HIV Screening

Bright Futures (April 2023):

HIV Risk Assessment: Conduct risk assessment at age 11 years, 12 years, 13 years, 14 years, 19 years, 20 years, and 21 years.

HIV Screening Lab Work: Conduct at least once between age 15-21 years. Also recommended anytime between ages 11-14 years, when a risk assessment is positive. And after initial screening, youth at increased risk of HIV infection should be retested annually or more frequently if at high risk.

Code(s)

See the codes in the <u>HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults</u> row above.

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row above.

Sudden Cardiac Arrest (SCA) and Sudden Cardiac Death (SCD) – Risk Assessment and ECG Screening

Bright Futures (July 2022):

All children should be evaluated for conditions predisposing to SCA and SCD in the course of routine health care. A thorough and detailed history, family history, and physical examination are necessary to begin assessing SCA and SCD risk. The ECG should be the first test ordered when there is a concern for SCA risk. The ECG should be interpreted by a physician trained in recognizing electrical heart disease (i.e., a pediatric cardiologist or pediatric electrophysiologist).

ECG Screening for those at Risk

Procedure Code(s):

93000, 93005, 93010

Diagnosis Code(s):

Required Screening Diagnosis Codes (requires at least one):

Adult: Z00.00, Z00.01 Child: Z00.121, Z00.129

And requires one of the following Additional Diagnosis Codes (requires at least one):

I42.0, I42.1, I42.2, I45.81, I49.8, I49.9, R55, R06.00, R06.09, R53.83, R00.2, R01.0, R01.1, R03.0, Q87.40, Q87.410, Q87.418, Q87.42, Q87.43, Q87.85, Q93.52, Z82.41, Z84.81, Z82.49

Risk Assessment

A risk assessment is included in the code for a wellness examination visit; see the codes in the Wellness Examinations row above.

ECG Screening for those at Risk:
Limited to ages 11 years to 21 years
(ends on 22nd birthday). Requires one of
the Screening Diagnosis Codes listed in
this row and one of the Additional
Diagnosis Codes listed in this row.

screening between the ages 0-21 years (perform risk assessment for hepatitis B virus (HBV) infection).

Also see the Expanded Women's Preventive Health section.

Certain codes may not be payable in all circumstances due to other policies or guidelines.

For preventive care medications, refer to the pharmacy plan administrator.

Service		
A date in this column is when the listed rating was released, not when the benefit is effective.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
Hepatitis B Virus Infection Screening	See the codes in the <u>Hepatitis B</u> <u>Virus Infection Screening</u> row above.	See the <u>Hepatitis B Virus Infection</u> <u>Screening</u> row above.
Bright Futures (July 2022): Bright Futures recommends		

Expanded Women's Preventive Health

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
Well-Woman Preventive Visits HRSA Requirement (Dec. 2021): WPSI Recommends that women receive at least one preventive	Procedure Code(s): Well-Woman Visits: See the Wellness Examinations row in the Preventive Care Services section.	Well-Woman Visits: See the Wellness Examinations row in the Preventive Care Services section.
care visit per year beginning in adolescence and continuing across the lifespan to ensure the provision of all recommended preventive services, including preconception and many services	Prenatal Office Visits: Evaluation and Management (Office Visits): 99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463	Prenatal Office Visits: Requires a Pregnancy Diagnosis Code.
necessary for prenatal and interconception care, are obtained. The primary purpose of these visits should be the delivery and coordination of recommended preventive services as determined by age and risk factors. These services may be completed at a single or as part of a series of visits that take place over time to obtain all necessary services depending on a woman's age, health status, reproductive health needs, pregnancy status, and risk factors. Well-women visits also include pre-pregnancy, prenatal, postpartum and interpregnancy	Pelvic Examination (add-on code): 99459	Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.
	Physician Prenatal Education, Group Setting: 99078	Physician Prenatal Education, Group Setting: Requires a Pregnancy Diagnosis Code.
	Prenatal Care (Antepartum) Visits: 59425, 59426	Prenatal Care (Antepartum) Visits: Does not have diagnosis code requirements for the preventive benefit to apply.
visits. Also see Wellness Examinations and other USPSTF	Global Obstetrical Codes: 59400, 59510, 59610, 59618	Global Obstetrical Codes: The routine, low-risk, prenatal visits portion of the code is covered as preventive.

Preventive Care Services

Page 35 of 46 Effective 10/01/2025

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or quidelines.

Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service A date in this column reflects when the listed rating was issued.	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
recommendations during pregnancy in the <i>Preventive Care Services</i> section.		Does not have diagnosis code requirements for the preventive benefit to apply.	
	Postpartum Care Visits (outpatient): 59430	Postpartum Care Visits (outpatient): Does not have diagnosis code requirements for the preventive benefit to apply.	
	Diagnosis Code(s): Pregnancy Diagnosis Codes	See above services that require a pregnancy diagnosis code.	
Screening for Diabetes in Pregnancy HRSA Requirement (Jan. 2023): Recommends screening pregnant	Procedure Code(s): Diabetes Screening: 82947, 82948, 82950, 82951, 82952, 83036	Diabetes Screening: Requires a Pregnancy Diagnosis Code (regardless of gestational week). Blood Draw:	
women for gestational diabetes mellitus after 24 weeks of gestation (preferably between 24 and 28 weeks of gestation) to prevent adverse birth outcomes, WPSI	Blood Draw: 36415, 36416 Diagnosis Code(s): Pregnancy Diagnosis Codes	Requires one of the diabetes screening procedure codes listed in this row and one of the <u>Pregnancy Diagnosis Codes</u> . Note : If a diabetes diagnosis code is	
recommends screening pregnant women with risk factors for type 2 diabetes or GDM before 24 weeks of gestation – ideally at the first prenatal visit.		present in any position, the preventive benefit will not be applied. See the <u>Diabetes Diagnosis Code List</u> .	
Also see the Screening for Pre- Diabetes and Type 2 Diabetes and Gestational Diabetes Screening sections of the Preventive Care Services section, and the Screening for Diabetes After Pregnancy section.			
Screening for Diabetes	Procedure Code(s):	Diabetes Screening:	
After Pregnancy HRSA Requirement (Jan. 2023): Recommends screening for type 2	Diabetes Screening: 82947, 82948, 82950, 82951, 82952, 83036	Requires one of the Required Screening diagnosis codes listed in this row and Z86.32.	
diabetes in women with a history of gestational diabetes mellitus	Blood Draw: 36415, 36416	No age limit.	
(GDM) who are not currently pregnant and who have not previously been diagnosed with type 2 diabetes. Initial testing should ideally occur within the first year postpartum and can be	Diagnosis Code(s): Required Screening Diagnosis Codes (requires at least one): Z00.00, Z00.01, Z13.1	Blood Draw: Requires one of the Diabetes Screening procedure codes listed in this row and one of the Required Screening diagnosis codes listed in this row and Z86.32.	
conducted as early as 4-6 weeks postpartum. Women who were not screened in the first year postpartum or those with a	And requires the following additional code: Additional Diagnosis Code Required: Z86.32 (personal history of	Note: If a diabetes diagnosis code is present in any position, the preventive benefit will not be applied. See the	

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

negative initial postpartum

Page 36 of 46 Effective 10/01/2025

Diabetes Diagnosis Code List.

gestational diabetes)

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Certain codes may not be payable in	all circumstances due to other policies o	r guidelines.
Service		Preventive Benefit Instructions
A date in this column reflects when the		Refer to Coverage Rationale, and FAQ's
listed rating was issued.	Code(s)	sections above for additional instructions.
screening test result should be		
screened at least every 3 years for		
a minimum of 10 years after		
pregnancy. For those with a		
positive screening test result in the		
early postpartum period, testing		
should be repeated at least 6		
months postpartum to confirm the		
diagnosis of diabetes regardless of		
the type of initial test (e.g., fasting		
plasma glucose, hemoglobin A1c, in the first 6 months postpartum		
regardless of whether the test		
results are positive or negative		
because the hemoglobin A1c test is		
less accurate during the first 6		
months postpartum.		
Also see <u>Gestational Diabetes</u>		
Screening and Screening for Pre-		
<u>Diabetes and Type 2</u> in the		
Preventive Care Services section,		
and the <u>Screening for Diabetes in</u>		
Pregnancy section.		
Screening for Urinary	See the Wellness Examinations row	See the Wellness Examinations row in
Incontinence	in the Preventive Care Services	the Preventive Care Services section
HRSA Requirement (Jan. 2024)	section above.	above.
The Women's Preventive Services		
Initiative recommends screening		
women for urinary incontinence		
annually. Screening should		
address whether women		
experience urinary incontinence		
and whether it impacts their		
activities and quality of life. If		
indicated, facilitating further		
evaluation and treatment is		
recommended.		
Counseling for Sexually	See the Wellness Examinations row	See the Wellness Examinations row in
Transmitted Infections	in the Preventive Care Services	the Preventive Care Services section
(STIs)	section above.	above.
HRSA Requirement (Dec. 2021):		
WPSI recommends directed		
behavioral counseling by a health		
care clinician or other appropriately		
trained individual for sexually active		
adolescent and adult women at an increased risk for STIs. WPSI		
IIIOIGASCU IISK IUI STIS. WESI		
recommends that clinicians review		

Preventive Care Services
United Healthcare Commercial and Individual Exchange Medical Policy

Page 37 of 46 Effective 10/01/2025

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service A date in this column reflects when the listed rating was issued. a woman's sexual history and risk	Code(s)	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.	
factors to help identify those at an increased risk of STIs. Risk factors include, but are not limited to, age younger than 25, a recent history of an STI, a new sex partner, multiple partners, a partner with concurrent partners, a partner with an STI, and a lack of or inconsistent condom use.			
For adolescents and women not identified as high risk, counseling to reduce the risk of STIs should be considered, as determined by clinical judgment.			
Screening for Human	Education and Risk	Education and Risk Assessment	
Immunodeficiency Virus Infection (HIV)	Assessment See the Wellness Examinations row in the Preventive Care Services	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.	
HRSA Requirement (Dec. 2021): The Women's Preventive Services Initiative (WPSI) recommends all adolescent and adult women, ages 15 and older, receive a screening test for human immunodeficiency virus (HIV) at least once during	See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the Preventive Care Services section	Screening Tests See the HIV (Human Immunodeficiency Virus) Screening for Adolescents and Adults row in the Preventive Care Services section above.	
their lifetime. Earlier or additional screening should be based on risk, and rescreening annually or more often may be appropriate beginning at age 13 for adolescent and adult women with an increased risk of HIV infection. The WPSI recommends risk assessment and	above		
prevention education for HIV infection beginning at age 13 and continuing as determined by risk. A screening test for HIV is recommended for all pregnant women upon initiation of prenatal care with rescreening during pregnancy based on risk factors. Rapid HIV testing is recommended for pregnant women who present in labor with an undocumented HIV			
status.	Code Creum 4 Bresonderer Code ()	Code Crown 4:	
Contraceptive Methods (Including Sterilizations)	Code Group 1 Procedure Code(s): Sterilizations:	Code Group 1: Does not have diagnosis code requirements for preventive benefits to apply.	

Preventive Care Services

United Healthcare Commercial and Individual Exchange Medical Policy

Page 38 of 46 Effective 10/01/2025

Expanded Women's Preventive Health These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the Preventive Care Services section above. Certain codes may not be payable in all circumstances due to other policies or guidelines. **Preventive Benefit Instructions** Service A date in this column reflects when the Refer to Coverage Rationale, and FAQ's Code(s) listed rating was issued. sections above for additional instructions. HRSA Requirement (Dec. 2021): Tubal Ligation, Oviduct Occlusion:58600, 58605, 58611, WPSI recommends that adolescent 58615, 58670, 58671, A4264 and adult women have access to the full range of contraceptives and (See Code Group 4 below for Tubal contraceptive care to prevent Ligation Follow-Up) unintended pregnancies and Contraceptive Methods: improve birth outcomes. Diaphragm or Cervical Cap: 57170. Contraceptive care includes A4261, A4266 screening, education, counseling, IUD (Paragard®): J7300 and provision of contraceptives (including in the postpartum period). IUD (Skyla®): J7301 Contraceptive care also includes IUD (Liletta®): J7297 follow-up care (e.g., management, IUD (Kyleena®): J7296 evaluation, and changes, including (See Code Group 2 below for the removal, continuation, and additional IUD codes) discontinuation of contraceptives). **Code Group 2 Procedure Code(s):** Code Group 2: WPSI recommends that the full range of U.S. Food and Drug Contraceptive Methods: Administration (FDA)-approved, -Implantable Devices: granted, or -cleared contraceptives, J7306, J7307 effective family planning practices, 11976 (capsule removal) and sterilization procedures be 11981 (implant insertion) available as part of contraceptive 11982 (implant removal) care. The full range of 11983 (removal with reinsertion) contraceptives includes those IUDs: currently listed in the FDA's Birth

J7298 (Mirena®)

S4989

Control Guide: (1) sterilization

rods, (3) copper intrauterine

doses), (5) injectable contraceptives, (6) oral

continuous use), (9) the

contraceptive rings, (11)

emergency contraception

(levonorgestrel), and (17)

or cleared by the FDA.

surgery for women, (2) implantable

devices, (4) intrauterine devices

with progestin (all durations and

contraceptives (combined pill), (7)

oral contraceptives (progestin only),

(8) oral contraceptives (extended or

contraceptive patch, (10) vaginal

diaphragms, (12) contraceptive

sponges, (13) cervical caps, (14)

condoms, (15) spermicides, (16)

emergency contraception (ulipristal acetate), and any additional

contraceptives approved, granted.

58300, S4981 (insertion)

58301 (removal)

(See Code Group 1 above for additional IUD codes)

Injections:

96372 (administration)

J1050 (injection)

Code Group 2 Diagnosis Code(s):

These are required for Code Group 2. Contraceptive Management: Z30.012, Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9

Code Group 3 Procedure Code(s):

Anesthesia for Sterilization: 00851, 00940, 00942, 00950, 00952, 01960, 01961, 01965, 01966, 01967, Requires one of the Code Group 2 diagnosis codes listed in this row.

Code Group 3:

Requires one of the Code Group 3 diagnosis code listed in this row.

Preventive Care Services UnitedHealthcare Commercial and Individual Exchange Medical Policy

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above Certain codes may not be payable in all circumstances due to other policies or quidelines.

For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.			
Service	an circumstances due to other policies of	Preventive Benefit Instructions	
A date in this column reflects when the		Refer to Coverage Rationale, and FAQ's	
listed rating was issued.	Code(s)	sections above for additional instructions.	
Notes: Coverage includes member reimbursement for the cost of FDA-approved, cleared, or granted mobile device applications for use as contraception consistent with the FDA-approved, cleared, or granted indication. For counseling and follow-up care, see the Wellness Examinations row in the Preventive Care Services section above. Certain employers may qualify for an exemption from covering contraceptive methods and sterilizations on account of religious objections. Refer to the plan's pharmacy benefit plan administrator for details on pharmacy contraceptives available under the plan's preventive benefit. Also refer to the Medical Policy titled Outpatient Surgical Procedures — Site of Service.	Sterilization - Laparoscopy with Removal of Adnexal Structures: 58661 Code Group 3 Diagnosis Code(s): This code is required for all Code Group 3 Procedure Codes: Sterilization: Z30.2 Tubal Ligation Follow-Up Hysterosalpingogram Code Group 4 Procedure Code(s): Catheterization and Introduction of Saline or Contrast Material: 58340	Code Group 4: Requires one of the Code Group 4 diagnosis code listed in this row.	
	Hysterosalpingography: 74740 Contrast Material: Q9967 Code Group 4 Diagnosis Code(s): Tubal Ligation Status: Z98.51		
	Code Group 5 Procedure Code(s): IUD Follow-Up Evaluation and Management (Office Visit): 99211, 99212 Pelvic Examination (add-on code): 99459 Refer to Code Group 7, Related Visits section below, for additional coding for Evaluation and Management (Office Visits). Code Group 5 Diagnosis Code(s): Encounter for routine checking of intrauterine contraceptive device: Z30.431	Code Group 5: Requires one of the Code Group 5 diagnosis code listed in this row. Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.	
	Code Group 6 Procedure Code(s): Impacted IUD removal 58562 Code Group 6 Diagnosis Code(s):	Code Group 6: Requires one of the Code Group 6 diagnosis codes listed in this row.	

Z30.432, Z30.433

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service

A date in this column reflects when the listed rating was issued.

Code(s)

Code Group 7 Procedure Code(s):

Related Visits: Related Evaluation and Management Office/Outpatient Visits for Contraception or Sterilization:

99202, 99203, 99204, 99205, 99211, 99212, 99213, 99214, 99215, 99417, G0463

Pelvic Examination (add-on code): 99459

Also see coding in the <u>Wellness</u> <u>Examinations</u> row above.

Related Pregnancy Tests:
Pregnancy Tests When Related to
Contraception or Sterilization: 81025,
84702, 84703

Code Group 7 Diagnosis Codes:

Tubal Ligation Status: Z98.51 Sterilization: Z30.2 Contraceptive Management: Z30.012, Z30.013, Z30.014, Z30.017, Z30.018

Z30.013, Z30.014, Z30.017, Z30.018, Z30.019, Z30.09, Z30.40, Z30.42, Z30.430, Z30.431, Z30.432, Z30.433, Z30.46, Z30.49, Z30.8, Z30.9

Preventive Benefit Instructions

Refer to Coverage Rationale, and FAQ's sections above for additional instructions.

Code Group 7:

Requires one of the Code Group 7 diagnosis codes listed in this row.

Pelvic Examination add-on code 99459: Preventive care services benefits may apply to 99459 when the related evaluation and management (office visit) code is applied to the preventive care services benefit. CPT code 99459 may not be payable in all circumstances due to other policies or guidelines.

Breastfeeding Services and Supplies

HRSA Requirement (Dec. 2021):

WPSI recommends comprehensive lactation support services including consultation; counseling; education by clinicians and peer support services; and breastfeeding equipment and supplies) during the antenatal, perinatal, and postpartum periods to optimize the successful initiation and maintenance of breastfeeding. Breastfeeding equipment and supplies include, but are not limited to, double electric breast pumps (including pump parts and maintenance) and breast milk storage supplies. Access to double electric pumps should be a priority to optimize breastfeeding and

Counseling and Education Procedure Code(s):

98960, 98961, 98962, 99242*, 99243*, 99244*, 99245*, 99341, 99342, 99344, 99345, 99347, 99348, 99349, 99350, S9443

Also see the codes in the <u>Wellness</u> <u>Examinations</u> row in the *Preventive Care Services* section above.

Diagnosis Code(s):

B37.89, N61.1, N64.4, N64.51, N64.52, N64.53, N64.59, N64.89, O91.011, O91.012, O91.013, O91.019, O91.02, O91.03, O91.111, O91.112, O91.113, O91.119, O91.13, O91.211, O91.212, O91.213, O91.219, O91.22, O91.23, O92.011, O92.012, O92.013, O92.019, O92.02, O92.03, O92.111, O92.112, O92.113, O92.119, O92.12, O92.13, O92.20, O92.29, O92.3, O92.4, O92.5,

Counseling and Education

Requires one of the diagnosis codes listed in this row for 98960-98962, 99242-99245, 99341-99345, and 99347-99350.

Does not have diagnosis code requirements for preventive benefits to apply for S9443.

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or quidelines.

Certain codes may not be payable in all circumstances due to other policies or guidelines.				
Service A date in this column reflects when the listed rating was issued. should not be predicated on prior failure of a manual pump. Breastfeeding equipment may also include equipment and supplies as clinically indicated to support dyads with breastfeeding difficulties and those who need additional services	Code(s) O92.70, O92.79, Q83.1, Q83.2, Q83.3, Q83.8,Z39.1, Z39.2 *For additional information on the reimbursement of consultation codes 99242-99245, refer to the Reimbursement Policy titled Consultation Services.	Preventive Benefit Instructions Refer to Coverage Rationale, and FAQ's sections above for additional instructions.		
services.	Breastfeeding Equipment & Supplies Procedure Code(s): Personal Use Electric Breast Pump: E0603 Breast Pump Supplies: A4281, A4282, A4283, A4284, A4285, A4286, A4287, A4288 Diagnosis Code(s): Pregnancy Diagnosis Codes or Z39.1.	Breastfeeding Equipment & Supplies E0603 is limited to one purchase per birth. E0603 and A4281-A4287 require at least one of the diagnosis codes listed in this row.		
Screening and Counseling for Interpersonal and Domestic Violence HRSA Requirement (Dec. 2016): Recommends screening adolescents and women for interpersonal and domestic violence, at least annually, and, when needed, providing or referring for initial intervention services. Interpersonal and domestic violence includes physical violence, sexual violence, stalking and psychological aggression (including coercion), reproductive coercion, neglect, and the threat of violence, abuse, or both. Intervention services include, but are not limited to, counseling, education, harm reduction strategies, and referral to appropriate supportive services. Also see the Screening for Intimate Partner Violence row in the Preventive Care Services section above.	See the Wellness Examinations row in the Preventive Care Services section above.	See the Wellness Examinations row in the Preventive Care Services section above.		

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Certain codes may not be payable in	all circumstances due to other policies o	r guidelines.
Service		Preventive Benefit Instructions
A date in this column reflects when the listed rating was issued.	Code(s)	Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
Breast Cancer Screening for Average-Risk Women	See the <u>Screening Mammography</u> row in the <i>Preventive Care Services</i> section above.	See the <u>Screening Mammography</u> row in the <i>Preventive Care Services</i> section above.
HRSA Requirement (Dec. 2016): Recommends that average-risk women initiate mammography screening no earlier than age 40 and no later than age 50. Screening mammography should occur at least biennially and as frequently as annually. Screening should continue through at least age 74 and age alone should not be the basis to discontinue screening. These screening recommendations are for women at average risk of breast cancer. Women at increased risk should also undergo periodic mammography screening; however, recommendations for additional services are beyond the scope of this recommendation.		
Screening for Cervical	Human Papillomavirus DNA	Human Papillomavirus DNA
Cancer HRSA Requirement (Dec. 2016): Recommends cervical cancer screening for average-risk women aged 21 to 65 years. For women aged 21 to 29 years recommends cervical cancer screening using cervical cytology (Pap test) every 3 years. Co-testing with cytology and human papillomavirus testing is not recommended for women younger than 30 years. Women aged 30 to 65 years should be screened with cytology and human papillomavirus testing every 5 years or cytology alone every 3 years. Women who are at average risk should not be	Testing (HPV) See the Cervical Cancer Screening row in the Preventive Care Services section above. Cervical Cytology (Pap Test) See the Cervical Cancer Screening row in the Preventive Care Services section above.	Testing (HPV) See the Cervical Cancer Screening row in the Preventive Care Services section above. Cervical Cytology (Pap Test) See the Cervical Cancer Screening row in the Preventive Care Services section above.
screened more than once every 3 years.		
Screening for Anxiety	Procedure Code(s):	Requires the diagnosis code listed in this row.
HRSA Requirement (Dec. 2019): The Women's Preventive Services Initiative recommends screening for anxiety in adolescent and adult	96127	uns iow.

Preventive Care Services

These are the requirements of the Health Resources and Services Administration (HRSA). For additional services covered for women, see the <u>Preventive Care Services</u> section above. Certain codes may not be payable in all circumstances due to other policies or guidelines.

Service		Preventive Benefit Instructions
A date in this column reflects when the listed rating was issued.	Code(s)	Refer to Coverage Rationale, and FAQ's sections above for additional instructions.
women, including those who are pregnant or postpartum. Optimal screening intervals are unknown and clinical judgement should be used to determine screening frequency. Given the high prevalence of anxiety disorders, lack of recognition in clinical practices, and multiple problems associated with untreated anxiety, clinicians should consider screening women who have not been recently screened.	Diagnosis Code(s): Encounter for Screening Examination for Other Mental Health and Behavioral Disorders: Z13.39	
Also see the rows for Anxiety Disorders in Adults Screening (USPSTF); Screening for Anxiety in Children and Adolescents (USPSTF); Screening for Depression in Adults (USPSTF); Depression in Children and Adolescents (Screening) (USPSTF); Perinatal Depression — Preventive Interventions (Counseling) (USPSTF); and Depression Screening (Bright Futures) in the Preventive Care Services section above.		
Obesity Prevention in Midlife Women (Counseling)	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.	See the Wellness Examinations row in the <i>Preventive Care Services</i> section above.
HRSA Requirement (Dec. 2021): WPSI recommends counseling midlife women aged 40 to 60 years with normal or overweight body mass index (BMI) (18.5-29.9 km/m2) to maintain weight or limit weight gain to prevent obesity. Counseling may include individualized discussion of healthy eating and physical activity.		

Diagnosis Codes

Preventive Care Services: ICD-10 Diagnosis Codes

Preventive Vaccine Codes

Preventive Care Services: Vaccine Codes

Benefit Considerations

Certain plans are not required to include coverage for the services identified by the federal Patient Protection and Affordable Care Act (PPACA). Refer to the member specific benefit plan document for coverage details.

Clinical Evidence

Refer to the Service column in the Applicable Codes section for the recommendation statements supporting this policy.

References

ACIP Vaccine-Specific Recommendations: https://www.cdc.gov/acip-recs/hcp/vaccine-specific/?CDC AAref Val=https://www.cdc.gov/vaccines/hcp/acip-recs/index.html. Accessed June 18, 2025.

American Academy of Family Physicians (AAFP) Clinical Preventive Services Recommendations: https://www.aafp.org/family-physician/patient-care/clinical-recommendations/clinical-practice-guidelines/clinical-preventive-services-recommendations.html. Accessed June 18, 2025.

American Academy of Pediatrics, Bright Futures Guidelines, 4th edition, Evidence and Rationale chapter: https://www.aap.org/en/practice-management/bright-futures/bright-futures-materials-and-tools/bright-futures-guidelines-and-pocket-guide/. Accessed June 18, 2025.

American Academy of Pediatrics/Bright Futures/Recommendations for Pediatric Preventive Healthcare. (For ages 0-21): https://downloads.aap.org/AAP/PDF/periodicity schedule.pdf. Accessed June 18, 2025.

American Academy of Pediatrics: http://www.aap.org/. Accessed June 18, 2025.

Centers for Disease Control and Prevention/Immunization Schedules: https://www.cdc.gov/vaccines/hcp/imz-schedules/index.html. Accessed June 18, 2025.

Grade Definitions for USPSTF Recommendations: http://www.uspreventiveservicestaskforce.org/Page/Name/grade-definitions. Accessed June 18, 2025.

July 19, 2010 IRS Interim Rules: http://www.irs.gov/irb/2010-29 IRB/index.html. Accessed June 18, 2025.

Published Recommendations, U.S. Preventive Services Task Force:

http://www.uspreventiveservicestaskforce.org/BrowseRec/Index/browse-recommendations. Accessed June 18, 2025.

U.S. Food and Drug Administration (FDA), Vaccines Licensed for Use in the United States: http://www.fda.gov/BiologicsBloodVaccines/Vaccines/ApprovedProducts/UCM093833. Accessed June 18, 2025.

U.S. Vaccine Names Information Page: https://www.cdc.gov/vaccines/hcp/vaccines-us/index.html. Accessed June 18, 2025.

Women's Preventive Services Guidelines (HRSA): https://www.hrsa.gov/womens-guidelines. Accessed June 18, 2025.

Women's Preventive Services Initiative (WPSI): https://www.womenspreventivehealth.org/recommendations/. Accessed June 18, 2025.

Policy History/Revision Information

Date	Summary of Changes
10/01/2025	Applicable Codes Preventive Care Services Statin Use for the Primary Prevention of Cardiovascular Disease in Adults - Cholesterol Screening (Lipid Disorders Screening) Updated list of applicable ICD-10 diagnosis codes for lipid disorders not covered under the preventive benefit to reflect annual edits: Added E78.010, E78.011, and E78.019 Removed E78.01
	Colorectal Cancer Screening: Colonoscopy, Sigmoidoscopy, Pathology and Anesthesia (for Colonoscopy or Sigmoidoscopy), Fecal Occult Blood Testing (FOBT) and Fecal Immunochemical Test (FIT), and Pre-Op Consultation

Date	Summary of Changes
	 Updated list of applicable ICD-10 diagnosis codes to reflect annual edits; added Z15.060 and Z15.068
	Healthy Diet and Physical Activity for Cardiovascular Disease Prevention in Adults with Cardiovascular Risk Factors: Behavioral Counseling Interventions: Hyperlipidemia/Dyslipidemia Updated list of applicable ICD-10 diagnosis codes to reflect annual edits: Added E78.010, E78.011, and E78.019 Removed E78.01
	Expanded Women's Preventive Health Contraceptive Methods (Including Sterilizations) Updated list of applicable HCPCS codes to reflect quarterly edits; revised description for J7300 Breastfeeding Services and Supplies: Breast Pump Supplies Updated list of applicable HCPCS codes to reflect quarterly edits; added A4288 Diabetes Diagnosis Code List Diabetes Mellitus: Type 2 Updated list of applicable ICD-10 diagnosis codes to reflect annual edits; added E11.A Supporting Information Archived previous policy version MP.016.55

Instructions for Use

This Medical Policy provides assistance in interpreting UnitedHealthcare standard benefit plans. When deciding coverage, the member specific benefit plan document must be referenced as the terms of the member specific benefit plan may differ from the standard plan. In the event of a conflict, the member specific benefit plan document governs. Before using this policy, please check the member specific benefit plan document and any applicable federal or state mandates. UnitedHealthcare reserves the right to modify its Policies and Guidelines as necessary. This Medical Policy is provided for informational purposes. It does not constitute medical advice.

This Medical Policy may also be applied to Medicare Advantage plans in certain instances. In the absence of a Medicare National Coverage Determination (NCD), Local Coverage Determination (LCD), or other Medicare coverage guidance, CMS allows a Medicare Advantage Organization (MAO) to create its own coverage determinations, using objective evidence-based rationale relying on authoritative evidence (Medicare IOM Pub. No. 100-16, Ch. 4, §90.5).

UnitedHealthcare may also use tools developed by third parties, such as the InterQual® criteria, to assist us in administering health benefits. UnitedHealthcare Medical Policies are intended to be used in connection with the independent professional medical judgment of a qualified health care provider and do not constitute the practice of medicine or medical advice.