

27. Pathophysiology of Acute Kidney Injury · assets 1-9 / 13




Fig. 21 The pathophysiology of cisplatin nephrotoxicity. Uptake of cisplatin in the PTCs (proximal tubule cells) is mediated by solute carrier family 22 member 2 (SLC22A2), SLC22A6, and SLC22A8 and its extrusion is mediated by CLC4A11 (MATE1). Upon entering PTCs, cisplatin is hydrated and accumulated to the toxic form by a γ -glutamyl transpeptidase (GGT) and kynurenine aminotransferase 1 (KYAT1)-dependent pathway. Cisplatin accumulation to toxic levels in PTCs induces DNA damage, mitochondrial damage, oxidative stress, endoplasmic reticulum (ER) stress, autophagy, and cell cycle regulation. Ultimately, these processes lead to cell death, acute kidney injury, and chronic kidney disease. [Created in BioRender; Okusa M. 2025. <https://BioRender.com/bbf7dxk>.]

Fig 27 1




Fig. 7.4 Overview of the pathways of regulated cell death. In general, two systems may be best differentiated when regulated cell death is considered. The caspase-controlled system includes apoptosis, necroptosis, and pyroptosis. In contrast, the peroxidation-controlled system of ferroptotic cell death functions entirely independently of the caspase-controlled network. [From Tonnus W, Gembardt F, Latk M, et al. The clinical relevance of neuroinflammation highlighting the importance of acute kidney injury and the adrenal glands. *Cell Death Differ.* 2019;26(11):68–82.)

Fig 27 4




Fig. 2.7. Tubular regeneration following acute kidney injury. Top panel, Tubular regeneration by differentiated cells. (A) Healthy tubules consist of nonproliferative mature epithelial cells that express markers of differentiation. (B) After injury, the epithelium is lost through apoptosis and necrosis. (C) Surviving epithelial cells differentiate, either in response to sublethal injury signals or owing to signals from other injured cells, and acquire a proliferative phenotype. (D) The surviving differentiated cells reconstitute the nephron epithelium. (E) Ultimately, most tubules are reconstituted by cells that have proliferated from the surviving differentiated cells. Bottom panel, Tubular regeneration by scattered progenitor epithelial cells. (F) Scattered tubular epithelial cells express signal transducer CD24, prominin-1 (CD133), and other genes that are characteristic of proximal tubule differentiation, such as vimentin and KIM-1. (G) After injury, mature tubular cells, but not scattered cells, undergo apoptosis. (H) Scattered tubular cells expand in response to injury and their progeny reconstitute the tubule. (I) The small subpopulation of scattered cells that survive after expansion. (From Chang-Fessenden M, Humphreys BD. Cellular plasticity in kidney injury repair. *Nat Rev Nephrol* 2017;13(1):39–46).

Fig_27_7




Fig. 27.2 Overview of pathophysiology of acute kidney injury (AKI). In the kidney, mechanisms pertaining to microvascular compartment, innate immunity, and ATN result in temporary, partial, or permanent loss of the glomerular filtration rate. Systemic responses may influence the extent of injury. Hemodynamic alterations may initiate AKI or exacerbate intrinsic microenvironmental mechanisms of AKI. Systemic immunologic mechanisms of proinflammatory or anti-inflammatory conditions may affect AKI, and neural mechanisms may attenuate AKI (see text for details).




Fig. 2/3 Morphology of human acute tubular necrosis. (A) Human biopsy specimens reveal significant proximal tubular cell damage, with intraluminal accumulation of apical membrane fragments and detached cells (b); thinning of proximal tubular cells to maintain monolayer tubule integrity (arrow), and dividing cells and accumulation of white cells within the microvasculature in the peritubular area (arrowheads). (B) Electron micrograph of a regenerating epithelial cell. Shown are small fragmented mitochondria (r). (C) Electron micrograph of renal epithelial cells showing nonreplicative state [black arrow] that morphologically supports the concept of back leak in the pathophysiology of AKI. (A, Courtesy M. Verma; B & C, Courtesy M. Verma and J. A. Verma). Tubular ultrastructure in acute renal failure in man: epithelial necrosis and regeneration. *Virchows Arch A Pathol Anat Histophysiol.* 1985;40(6):775-789.

Fig 27 3




Fig. 27.5 Autophagy. The forms of autophagy are macroautophagy, microautophagy, and chaperone-mediated autophagy. Macroautophagy starts with the de novo formation of a cup-shaped isolation double membrane that engulfs a portion of cytoplasm. Microautophagy involves the engulfment of cytoplasm instantly at the lysosomal membrane by invagination, protrusion, and separation. Chaperone-mediated autophagy is a process of direct transport of unfolded proteins via the lysosomal chaperonin hsc70 and LAMP-2. All forms of autophagy subsequently lead to the degradation of intra-autophagosomal components by lysosomal hydrolases. PE, Phosphatidylethanolamine. (From Periyasamy-Thandavesan S, Jiang M, Schoenlein P, Dong Z. Autophagy: molecular machinery, regulation, and implications for renal pathophysiology. *Am J Physiol Renal Physiol.* 2009;297(2):F244–F256.)

Fig 27 5


[illegible]

Fig 27 6




Fig. 27.8 Ex vivo microcomputed tomography (CT) imaging and three-dimensional quantification of the renal vasculature reveal significant alterations in renovascularities in progressive ischemia-reperfusion-induced (IR) renal injury. Lower panel: Representative high-resolution, ex vivo, micro-CT images of sham control and IR kidneys from days 14, 21, and 56 after Microfil perfusion (two-dimensional cross-sectional images in transverse [b], coronal [c], and sagittal [d] planes, as well as three-dimensional volume renderings). Note the progressive rarefaction of functional vessels between the continuous stretching of fibrotic kidneys over time. Scale bar: 200 μ m. (Upper panel) Quantitative CT-based quantification of vascular branching points per increasing vessel order (from hilus to periphery). A continuous reduction in vessel branching points was observed between the sham control and IR kidneys during the first 2 weeks after ischemia, followed by a partial recovery. IR injury. (From Ehrlich J, Babitskaya J, Griesma F, et al. Quantitative micro-computed tomography imaging of vascular dysfunction in progressive kidney disease. *J Am Soc Nephrol*. 2016;27(2):502-532.)

Fig_27_9







Fig. 27.3 Microvascular dysfunction in septic acute kidney injury. Multiple mechanisms are involved in the development of sepsis-related microvascular dysfunction, among them endothelial dysfunction related partly to increased toxin-derived and pathogen-derived mediators, as well as to reduced nitric oxide bioavailability. These factors lead to endothelial dysfunction, which in turn causes microvascular dysfunction, resulting in an altered glycocalyx. The glycocalyx is a layer of glycoproteins that covers the endothelial surface, facilitating the flow of red blood cells and limiting the adhesion of leukocytes and platelets to the endothelium. The glycocalyx may be substantially altered during sepsis, so that adherence between the vascular endothelium and circulating cells (e.g., leukocytes) is increased. Microvascular dysfunction is also caused by increased leukocyte adhesion to the endothelium, which in turn causes microvascular dysfunction. Microvascular dysfunction might also participate in sepsis-induced microvascular alterations, as well as alterations in leukocyte deformability and their adhesion to the endothelium. All these phenomena might cause heterogeneity in microvascular blood flow, with a decrease in vascularity and nonperfused capillaries, resulting in an increased diffusion distance for oxygen and in alterations in oxygen extraction. *endothelium*, endothelial cells; *glycocalyx*, glycocalyx; *leukocytes*, white blood cells; *microvascular dysfunction*, microvascular dysfunction; *nitric oxide*, nitric oxide; *platelets*, platelets; *sepsis*, sepsis; *vascular endothelium*, vascular endothelium; *white blood cells*, white blood cells.

Fig_27_8

27. Pathophysiology of Acute Kidney Injury · assets 10-13 / 13



Fig_27_10

[illegible]

eFig. 27.6 The impact of acute kidney injury (AKI) on distant organs. AKI causes hemodynamic, humoral, and immunologic changes, which lead to dysfunction of distant organs, including lung, heart, brain, liver, intestine, and the immune system. (From Lee SA, Cozzoli M, Bush LE, Rabb H. Distant organ dysfunction in acute kidney injury: a review. *Am J Kidney Dis.* 2018;72[6]:846–856.)

Fig_e27_6

Organ	Mechanism	Species	Reference
Lung	CXCL1	Mouse	638
	TNF- α	Mouse	639
	IL-10	Mouse	611,640
	Neutrophil elastase	Mouse	641
	Lung permeability	Mouse	610
	Osteopontin and lung injury	Mouse	612
	Pulmonary edema, inflammatory cytokines	Rat	642
	Oxidant stress	Rat	643
	Systemic cytokines and HMGB1	Human (ex vivo)	644
	Heart	TNF- α	Rat
	Mitochondrial fission protein	Mouse	645
Brain	RAAS	Mouse	646
Liver	Administration of glutathione	Rat	647
	Hepatic oxidant stress	Mouse	648,649
Intestine	Hepatic oxidant stress	Mouse	650

Table 27_1