



Fig. 78.1 Countries and territories that constitute the Far East. Modified from Wikipedia, https://en.wikipedia.org/wiki/Geography_of_Asia. Map lines delineate study areas and do not necessarily depict accepted national boundaries.

Fig_78_1




Fig. 78.2 Prevalence of immunoglobulin A nephropathy as indicated by percentage of biopsy-proven primary glomerulonephritis in different parts of the world. (Reprinted by permission from Macmillan Publishers Ltd. From Lai KN, Tang SCW, Schena FP, et al. IgA nephropathy. *Nat Rev Dis Primers*. 2016;2:16001.) Map lines delineate study areas and do not necessarily depict accepted national boundaries.

Fig_78_2




Fig. 78.3 Proportion of primary diagnoses of end-stage renal disease in prevalent patients on dialysis in Japan. GN, Glomerulonephritis; PKD, polycystic kidney disease. (Reprinted with permission from Elsevier. From Hanafusa N, Nakai S, Iseki K, et al. Japanese Society for Dialysis Therapy Renal Data Registry – a window through which we can view the details of Japanese dialysis population. *Kidney Int Suppl*. 2015;5:15–22.)

Fig_78_3

Table 78.1 Secondary Renal Diseases Presenting as Acute Kidney Injury in East, Southeast, and Northeast Asia	
Causes	Examples
Autoimmunity	- Lupus nephritis - ANCA vasculitis - Antiglomerular basement membrane disease - Miscellaneous glomerulonephritis
Infections	- Leptospirosis - Dengue fever - Hantaviruses (hemorrhagic fever with renal syndrome) - Scrub typhus - Malaria - Severe fever with thrombocytopenia syndrome virus
Toxins	- From plant (e.g., aristolochic acid and djenkolic acid) - From animal or insect (e.g., snake venom and grass carp fish gallbladder) - Chemical (e.g., ketamine abuse presenting with CKD or AKI due to obstructive uropathy)
AKI, Acute kidney injury; ANCA, antineutrophil cytoplasmic antibody; CKD, chronic kidney disease.	

Table_78_1

Table 78.2 Chinese Medicinal Herbs that Contain Aristolochic Acids	
Name of Chinese Herb	Botanical Name
Guang Fang Ji (Fangchi)	<i>Aristolochia fangchi</i>
Xixin	Radix et Rhizoma Asari
Guan Mu Tong	<i>Aristolochia manshuriensis</i>
Qing Mu Xiang	<i>Aristolochia cucurbitifolia</i>
Ma Dou Ling	<i>Aristolochia debilis</i>
Tian Xian Teng	<i>Aristolochia contorta</i>
Modified from Yang HY, Chen PC, Wang JD. Chinese herbs containing aristolochic acid associated with renal failure and urothelial carcinoma: a review from epidemiologic observations to causal inference. <i>Biomed Res Int</i> . 2014;2014:569325.	

Table_78_2

Table 78.3 Typical Clinical Phases of Hemorrhagic Fever with Renal Syndrome due to Hantavirus Infection				
Phase of Illness	Time of Occurrence	Principal Features	Signs and Symptoms	Complications
1. Febrile	1–7 days	Fever	- Headache - Vomiting - Abdominal pain - Back pain - Visual disturbance - Capillary leak - Pulmonary symptoms	- Acute encephalomyelitis - Bleeding - Shock - Multiorgan failure - Pituitary hemorrhage - Renal - Pulmonary edema
2. Hypotensive	1–3 days	Hypotension	Severe oliguria can occur	Acute respiratory distress syndrome
3. Oliguric	2–6 days	↓ Urine output	Fluid retention	Disseminated intravascular coagulation
4. Polyuric	2 weeks	↑ Urine output	Decrease of body weight	
5. Convalescent	3–6 months		- Weakness - Fatigue	
Modified from Jiang H, Du H, Wang LM, et al. Hemorrhagic fever with renal syndrome: pathogenesis and clinical picture. <i>Front Cell Infect Microbiol</i> . 2016;6:1.				

Table_78_3

Table 78.4 HBsAg Seroprevalence and the Number of People Living with Chronic Hepatitis B Virus in the General Population in Some Asian Countries			
Country	Prevalence Estimates (%; 95% CI)	Population Size Per Country	HBsAg-Positive Population
China	5.49 (5.47–5.50)	1,359,821,465	74,601,204
Japan	1.02 (1.01–1.02)	127,352,833	1,294,431
Malaysia	0.74 (0.70–0.77)	28,275,935	208,540
Mongolia	9.07 (8.41–9.78)	2,712,738	246,070
The Philippines	4.63 (4.53–4.73)	93,444,322	4,326,212
Singapore	4.09 (3.87–4.33)	5,078,969	207,943
South Korea	4.36 (4.36–4.37)	48,453,931	2,111,914
Thailand	6.42 (6.37–6.47)	66,402,316	4,260,008
Vietnam	10.79 (10.29–11.31)	89,047,397	9,607,438
CI, Confidence interval; HBsAg, hepatitis B surface antigen; HBV, hepatitis B virus. Modified from Schweitzer A, Horn J, Mikolajczyk RT, et al. Estimations of worldwide prevalence of chronic hepatitis B virus infection: a systematic review of data published between 1965 and 2013. <i>Lancet</i> . 2015;386:1546–1555.			

Table_78_4