

Chronic Kidney Disease–Mineral Bone Disorder

Anna L. Zisman | Stuart M. Sprague

CHAPTER OUTLINE

PATHOPHYSIOLOGY OF CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER, 1533

DIAGNOSIS OF CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER, 1550

CLINICAL CONSEQUENCES OF THE ABNORMALITIES IN CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER, 1558

CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER IN KIDNEY TRANSPLANT RECIPIENTS, 1561

SUMMARY, 1563

KEY POINTS

- Changes in mineral metabolism parameters occur as early as stage 2 chronic kidney disease (CKD), most commonly with increased serum fibroblast growth factor 23 concentrations.
- Pi bioavailability depends on the food source, with plant-based sources such as legumes having less bioavailable Pi due to human's inability to digest phytate, the major Pi storage form in plants.
- The commonly used albumin-corrected total calcium values misclassify the true hypercalcemia or hypocalcemia in 20% of patients with CKD; thus whenever possible, ionized calcium concentrations should also be obtained, particularly when starting or modifying therapies for CKD-MBD (mineral bone disorder).
- Vascular calcification of the medial layer of elastic arteries is extremely common in patients with CKD. The pathophysiology may be similar to that of intimal calcifications associated with atherosclerosis.
- There is no definitive noninvasive way (by imaging or bloodwork) to assess bone physiology parameters such as turnover and mineralization.
- The newer system for renal osteodystrophy classification focuses on abnormalities in bone turnover, mineralization, and volume, all of which affect bone strength.
- Dual-energy x-ray absorptiometry may be used for assessment of fracture risk in patients with CKD, although clinical trials linking changes in dual-energy x-ray absorptiometry to relevant outcomes including fracture are lacking.
- There are no randomized controlled trials in CKD or end-stage kidney disease to show that treatment to a specific parathyroid hormone target leads to improved outcomes.
- Patients on dialysis have higher rates of hip fracture than in the general population in all age groups and have a higher mortality when they experience a fracture.
- Fractures, especially in the appendicular skeleton, are common in kidney transplant recipients.

In persons with healthy kidneys, normal serum concentrations of calcium and phosphorus are maintained through the interaction of three hormones: parathyroid hormone (PTH); calcitriol (1,25-dihydroxyvitamin D₃ 1,25[OH]₂D), the active hormonal metabolite of vitamin D; and fibroblast growth factor 23 (FGF-23). Circulating or soluble klotho also plays a role in mineral homeostasis.

These hormones act on four primary target organs: bone, kidney, intestine, and parathyroid glands. The kidneys play a critical role in the regulation of normal serum calcium and phosphorus concentrations and of the three hormones. Thus derangements in mineral homeostasis are common in patients with chronic kidney disease (CKD). Abnormalities begin in early CKD and are nearly universally

observed at glomerular filtration rates (GFR) <30 mL/min. With progression of CKD, the body attempts to maintain normal serum concentrations of calcium and phosphorus by altering the production of calcitriol, PTH, FGF-23, and klotho. Eventually these compensatory responses become unable to maintain normal mineral homeostasis, resulting in 1. altered serum concentrations of calcium, phosphorus, PTH, calcitriol, FGF-23, and klotho; 2. disturbances in bone remodeling and mineralization (often referred to as “renal osteodystrophy”) and/or impaired linear growth in children; and 3. extraskeletal calcification in soft tissues and arteries. In 2006, the term chronic kidney disease–mineral and bone disorder (CKD-MBD) was developed to describe this triad of abnormalities in biochemical measures, skeletal abnormalities, and extraskeletal calcification (Table 52.1).¹ These abnormalities that constitute CKD-MBD are interrelated in both the pathophysiology of the disease and response to treatment. All three components of CKD-MBD are associated with increased risk of fractures, cardiovascular disease, and mortality in patients with CKD stages 4 through 5D. However, to enhance understanding of the complex integration of these abnormalities in CKD, each component is first discussed independently.

PATHOPHYSIOLOGY OF CHRONIC KIDNEY DISEASE—MINERAL AND BONE DISORDER

PHOSPHORUS AND CALCIUM HOMEOSTASIS

PHOSPHORUS BALANCE AND HOMEOSTASIS

Inorganic phosphorus is critical for numerous physiologic functions, including skeletal development, mineral

metabolism, cell membrane phospholipid content and function, cell signaling, platelet aggregation, and energy transfer through mitochondrial metabolism. Because of the importance of phosphorus in these functions, normal homeostasis maintains serum phosphorus concentrations between 2.5 and 4.5 mg/dL (0.81 and 1.45 mmol/L). Serum concentrations are highest in infants and decrease throughout growth, reaching adult levels in the late teens. Total adult body stores of phosphorus are approximately 700 g, of which 85% is contained in bone in the form of hydroxyapatite ($[\text{Ca}]_{10}[\text{PO}_4]_6[\text{OH}]_2$). Of the remainder, 14% is intracellular and only 1% is extracellular. Of this extracellular phosphorus, 70% is organic (phosphate) and contained within phospholipids and 30% is inorganic. The inorganic fraction is 15% protein bound, and the remaining 85% is either complexed with sodium, magnesium, or calcium or circulates as the free monohydrogen or dihydrogen form. It is this inorganic fraction that is freely circulating and measured. At a pH of 7.4, inorganic phosphates are in a ratio of about 4:1 HPO_4^{-2} to $\text{H}_2\text{PO}_4^{-1}$. For that reason, the serum phosphorus concentration is usually expressed in mmol/L rather than mEq/L. Thus serum measurements reflect only a minor fraction of total body phosphorus and therefore do not accurately reflect total body stores in the setting of the abnormal homeostasis that occurs in CKD. Furthermore, there is considerable diurnal variation in serum phosphorus concentrations in healthy persons,² as well as in patients with advanced CKD.³ The terms “phosphorus” and “phosphate” are often used interchangeably, but strictly speaking, phosphate means the inorganic freely available form (HPO_4^{-2} and $\text{H}_2\text{PO}_4^{-1}$). However, most laboratories report phosphate, the measurable inorganic component of total body phosphorus, as phosphorus. For simplicity, we use the abbreviation Pi to represent phosphate and/or phosphorus throughout this chapter.

Pi is contained in almost all foods and is generally associated with a food’s protein content or phosphate-containing additives. In most commonly ingested foods without additives, the mean Pi content ranges from 9.0 to 14.6 mg per gram of protein, with many foods having up to a 28% higher Pi content because of additives, which are used as preservatives, acidifying agents, acidity buffers, and emulsifying agents.⁴ Although the recommended daily allowance for Pi is 800 mg/day, the average American diet contains approximately 1000 to 1400 mg Pi and that amount does not necessarily include inorganic Pi added as a preservative, a common practice.⁵ The source of Pi has a significant impact on bioavailability. Pi in the form of preservatives or additives is nearly 100% bioavailable, whereas Pi bound to phytate, as in legumes, is less bioavailable owing to the lack of the enzyme phytase in humans.⁶ In studies, the source of Pi directly affects Pi homeostasis.³ As a result, it is challenging to balance dietary Pi restriction against the need for adequate protein intake in patients with CKD, especially with malnutrition present in up to 50% of patients undergoing dialysis. Pi balance in earlier stages of CKD (stage G3a, G3b, and in most cases G4) is generally neutral because of the phosphaturic effects of PTH and FGF-23⁷; as these compensatory mechanisms begin to fail and particularly when patients have little to no residual kidney function, however, positive Pi balance likely ensues.

Table 52.1 Kidney Disease Improving Global Outcomes Classification of Chronic Kidney Disease–Mineral Bone Disorder and Renal Osteodystrophy

Definition of chronic kidney disease–mineral bone disorder (CKD-MBD)	A systemic disorder of mineral and bone metabolism due to CKD manifested by one or a combination of the following: Abnormalities of calcium, phosphorus, parathyroid hormone, or vitamin D metabolism Abnormalities in bone turnover, mineralization, volume, linear growth, or strength Vascular or other soft tissue calcification
Definition of renal osteodystrophy	Renal osteodystrophy is an alteration of bone morphology in patients with CKD. It is one measure of the skeletal component of the systemic disorder of CKD-MBD that is quantifiable by histomorphometry of bone biopsy.

From Moe S, Drüeke T, Cunningham J, et al. Definition, evaluation, and classification of renal osteodystrophy: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int.* 2006;69:1945–1953.

Approximately 60% to 70% of dietary Pi is absorbed by the gastrointestinal tract, predominantly in the small intestine, although transport can occur in all intestinal segments. Pi absorption occurs via passive sodium-independent transport and active sodium-dependent transport. Although it can vary depending on the study and experimental design, active transport is approximately 50% of total transport (Fig. 52.1).⁸ Passive transport occurs down electrochemical gradient through paracellular tight junctions; claudins and occludins appear to be involved and control transport rates and ion specificity.⁹ Active absorption occurs via the epithelial brush border type II (solute carrier A34) transporters, specifically the sodium-Pi cotransporter (NaPi-IIb) using energy from the basolateral sodium-potassium ATPase transporter. Complete ablation of the NaPi-IIb gene in mice demonstrates that this transporter is responsible for 90% of sodium-dependent transport but only 50% of total intestinal Pi transport.¹⁰ In animals with CKD induced by adenine, ablation of the NaPi-IIb gene lowers serum Pi, with additional lowering by the Pi binder sevelamer, suggesting that both active transport and passive transport are important in CKD.¹¹ The NaPi-IIb transporter is predominantly stimulated by high dietary Pi and calcitriol.⁸ In addition, studies suggest that the phosphatonins, matrix extracellular phosphoglycoprotein,¹² and FGF-23¹³ may play

a role in intestinal transport. Tenapanor, an inhibitor of the gut sodium/hydrogen exchanger, also affects Pi transport in the intestine. Tenapanor decreases paracellular absorption by increasing resistance to Pi transport at tight junctions and decreasing NaPi-IIb expression, thereby decreasing active Pi transport.¹⁴ However, dietary Pi appears the most important regulator of intestinal absorption.

The kidneys are responsible for maintaining Pi balance by excreting the net amount of Pi absorbed (see Chapter 7). Most inorganic Pi is freely filtered by the glomerulus with approximately 70% to 80% reabsorbed in the proximal tubule, which serves as the primary regulated site of the kidney. The remaining 20% to 30% is reabsorbed in the distal tubule. Pi transport across the apical lumen occurs via an active transport process that is driven by active sodium transport on the basolateral side by the sodium-potassium adenosine triphosphatase (Na⁺-K⁺-ATPase). The primary transporters on the luminal surface are NaPi-IIa (SLC34A1) and NaPi-IIc (SLC34A3), with a minor component via the type III sodium-dependent Pi cotransporter Pit-2 (SLC20A2). PTH and FGF-23 both downregulate these NaPi transporters but through different signaling mechanisms. FGF-23 stimulates endocytosis of the transporters after signaling through the FGF receptor (FGFR)–klotho complex, described later. PTH, after binding to PTHR1 receptor, leads to increased cyclic




Fig. 52.1 Intestinal phosphate transport. Approximately 50% of phosphate (Pi) transport is sodium (Na⁺) dependent, due to active transport, and regulated by a number of factors. The remaining phosphate transport is sodium independent and due to paracellular or transcellular transport. *FGF-23*, Fibroblast growth factor 23; *MEPE*, matrix extracellular phosphoglycoprotein; *Na⁺/K⁺ ATPase*, sodium-potassium adenosine triphosphatase; *NaPi-IIb*, the sodium-Pi cotransporter. (From Lee GJ, Marks J. Intestinal phosphate transport: a therapeutic target in chronic kidney disease and beyond? *Pediatr Nephrol*. 2015;30:363–371. With permission.)

adenosine monophosphate/protein kinase A (cAMP/PKA) signaling with the scaffolding protein $\text{Na}^+\text{-H}^+$ exchanger regulatory factors 1 and 3 (NHERF1 and NHERF3) playing an important role. Pi has been shown to stimulate FGF23 release from osteocytes via induction of kidney-specific glycolysis and synthesis of glycerol-3-phosphate (G-3-P) by the proximal tubule.¹⁵ The enzyme required for glycolysis, G-3-P dehydrogenase, is Pi dependent and reabsorption of Pi via NaPi-IIa is critical for the G-3-P production. FGF-23 also decreases calcitriol production by reducing the conversion of 25(OH)D to calcitriol and increasing its catabolism. The reduction in calcitriol results in further reduction of intestinal Pi absorption.

CALCIUM BALANCE AND HOMEOSTASIS

Serum calcium concentrations are tightly controlled within a relatively narrow range, usually 8.5 to 10.5 mg/dL (2.1 to 2.6 mmol/L). However, the serum calcium concentration is a poor reflection of overall total body calcium because serum concentrations account for <1% of total body calcium; the remainder of total body calcium is stored in bone. Ionized calcium, generally 40% of total serum calcium, is physiologically active, whereas nonionized calcium is bound to albumin or anions such as citrate, bicarbonate, and Pi. In the presence of hypoalbuminemia, there is a relative increase in the ionized calcium relative to the total calcium; thus total serum calcium measurement may underestimate the physiologically active (ionized) serum calcium. A commonly used formula for estimating the ionized calcium from the total calcium value is to add 0.8 mg/dL for every 1-mg decrease in serum albumin below 4 mg/dL. However, in a study in patients with CKD stages 3 to 5 not undergoing dialysis, total calcium concentration and albumin-corrected total calcium values failed to correctly classify 20% of patients as either hypocalcemic or hypercalcemic whose state was documented by ionized calcium measurement.¹⁶ The sensitivity to detect true hypocalcemia or hypercalcemia was only 40% and 21% for total serum calcium and 36% and 21% for albumin-corrected serum calcium, respectively.¹⁶ The primary causes of this discordance are thought to be albumin, PTH, and pH, although the latter has been questioned.¹⁷ Thus whenever possible, ionized calcium measurement should be used. Serum concentrations of ionized calcium are maintained within the normal range by the secretion of PTH, as discussed later.

In normal individuals, the net calcium balance (intake–output) varies with age. Children and young adults are usually in a slightly positive net calcium balance to enhance linear growth; beyond ages 25 to 35 years, when bones stop growing, calcium balance tends to be neutral. Individuals with normal kidney function have protection against calcium overload by virtue of their ability to increase renal excretion of calcium and reduce intestinal absorption of calcium through the actions of PTH and calcitriol. However, in CKD the ability to maintain normal homeostasis, including a normal serum ionized calcium level and appropriate calcium balance for age, is diminished, and lost completely when kidney function is absent. Two studies in patients with CKD stages G3b to G4 demonstrate that 1000 mg per day of dietary or calcium supplement/binder leads to near-neutral calcium balance (reviewed in [Chapter 17](#)).^{7,18} Concerns about inadequate calcium intake in the CKD population

contributing to worsening bone health have led to updated guidelines, emphasizing the need for 800 to 1000 mg per day of calcium in adults and age-appropriate normal calcium intake in children.¹⁹

Calcium absorption across the intestinal epithelium occurs via a vitamin D–dependent, saturable (transcellular) pathway and a vitamin D–independent, nonsaturable (paracellular) pathway. In states of adequate dietary calcium, the paracellular mechanism prevails, but the vitamin D–dependent pathways are critical in calcium-deficient states. The transcellular absorption occurs via three steps: 1. calcium enters from the lumen into the cells via transient receptor potential vanilloid (TRPV) channels, of which TRPV6 is most important in the intestine²⁰; 2. the intracellular calcium associates with calbindin-D9K to be “ferried” to the basolateral membrane; and 3. calcium is removed from the enterocytes predominantly via the calcium-ATPase, with the $\text{Na}^+\text{-Ca}^{2+}$ exchanger playing a minor role. The duodenum is the major site of calcium absorption, although the other segments of the small intestine and the colon also contribute to net calcium absorption. All of the key regulatory components of active calcium transport—TRPV, calbindin, the $\text{Ca}^{2+}\text{-ATPase}$ (PMCA1b), and the $\text{Na}^+\text{-Ca}^{2+}$ exchanger (NCX1)—are upregulated by calcitriol.²¹ Mice with intestinal knockdown of the vitamin D receptor (VDR) are still able to maintain normal calcium levels because of PTH-induced increased bone resorption; by contrast, global knockdown of VDR leads to hypocalcemia. Hypocalcemia with VDR knockdown can be corrected with either a high-calcium diet (and presumed intestinal paracellular calcium transport) or the administration of calcitriol.²² Thus bone and kidney can compensate for impaired responsiveness of the intestinal VDR, and diet alone can compensate for a total lack of vitamin D. The 1α -hydroxylase enzyme (CYP27B1) is also located throughout the intestine; to date, however, conversion of 25(OH)D to calcitriol has been identified only in the colonic epithelial cells in inflammation.²³

The renal transport of calcium is further detailed in [Chapter 7](#). In the kidney, the majority (60%–70%) of calcium is reabsorbed passively in the proximal tubule, a process driven by a transepithelial electrochemical gradient that is generated by sodium and water reabsorption. Paracellular movement of calcium occurs via pore-forming claudins 5, 10a, and 17.²⁴ In the thick ascending limb, another 10% of calcium is reabsorbed via paracellular transport. Calcium-sensing receptor (CaSR) activation in this segment inhibits calcium absorption. This paracellular reabsorption occurs via claudins 14, 16 (paracellin), and 19,²⁵ with genetic defects in claudins resulting in syndromes of hypercalciuria and hypomagnesemia.²⁶ However, the more regulated aspect of calcium reabsorption occurs via transcellular pathways in the distal convoluted tubule and connecting tubule ([Fig. 52.2](#)). The mechanism is similar to intestinal transport: Calcium enters these cells via TRPV5 calcium channels down electrochemical gradients. In the cells, calcium binds with calbindin-D28k and is transported to the basolateral membrane, where calcium is actively reabsorbed by the $\text{Na}^+\text{-Ca}^{2+}$ exchanger (NCX1), and the plasma membrane adenosine triphosphatase (ATPase; PMCA1b). As in the intestinal epithelial cell, calcitriol upregulates all of these transport proteins.²⁷ PTH has an indirect effect on renal calcium handling via stimulating the synthesis of




Fig. 52.2 Epithelial calcium active transport. The late part of the distal convoluted tubule (DCT) and connecting tubule (CNT) play an important role in fine-tuning renal excretion of Ca^{2+} . The epithelial Ca^{2+} channel (TRPV5) is primarily expressed apically in these segments and colocalizes with calbindin- $\text{D}_{28\text{K}}$ (28K), $\text{Na}^{+}/\text{Ca}^{2+}$ exchanger (NCX1), and the plasma membrane adenosine triphosphatase (ATPase; PMCA1b). Upon entry via TRPV5, Ca^{2+} is buffered by 28K and diffuses to the basolateral membrane, where it is released and extruded by a concerted action of NCX1 and PMCA1b. In addition, the basolateral membrane exposes a parathyroid hormone (PTH) receptor (PTHr) and the $\text{Na}^{+}/\text{K}^{+}$ -ATPase consisting of the α , β , and γ subunits. PTHr activation by PTH stimulates TRPV5 activity and entered Ca^{2+} can subsequently control the expression level of the Ca^{2+} transporters. At the apical membrane, there is a bradykinin receptor (BK2) that is activated by urinary tissue kallikrein (TK) to activate TRPV5-mediated Ca^{2+} influx. In the cell, entered Ca^{2+} acts as negative feedback on channel activity, and 28K plays a regulatory role by association with TRPV5 under low intracellular Ca^{2+} concentrations. Extracellular urinary klotho directly stimulates TRPV5 at the apical membrane by modification of the N-glycan, whereas intracellular klotho enhances $\text{Na}^{+}/\text{K}^{+}$ -ATPase surface expression, which in turn activates NCX1-mediated Ca^{2+} efflux. ADP, Adenosine diphosphate; DCT1, early part of distal convoluted tubule; PT, proximal tubule; TAL, thick ascending limb of Henle. (From Boros S, Bindels RJM, Hoenderop JGJ. Active Ca^{2+} reabsorption in the connecting tubule. *Pflügers Arch Eur J Physiol.* 2009;458:99–109.)

calcitriol and increases TRPV5 activity.²⁸ The enzymatic activity of circulating klotho has been shown to cleave the extracellular domain of TRPV5 channels, keeping these channels at the cell membrane and thereby facilitating calcium reabsorption.²⁹

CALCIUM-SENSING RECEPTOR

Physiologic studies in animals and humans in the 1980s demonstrated the rapid release of PTH in response to small reductions in blood-ionized calcium, lending support to the existence of a CaSR in the parathyroid gland that was subsequently cloned in 1993.³⁰ The CaSR was shown to belong to the superfamily of G protein-coupled receptors and is a glycosylated protein with a large extracellular domain, seven membrane-spanning segments, and a relatively large cytoplasmic domain. The primary ligand for the CaSR is Ca^{2+} , but it also senses other divalent and polyvalent cations including Mg^{2+} , Be^{2+} , La^{3+} , Gd^{3+} , and polyarginine.³¹ Extracellular calcium binds to multiple sites, leading to conformational changes that result in activation of phospholipases C, A_2 , and D, as well as inhibition of cAMP production.³² Activation of the CaSR stimulates phospholipase C, leading to an increase in inositol 1,4,5-triphosphate (IP_3), which mobilizes intracellular

calcium and decreases PTH secretion (Fig. 52.3). By contrast, inactivation of the CaSR reduces intracellular calcium and increases PTH secretion. CaSR messenger RNA (mRNA) is widely expressed in multiple tissues including organs responsible for CKD-MBD (parathyroid, kidney, thyroid, bone, intestine, vasculature). Studies have demonstrated a diverse role for the CaSR in disease including in the gastrointestinal tract, where it regulates gastrin, glucagon-like peptide-1, acid, and hormone secretion and is involved in taste, gastrointestinal fluid transport, and cell turnover.³³

$\text{CaSR}^{-/-}$ mice die shortly after birth owing to hypercalcemia, hypocalciuria, and hyperparathyroidism. If the PTH gene is also ablated, the mice survive and most organs appear histologically normal, with healing of bone mineralization defects but no change in hypocalciuria.³⁴ However, these $\text{CaSR}^{-/-}/\text{PTH}^{-/-}$ mice demonstrate hypercalcemia in response to oral calcium, infusion of PTH, or administration of calcitriol, whereas the $\text{CaSR}^{+/+}/\text{PTH}^{-/-}$ mice are able to decrease gastrointestinal calcium absorption and increase renal calcium excretion to maintain normal levels of serum calcium.^{35,36} These data indicate that CaSR activation corrects hypocalcemia by increasing PTH, whereas in hypercalcemia, the CaSR acts independent of PTH by increasing renal calcium excretion. In uremic




Fig. 52.3 Calcium-sensing receptor (CaR). Activation of the CaR by calcium stimulates phospholipase C, leading to increased inositol 1,4,5-trisphosphate (IP_3), which mobilizes intracellular calcium and inhibits parathyroid hormone (PTH) synthesis. A decrease in serum calcium (Ca^{2+}) inhibits intracellular signaling, leading to increased PTH synthesis and secretion. ER, Endoplasmic reticulum. (From Friedman PA, Goodman WG. PTH(1-84)/PTH(7-84): a balance of power. *Am J Physiol Renal Physiol*. 2006;290:F975–F984.)

animals, the expression of CaSR in the parathyroid gland is downregulated by a high-Pi diet and upregulated by magnesium³⁷ and calcimimetics. In parathyroid glands from patients with secondary hyperparathyroidism, the expression of the CaSR is downregulated in comparison with expression in nonuremic patients³⁸ but can be upregulated with the administration of calcimimetics.³⁹

The CaSR is expressed throughout the kidney, where it is found in diverse locations and performs multiple physiologic functions: podocyte (cytoskeleton changes); proximal tubule (phosphate reabsorption, calcitriol synthesis, acidification/fluid reabsorption); macula densa (renin secretion); thick ascending loop of Henle (calcium, sodium, potassium, and chloride handling); distal convoluted tubule/connecting tubule (calcium transport); and collecting duct (acid/base, water handling).⁴⁰ Diverse functions in the kidney signify how important calcium homeostasis is to normal kidney function and vice versa. Furthermore, many of these functions avoid renal calcium precipitation. Most notably, activation of the luminal CaSR in the collecting duct by elevated urine calcium values leads to urinary acidification and polyuria,⁴¹⁹ a common clinical symptom in hypercalcemia, and prevents calcium-Pi precipitation.

There is some disagreement as to the role of the CaSR in bone and whether this role depends on PTH. Clearly bone cells respond to calcium. The CaSR is important in fetal bone development; conditional deletion of the CaSR in early osteoblasts leads to altered bone phenotype, although the results vary depending on the construct used.⁴² Studies suggest that the CaSR modulates both bone resorption and formation induced by PTH.⁴³ In vitro, mesenchymal stem cells appear to require CaSR for differentiation,⁴⁴ but in more differentiated cells, calcium channels appear more involved in calcium transport.⁴⁵ An initial report found that calcium also regulates FGF-23 synthesis in bone, although this effect does not appear to be mediated via the CaSR.⁴⁶ Calcimimetics, allosteric activators of the CaSR, are used to treat secondary hyperparathyroidism as discussed in Chapter 62.


Data also suggest a role for the CaSR in vascular calcification. Immunohistochemical staining demonstrates expression of CaSR on normal human arteries with downregulation in areas of calcification.⁴⁷ The CaSR is expressed on cultured vascular smooth muscle cells (VSMCs), and calcimimetics inhibit in vitro calcification.^{47,48} The calcimimetic R-568 reverses calcitriol-induced arterial calcification⁴⁹ and inhibits proliferation of VSMCs and endothelial cells in the 5/6 nephrectomy rat model.⁵⁰ Calcimimetics also retard uremia-enhanced vascular calcification and atherosclerosis in the ApoE^{-/-} mouse⁵¹ and prevent arterial and myocardial calcification in the Cy/+ model of slowly progressive CKD-MBD.⁵² Calcimimetics also upregulate a potential local inhibitor of arterial calcification, matrix gla protein.⁵³ A trial of cinacalcet in patients with end-stage kidney disease (ESKD) treated with calcium-based phosphate binders showed amelioration in calcification relative to placebo (and active vitamin D analogs).⁵⁴ These data support a role for the CaSR in all three components of CKD-MBD.

HORMONAL REGULATION OF CHRONIC KIDNEY DISEASE—MINERAL AND BONE DISORDER

PARATHYROID HORMONE

The primary function of PTH is to maintain calcium homeostasis (Fig. 52.4) by 1. increasing bone mineral dissolution, thus releasing calcium and Pi; 2. increasing renal reabsorption of calcium and excretion of Pi; 3. increasing the activity of the renal CYP27B1 enzyme to convert 25(OH)D to calcitriol; and 4. enhancing the gastrointestinal absorption of both calcium and Pi indirectly through its effects on the synthesis of calcitriol. In healthy persons, the increase in serum PTH concentration in response to hypocalcemia effectively restores serum calcium and maintains serum Pi concentrations. The kidneys are critical to this normal homeostatic response; thus patients with more severe CKD may lose the capacity to maintain calcium homeostasis.

PTH is cleaved to an 84-amino acid protein in the parathyroid gland, where it is stored as fragments in secretory granules for release. Once released, the circulating 1–84-amino acid protein has a half-life of 2 to 4 minutes and is further metabolized in the liver and kidney. PTH secretion occurs in response to hypocalcemia, hyperphosphatemia, and calcitriol deficiency. The extracellular concentration of ionized calcium is the most important determinant of minute-to-minute secretion of PTH from stored secretory granules. The CaSR mediates the rapid response in PTH secretion, which occurs within seconds of changes in ionized calcium concentration. Inactivating mutations of the CaSR have been associated with neonatal severe hyperparathyroidism and benign familial hypocalciuric hypercalcemia. Affected patients have asymptomatic elevations of serum calcium in the presence of nonsuppressed PTH. Activating mutations have been found in patients with autosomal dominant hypocalcemia resulting in inhibition of PTH secretion at relatively lower serum calcium concentrations. PTH is released as both an intact (1–84) protein and carboxy (C)-terminal fragments (often called PTH [7–84]; see Fig. 52.3). C-terminal PTH has the opposite effect on calcium release from bone in animals and cultured calvariae from that of PTH with an intact N terminus.⁵⁵ In addition, the C-terminal PTH inhibits apoptosis in osteoblasts, whereas



Note: Items shown in blue are directly affected by decrease in renal function.

Fig. 52.4 Normalization of serum calcium by multiple actions of parathyroid hormone (PTH). Serum levels of ionized calcium (Ca) are maintained in the normal range by induction of increases in the secretion of PTH. PTH acts to increase bone resorption, renal calcium reabsorption, and the conversion of 25(OH)D to calcitriol in the kidney, thereby increasing gastrointestinal calcium absorption. The blue boxes indicate processes that are abnormal in chronic kidney disease, leading to altered calcium homeostasis. PO_4 , Phosphate. (From Moe SM. Calcium, phosphorus, and vitamin D metabolism in renal disease and chronic renal failure. In: Kopple JD, Massry SG, eds. *Nutritional Management of Renal Disease*. Philadelphia: Lippincott Williams & Wilkins; 2004:261–285.)

the N-terminal PTH induces apoptosis.⁵⁵ Regulators of PTH secretion act by changing the proportion of intact 1–84 and C-terminal fragments.

PTH binds to the PTH1 receptor (PTH1R), which is a member of the G protein–linked seven membrane–spanning receptor family and is widely expressed. PTH-related peptide (PTHrp) shares homology with the first few amino acids of PTH and also binds the PTH1R. Activation of the PTH1R stimulates heterodimeric G proteins G_s (leading to stimulation of cAMP and PKA signaling) and $G_{\alpha q}$ (leading to activation of IP_3 and protein kinase C), ultimately resulting in changes in intracellular calcium.⁵⁶ PTH1R activation may vary in response to time exposure, secondary conformational changes after binding, and which cell signaling mechanism is preferentially activated. In general, the effects of PTH are systemic and those of PTHrp are autocrine.

The relations among calcium, Pi, FGF-23, and calcitriol in the development of secondary hyperparathyroidism in CKD are complex and nearly impossible to fully evaluate in humans because changes in one lead to rapid changes in the others. The response to a decrease in ionized calcium mediated by the CaSR is likely the most potent stimulus for PTH release. Pi increases PTH production by enhancing the stability of PTH mRNA.⁵⁷ FGF-23 directly stimulates

PTH release in a klotho-independent manner.⁵⁸ Calcitriol suppresses PTH release via the VDR to lead to direct suppression of the gene. Other vitamin D compounds that bind to the VDR with lower affinity still reduce PTH release if given in high enough quantities.⁵⁹ Although PTH-induced signaling predominantly affects mineral metabolism, there are also many extraskelletal manifestations of PTH excess in CKD. These include encephalopathy, anemia, extraskelletal calcification, peripheral neuropathy, cardiac dysfunction, hyperlipidemia, bone and muscle pain, pruritus, and impotence.⁶⁰

In the kidney, PTH facilitates calcium reabsorption and Pi excretion, as noted earlier. In bone, PTH receptors are located on osteoblasts, with a time-dependent effect. PTH administered long term inhibits osteoblast differentiation and mineralization. By contrast, the administration of PTH to osteoblasts in a pulse rather than a continuous manner stimulates osteoblast proliferation, forming the basis for the administration of PTH as an anabolic therapy for osteoporosis.⁶¹ PTH also interacts with wnt/ β -catenin signaling, as discussed in the bone section.

VITAMIN D

Cholesterol is synthesized to 7-dehydrocholesterol, which in turn is metabolized in the skin to vitamin D_3 (Fig. 52.5). This reaction is facilitated by ultraviolet light (ultraviolet B) and higher temperature and is therefore reduced in individuals with high skin melanin content and inhibited by sunscreen containing sun protection factor 8 or higher. In addition, there are dietary sources of vitamin D_2 (ergocalciferol) and vitamin D_3 (cholecalciferol). The difference between D_2 (plant source) and D_3 (animal source) compounds is the presence of a double bond (D_2) between carbon numbers 22 and 23 in the side chain. Once in the blood, both D_2 and D_3 bind with vitamin D–binding protein (DBP) and are carried to the liver, where they are hydroxylated by CYP27A1 (25-hydroxylase) in an essentially unregulated manner to yield 25(OH)D, often called calcidiol. Once they are converted to calcidiol, there appears to be no difference between the biologic activities of D_2 and D_3 . Calcidiol is then converted in the kidney (or other cells) to calcitriol by the action of CYP27B1. This active metabolite is also degraded by other kidney enzymes, 24,25-hydroxylase (CYP24A1), and CYP3A4, providing the primary metabolism of the active compound.

Vitamin D–binding protein is a 58-kDa protein synthesized in the liver. Its serum levels in humans are between 4 and 8 mM, and the protein has a half-life of 3 days. Both the parent vitamin D, 25(OH)D, and calcitriol are carried in the circulation by DBP, but its greater affinity is for 25(OH)D. Targeted gene disruption studies show that DBP-null mice have a marked reduction in both circulating and tissue distributions of calcitriol and yet are normocalcemic, indicating that the primary role of DBP is to maintain stable serum stores of vitamin D metabolites.⁶² At the cellular level, both 25(OH)D and calcitriol are endocytosed. Inside the cell, calcitriol can be inactivated by mitochondrial 24-hydroxylase (CYP24A1) or can bind to the VDR in the cytoplasm. Once the VDR–ligand binding has occurred, the VDR translocates to the nucleus, where it heterodimerizes with the retinoid X receptor. This complex binds the vitamin D response element of target genes and




Fig. 52.5 Overview of vitamin D metabolism. Vitamin D is obtained from dietary sources and is metabolized via ultraviolet light (ultraviolet B [UVB]) from 7-dehydrocholesterol in the skin. Both sources (diet and skin) of vitamin D₂ and vitamin D₃ bind to vitamin D-binding protein (DBP) and circulate to the liver. In the liver, vitamin D is hydroxylated by CYP27A1 (25-hydroxylase) to 25(OH)D, commonly referred to as calcidiol. Calcidiol is then further metabolized to calcitriol by the 1 α -hydroxylase enzyme (CYP27B1) at the level of the kidney. The active metabolite 1,25(OH)₂D (calcitriol) acts principally on the target organs of intestine, parathyroid (PTH) gland, bone cell precursors, and the kidney. Calcitriol is metabolized to the inert 1,24,25(OH)₃D through the action of the 24,25-hydroxylase enzyme (CYP24). Calcidiol is similarly hydroxylated to 24,25(OH)₂D. (Modified from Moe SM. Renal osteodystrophy. In: Pereira BJJ, Sayegh M, Blake P, eds. *Chronic Kidney Disease: Dialysis and Transplantation*. 2nd ed. Philadelphia: Elsevier Saunders; 2004.)

recruits transcription factors and corepressors/coactivators that modulate the transcription.⁶³ These corepressors and coactivators appear to be specific for the ligand, and thus different forms and analogs of vitamin D may produce different effects at each tissue, forming the basis for the pharmacologic development of analogs. Degradation of calcitriol is believed to occur principally in the kidney, from side cleavage and oxidation, to form 24,25(OH)₂D.⁶³

It is generally accepted that the major source of the circulating levels of calcitriol in the kidney. There is evidence that both 25(OH)D and calcitriol have local tissue effects because the VDR, CYP27B1, and CYP24A1 are found in many cells throughout the body.⁶⁴ While the kidneys are the predominant site of conversion of 25(OH)D to calcitriol, there is evidence of conversion in other organs, associated with CYP27B1 expression and/or activity in normal and abnormal cells. Other cell types active in conversion of 25(OH)D to calcitriol include osteoblasts, breast epithelial cells (normal and cancerous), prostate gland (normal and cancerous), alveolar and circulating macrophages, pancreatic islet cells, synovial cells, and arterial endothelial cells. Some of these cells may directly take up calcitriol, and others may endocytose the DBP-25(OH)D complex in a megalin-mediated manner (Fig. 52.6), after which the 25(OH)D is hydroxylated by CYP27B1 to act on that specific cell. The presence of CYP24A1 in cells also indicates that the

metabolism of calcitriol may be regulated at a cellular level. Circulating concentrations of 25(OH)D are 1000 times higher than those of calcitriol, and thus 25(OH)D exhibits a local, or autocrine/paracrine, effect on many cell types.⁶⁴

Circulating calcitriol mediates its cellular function via nongenomic and genomic mechanisms. Calcitriol facilitates the uptake of calcium in intestinal and renal epithelium by increasing the activity of the voltage-dependent calcium channels TRPV5 and TRPV6. Calcitriol then enhances the transport of calcium through and out of cells, by upregulating the calcium transport protein calbindin (calbindin-D9k in intestine, and calbindin-D28k in kidney) and the basolateral calcium-ATPase as detailed earlier. The CYP27B1 in the kidney is the site of regulation of calcitriol synthesis by numerous other factors including serum calcium, serum Pi, estrogen, prolactin, growth hormone, FGF-23, and calcitriol itself. Studies show that FGF-23 and inflammatory mediators such as interferon regulate CYP27B1 at nonrenal sites.⁶⁵ In vitamin D knockout animals, parathyroid gland hyperplasia is consistently observed despite normalization of serum calcium. However, gland growth can be blunted by exogenous administration of calcitriol even in the absence of VDR, demonstrating a role for calcitriol in regulation of parathyroid gland growth.⁶⁶ In the rat, a single small dose of calcitriol decreases PTH secretion by nearly 100%. Studies in the 1970s demonstrated that oral calcitriol, but


ROLE OF EXTRARENAL 1α -HYDROXYLASE

Fig. 52.6 Concept of the role of the extrarenal 1α -hydroxylase. The metabolism of vitamin D in the context of the cells involved is shown. *Upper right*, Proximal tubular cell showing the key elements in the uptake of 25-(OH)D_3 and its conversion to $1\alpha,25\text{-(OH)}_2\text{D}_3$. Megalin/cubilin are cell surface receptors that execute endocytosis of the vitamin D-binding protein (DBP)- 25-(OH)D_3 complex, and CYP27B1 is the main component of the 1α -hydroxylase, responsible for making $1\alpha,25\text{-(OH)}_2\text{D}_3$. *Middle left*, Simple target cell that takes up $1\alpha,25\text{-(OH)}_2\text{D}_3$ as the free ligand originally ferried to the target cell bound to DBP. The picture shows the key elements of the transcriptional machinery, as well as some representative gene products, including the cell division protein p21, the bone matrix protein osteopontin, the calcium transport protein calbindin, and the autoregulatory protein CYP24A1. *Lower right*, Target cell expressing extrarenal 1α -hydroxylase, which possesses megalin/cubilin machinery to take up the DBP- 25-(OH)D_3 complex and also expresses CYP27B1, enabling it to make $1\alpha,25\text{-(OH)}_2\text{D}_3$ intracellularly and also to respond in a likewise manner to the simple target cell because it also possesses the vitamin D receptor (VDR) and other transcriptional machinery. The expectation is that cells involved in cell differentiation or in control of cell division require higher concentrations of $1\alpha,25\text{-(OH)}_2\text{D}_3$ in order to modulate a different set of genes and that the CYP27B1 boosts local production to augment “circulating” $1\alpha,25\text{-(OH)}_2\text{D}_3$ arriving from the kidney in the bloodstream. With normal physiologic processes, locally produced $1\alpha,25\text{-(OH)}_2\text{D}_3$ would not enter the general circulation, although in pathologic conditions (e.g., sarcoidosis) it might. At this time, it is not clear how many cell types can be considered simple target cells and how many possess the CYP27B1 and megalin/cubilin to allow for local production of hormones. mRNA, Messenger RNA; RXR, retinoid X receptor. (From Jones G. Expanding role for vitamin D in chronic kidney disease: importance of blood 25-OH-D levels and extra-renal 1α -hydroxylase in the classical and nonclassical actions of $1\alpha,25\text{-dihydroxyvitamin D}_3$. *Semin Dial.* 2007;20:316–324. With permission.)

not the precursor hormone vitamin D_3 , suppressed PTH in patients undergoing dialysis, leading to widespread use of calcitriol or its analogs in the management of secondary hyperparathyroidism. However, later studies in animals have demonstrated efficacy of 25-(OH)D in suppression of PTH, but the doses required are much greater than for calcitriol. Studies in humans have shown efficacy of 25-(OH)D in suppressing PTH in patients with advanced CKD, but direct comparison studies are lacking.⁶⁷

Calcitriol has multiple effects on many cells that are important in bone remodeling; therefore it is not surprising that bone defects are well described in vitamin D-deficient states. However, the direct effects of the vitamin D system on bone have been difficult to differentiate from the secondary effects of hypocalcemia and hyperparathyroidism in vitamin D-deficient models. Transgenic animals including 1α -hydroxylase^{-/-}/VDR^{-/-} and 1α -hydroxylase^{-/-}/VDR^{-/-}, have impaired bone mineralization. In these animals, mineralization can be corrected with normalization of serum

calcium; even exogenous calcitriol does not fully correct mineralization in the 1α -hydroxylase^{-/-} animals unless serum calcium concentrations are also restored. Studies evaluating bone remodeling also demonstrate an important role for the calcitriol/VDR system. If hypocalcemia is not corrected (leading to secondary hyperparathyroidism), there is increased osteoblast activity and bone formation from the anabolic effects of PTH. The activation of osteoclasts by PTH is blunted, suggesting a synergistic effect of vitamin D and PTH. Supporting this suggestion is the finding that when serum calcium concentrations are corrected by “rescue” diets and secondary hyperparathyroidism is prevented, osteoblast numbers, mineralization activity, and bone volume are still reduced. Comparison studies of 1α -hydroxylase^{-/-} and PTH^{-/-} mice demonstrate a predominant role for PTH in appositional bone growth and for vitamin D in endochondral bone formation. Thus the calcitriol/VDR system has anabolic bone effects that are necessary for bone formation and are supplemental to the effect of PTH.⁶⁸

FGF-23 AND KLOTHO

“Phosphatonins” are circulating factors that regulate urinary Pi excretion. Two main phosphatonins have been described: FGF-23 and matrix extracellular phosphoglycoprotein. Various forms of rickets have now been found to be due to abnormalities in FGF-23. Autosomal dominant hypophosphatemic rickets is rare and is associated with a mutation that limits normal degradation of FGF-23. Autosomal recessive hypophosphatemic rickets is also rare and is due to a mutation in dentin matrix protein (DMP), a locally produced inhibitor of FGF-23. X-linked hypophosphatemic rickets is the most common form of rickets due to a mutation in *PHEX* (phosphate-regulating gene with homologies to endopeptidases located on the X chromosome). Mutations in *PHEX* have been found to have deficient degradation of FGF-23 in the osteocyte, leading to inappropriately high serum concentrations of FGF-23.⁶⁹ Thus what previously was thought to be disorders of different etiologies are now all linked to FGF-23.

FGF-23 is a 251-amino acid hormone predominantly produced from bone cells (osteocytes and osteoblasts) during active bone remodeling, but its mRNA is also found in the heart, liver, thyroid/parathyroid, intestine, and skeletal muscle.⁷⁰ FGF-23 production in the osteocyte is stimulated by PTH⁷¹ and calcitriol.⁷² Elevated Pi or Pi load and hypercalcemia may also stimulate FGF-23, but this stimulation appears to be indirect. In the osteocyte, both DMP1 and PHEX proteins degrade FGF-23 such that

mutations in their corresponding genes lead to excess FGF-23.⁶⁹ In turn, calcitriol increases PHEX and FGF-23 inhibits calcitriol, completing a feedback loop. Regulation of NaPi-IIa by FGF-23 is independent of PTH; FGF-23 also inhibits the conversion of 25(OH)D to calcitriol by inhibition of CYP27B1 in the renal tubules⁷³ and at extrarenal sites,⁶⁵ and it increases catabolism of calcitriol by activation of CYP24,⁷³ leading to hypophosphatemia and inappropriately normal or low serum calcitriol concentrations. An overview of the FGF-23–klotho axis is shown in Fig. 52.7.

FGF-23 is a member of a diverse family of 18 FGFs that bind to one of four receptors (FGFRs) via a heparan sulfate cofactor— or klotho coreceptor—dependent manner, leading to diverse biologic effects.⁷⁴² Identification of klotho as a coreceptor for FGF-23 was due to nearly identical phenotypes of the knockout mice including hyperphosphatemia, hypercalcemia, and excess calcitriol levels associated with early mortality, growth retardation, vascular calcification, cardiac hypertrophy, and osteopenia.⁷⁵ Klotho was originally identified as an aging suppressor gene. α -Klotho is expressed in the kidney, skin, choroid plexus, pancreas, and parathyroid gland and forms complexes with FGFR1 and FGFR4 to enhance FGF-23 signaling. β -Klotho is expressed in the liver and fat, forms complexes with FGFR1 and FGFR4, and supports FGF-15/19 and FGF-21 signaling. γ -Klotho increases FGF-19 activity and is expressed in the eye, fat, and kidney. All three klothos are transmembrane proteins, with short intracellular domains and large extracellular domains




Fig. 52.7 The bone–kidney–parathyroid endocrine axes mediated by fibroblast growth factor 23 (FGF-23) and klotho. Active form of vitamin D (calcitriol) binds to vitamin D receptor (VDR) in the bone (osteocytes). The ligand-bound VDR forms a heterodimer with a nuclear receptor (RXR) and transactivates expression of the FGF-23 gene. FGF-23 secreted from bone acts on the klotho–FGF receptor (FGFR) complex expressed in the kidney (the bone–kidney axis) and parathyroid gland (the bone–parathyroid axis). In the kidney, FGF-23 suppresses synthesis of active vitamin D by downregulating expression of the *Cyp27b1* gene and promotes its inactivation by upregulating expression of the *Cyp24* gene, thereby closing a negative feedback loop for vitamin D homeostasis. In the parathyroid gland, FGF-23 suppresses production and secretion of parathyroid hormone (PTH). PTH binds to the PTH receptor (PTHr) expressed on renal tubular cells, leading to upregulation of *Cyp27b1* gene expression. Thus suppression of PTH by FGF-23 reduces expression of the *Cyp27b1* gene and serum levels of calcitriol. This step closes another long negative feedback loop for vitamin D homeostasis. (From Kuro-o M. Overview of the FGF23–Klotho axis. *Pediatr Nephrol*. 2010;25:583–590. With permission.)

that have β -glucosidase cleavage sites. In animals, α -klotho expressed in the distal tubule can be cleaved to release the extracellular domain into the circulation. Further, alternative splicing of the α -klotho gene produces only the extracellular domain that is released from cells and is called circulating or soluble klotho.⁷⁶ The soluble α -klotho acts as a coreceptor at nonrenal sites and prevents FGF-23–induced cardiac hypertrophy.⁷⁵ The kidney regulates both renal production of soluble klotho and renal excretion.⁷⁶

In the kidney, tissue klotho is downregulated⁷⁶ and FGF-23 is upregulated early in the course of CKD.⁷⁷ Klotho is shed from the distal convoluted tubule to serve as a coreceptor with FGF-23 on the proximal tubule, where it inhibits reabsorption of Pi that is regulated by NaPi-IIa, NaPi-IIc, and Pit-1 (similar to PTH but via different signaling mechanisms) and suppresses CYP27B1 to inhibit calcitriol production (opposite of PTH).⁷⁸ α -Klotho also increases calcium reabsorption by stimulating TRPV5,⁷⁹ decreases potassium excretion through effects on the ROMK1 (renal outer medullary potassium 1) channel,⁸⁰ protects against kidney injury and fibrosis,⁸¹ and decreases insulin resistance.⁷⁸ Table 52.2 demonstrates the parallel changes in klotho deficiency and CKD.

In recent years a two-way relationship between iron deficiency and FGF23 has been described. Initially identified in patients with autosomal dominant hypophosphatemic rickets in whom iron deficiency exacerbated hypophosphatemia,⁸² iron deficiency has now been confirmed to increase FGF23 production.⁸³ Because this rise is typically balanced by increased cleavage of FGF23, perturbations in serum Pi are not classically noted. Severe hypophosphatemia can result when the balance of production and cleavage is disturbed, such as with administration of specific IV iron formulations (ferric caboxymaltose) that may impair FGF23 cleavage by mechanisms that are not yet clear.⁸⁴ Inflammation and erythropoietin administration are also potent drivers of FGF23 production, but these factors are classically coupled with increased cleavage.⁸⁵ In CKD, the balance of production and cleavage is disturbed, with a concomitant decrease in the ability of the kidney to excrete Pi.⁸⁶ Notably, FGF23 itself can promote inflammation, iron deficiency, and anemia via secretion of inflammatory cytokines through FGFR4 activation.⁸⁷ In a mouse model of CKD, inhibition of FGF23 signaling reverses iron deficiency, anemia, and markers of inflammation.⁸⁸

In addition to klotho's role in mineral metabolism, klotho and FGF-23 are implicated in cardiovascular disease. FGF-23 can induce cardiac hypertrophy and increases intracellular calcium in a klotho-independent manner.^{75,89,90} In humans, diminished kidney function and the associated inflammation may be required to manifest the negative cardiac effects of FGF23.⁹¹ Klotho is localized in the heart at the sinoatrial node, and klotho deficiency may lead to arrhythmias.⁹² Klotho suppresses cardiomyocyte apoptosis⁹³ and downregulates TRPC6 (transient receptor potential cation 6) calcium channel in the cardiomyocyte.⁹⁴ Both klotho and FGFR1 and FGFR3, but not FGF-23 and FGFR4, are expressed in human arteries.^{95,96} In human arteries from patients with CKD, klotho, FGFR1, and FGFR3 are downregulated in the presence of calcification. VDR activators upregulate klotho, leading to an anticalcific effect on FGF-23-induced calcification.⁹⁶ Decreased klotho impairs

Table 52.2 Comparison of Phenotypes of Klotho Deficiency and Chronic Kidney Disease

	Klotho Deficiency	Chronic Kidney Disease
Blood Chemistry		
Phosphate	↑↑↑↑	↑ or ↑↑↑ ^a
Calcium	↑	↔ or ↓↓
Creatinine	↑	↑↑↑
Calcitriol	↑↑↑	↓↓↓
Parathyroid hormone	↔ or ↓	↑↑
Fibroblast growth factor 23	↑↑↑	↑↑
Klotho	↓↓↓ or disappears	↓↓ at ESKD ^b
Gross Phenotypes		
Body weight	↓↓↓	↓↓
Growth retardation	↓↓↓	↓↓ in children
Physical activity	↓↓↓	↓
Fertility	↓↓↓	↓↓
Life span	↓↓↓	↓↓
Cardiovascular Disease		
Cardiac hypertrophy	↑↑	↑↑↑
Cardiac fibrosis	↑↑	↑↑↑
Vascular calcification	↑↑↑↑	↑↑
Atherosclerosis	↑↑	↑↑↑
Blood pressure	↑	↑↑↑↑
Hematocrit levels	↓	↓↓↓
Bone disease	↓↓↓	↓↓↓

^aDuring early chronic kidney disease, blood phosphate level is in the normal range.

^bBlood klotho may be increased in early chronic kidney disease. ESKD, End-stage kidney disease; ↓, decreases; ↑, increases; ↔, unchanged.

Modified from Hu MC, Kuro-o M, Moe OW. Renal and extrarenal actions of Klotho. *Semin Nephrol*. 2013;33:118–129.

endothelial function.⁹⁷ Activation of the renin–angiotensin–aldosterone system reduces renal klotho expression.⁹⁸ In summary, discovery of the many systemic effects of FGF-23 and klotho has revolutionized our understanding of CKD-MBD.

BONE BIOLOGY

The majority of the total body stores of calcium and Pi is located in bone, and therefore bone plays an integral role in homeostasis. Trabecular (cancellous) bone is located predominantly in the epiphyses of the long bones, is 15% to 25% calcified, and serves a metabolic function, with a relatively short turnover time as shown by calcium⁴⁷ studies. By contrast, cortical (compact) bone is located in the shafts of long bones and is 80% to 90% calcified. This bone serves primarily a protective and mechanical function and has a calcium turnover time of months. Bone consists principally (90%) of highly organized cross-linked fibers of type I collagen; the remainder consists of proteoglycans and

“noncollagen” proteins such as osteopontin, osteocalcin, osteonectin, and alkaline phosphatase. Hydroxyapatite— $\text{Ca}_{10}(\text{PO}_4)_6(\text{OH})_2$ —is the primary bone crystal.

The cellular components of bone are cartilage cells, which are critical to bone development; osteoblasts, which are the bone-forming cells; and osteoclasts, which are the bone-resorbing cells. Osteoblasts are derived from progenitor mesenchymal cells located in the bone marrow. They are then induced to become osteoprogenitor cells, then endosteal or periosteal progenitor cells, and then mature osteoblasts. The control of this differentiation pathway is complicated and involves integration of circulating hormones, locally produced factors from the mesenchymal-hematopoietic cell niche, and transcription factors. Once bone formation is complete, osteoblasts may undergo apoptosis or may become quiescent cells trapped within the mineralized bone in the form of osteocytes.⁹⁹ The osteocytes are interconnected through a series of canaliculi and serve as mechanoreceptors. Osteocytes detect and respond to mechanical loading and initiate bone remodeling by regulating local osteoclastogenesis via paracrine signals. Osteoclasts are derived from hematopoietic precursor cells that differentiate and are signaled to arrive at a certain place in the bone through the osteoprotegerin (OPG)/RANKL (receptor activator of nuclear factor κB ligand) system detailed later. Once there, they fuse to form the multinucleated cells known as osteoclasts, which become highly polarized, reabsorbing bone through the release of derivative enzymes. These cells move along a resorption surface via changes in the cytoskeleton. PTH, cytokines, and calcitriol are all important in inducing the fusion of the committed osteoclast precursors.

The control of bone remodeling is highly complex but appears to occur in distinct phases, as follows: 1. osteoclast recruitment and activation, 2. osteoclast resorption, 3. preosteoblast migration and differentiation, 4. osteoblast deposition of matrix (osteoid or unmineralized bone), 5. mineralization, and 6. quiescence. At any one time, <15% to 20% of the bone surface is undergoing remodeling, and this process in a single bone remodeling unit can take 3 to 6 months.¹⁰⁰ How or why a certain segment of bone undergoes a remodeling cycle is not completely clear. The three main systems that interact to regulate remodeling are OPG/RANKL, sclerostin/Wnt/ β -catenin, and PTH/PTHrP, which are discussed separately.

OPG/RANKL

The identification of the OPG and RANK system in the 1980s sheds new light on the control of osteoclast function and the long-observed coupling of osteoblasts and osteoclasts. RANK is located on osteoclasts, and RANK ligand (RANKL) is secreted by osteoblasts. Osteoblasts also synthesize the decoy protein OPG, which can bind to RANK ligand and inhibit the subsequent binding of RANKL to RANK on osteoclasts, thus inhibiting bone resorption (Fig. 52.8). Alternatively, if OPG production is decreased, RANKL can bind with RANK on osteoclasts and induce osteoclastic bone resorption. This control system is regulated by nearly every cytokine and hormone thought important in bone remodeling including PTH, calcitriol, estrogen, glucocorticoids, interleukins, prostaglandins, and members of the transforming growth factor- β superfamily of cytokines.¹⁰¹ OPG has been successful in preventing bone resorption in models of osteoporosis, as




Fig. 52.8 Role of osteoprotegerin (OPG)/receptor activator of nuclear factor κB ligand (RANKL) in bone remodeling. Mechanisms of action for OPG, RANKL, and RANK (receptor activator of nuclear factor κB) are depicted in this diagram. RANKL is produced by osteoblasts, bone marrow stromal cells, and other cells under the control of various proresorptive growth factors, hormones, and cytokines. Osteoblasts and stromal cells produce OPG, which binds to and thereby inactivates RANKL. The major binding complex is likely to be a single OPG homodimer interacting with high affinity with a single RANKL homotrimer. In the absence of OPG, RANKL activates its receptor, RANK, found on osteoclasts and preosteoclast precursors. RANK–RANKL interactions lead to preosteoclast recruitment, fusion into multinucleated osteoclasts, osteoclast activation, and osteoclast survival. Each of these RANK-mediated responses can be fully inhibited by OPG. *CFU-M*, Macrophage colony-forming unit. (From Kearns AE, Khosla S, Kostenuik PJ. Receptor activator of nuclear factor κB ligand and osteoprotegerin regulation of bone remodeling in health and disease. *Endocr Rev*. 2008;29:155–192.)

well as hormone- and cytokine-induced bone resorption,¹⁰¹ and denosumab, an anti-RANKL antibody, is an approved anabolic drug for the treatment of osteoporosis.¹⁰² Interestingly, abnormalities in the OPG/RANKL system have been found in kidney disease,¹⁰³ and early animal models suggest that treatment with OPG may have a

protective role in hyperparathyroid bone disease.¹⁰⁴ Initial studies in patients receiving dialysis have demonstrated hypocalcemia as a severe adverse effect of denosumab as an inhibitor of osteoclastic resorption.¹⁰⁵ More information is required to understand how this system regulates bone remodeling in the context of CKD.




Fig. 52.9 Parathyroid hormone (PTH) and β -catenin signaling in bone remodeling. Osteocytes control bone formation through the secretion of the Wnt antagonists sclerostin (SOST) and Dickkopf Wnt signaling pathway inhibitor 1 (*dkk-1*), the expression of which is regulated by mechanosignals and by signaling of PTH and bone morphogenetic protein (BMP). PTH represses expression of these antagonists, whereas BMP signaling, which is mediated by BMP receptor 1A (*BMPR1A*), induces their expression. Moreover, Wnt signaling in osteocytes controls the production of osteoprotegerin (OPG), which is the decoy receptor for the key osteoclast differentiation factor RANKL (receptor activator of nuclear factor κ B ligand). Osteoblast-expressed WNT5a stimulates differentiation of osteoclast precursors as a result of binding to the FZD–ROR2 (frizzled and receptor tyrosine kinase–like orphan receptor 2) receptor complex. In a feedback loop for bone remodeling, osteoclasts stimulate the local differentiation of osteoblasts at the end of the resorption phase by secreting Wnt ligands. In addition, activation of parathyroid hormone 1 receptor (*PTH1R*)-mediated signaling in osteoblasts and osteocytes leads to stabilization of β -catenin and thus activation of Wnt signaling. LRP5/6, Low-density lipoprotein (LDL) receptor–related proteins 5 and 6; PKA, protein kinase signaling. (From Baron R, Kneissel M. Wnt signaling in bone homeostasis and disease: from human mutations to treatments. *Nat Med.* 2013;19:179–192. With permission.)

SCLEROSTIN/ WNT/ β -CATENIN

Genetic defects in the gene *SOST* have been identified in rare bone disorders. Sclerostin, the protein product of this gene, binds to low-density lipoprotein receptor–related proteins 5 and 6 (LRP5/LRP6) on the osteocyte to competitively inhibit the binding of the protein Wnt (Fig. 52.9). Normally, Wnt binding to LRP5/LRP6 leads to stabilization of β -catenin (canonical pathway) and regulation of normal bone accrual via osteoblast differentiation. In the presence of sclerostin, the β -catenin is degraded, and mesenchymal stem cell differentiation to mature bone cells is inhibited. In animal models, sclerostin deletion enhances bone accrual,¹⁰⁶ and in early human trials, treatment with an antibody to sclerostin was found to be anabolic.^{107,108} Sclerostin concentrations are elevated in the blood and bone of patients with CKD¹⁰⁹ and bone in animals with CKD¹¹⁰ and are associated with decreased BMD and increased fracture risk.¹¹¹ Increased cardiovascular disease risk and mortality have been correlated with sclerostin levels in some studies,^{111,112} though not confirmed by others.¹¹³ The anabolic agent, romosozumab, an antisclerostin antibody, may be efficacious in the treatment of renal osteodystrophy.¹¹⁴ However, initial studies in animals found that romosozumab was not efficacious in the setting of elevated PTH, although it did improve bone volume when PTH was suppressed.¹¹⁵ Dickkopf-related protein 1 (dkk-1) also inhibits Wnt binding to LRP5/LRP6, and an antibody to this circulating inhibitor of Wnt signaling improved bone remodeling in a model of early CKD.¹¹⁶ In osteocytes, PTH directly suppresses sclerostin and dkk-1 secretion^{71,117} and thus inhibits the production of circulating inhibitors of Wnt signaling.

PTH/PTH1R

In bone, PTH binds to its receptor, PTH1R, and activates β -catenin signaling via multiple mechanisms (Fig. 52.9): 1. direct activation through cAMP signaling; 2. indirect activation via osteoclast activation, which then increases β -catenin activity in osteoblasts; and 3. by binding to LRP6 to activate LRP5/LRP6 signaling even in the absence of Wnt ligands.¹⁰⁶ Thus PTH can activate β -catenin through non-Wnt-mediated pathways and through pathways not regulated by sclerostin or dkk-1. There are also differences in responses to continuous and intermittent PTH exposures. Mice expressing a constitutively active PTH1R or animals receiving continuous infusion of PTH(1–84) (analogous to secondary hyperparathyroidism) also have Wnt-dependent remodeling with increased osteoclast bone resorption via the OPG/RANKL system, leading to osteoblast activation and β -catenin activation.^{118,119} Hyperphosphatemia also activates β -catenin signaling.¹²⁰ Thus in CKD with hyperphosphatemia and secondary hyperparathyroidism, there is activation of β -catenin by PTH-mediated inhibition of circulating inhibitors of Wnt signaling (sclerostin and dkk-1), PTH-mediated effects independent of Wnt signaling, and phosphorus-mediated effects, all leading to enhanced mesenchymal differentiation to osteoblasts, increased RANKL-induced osteoclast activation, and increased bone resorption.

PATHOPHYSIOLOGY OF VASCULAR CALCIFICATION

Vascular disease may be due to a variety of different pathologic processes in different arterial segments, all of which can be

calcified. Atherosclerotic disease is characterized by fibro-fatty plaque formation, and on the basis of autopsy data and animal models, calcification had been thought to occur late in the disease course. These plaques can protrude into the arterial lumen, leading to a filling defect on angiography (Fig. 52.10A). However, advances in imaging, especially intravascular ultrasonography, have demonstrated that atherosclerosis can also be a circumferential lesion (without an obstructed lumen) with calcification earlier in the course of the disease.¹²¹ The medial layer may also be affected in arteriosclerosis, leading to thickening commonly found in elastic arteries (Fig. 52.10B). In addition to the larger elastic arteries, smaller elastic arteries may be affected by medial thickening and calcification, classically described as Mönckeberg calcification or medial calcinosis. This condition is more common in advanced age and in patients with diabetes mellitus and/or CKD, and it is associated with all-cause and cardiovascular mortality in patients with diabetes with and without CKD, as well as in patients with CKD with or without diabetes.

Although initially believed to be related to spontaneous precipitation in the setting of high serum concentrations of calcium and Pi, vascular calcification is now known to be a tightly regulated process that resembles mineralization in bone, a process kept “in check” through the actions of inhibitors of calcification. The current hypothesis accepted by most investigators is that VSMCs dedifferentiate or transform to osteocyte/chondrocyte-like cells (Fig. 52.11). These cells then lay down an extracellular matrix of collagen and noncollagenous proteins and make matrix vesicles that attach to the extracellular matrix to initiate and propagate mineralization. This process is regulated by cells, extracellular matrix proteins, and inhibitors that may act locally or systemically. In advanced CKD, there is abnormal bone remodeling and reduced renal clearance of phosphate, generating a positive calcium and Pi balance that “feeds” the mineral composition of matrix vesicles and augments the ability of existing calcification to expand. The evidence for each of these steps is discussed in the following sections.

CELLULAR TRANSFORMATION

VSMCs, osteoblasts, chondrocytes, and adipocytes differentiate from mesenchymal precursors with normal differentiation and transformation (dedifferentiation), which are controlled by various transcription factors (Fig. 52.12). Expression of the osteoblast differentiation factor core-binding factor α -1 (Cbfa1), now called Runx-2, has been identified in the inferior epigastric artery of adults undergoing kidney transplantation¹²² and in sections from the brachial arteries of children undergoing dialysis.¹²³ Runx-2 is essential for normal bone development, in that Runx-2 knockout mice fail to form a skeleton.¹²⁴ Genetic techniques have confirmed that VSMCs give rise to osteochondrogenic-like cells in calcified blood vessels (as opposed to circulating cells).¹²⁵ Osteoclast-like cells can also be seen, more commonly in intimal lesions, and as in bone, they appear to arise from circulating precursors.¹²⁶

In vitro, VSMCs upregulate Runx-2 in response to elevated Pi mediated by the type III sodium-dependent Pi cotransporters Pit-1 and Pit-2.¹²⁷ In addition, VSMCs incubated with uremic serum (pooled from anuric patients




Fig. 52.10 Arterial calcification. Histologic differences between atherosclerotic, or intimal, calcification (A) and medial calcification (B). *Int.*, Internal.

undergoing dialysis), in comparison with normal serum, express Runx-2 and its downstream protein osteopontin via a non-Pi-mediated mechanism.¹²⁸ Excess calcium can also induce mineralization *in vitro*, and the effects of calcium are additive to those of increased Pi.¹²⁹ FGF-23 enhances Pi-induced vascular calcification in rat aortic rings and rat aorta VSMCs by promoting osteoblastic differentiation.¹³⁰ Most recently, calprotectin has been identified as a critical contributor to calcification in human VSMC and mouse aortic rings and was separately associated with cardiovascular outcomes and mortality in a diverse international cohort.¹³¹ Numerous traditional and nontraditional cardiovascular risk factors in CKD can induce the transformation of VSMCs into osteoblast-like cells, with subsequent calcification *in vitro* (see Fig. 52.12). Given these data, it is not surprising that arterial calcification is so common in patients with CKD.

In animal models of CKD, secondary hyperparathyroidism develops spontaneously with loss of kidney function and can be associated with vascular calcification.^{52,132,133} Unfortunately, it is difficult to distinguish between the effects of Pi and PTH. However, one study found that vascular calcification developed in nephrectomized animals achieving supraphysiologic PTH levels by infusion, regardless of the Pi intake.¹³⁴ In a similar study, Runx-2 was upregulated in animals in three of the four Pi/PTH intake groups—those fed normal PTH + high Pi, high PTH + high Pi, and high PTH + low Pi—indicating that both Pi and PTH may lead to Runx-2 upregulation in arteries. However, the animals given high PTH + low Pi did not demonstrate arterial calcification,¹³⁵ suggesting that Pi is needed to provide the substrate for calcification to progress.

Arterial calcification and mineralization of bone are inversely related. In animal models of excessive bone resorption, treatments aimed at decreasing bone remodeling by inhibition of osteoclast activity (i.e., bisphosphonates

and calcimimetics) have been found helpful in preventing vascular calcification in some¹³⁶⁻¹³⁸ but not all studies.¹³⁹ Correction of low-turnover bone disease also appears to improve arterial calcification in animals.^{140,141} The role of vitamin D has been controversial, but more recent data suggest that it is only when the circulating levels of calcitriol are increased (and thus induce hypercalcemia or hyperphosphatemia) that calcification is observed.¹⁴²

MATRIX VESICLES AND APOPTOSIS

In chondrocytes and osteoblasts, normal mineralization is initiated when matrix vesicles are released into the extracellular space from the cell surface via polarized budding with subsequent attachment to extracellular matrix proteins. Matrix vesicles are characterized by both their appearance as small (50–200 nm), electron-dense spherical particles on electron microscopy and the biochemical presence of calcium and Pi, alkaline phosphatase, and the membrane protein annexins. Matrix vesicles have been identified in nearly all forms of mineralization/calcification in human tissues including bone, cartilage, tendon, calciphylaxis, and atherosclerosis. Cultured VSMCs incubated with elevated concentrations of calcium and Pi release matrix vesicles into the media, and the presence of fetuin-A (AHSG or α -2-Heremans-Schmid glycoprotein), the circulating inhibitor of mineralization (see later), decreases calcium uptake of the matrix vesicles.¹⁴³ Collagenase digestion of VSMCs has been found to lead to two populations of matrix vesicles, a secreted form in the media that had high fetuin-A and low annexin II content and could not mineralize type I collagen, and cellular matrix vesicles that had low fetuin-A and high annexin II content and could mineralize.¹⁴⁴ Matrix vesicles are similar to exosomes, which are known to transfer microRNAs from cell to cell and thus may play a role in calcification.¹⁴⁵ MicroRNAs have been found




Fig. 52.11 Overview of the pathophysiology of vascular calcification. Normally, mesenchymal stem cells differentiate to adipocytes, osteoblasts, chondrocytes, and vascular smooth muscle cells (VSMCs). In the setting of chronic kidney disease (CKD), diabetes, aging, inflammation, and the presence of multiple other toxins, these VSMCs can dedifferentiate or transform to chondrocyte/osteoblast-like cells by upregulation of transcription factors such as runt-related transcription factor 2 (*RUNX-2*) and homeobox protein *MSX2*. These transcription factors are critical for normal bone development, and thus their upregulation in VSMCs is indicative of a phenotypic switch. These osteocyte/chondrocyte-like VSMCs then become calcified in a process similar to bone formation. The cells lay down collagen and noncollagenous proteins in the intima or media and incorporate calcium (Ca) and phosphorus (Pi) into matrix vesicles to initiate mineralization and further grow the mineral into hydroxyapatite. The overall positive calcium and phosphorus balance of most patients undergoing dialysis feeds both the cellular transformation and generation of matrix vesicles (MVs). In addition, the extremes of bone turnover in CKD (low and high turnover or adynamic and hyperparathyroid bone, respectively) increase the available calcium and phosphorus by altering the bone content of these minerals. Ultimately, whether an artery calcifies or not depends on the strength of the army of inhibitors (Is) standing by in the circulation (fetuin-A) and in the arteries (e.g., pyrophosphate [PPi], matrix gla protein [MGP], and osteopontin [OP]). (From Moe SM, Chen NX. Mechanisms of vascular calcification in chronic kidney disease. *J Am Soc Nephrol*. 2008;19:213–216. With permission.)

to regulate the phenotypic switch from contractile to synthetic VSMCs¹⁴⁶ and are involved in the regulation of arterial calcification in vitro in animal models.^{145,147} These results suggest that the cellular regulation of the content of matrix vesicles may regulate the type and mineralizing capacity of the vesicle.

In addition to matrix vesicles, apoptotic bodies can induce calcification in vitro in VSMCs. Apoptotic bodies stimulated by calcium-Pi crystals of approximately 1 μm or less in diameter cause a rapid rise in intracellular calcium concentration and apoptosis, an effect triggered by lysosomal degradation.¹⁴⁸ Apoptosis has been identified in arterial segments with calcification obtained from children with ESKD.¹²³ Activation of the DNA damage response by prelamin A accelerates arterial calcification.¹⁴⁹ Autophagy, a regulated process of cell survival, counteracts Pi-induced calcification by reducing matrix vesicle release.¹⁵⁰ By contrast, atorvastatin protects against calcification by inducing autophagy via suppression of the β -catenin

pathway.¹⁵¹ Thus cells appear to guard against calcification when able via a number of pathways.

INHIBITORS OF VASCULAR CALCIFICATION

Vascular calcification, although prevalent in patients with CKD and particularly in patients receiving dialysis, is not uniform. Approximately 20% (depending on the published series) of patients undergoing dialysis have no vascular calcification and continue to have no calcification on follow-up despite risk factors similar to those in patients who do have calcification. These data support the concept of calcification inhibitors. Knockout animal models have demonstrated that selective deletion of many genes leads to vascular calcification.¹⁵² These studies imply that mineralization (or calcification) of arteries will occur, at least in some species or individuals, unless inhibited. This concept, that the regulation of calcification in blood vessels occurs principally via inhibition rather than promotion, may also be true in bone.¹⁵³ Inhibitors can be circulating




Fig. 52.12 Factors that regulate pathways involved in the pathogenesis of vascular calcification. Multiple factors regulate each step of extraskeletal calcification. ACE, Angiotensin-converting enzyme; All, angiotensin II; Ca, calcium; LRP5, low-density lipoprotein (LDL) receptor-related protein 5; MGP, matrix gamma-carboxyglutamate (Gla) protein; miRNA, microRNA; OPG, osteoprotegerin; Pi, phosphate; pOPN, osteopontin; PPI, pyrophosphate; RAGE, advanced glycosylation end product receptor; RANKL, receptor activator of nuclear factor κ B ligand; ROS, reactive oxygen species; Runx2, runt-related transcription factor 2; TGF- β , transforming growth factor β ; VSMCs, vascular smooth muscle cells; Wnt, wingless-type MMTV integration site family member. (From Wu M, Rementer C, Giachelli CM. Vascular calcification: an update on mechanisms and challenges in treatment. *Calcif Tissue Int.* 2013;93:365–373. With permission.)

or locally produced and site specific. Three inhibitors that have been well characterized in the arterial calcification of CKD are fetuin-A, matrix γ -carboxyglutamate (Gla) protein (MGP), and OPG.

Many other inhibitors of calcification exist. In aggregate, the data discussed in this section support the diversity and abundance of naturally occurring inhibitors of calcification. Thus vascular calcification in CKD represents a state of increased calcification promoters and decreased calcification inhibitors.

Fetuin-A

Fetuin-A is a circulating inhibitor of calcification, abundant in plasma, and mainly produced by the liver in adults. The transcription and synthesis of fetuin-A are downregulated during inflammation, and thus it is also a reverse acute-phase reactant, like serum albumin and several other hepatic proteins. Fetuin-A binds to both calcium and Pi in the serum, forming small “calciparticles” that are removed through the reticuloendothelial system. Fetuin-A inhibits the de novo formation and precipitation of the apatite precursor mineral basic calcium-Pi but does not dissolve it once the basic calcium-Pi is formed.¹⁵⁴ Therefore fetuin-A could be viewed as acting as a host defense to “cleanse the blood” of unwanted calcium and Pi and to prevent undesirable calcification in the circulation without causing bone demineralization. Fetuin-A has been found in matrix vesicles from VSMCs, and its presence renders the vesicles incapable of mineralization.^{143,155} Fetuin-A is abundant in serum and is a major factor in the calcification propensity

of serum,¹⁵⁶ a measure of which has been shown to reflect all-cause mortality in patients with CKD.¹⁵⁷

Targeted disruption of fetuin-A leads to diffuse and profound soft tissue calcification and to arteriolar calcification of muscle, kidney, and lung but not large arteries.¹⁵⁸ When Ahsg^{-/-} mice were crossed with ApoE^{-/-} mice, the latter known to have increased cholesterol and atherosclerosis, both aorta and coronary artery calcification developed in the double-deficient Ahsg^{-/-}/ApoE^{-/-} mice with high-Pi diet alone and were accelerated by CKD.¹⁵⁹ Thus extensive and multisite arterial calcification in this animal model required genetic predisposition to atherosclerosis (Apo E^{-/-}), a genetic defect in an inhibitor of mineralization (Ahsg^{-/-}), and hyperphosphatemia that was further accelerated by CKD. These data support the redundancy of the inhibitor system, in which multiple local regulators compensate for the absence of the circulating inhibitor fetuin-A. In patients with CKD and ESKD, lower serum fetuin-A concentrations are associated with mortality.¹⁶⁰⁻¹⁶² Serum concentrations of fetuin-A in patients undergoing dialysis were found to be inversely correlated with coronary artery calcification assessed by spiral computed tomography (CT),¹⁶³ with carotid artery plaques,¹⁶⁴ and, in children undergoing dialysis, arterial stiffness.¹²³ Similar findings have been shown in CKD.¹²³ Fetuin-A deficiency in CKD is likely due to chronic inflammation, although gene polymorphisms may also play a role. It is also possible that there is inappropriate upregulation, leading to a relative deficiency of fetuin-A in the setting of elevated calcium and Pi.

MATRIX GAMMA-CARBOXYGLUTAMATE PROTEIN

Matrix Gla protein (MGP) is a vitamin K–dependent protein expressed in many tissues but highly expressed in arteries and bone, where it acts predominantly as a local regulator of vascular calcification. MGP knockout mice have excessive cartilage and growth plate mineralization and arterial medial calcification, resulting in early mortality.¹⁶⁵ In MGP-deficient mice, calcification depends on elastin fragmentation due to increased elastase production.¹⁶⁶ Warfarin use and/or nutritional vitamin K deficiency result in undercarboxylation of MGP and impaired function.¹⁶⁷ Warfarin use is also a known risk factor for calcific uremic arteriolopathy (also referred to as calciphylaxis) and can induce calcification in an animal model of CKD.¹⁶⁸ Serum concentrations of carboxylated MGP were found to be nearly undetectable in patients undergoing dialysis and in patients with atherosclerotic disease.¹⁶⁷ Furthermore, lower levels of carboxylated MGP were associated with coronary artery calcification, arterial stiffness, and higher serum Pi in patients undergoing dialysis.¹⁶⁹ Lower serum concentrations of relative carboxylated MGP, as well as vitamin K deficiency, were also found in 20 patients with calcific uremic arteriolopathy compared with matched controls.¹⁷⁰ Progression of coronary calcification was more pronounced in patients taking warfarin.¹⁷¹ Supplementation of vitamin K in patients undergoing dialysis can restore carboxylated MGP¹⁷² and perhaps prevent progression of calcification,¹⁷³ but studies to determine whether such supplementation reduces calcification and cardiovascular events remain small and inconclusive.¹⁷⁴

Pyrophosphate

Another naturally occurring inhibitor of mineralization is pyrophosphate, which inhibits the formation of calcium-Pi crystals *in vitro*. Pyrophosphate is produced by VSMCs and inhibits arterial calcification. Pyrophosphate is inhibited by tissue-nonspecific alkaline phosphatase (TNAP), and TNAP activity is increased in calcified arteries from uremic animals¹⁷⁵ and patients with stage 5 CKD.¹²³ Pyrophosphate is also inhibited by another enzyme, ectonucleotide pyrophosphate/phosphodiesterase I (NPPI). Children deficient in NPPI have infantile arterial calcification.¹⁷⁶ Circulating levels of pyrophosphate are decreased in patients undergoing dialysis¹⁷⁷ and are negatively associated with arterial calcification in patients with CKD.¹⁷⁸ The intraperitoneal administration of pyrophosphate reduced vascular calcification in a rodent model of CKD.¹⁷⁹

Osteoprotegerin

Osteopenia and arterial calcification develop in mice null for OPG, implying that OPG is an important direct inhibitor of vascular calcification, but it is not clear whether this development was due to abnormalities in bone¹⁸⁰ or a direct arterial effect. Studies in the low-density lipoprotein receptor null mice, a model of atherogenesis, demonstrated that the administration of OPG did not prevent atherosclerotic lesions but did prevent calcification of those lesions.¹⁸¹ Bone marrow and vessel wall OPG reduces atherosclerosis and calcification,¹⁸² via regulation of the procalcific effects of RANKL on VSMCs.¹⁸³ Such procalcification effects appear related to inflammation¹⁸⁴ and may explain mechanisms of calcification in areas of macrophage-laden atherosclerotic plaques.

INTEGRATED REGULATION OF PHOSPHORUS AND CALCIUM

The four hormones PTH, FGF-23, calcitriol, and klotho work together to maintain normal Pi and calcium homeostasis to achieve appropriate balance in the blood and urine of these ions to avoid extraskeletal calcification and ensure adequate availability of these ions for bone that is growing (modeling) or remodeling. A summary of the integrated physiologic response to hyperphosphatemia is depicted in Fig. 52.13. This response is a complex system of multiple integrated feedback loops and is easier to understand if broken into loops that regulate calcitriol, Pi, and calcium.

PTH-FGF-23–Calcitriol Loop

PTH and FGF-23 have similar effects in stimulating Pi excretion. However, these hormones differ in their effects on the vitamin D axis. PTH stimulates CYP27B1 activity, thus increasing the production of calcitriol, which in turn negatively feeds back on the parathyroid gland to decrease PTH secretion. By contrast, FGF-23 inhibits CYP27B1 and stimulates CYP24, thereby decreasing the production of calcitriol and feeding back to limit further secretion of FGF-23, as normally calcitriol stimulates FGF-23 production.

Pi–PTH–FGF-23 Loop

As Pi levels increase (or more likely there is a long-term Pi load), both PTH and FGF-23 are increased, the latter from bone. Both the elevated PTH and FGF-23 increase urinary Pi excretion through downregulation of NaPi transporters. The effect of FGF-23 in the kidney is klotho dependent.




Fig. 52.13 Regulation of serum phosphorus levels. As phosphorus levels increase (or there is a long-term phosphorus load), levels of both parathyroid hormone (PTH) and fibroblast growth factor 23 (FGF-23) are increased. Both of these elevations in turn increase urinary phosphate (Pi) excretion. The two hormones differ in respect to their effects on the vitamin D axis. PTH stimulates 1 α -hydroxylase activity, thereby increasing the production of calcitriol, which in turn negatively feeds back on the parathyroid gland to decrease PTH secretion. By contrast, FGF-23 inhibits 1 α -hydroxylase activity, thereby decreasing the production of calcitriol, thus feeding back to stimulate further secretion of FGF-23. FGF-23 and PTH also regulate each other. Finally, low calcium levels stimulate PTH, whereas high calcium levels stimulate FGF-23. Lastly, there is some evidence that FGF-23 also inhibits PTH secretion.

PTH increases renal calcium reabsorption, minimizing the possibility of high calcium and Pi concentrations in urine at a time when there is a desire to increase Pi urinary excretion. PTH stimulates the secretion of FGF-23 from osteocytes, and increased FGF-23 inhibits PTH by decreasing both *PTH* gene expression and PTH secretion.^{185,186}

Calcium-PTH-FGF-23 Loop

Hypocalcemia, a potent stimulator of PTH, blunts FGF-23 release.¹⁸⁷ The latter would therefore “remove” both the FGF-23 inhibition of PTH and the FGF-23 inhibition of calcitriol synthesis during times of hypocalcemia. This process would maximize both the PTH effects to increase renal calcium reabsorption, increase bone resorption, and enhance calcitriol stimulation of intestinal calcium absorption with the goal of normalizing serum calcium concentrations. Hypercalcemia has opposing effects: It stimulates FGF-23¹⁸⁸ (which reduces PTH and calcitriol synthesis) and directly inhibits calcitriol synthesis and PTH secretion. The result is decreased intestinal calcium absorption, renal reabsorption of calcium, and bone resorption.

DIAGNOSIS OF CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER

MEASUREMENT OF THE BIOCHEMICAL ABNORMALITIES IN CKD-MBD

The measurement of calcium and Pi was discussed earlier in this chapter. Table 52.3 summarizes currently measured biomarkers used for the diagnosis and management of CKD-MBD.

PARATHYROID HORMONE

PTH concentration in plasma or serum serves as not only an indicator of abnormal mineral metabolism in CKD-MBD but also a noninvasive biochemical sign for the initial diagnosis of osteitis fibrosa cystica, the most common form

of renal osteodystrophy in CKD-MBD. PTH measurement also can be a useful index for monitoring the evolution of renal osteodystrophy and can serve as a surrogate measure of bone turnover in patients with CKD. Although the sensitivity and specificity of PTH as a marker of bone remodeling are not ideal, it is the best single marker available at the current time.¹⁸⁹ The combination of PTH, bone-specific alkaline phosphatase, and amino-terminal propeptide of type 1 collagen discriminates high versus low bone turnover in patients receiving dialysis.¹⁸⁹ However, the definitive method for establishing the specific type of renal osteodystrophy in individual patients requires bone biopsy, an invasive diagnostic procedure, and access to specialized laboratory personnel and equipment capable of providing assessments of bone histology.

PTH circulates not only in the form of the intact 84-amino acid peptide but also as multiple fragments of the hormone, particularly from the middle and C-terminal regions of the PTH molecule. These PTH fragments arise from direct secretion from the parathyroid gland as well as from metabolism of PTH(1–84) by peripheral organs, especially liver and kidney. The biologically active hormone produced (PTH[1–84]) exerts its effects through the interaction of its first 34 amino acids with PTHR1. PTH(1–84) has a plasma half-life of 2 to 4 minutes. In comparison, the half-life of C-terminal fragments, which are cleared principally by the kidney, is 5 to 10 times longer with normal kidney function and even longer in the presence of CKD. There is also a diurnal variation in the secretion of PTH and the release is oscillatory, further complicating measurement.

Assays for PTH have undergone many improvements over the years (Fig. 52.14). In the early 1960s, radioimmunoassays were developed for measurement of PTH. However, these assays proved to be unreliable owing to different characteristics of the antisera used and are referred to as “first-generation” assays; consequently, two-site immunometric assays (IMAs) are referred to as “second- and third-generation” assays. The typical second-generation IMAs (known as intact PTH assays) measure PTH(1–84) and other large C-terminal PTH

Table 52.3 Biomarkers for Chronic Kidney Disease–Mineral Bone Disorder

	Affected by Sample Processing	Assay Validity	Renally Excreted	Diurnal Variation	Seasonal Variation	Variation With Meals	Variation With Dialysis Time
Parathyroid hormone	Yes	No; some assays pick up fragments	No	Yes	No	No	Yes
25(OH)D (calcidiol)	No	Good (uncertain importance of differentiating D ₂ from D ₃)	No	No	Yes	No	No
1,25(OH) ₂ D (calcitriol)	No	Good	No	Yes	No	No	?
Fibroblast growth factor 23	No	Intact versus C terminal	No	?	?	Yes	No
Soluble α -klotho	?	Uncertain	Yes	?	?	?	?
Sclerostin	?	Uncertain, likely valid	No	?	?	?	?
Bone-specific alkaline phosphatase	No	Good	No	No	No	?	No

“Assay validity” indicates that the measurement is of the biologically active hormone or marker, not fragments.

?, Indicates insufficient data.




Fig. 52.14 Parathyroid hormone (PTH) assays. Schematic presentation of PTH(1–84) and the relationship between PTH assays, PTH assay epitopes, and PTH molecular forms detected in the circulation. The *upper panel* depicts the structure of human PTH and the epitopes detected by various PTH assays. First-generation PTH assays detect full-length PTH (1–84) in addition to PTH fragments. These assays include radioimmunoassays (RIAs) that use antisera specific to the amino-terminal (N-RIA), middle (MID-RIA), or carboxyl-terminal (C-RIA) regions of PTH. Second-generation “intact PTH” assays detect full-length PTH (1–84) and non(1–84)-PTH fragments. Third-generation PTH assays (bioactive PTH) detect only full-length PTH (1–84). The *bottom panel* depicts PTH molecular forms present in the circulation. IMA, Immunometric assay; N-PTH, amino-terminal PTH; term., terminal. (From Henrich LM, Rogol AD, D’Amour P, et al. Persistent hypercalcemia after parathyroidectomy in an adolescent and effect of treatment with cinacalcet HCl. *Clin Chem*. 2006;52:2286–2293.)

fragments because the antibodies do not bind to amino acid 1. These assays are commonly used in clinical practice. By contrast, third-generation assays (bioactive, whole, or bio-intact PTH assays) use capture antibodies similar to that of the intact PTH assays but also use detection antibodies directed against epitopes at the extreme N-terminal end (epitopes 1–4) of the molecule, and therefore they are believed to detect exclusively the biologically active PTH(1–84) (Fig. 52.14). This difference may be important because C-terminal fragments (lacking small or large portions of the N terminus) are most abundant, representing approximately 80% of circulating PTH in healthy persons and 95% in patients with CKD.¹⁹⁰ This finding may in part explain why elevated PTH concentrations are “normal” in CKD yet are increased relative to values observed in patients without CKD. The second-generation intact assays are commonly used on automated platforms. Although each assay has a reasonable coefficient of variation, the standards for the commercially available assays are not uniform and the detection antibodies do not all bind at the same sites. Thus the assay kit-to-kit variability can be high, with as much as a fivefold difference depending on the kit used.¹⁹¹ This is the reason the Kidney Disease Improving Global Outcomes (KDIGO) guidelines recommended using the same assay every time and evaluating trends rather than targeting precise PTH values.¹⁹²

VITAMIN D

Serum calcitriol concentrations are generally measured by immunoassay, although the gold standard for calcitriol measurement is high-performance liquid chromatography, which is not widely available clinically. Unlike in PTH

assays, sample handling in calcitriol immunoassays has little effect on results. However, vitamin D circulates as both D₂ and D₃, and some laboratory kits measure only D₂, others measure only D₃, and still others measure both (expressed as 25-hydroxyvitamin D). The rationale for distinguishing D₂ from D₃ is controversial because it is unclear how differentiating the forms of vitamin D affects management- or patient-level outcomes. As with PTH, there is some assay-to-assay variability, which could affect the classification of insufficiency/deficiency or sufficient levels of calcitriol.¹⁹³ Fortunately, current initiatives are under way to standardize these assays.¹⁹⁴ The half-life of calcitriol is long and thus represents total body stores.

By contrast, calcitriol levels are generally measured only in the setting of hypercalcemia. The half-life is comparatively short, and the assay more expensive and difficult. Interpretation also requires consideration of clinical context. In other words, a patient with stage G4 or G5 CKD and hypercalcemia may have high normal serum calcitriol, which may be distinctly abnormal given the CKD stage and serum calcium concentration and should prompt consideration of an extrarenal source of calcitriol.

FGF-23

FGF-23 is currently measured primarily with two different assays (Fig. 52.15). The first uses two antibodies directed against the C-terminal end and thus measures the intact, as well as C-terminal, fragments (results are reported in relative unit/mL). The second assay uses one antibody directed against an epitope within the N-terminal region and a second antibody directed against an epitope within




Fig. 52.15 Fibroblast growth factor 23 (FGF-23) assays. (A) FGF-23 O-glycosylation site and epitopes recognized by antibodies used in current assays. (B) Spectrum of serum FGF-23 levels in early chronic kidney disease and end-stage kidney disease (ESKD) compared with the normal reference range and levels associated with different disorders affecting FGF-23. *ADHR*, Autosomal dominant hypophosphatemic rickets; *ARHP*, autosomal recessive hypophosphatemia; *TIO*, tumor-induced osteomalacia; *XLH*, X-linked hypophosphatemia. (From Block GA, Ix JH, Ketteler M, et al. Phosphate homeostasis in CKD: report of a scientific symposium sponsored by the National Kidney Foundation. *Am J Kidney Dis.* 2013;62:457–473. With permission.)

the C-terminal region of the molecule, and thus it detects intact molecules (results are reported in picograms per milliliter).⁸⁵ Although these two assays appear comparable in the association with clinical events, they have poor agreement because of differences in FGF-23 fragment detection, antibody specificity, and calibration. Such analytic variability does not permit direct comparison of FGF-23 measurements made with different assays, a fact that probably, at least in part, accounts for some of the inconsistencies noted among observational studies.¹⁹⁵ FGF-23 can be detected in the urine, although at this time it is unclear how much, if any, of the hormone is cleared by the kidneys.^{196,197} From a clinical perspective, more data are required before the use of FGF-23 measurements for routine clinical management.

SOLUBLE KLOTHO

It is unclear whether the circulating or soluble α -klotho levels reflect tissue-level expression of klotho. Some studies have found that low circulating levels were associated with progression of CKD,¹⁹⁸ but other studies have failed to confirm this finding.¹⁹⁹ A decrease in soluble klotho is detectable before aberrations in FGF23, PTH, or Pi, which may herald onset of CKD-MBD.¹⁹⁸ Klotho can be detected in urine,²⁰⁰ suggesting its levels may be altered by residual renal function. Different assay kits give variable results.^{76,201}

SCLEROSTIN

Circulating sclerostin concentrations are elevated in CKD²⁰² and rise with progressive disease. However, sclerostin does

not appear to be renally excreted,²⁰³ suggesting that the rising levels reflect underlying biology. The role of sclerostin in clinical diagnosis of CKD remains exploratory, although elevated sclerostin values are associated with arterial calcification²⁰⁴ and increased osteoblast number in human bone biopsy specimens.²⁰⁵

BONE-SPECIFIC ALKALINE PHOSPHATASE

Bone-specific alkaline phosphatase (BALP) is not cleared renally. BALP concentration has relatively good correlation with bone formation in CKD and may be additive to the interpretation of PTH measurements.¹⁹² However, its concentration has limited ability as an independent measurement.^{188,206}

COLLAGEN-BASED BONE BIOMARKERS

Osteoblasts secrete C- and N-terminal cleavage products of type I procollagen called secreted procollagen type IN propeptide (s-PINP) and secreted procollagen type IC propeptide (s-PICP), which are markers for bone formation. By contrast, serum C-terminal cross-linking telopeptide of type 1 collagen (s-CTX) and serum N-terminal cross-linking telopeptide of type I collagen (s-NTX) are measured as fragments of cross-links that are released when bone is resorbed. With the exception of the s-PICP, all of these markers are renally excreted, making interpretation of their measurements difficult.²⁰⁷ In cross-sectional analyses, higher serum concentrations are associated with higher odds of fracture.²⁰⁸




Fig. 52.16 Determinants of bone strength. Bone strength comprises both bone density and bone quality. “Bone quality” refers to bone turnover, microarchitecture, microfractures, mineralization, and composition of the mineral matrix. “Trabecular microarchitecture” involves trabecular thickness, the ratio of plates and rods, and their connectivity and spacing. “Cortical microarchitecture” consists of cortical thickness, porosity, and bone size. Composition of the mineral matrix includes changes in the cross-linking of type I collagen and alterations in the size and structure of bone mineral. Bones accumulate microfractures over time even with normal physical activity. The ability to repair them affects bone quality. *BMD*, Bone mineral density. (From Moorthi R, Moe S. Recent advances in the noninvasive diagnosis of renal osteodystrophy. *Kidney Int.* 2013;84:866–894. With permission.)

TARTRATE-RESISTANT ACID PHOSPHATASE 5B

Tartrate-resistant acid phosphatase 5b is released by osteoclasts during bone resorption and thus may be a good marker of bone resorption.²⁰⁹ Use of this biomarker is growing because, in concert with others, it may be useful to discriminate low versus high bone turnover in CKD-MBD.²¹⁰

BONE BIOPSY ASSESSMENT OF BONE IN CHRONIC KIDNEY DISEASE—MINERAL AND BONE DISORDER

Abnormalities of bone quality and quantity are common in CKD-MBD (Fig. 52.16), leading to fractures and impaired growth in children. “Renal osteodystrophy” is defined as an alteration of bone morphology in patients with CKD that is quantifiable by bone histomorphometry.¹

HISTOMORPHOMETRY IN PATIENTS WITH CHRONIC KIDNEY DISEASE

Clinical assessment of bone remodeling is best performed with a bone biopsy of the trabecular bone, usually at the iliac crest. The patient is given a tetracycline derivative approximately 3 to 4 weeks before the bone biopsy and a different tetracycline derivative 3 to 5 days prior. Tetracycline binds to hydroxyapatite and emits fluorescence, thereby serving as a label for the bone. A core of predominantly trabecular bone is collected and embedded in plastic material and then sectioned. The sections can be visualized with special stains under fluorescent microscopy to

determine the amount of bone between administrations of the two tetracycline labels or that formed in the interval. This dynamic parameter assessed with bone biopsy is the basis for evaluating bone turnover, which is key in discerning types of renal osteodystrophy. In addition to dynamic indices, bone biopsies can be analyzed by quantitative histomorphometry for static parameters as well. The nomenclature for these assessments has been standardized.²¹¹

Clinically, bone biopsies are most useful for differentiating bone turnover, as well as bone volume and mineralization. However, with the advent of several new markers of bone turnover, the use of bone biopsy has been reserved primarily for the diagnosis of renal osteodystrophy and for research purposes. Sherrard and colleagues²¹² proposed a classification system for renal osteodystrophy that used the parameters of osteoid (unmineralized bone) area as a percentage of total bone area and fibrosis. These two static parameters, together with the dynamic bone turnover assessed by bone formation rate or activation frequency, have been used to distinguish the various forms of renal osteodystrophy over the past 30 years²¹³; however, this evaluation has been replaced by the KDIGO TMV (turnover, mineralization, volume) system (discussed later).¹

Fig. 52.17 illustrates bone histology using the original classification scheme. Normal bone is illustrated in Fig. 52.17A. Histologic features of high-turnover disease (predominant hyperparathyroidism or osteitis fibrosa cystica) are characterized by an increased rate of bone formation, increased bone resorption, extensive osteoclastic and osteoblastic activity, and progressive increase in endosteal peritrabecular fibrosis (Fig. 52.17B). High osteoblast activity is manifested by an increase in unmineralized bone matrix. The number of osteoclasts is also increased, as well as total resorption surface. There may be numerous dissecting cavities through which the osteoclasts tunnel into individual trabeculae. In osteitis fibrosa cystica, the alignment of strands of collagen in the bone matrix has an irregular woven pattern, unlike the normal lamellar (parallel) alignment of strands of collagen in normal bone. Although woven bone may appear to be thicker, the disorganized collagen structure may render the bone physically more vulnerable to stress.

Low-turnover (adynamic) bone disease (Fig. 52.17C) is characterized histologically by absence of cellular (osteoblast and osteoclast) activity, osteoid formation, and endosteal fibrosis. It appears to be essentially a disorder of decreased bone formation accompanied by a secondary decrease in bone mineralization. Although low-turnover disease is common in the absence of aluminum, it was initially described as a result of aluminum toxicity. Aluminum bone disease is diagnosed with special staining that demonstrates the presence of aluminum deposits at the mineralization front (Fig. 52.17D).²¹² Frequently, aluminum disease is associated with osteomalacia. Osteomalacia is characterized by an excess of unmineralized osteoid, which manifests as wide osteoid seams and a markedly decreased mineralization rate (Fig. 52.17E). The presence of increased unmineralized osteoid per se does not necessarily indicate a mineralizing defect because larger quantities of osteoid appear in conditions associated with high rates of bone formation when mineralization lags behind the increased




Fig. 52.17 Bone histology. (A) Normal bone. (B) Hyperparathyroid bone (increased osteoclast and osteoblasts and fibrosis). (C) Adynamic bone (no cellular activity and no osteoid). (D) Aluminum bone disease (left aluminum staining at mineralization front) and right two panels show accumulation of osteoid (*orange-red stain*). (E) Osteomalacia (increased unmineralized osteoid in *pink/red*). (F) Mixed uremic osteodystrophy presence of increased osteoid (*orange red*) indicating mineralization defect and increased osteoclast activity. (A, B, D [*right*], and E, courtesy S. L. Teitelbaum, M.D.; D [*left*], courtesy D. J. Sherrard, M.D.)

synthesis of matrix. Other features of osteomalacia include the absence of cellular activity and absence of endosteal fibrosis.

Clinical Relevance

Patients with adynamic or low-turnover bone disease are at risk for fractures and vascular calcification. Common therapies for osteoporosis such as bisphosphonates and denosumab inhibit bone turnover and can result in adynamic bone disease in patients with CKD and end-stage kidney disease. Therefore these drugs are not recommended for such patients unless work-up reveals a high-turnover state. In fact, in a study of osteoporotic postmenopausal women on dialysis who received denosumab, 10.7% of women required hospitalization for severe hypocalcemia and 1.3% died.²¹⁴

“Mixed uremic osteodystrophy” is the term used to describe bone biopsies that have features of secondary hyperparathyroidism together with evidence of a mineralization defect (Fig. 52.17F). There is extensive osteoclastic and osteoblastic activity and increased endosteal peritrabecular fibrosis coupled with more osteoid than expected, and tetracycline labeling uncovers a concomitant mineralization defect. Unfortunately, mixed uremic osteodystrophy, in particular, and high- and low-turnover bone diseases have been inconsistent and poorly defined.

THE SPECTRUM OF BONE HISTOMORPHOMETRY IN CKD

The prevalence of different forms of renal osteodystrophy has changed over the past decade. Whereas osteitis fibrosa cystica due to severe hyperparathyroidism had previously been the predominant lesion, the prevalence of mixed uremic osteodystrophy and adynamic bone disease has increased. However, the overall percentage of patients with high bone formation compared with low bone formation has not changed dramatically over the past 20 to 30 years, although osteomalacia has been essentially “replaced” by adynamic bone disease. There are differences in prevalence of mixed uremic osteodystrophy in patients not yet undergoing dialysis, which appears to depend on the level of GFR and the country in which the study was performed.²¹³ Two large analyses of patients undergoing long-term dialysis revealed a relatively high incidence of low bone turnover; one study of 489 biopsy specimens in predominantly white patients revealed low turnover in 59%,¹⁸⁹ whereas in another study of 630 patients, low-turnover disease was noted in 62% of White but only 32% of Black patients.²¹³ The prevalence of mineralization defect or osteomalacia was relatively low at only 3%.²¹³ Thus these data demonstrate that histologic abnormalities of bone begin early in the course of CKD and that differences in bone turnover may differ on the basis of race or country/region of origin.

By contrast, low-turnover bone disease has diverse pathophysiology. In the 1980s, aluminum-induced osteomalacia was common. The potential toxicity of aluminum was initially recognized by Alfrey, who identified a fatal neurologic syndrome in patients receiving dialysis consisting of dyspraxia, seizures, and electroencephalographic abnormalities in association with high brain aluminum levels on autopsy.²¹⁵ The source of aluminum in these severe cases was believed to be elevated

aluminum concentrations in dialysate water. Subsequently, aluminum-containing phosphate binders were also identified as a source. The additional symptoms of fractures, myopathy, and microcytic anemia were described several years after the initial reports of the neurologic syndrome.²¹⁵ Fortunately, exposure to aluminum is limited in the modern era, and the incidence of aluminum bone disease is relatively rare. However, the diagnosis of aluminum-induced bone disease can be difficult because aluminum toxicity is due to tissue burden, not serum concentrations. Thus if aluminum bone disease is suspected, bone biopsy remains the gold standard for making the diagnosis.^{216,217}

In adynamic bone disease, there is a paucity of cells with resultant low bone turnover (Fig. 52.17C). Unlike in osteomalacia, in adynamic bone, there is no increase in osteoid or unmineralized bone. The lack of bone cell activity led to the initial description of the disease as “aplastic” bone disease. Early investigators believed that the disease was due to aluminum, but it was later identified in the absence of aluminum. The etiology of adynamic bone disease is likely multifactorial, and major contributory factors include diabetes mellitus, aging, and malnutrition.²¹⁸

Proposed pathophysiologic mechanisms of low bone turnover are listed in Table 52.4. Increases in both sclerostin and dkk-1, which are soluble inhibitors of Wnt signaling that inhibit osteoblastic bone formation, likely play a role in development of adynamic bone disease.^{109,219,220} Circulating fragments of PTH (7–84–amino acid fragments) may also be antagonists to PTH,²²¹ resulting in an effective resistance to 1–84 amino acid at the level of bone. There is evidence that markedly elevated concentrations of FGF-23 may be associated with decreased osteoblastic activity.²²² Furthermore, abnormal regulation of cell differentiation in the presence of kidney failure may explain, in part, the relative paucity of cells in adynamic bone, although this possibility remains to be proven. In rats, the administration of bone morphogenetic protein-7 can restore normal cell function, supporting that a failure of normal cell differentiation, likely due to a number of causes, may be critical.²²³ Although most patients with low-turnover bone disease are asymptomatic, they are at increased risk of fracture owing to impaired remodeling^{222,224,225} and at risk of vascular calcification because of the inability of bone to buffer both acute and chronic calcium loads.^{222,226,227}

TURNOVER, MINERALIZATION, VOLUME CLASSIFICATION

As previously noted, KDIGO recommends that the definition of renal osteodystrophy be limited to describing the alterations of bone morphology in patients with CKD and is one measure of the skeletal component of the systemic disorder of CKD-MBD that can be quantifiable by histomorphometry.¹ Historically, “renal osteodystrophy” often included disorders of bone mineral metabolism in addition to histomorphometric changes in bone. As our understanding of bone biology progresses, there is greater appreciation of the diverse physiologic processes leading to similar bone biopsy findings. In addition, information on bone volume as an independent parameter is available.²²⁶ Thus the previous classification was updated by KDIGO to use three key histologic descriptors: bone turnover, mineralization, and volume (referred to as the

Table 52.4 Causes and Proposed Mechanisms of Decreased Bone Formation in Patients With Chronic Kidney Disease

	Mechanism of Decreased Osteoblast Activity
Low serum calcitriol	↓ Osteoblast differentiation ↓ Osteoblast life span
Metabolic acidosis	↓ Calcitriol production ↓ Collagen synthesis
High serum phosphate	↓ Calcitriol production
Calcium loading/ hypercalcemia	↓ Calcitriol production and ↑ calcitriol degradation (mediated by calcium- sensing receptor)
High serum interleukins 1 and 6, tumor necrosis factor	↓ Osteoblast life span
Low serum insulinlike growth factor-I (IGF-I) activity	↓ IGF-I and IGF binding protein 5 (IGFBP5) levels ↑ Inhibitory IGFBP (2, 4, 6) levels
Sclerostin	↓ Osteoblast life span ↓ Wnt/β-catenin signaling
Dickkopf-1 (dkk-1)	↓ Osteoblastic activity ↓ Wnt/β-catenin signaling
Malnutrition, proteinuria	↓ Osteoblastic activity
Diabetes	↓ IGF-I; 25-hydroxyvitamin D levels ↓ 25-Hydroxyvitamin D and calcitriol levels ↑ Advanced glycation end products (AGEs)
Age-related	↓ Osteoblast life span ↑ AGEs ↓ Osteoblast life span
Hypogonadal	
Women (↓ estrogen and ↑ sex hormone-binding globulin [SHBG])	↓ Osteoblast life span
Men (↓ testosterone and ↑ SHBG)	↓ Osteoblast life span
Uremic toxins (uric acid)	↓ Calcitriol production ↓ Vitamin D receptor activity ↓ Osteoblast proliferation
Aluminum toxicity	↓ Osteoblast activity

↓, Decreased; ↑, increased.

Table 52.5 Turnover, Mineralization, Volume Classification System for Renal Osteodystrophy

Turnover	Mineralization	Volume
Low	Normal	Low
Normal		Normal
High	Abnormal	High

From Moe S, Drüeke T, Cunningham J, et al. Definition, evaluation, and classification of renal osteodystrophy: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int.* 2006;69:1945–1953.

acceptable parameters for assessing bone turnover. Bone turnover is affected mainly by hormones, cytokines, mechanical stimuli, and growth factors that influence the recruitment, differentiation, and activity of osteoclasts and osteoblasts. It is important to clarify that although bone formation rate is frequently similar to bone resorption rate, which cannot be measured directly, it is not always so. Imbalance in these processes can affect bone volume. For example, if resorption exceeds formation, negative bone balance and decreased bone volume result.

Mineralization reflects how well bone collagen becomes calcified during the formation phase of skeletal remodeling. It is assessed with histomorphometry by static measurements of osteoid volume and osteoid thickness and by dynamic, tetracycline-based measurements of mineralization lag time and osteoid maturation time. Causes of impaired mineralization include inadequate vitamin D nutrition, mineral (calcium or Pi) deficiency, acidosis, and aluminum toxicity.

Volume indicates the amount of bone per unit volume of tissue. It is assessed with histomorphometry by static measurements of bone volume in cancellous bone. Determinants of bone volume include age, sex, race, genetic factors, nutrition, endocrine disorders, mechanical stimuli, toxicities, neurologic function, vascular supply, growth factors, and cytokines.

The KDIGO classification is consistent with the historically used classification system²¹² but provides more information on parameters other than turnover. Two large-scale analyses using the updated TMV classification scheme suggested that it provides clinically relevant information.^{189,213} Low bone volume and low bone turnover are more common than heretofore appreciated, whereas defective mineralization is relatively rare in adult patients with ESKD.

Clinically, serum PTH concentration is used as a surrogate biomarker to predict bone turnover. However, as detailed earlier, studies evaluating the ability of the serum concentration of (intact) PTH to predict low- and/or high-turnover bone diseases have been disappointing. The ability to reliably predict the presence of high-turnover bone disease is poor until serum PTH concentrations exceed 500 pg/mL.^{189,213} Primarily on the basis of earlier studies using the Allegro intact PTH assay, the Kidney Disease Outcomes Quality Initiative (KDOQI) guidelines

TMV classification), with any combination of each of the descriptors possible in a given specimen (Table 52.5).¹ The TMV classification scheme provides a clinically relevant description of the underlying bone pathology as assessed by histomorphometry, which in turn helps define the pathophysiology and thereby guide therapy.

Turnover reflects the rate of skeletal remodeling, which is normally the coupled process of bone resorption and bone formation. Turnover is assessed with histomorphometry by dynamic measurements of osteoblast function using double-tetracycline labeling, as previously described. Bone formation rate and activation frequency represent

recommend a target-intact PTH level of 150 to 300 pg/mL.²¹⁶ Unfortunately, studies that correlate intact PTH with bone histology were performed with an assay no longer available. Use of currently available assays has shown that the intact PTH value between 150 and 300 pg/mL is not predictive of underlying bone histology.²²⁸ However, an analysis of 610 biopsy specimens found that although the intact PTH value could not predict underlying bone histology, it was able to discriminate low-turnover from non-low-turnover bone diseases and high-turnover from non-high-turnover bone diseases.¹⁸⁹

NONINVASIVE ASSESSMENT OF BONE

DUAL-ENERGY X-RAY ABSORPTIOMETRY

Dual-energy X-ray absorptiometry (DXA) measures areal bone mineral density (aBMD or simply BMD) in g/cm² using minimal radiation and rapid scan times. BMD assessment by DXA has good reproducibility (<1%–2% coefficient variation) and reliable reference ranges for age, sex, and race. In the general population, BMD measured by DXA can be used clinically to define osteoporosis and is an accepted surrogate endpoint after prospective studies demonstrated an age-dependent predictive value of DXA for fractures.²²⁹ However, the discordance between changes in DXA findings and the efficacy of drugs at preventing fracture has led to appreciation of the importance of bone “quality,” which is not assessed by DXA. This observation has forced the use of fractures as endpoints for approval of new therapeutics for the treatment of osteoporosis.

The KDIGO CKD-MBD guideline recommends DXA to assess fracture risk in patients with stage 1 through early stage 3 CKD, as long as biochemical testing does not suggest CKD-MBD.¹⁹² However, for patients with CKD stages 3b through 5, the guideline did not recommend DXA owing to the lack of definitive data demonstrating that DXA predicts fracture in CKD-MBD. A meta-analysis was performed to determine whether DXA measurements at the femoral neck, spine, and radius were associated with spine and/or nonspine fractures in patients undergoing dialysis.^{230,232} On the basis of results of 6 studies with 683 patients, lower BMD in the spine and distal radius, but not in the femoral neck, was associated with fractures. Although BMD may have been lower in patients with CKD and a history of fracture, there is considerable overlap in BMD such that BMD provides poor fracture discrimination in individuals. In addition, for lumbar spine assessment of BMD by DXA, existing aortic calcification may confound the measurement. Since the publication of 2017 KDIGO guidelines, studies have demonstrated fracture prediction value of DXA in patients with CKD at least equivalent to that in the general population.^{232,233} These data therefore support that DXA can predict fracture risk in patients with CKD. The addition of the Trabecular Bone Score (TBS) to DXA may further improve fracture prediction in the general population and in patients with CKD, ESKD, and post transplantation.²³⁴⁻²³⁶ However, DXA cannot make a specific diagnosis as to why there is low BMD. Unlike patients with normal or near-normal kidney function and low BMD by DXA being classified as having osteoporosis, patients with CKD and low bone density should not be routinely treated with antiosteoporosis therapies.²³⁷ DXA is an inexpensive and widely available technique that can be

easily standardized among sites, so it may be a good tool in longitudinal CKD research studies for the serial assessment of BMD in response to interventions. Unfortunately, to date, treatment studies using DXA as an endpoint in patients with CKD are limited.

QUANTITATIVE COMPUTED TOMOGRAPHY TECHNIQUES

Quantitative CT (QCT) allows three-dimensional imaging of cross sections of the central and axial skeleton to provide spatial or volumetric BMD (vBMD). It also allows distinction between cortical and trabecular compartments. In CKD, QCT measures of trabecular bone density at the spine have been correlated with trabecular bone volume histomorphometry.²³⁸ Peripheral QCT (pQCT) avoids the large dose of ionizing radiation exposure for patients by focusing on the tibia and distal radius, and in one study it was associated with fracture.²³⁹ Although derived from a single study, these results are in line with expected associations between loss of bone quality and fracture risk in patients undergoing dialysis.

High-resolution peripheral computed tomography (HRpQCT) has greater resolution than pQCT and allows evaluation of trabecular microarchitecture (bone volume fraction, trabecular thickness, separation, and number). HRpQCT of the radius and tibia was found to discriminate among patients with CKD with and without fracture.^{241,242} However, the discriminatory power of the various HRpQCT parameters to individually discriminate between those with and without fractures by receiver operating curve analyses was <0.75. When patients with the longest duration of CKD were considered, the area under the receiver operating curve improved to more than 0.8 for multiple parameters including radial cortical thickness, radial total vBMD, and cortical vBMD. Interestingly, aBMD by DXA of the ultradistal radius performed similarly in its ability to discriminate prevalent fractures in this subpopulation with the longest duration of kidney disease.²⁴² In another study, areal BMD by DXA at the ultradistal radius was superior to HRpQCT measures for fracture discrimination.²⁴³ One study demonstrates the superiority of HRpQCT in identifying changes associated with acidosis in CKD.²⁴⁴ In a group of patients with CKD followed longitudinally with DXA and HRpQCT, Levy and colleagues²⁴⁴ noted that more severe acidosis at baseline and worsening acidosis over time were both related to lower trabecular number, while more severe acidosis at baseline was associated with increases in trabecular separation and network heterogeneity. Concurrently, no changes in DXA scores were noted relative to acidosis.²⁴⁴

MICROCOMPUTED TOMOGRAPHY AND MICROMAGNETIC RESONANCE IMAGING

Resolution of microtechniques is as low as 8 mm, compared with about 100 μm for HRpQCT, thus providing spatial resolution almost that of an actual bone biopsy. Micro-CT was used to assess the effects of severe high-turnover renal osteodystrophy on trabecular and cortical architecture in growing rats. The technique demonstrated irregular trabecular thickening and loss of trabecular connectivity in the femoral neck and greater endocortical porosity in the femoral shaft, consistent with biopsy findings in high-turnover osteodystrophy.²⁴⁵ Unfortunately, this technique

is currently limited to in vitro studies. In an evaluation of 17 patients undergoing hemodialysis who had secondary hyperparathyroidism, micromagnetic resonance imaging demonstrated disruptions of the distal tibial trabecular network.²⁴⁶ There have not been subsequent confirmatory studies or studies comparing this technique with other imaging methods in the ability to predict fractures in CKD.

ASSESSMENT OF VASCULAR CALCIFICATION

Arterial calcification can be detected in humans through numerous techniques. Plain radiographs can be used to assess the presence or absence, and thus prevalence, of vascular calcification. Scoring methods using the number of aorta segments calcified along the lumbar spine can also allow reproducible quantification during longitudinal follow-up,²⁴⁷ although the sensitivity is less than with CT-based imaging. Although some distinction can be made between medial and intimal calcification on plain radiographs,²⁴⁸ the reproducibility among multiple research sites of this method for differentiation of calcification type has not been evaluated. Ultrasonography of the carotid arteries can be used to assess intimal–medial thickness, which correlates well with atherosclerosis and cardiovascular events. In a later study, intravascular ultrasonography of the coronary arteries was able to detect atherosclerotic lesions and calcification. This technique, although invasive, can detect circumferential lesions and external remodeling of atherosclerosis—lesions that invade the internal elastic artery into the medial layer rather than protrude into the vessel lumen. The latter luminal lesion is all that is detected on angiography, leading some to question the use of angiography as the “gold standard.”²⁴⁹ Finally, technologic advances have led to ultrafast CT scans—electron beam CT (EBCT) and multislice CT—that use electrocardiography gating to allow imaging only in diastole, thus avoiding motion artifact of the heart. These techniques have allowed reproducible quantification of coronary artery and aorta calcification and therefore serve as excellent intermediate endpoints for CKD-MBD interventions that might alter progression of vascular calcification. Unfortunately, these techniques do not allow differentiation of medial from intimal calcification, and the radiation dose with their use is large.

A study in each of 140 prevalent patients undergoing hemodialysis compared a lateral radiograph of the lumbar abdominal aorta, an echocardiogram, and measurement of pulse pressure with EBCT results.²⁵⁰ Calcification of the abdominal aorta was scored as 0 through 24 divided into tertiles; echocardiograms were graded as 0 through 2 for absence or presence of calcification of the mitral and aortic valves; and pulse pressure was divided in quartiles. The researchers found that the likelihood ratio (95% confidence interval [CI]) of coronary artery calcification score on EBCT of 100 or higher was 1.79 (1.09–2.96) for calcification of either valve on echocardiography and 7.50 (2.89–19.5) for participants with a lateral abdominal radiographic score ≥ 7 . Verbeke and colleagues²⁵¹ confirmed this predictive value of lateral abdominal radiographs in more than 1000 patients receiving dialysis.

On the basis of observational studies, calcification detected in all of these imaging studies is associated with mortality in

patients with CKD. Bellasi and Raggi²⁵² analyzed multiple such studies and calculated the positive predictive value to be between 19% and 52% and the negative predictive value to be between 68% and 100%. The addition of the clinical variables age and dialysis vintage to abdominal calcification termed the “cardiovascular calcification index,” was linearly associated with death, with a 12% increase in hazard ratio for each point increase in the cardiovascular calcification index.²⁵³

CLINICAL CONSEQUENCES OF THE ABNORMALITIES IN CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER

BIOCHEMICAL ABNORMALITIES

PHOSPHORUS AND CALCIUM

Epidemiologic data suggest that serum Pi levels above the normal range are associated with increases in morbidity and all-cause and cardiovascular mortality in patients with CKD. These studies differ in their sample size, analyses, and chosen reference ranges, and most evaluate only patients undergoing dialysis. In patients with CKD stages 2 to 5, higher levels of serum Pi, even within the population reference range, have been associated with increased risk of all-cause or cardiovascular mortality in one study²⁵⁴ but not in other studies.^{255,256} However, in patients undergoing dialysis, nearly all studies demonstrate an association of elevated Pi with mortality, although there are slight differences in the inflection point or range at which Pi becomes significantly associated with increased all-cause mortality. Furthermore, several retrospective studies have demonstrated that treatment of patients undergoing dialysis with Pi binders is associated with a 20% to 40% lower risk of death.^{257–259} However, there are no prospective studies that have demonstrated a survival benefit of lowering serum phosphate concentrations, and there is no specific target serum phosphate that has been identified as conferring a clinically relevant benefit. In patients with CKD stages 3 to 5, there are no data to support an increased risk of mortality or fracture with elevated serum calcium concentrations. However, as with Pi, there are several studies in patients receiving dialysis demonstrating higher mortality with higher serum calcium concentration and in some studies, among patients with very low serum calcium concentrations, as the latter may reflect more severe vitamin D deficiency and secondary hyperparathyroidism.

PARATHYROID HORMONE

Although PTH has long been considered a surrogate marker for bone disease, it also has systemic effects including effects on endothelial, cardiac, and skeletal health. As with other biochemical measures of CKD-MBD, observational studies have found an association of all-cause mortality with various levels of PTH, with inflection points ranging from more than 400 to 600 pg/mL. On basis of these observational data and the assay limitations, the KDIGO guidelines consider serum concentrations of intact PTH within two and nine times the upper limit of normal for the PTH assay (<130 and >585 pg/mL for most kits with an upper normal limit of 65 pg/mL) as a reasonable target range, and that values within

that range should be interpreted by evaluation of trends. Intervention should occur if the trends are consistently going up or down.¹⁹² However, it is important to recognize that there are no randomized clinical trials demonstrating that treatment to achieve specific PTH concentrations results in improved outcomes, with the exception of the Evaluation Of Cinacalcet Hydrochloride (HCl) Therapy to Lower CardioVascular Events (EVOLVE) trial discussed in Chapter 62.

A Cochrane meta-analysis found that serum Pi and calcium concentrations, but not PTH, were associated with cardiovascular and all-cause mortality.²⁶⁰

COMBINATION OF ABNORMALITIES OF CALCIUM, PHOSPHORUS, AND PARATHYROID HORMONE

The relationship of calcium, Pi, and PTH with outcomes is further complicated by the clinical reality that these laboratory parameters do not respond in isolation from one another, but rather change depending on the levels of other parameters and treatments. For example, Stevens and colleagues²⁶¹ assessed various biochemical combinations and found that the relative risk for mortality was highest when serum concentrations of calcium and Pi were elevated and the serum PTH concentration was low. They discovered that mortality was lowest when serum concentrations of calcium and Pi were normal and the serum PTH concentration was high. In addition, results were modified by vintage (time since initiation of dialysis). A Dialysis Outcomes and Practice Patterns Study (DOPPS) trial also evaluated combinations of serum parameters of mineral metabolism and reached slightly different conclusions.²⁶² The DOPPS researchers found that in the setting of an elevated serum PTH (>300 pg/mL), hypercalcemia (>10 mg/dL) was associated with higher mortality risk even with normal serum Pi. Block and associates, analyzing average values over 4 months from more than 26,000 patients undergoing dialysis, categorized them according to whether serum concentrations of calcium, Pi, and PTH were below, within, or above target ranges. In only 20% of patients were all three variables within target ranges. Those patients with high PTH and high serum calcium had consistently higher mortality.²⁶³ It is important to remember that abnormalities in mineral metabolism are common. Ultimately, we need quality evidence (i.e., from prospective randomized controlled trials using approaches that target specific biochemical endpoints) to determine whether these treatments—despite having been commonly administered in some cases for decades—are truly safe and efficacious.

FIBROBLAST GROWTH FACTOR 23

Serum concentrations of FGF-23 (>100 RU/mL) were observed in 70% and 100% of patients with estimated GFRs of 50 and <20 mL/min/1.73 m², respectively, in the Chronic Renal Insufficiency Cohort (CRIC) study.⁷⁷ Serum FGF-23 concentrations typically rise as kidney function declines over time, presumably to maintain hemostasis vis-à-vis Pi and calcium. In CKD, elevations of serum FGF-23 are associated with death,²⁶⁴ progression of CKD,^{264,265} left ventricular hypertrophy,^{89,266,267} and cardiovascular events,^{269,270} independent of serum concentrations of Pi and PTH. Once patients reach levels of kidney function poor enough to require dialysis, serum concentrations of FGF-23 can be 1000-fold greater, and these have been associated with

poorer survival in incident patients in some studies²⁷⁰ but not others.^{271,272} In prevalent patients receiving dialysis, serum FGF-23 was also associated with mortality²⁷³ and directly correlated with left ventricular mass index.²⁷² In a secondary analysis of nearly 2000 hemodialysis patients treated with cinacalcet, a reduction of serum FGF-23 concentrations by 30% or more was associated with reduced risks of death and cardiovascular events.²⁷⁴

NUTRITIONAL VITAMIN D DEFICIENCY

Although traditionally the term “vitamin D” has been used to indicate the active metabolite, calcitriol, the correct use of the term is for the precursor molecule, calcidiol or 25(OH)D. In the general population, serum calcidiol concentrations are accepted as the standard measures of nutritional uptake because they correlate best with end-organ effects. The conversion of vitamin D₂ and vitamin D₃ to 25(OH)D by CYP27A1 is essentially unregulated, and therefore levels of 25(OH)D are a reliable indicator of the vitamin D status of a given individual. However, what constitutes adequate stores remains a controversy. Although there is no absolute level of calcidiol that defines deficiency, a level <10 ng/mL (25 nmol/L) is typically used because it is associated with rickets in children and osteomalacia in adults. The term “vitamin D insufficiency” has been used to describe less severe calcidiol-deficient states. Although controversial, the typical range of “insufficient” calcidiol levels is 10 to 30 ng/mL (25–75 nmol/L). In contrast to the severity of bone disease observed with deficiency, insufficiency is associated with elevated PTH and osteoporosis.²⁷⁵ Unfortunately, despite supplementation of calciferol in various foods such as milk, calcidiol insufficiency is relatively common in the general population, particularly in some at-risk groups such as African Americans, hospitalized patients, nursing home residents, and people living in northern climates.²⁷⁶ The Institute of Medicine convened a panel whose report, released in November 2010, recommended that serum vitamin D concentrations of 20 ng/mL are adequate and that the adult-recommended dietary allowance is 600 IU/day, with an upper-level intake of 4000 IU/day.²⁷⁷ The Endocrine Society has subsequently recommended a vitamin D level higher than 30 ng/mL, further suggesting that 40 to 60 ng/mL is ideal and that up to 100 ng/mL is safe.²⁷⁸ Either vitamin D₂ or vitamin D₃ can be used for vitamin D supplementation. Although there has been controversy regarding the use of D₃ or D₂ for achieving and maintaining higher serum vitamin D levels, most analyses have found them to be equally effective in raising and maintaining serum vitamin D levels in patients without CKD.²⁷⁹

Vitamin D deficiency (<20 ng/mL) and insufficiency (<30 ng/mL) are common in patients with CKD, with only 29% and 17% of those with stage 3 and 4 disease, respectively, having sufficient levels in one study.²⁸⁰ This prevalence of vitamin D deficiency and insufficiency in an ambulatory CKD population is similar to that in non-CKD nursing home residents, hospitalized patients, and elderly women with hip fractures. However, frank vitamin D deficiency (<10 ng/mL) was observed in 14% and 26% of patients with stage 3 and 4 CKD, respectively.²⁸⁰ Several other investigators have also

found widespread vitamin D insufficiency in patients with CKD, although not all investigators have identified a graded relation with CKD stage.²⁸⁰⁻²⁸⁴ Current targets for vitamin D sufficiency may be too low, as several studies have shown that serum levels of 40 to 50 were associated with appropriate PTH suppression in CKD without hypercalcemia or hyperphosphatemia.^{284,285}

Evidence from epidemiologic studies suggests that vitamin D deficiency and supplementation of vitamin D are associated with survival in patients with CKD regardless of dialysis status, although these analyses are heavily confounded, prohibiting causal inference.²⁸⁶⁻²⁹¹ A meta-analysis of prospective studies demonstrated a significant decrease, 14%, in mortality risk for every 10-ng/mL increase in serum vitamin D concentration.²⁹² Furthermore, the association of vitamin D supplementation and survival appears to be independent of changes in serum calcium, Pi, and PTH concentrations. There are several plausible mechanisms that could explain the connection between vitamin D deficiency and insufficiency and mortality; for example, serum vitamin D concentrations are inversely correlated with multiple cardiovascular risk factors including plasma renin activity, blood pressure, left ventricular mass, markers of inflammation, markers of insulin resistance and type 2 diabetes mellitus, and albuminuria.²⁹³⁻²⁹⁷

In animal studies, activated vitamin D inhibits renin production.^{298,299} Several trials in patients with CKD reported that the use of an active vitamin D analog, paricalcitol, resulted in a significant reduction in proteinuria.³⁰⁰⁻³⁰² Evaluation of patients with nondialysis-requiring CKD found an inverse correlation between serum vitamin D concentrations and brachial artery flow-mediated dilation as a marker of endothelial dysfunction.³⁰³ Other studies have demonstrated that patients undergoing hemodialysis who have received cholecalciferol supplements have lower inflammatory parameters, as indicated by higher serum albumin and lower C-reactive protein and interleukin-6 concentrations, as well as improved cardiac function, as reflected by lower brain natriuretic peptide and left ventricular mass index.^{304,305} In a prospective placebo-controlled trial in patients who had not started dialysis, paricalcitol failed to reduce left ventricular mass after 48 weeks in patients with mild-to-moderate left ventricular hypertrophy. There was, however, a decrease in hospitalizations, a secondary endpoint.³⁰⁶ The lack of effect may have been a function of the relatively high dose of paricalcitol, which may have increased serum FGF-23 concentrations, as multiple studies have shown a direct link between FGF-23 and left ventricular hypertrophy.^{88,266,306,307} Despite multiple observational studies and several clinical trials, additional controlled trials are required to determine the safety and efficacy of (nutritional and active) vitamin D supplementation in patients with CKD.

FRACTURE

The incidence of fractures was reported to be significantly higher in patients with nondialysis-requiring CKD compared with persons with normal or near-normal kidney function, although lower than the risk of fractures in patients receiving dialysis,^{192,308-310} suggesting that kidney disease is associated with progressive bone deterioration. However, not all studies have confirmed increased rates of hip, wrist, and

vertebral fractures in patients with CKD independent of age and sex.³¹² Newer studies have suggested that fracture rates in ESKD patients are declining compared with historical data.³¹²

In patients undergoing dialysis, the studies are consistent and demonstrate a markedly higher incidence of hip fracture among all age groups.^{224,313} Patients undergoing hemodialysis may have a 50% higher rate of hip fractures than patients undergoing peritoneal dialysis.³¹⁴ Furthermore, hip fractures in patients on dialysis were associated with a doubling of mortality observed after hip fractures in patients not on dialysis.^{224,315} Depending on the study, 10% to 52% of prevalent patients undergoing dialysis have had fractures.¹⁹² Unfortunately, most of these studies are of cross-sectional design or were cohort studies with limited follow-up.

Hyperparathyroidism is known to cause abnormal bone remodeling, especially of cortical bone. Serial studies using HRpQCT show a rapid decline in cortical bone in patients on dialysis.³¹⁶ In prevalent patients undergoing dialysis who have secondary hyperparathyroidism, clinical fractures are predominantly cortical in location.³¹⁷ However, the association of PTH with fracture has been inconsistent,¹⁹² perhaps because both high and low PTH values are associated with fractures. Other risk factors for hip fracture include older age, female sex, low serum albumin, prior kidney transplantation, and peripheral vascular disease.^{208,318-320} Falls are also more common in patients with CKD and predispose to fracture. Falls in patients undergoing dialysis may be related to peripheral vascular disease,³¹³ low muscle strength, impaired neuromuscular function,³²¹ and the administration of psychoactive medications.³¹⁹

In summary, the abnormalities of bone, in terms of both quality and quantity, variable biochemical abnormalities, and multiple comorbid conditions likely all contribute to the pathogenesis of fractures in ESKD. This complexity also may limit generalizability of therapies routinely used in the general population to patients with the disease.

EXTRASKELETAL CALCIFICATION

Arterial calcification has a multitude of clinical manifestations. Intimal calcification (in association with atherosclerotic disease) can lead to myocardial infarction from stenosis and acute thrombus or to ischemia in both coronary and peripheral arteries. Medial calcification (or circumferential calcification) can lead to arterial stiffening, with reduced compliance of the artery and an inability to appropriately dilate in the setting of increased stress. In coronary arteries, similar symptoms of ischemia can develop and in theory could lead to arrhythmias and sudden death. In larger arteries such as the aorta, calcification can lead to increased pulse wave velocity and elevated pulse pressure and is commonly associated with systolic hypertension in the elderly, a known risk factor for cardiovascular disease in the general population. In addition, the premature return of wave reflections during systole (instead of diastole) can lead to altered coronary perfusion, especially in the setting of left ventricular hypertrophy. Lastly, calcification of the arterioles of the skin and other organs can lead to localized infarction and ischemia, including ischemic bowel and calcific uremic arteriolopathy (calciphylaxis).

VASCULAR CALCIFICATION IN PATIENTS WITH CHRONIC KIDNEY DISEASE

The high prevalence of vascular calcification in patients with CKD is not a new observation. In 1979, Ibels and colleagues³²² demonstrated that both renal and internal iliac arteries of patients undergoing renal transplantation had greater atherogenic/intimal disease and increased calcification (detected by biochemical methods) than transplant donors. In addition, the medial layer was thicker and more calcified in uremic patients than in the donors. A study comparing histologic changes in coronary arteries from patients undergoing dialysis at autopsy were compared with nondialysis, age-matched patients who had died from a cardiac event found similar magnitudes of atherosclerotic plaque burden and intimal thickness, but with more calcification in the patients who had received dialysis.³²³ In addition, morphometry of the arteries demonstrated increased medial thickening. When these same investigators evaluated more distal segments of the coronary arteries, they found medial calcification.³²⁴ Studies of the inferior epigastric artery of patients undergoing kidney transplantation demonstrated calcification in 31% of patients, with histology demonstrating isolated medial calcification without evidence of atherosclerotic or intimal changes, and both medial and intimal calcification in the same artery without disruption of the internal elastic lamina.³²⁵ Thus there is histologic evidence of 1. increased arterial calcification in coronary, renal, and iliac arteries of patients who undergo dialysis than in those who do not; and 2. the presence of both intimal and medial calcification in patients undergoing dialysis, and this can occur independently of each other, suggesting unique inciting factors. As noted previously, it is likely that calcification of plaque (atherosclerosis) and calcification of medial layer occur by similar mechanisms. However, the initiating event is clearly different.

Braun and colleagues³²⁶ first demonstrated that coronary artery calcification as detected by EBCT increased with advancing age in patients on dialysis and that the calcification scores were twofold to fivefold higher in patients on dialysis than in age-matched persons with normal kidney function and angiographically proven coronary artery disease. Nearly 50% to 60% of patients starting hemodialysis have evidence of coronary artery calcification,³²⁷ with higher prevalence in patients with diabetes.³²⁸ The prevalence of detectable coronary or peripheral artery calcification is 51% to 93% in prevalent patients undergoing dialysis and 47% to 83% in patients with CKD stages 3 through 5. Valvular calcification is similarly common, occurring in 20% to 47% of patients on dialysis.¹⁹² Studies that have evaluated the natural history of coronary calcification demonstrate that once present, the course is progressive. By contrast, many patients without calcification remain free of calcification for several years.^{192,329,330} The presence of calcification in the coronary arteries, valves, and peripheral arteries is associated with mortality.²⁵² Given the high prevalence of calcification, it is currently uncertain whether routine imaging offers clinical utility or whether it can help stratify patients whose calcification will and will not respond to specific therapies.

There is an inverse relation between bone mineralization and vascular calcification in patients receiving dialysis. There are several distinct but not mutually exclusive mechanisms

by which disturbances in the bone metabolism may cause or accelerate vascular calcification.^{326,331} Interestingly, it appears to be low-turnover bone disease that leads to highest risk of vascular calcification. London and colleagues³³² evaluated patients on hemodialysis who underwent assessment of vascular calcification by ultrasonography with semiquantitative scoring and by bone biopsy with histomorphometry. Those patients with lowest bone formation rates and decreased osteoblast surfaces had the most prominent degree of peripheral artery calcification, and this relation held true in patients with and without previous parathyroidectomy.³³² Another study found that patients with low-turnover bone disease on biopsy were more likely to have progression of coronary artery calcification over 1 year as assessed by serial multislice CT than those with high-turnover bone disease.²²⁸ Abnormalities in cortical and trabecular bone on HRpQCT were associated with coronary artery calcification.³³³ The likely mechanism relates to a reduced ability to mineralize bone in ESKD, especially in the setting of low-turnover bone disease, leading to dystrophic calcification in the setting of calcium (and vitamin D) intake; actively remodeling bone, by contrast, is able to mineralize, as demonstrated by Kurz and associates.²²⁶ In contrast to evidence in low-turnover bone disease, data are less robust as to the effects of high-turnover bone disease on vascular calcification, perhaps because the treatments for high-turnover bone disease confound the effects of calcification.

CHRONIC KIDNEY DISEASE-MINERAL AND BONE DISORDER IN KIDNEY TRANSPLANT RECIPIENTS

Bone and mineral disorders are universal complications in patients with CKD before transplantation. Ideally, many of these complications would be improved with successful kidney transplantation. Unfortunately, in many cases, kidney transplantation achieves milder forms of CKD (as opposed to normal kidney function), and thus kidney transplant recipients still suffer from disorders associated with CKD-MBD. In addition, disorders of mineral metabolism occur following successful transplantation, including the effects of medications (steroids and calcineurin inhibitors), persistence of underlying disorders (hyperparathyroidism and vitamin D deficiency), and development of hyperphosphaturia with hypophosphatemia (especially with persistent hyperparathyroidism).

BIOCHEMICAL CHANGES AFTER TRANSPLANTATION

Biochemical changes in mineral metabolism are common in patients after kidney transplantation.³³⁴⁻³³⁷ Decreased calcidiol levels are common in kidney transplant recipients, with reported prevalence of up to 81%.³³⁵ The causes are multifactorial and may include nutritional deficiency, malabsorption, and decreased sun exposure. However, the improvement in kidney function results in an increase in CYP27B1 activity, resulting in greater conversion of calcidiol to calcitriol, which subsequently causes a gradual decrease in PTH concentrations during the first 3 to 6 months after transplantation. However, persistent hyperparathyroidism

exists in about 33% and 20% of patients at 6 and 12 months, respectively.³³⁸ Serum FGF-23 concentrations decrease immediately after transplantation, although they remain inappropriately elevated for the prevailing serum Pi concentration. By about 3 months after transplantation, serum FGF-23 concentrations are reduced by 89%, and at 12 months FGF-23 concentrations appear to be similar to those in patients without transplantation who have comparable stages of CKD.³³⁵ The elevated FGF-23 inhibits CYP27B1 and contributes to the low calcitriol concentrations observed in the early post-transplantation period, likely contributing to the persistent hyperparathyroidism seen after transplantation. Thus elevated FGF-23 and PTH concentrations combined with low calcitriol concentrations likely contribute to hypophosphatemia observed in many transplant recipients.^{338,339}

Serum calcium concentrations typically drop soon after transplantation, as a result of cessation of therapy with calcitriol and its analogs and/or calcium-containing phosphate binders. After that initial drop, however, calcium concentrations progressively increase, with hypercalcemia developing in a substantial number of patients during the first to third month after transplantation. Hypercalcemia can be transient, resolving by 6 to 8 months in some cases but lasting for many years in others. Hypercalcemia is typically associated with hyperparathyroidism and results from enhanced renal tubular reabsorption of calcium under the influence of PTH in the functioning allograft, the effects of calcitriol on the gastrointestinal absorption of calcium, and, potentially, a direct effect of PTH in causing calcium efflux from bone. Low bone turnover with relatively low PTH may also result in hypercalcemia because the bone is not able to act as a buffer for the circulating calcium pool.^{334,335}

BONE CHANGES AFTER TRANSPLANTATION

Disorders of mineral metabolism and associated bone disease lead to fractures following transplantation, which occur after up to 44% of successful kidney transplants.³³⁴ This fracture risk appears to be less than that observed with other solid organ transplants.³³⁴ The combination of kidney and pancreas transplantation raises fracture risk above that associated with kidney transplantation alone.^{340,341} In a retrospective analysis of 68,814 patients, fractures developed in 22.5% within 5 years.³⁴² In a study of first-time kidney transplant recipients between 1997 and 2010, the incidence of hip fracture was 3.8 per 1000 person-years. The risk of fracture in 2010 was 0.56 (95% CI, 0.47–0.77) in comparison with that in 1997, indicating a decrease over the 13-year period, presumably due to changes in immunosuppression therapy.³⁴³ Factors associated with higher fracture risk during the first 5 years included female sex, age older than 45 years, white race, recipient of a deceased donor kidney, increased human leukocyte antigen mismatches, diabetes, pretransplantation dialysis, steroid use, and an aggressive induction regimen. Importantly, even with the use of steroid-sparing regimens, total (all sites) fractures have occurred in up to 50% of kidney transplant recipients at 6 years.^{343,344} By contrast, a Canadian study evaluated 4821 kidney transplant recipients and found that the 10-year cumulative incidence of hip fracture was 1.7%.³⁴⁵

Studies evaluating bone histology in recipients of kidney transplants are limited. The main alteration is an uncoupling of bone remodeling, which results in a decrease in bone formation with persistent bone resorption and net bone loss. In a small cohort of young patients who underwent preemptive transplantation and were treated predominantly with corticosteroids, bone biopsies revealed a mineralization defect as early as 6 months after transplantation.³⁴⁶ Another study evaluating the early histologic changes after transplantation revealed that osteoid volume, osteoid thickness, osteoid resorption surface, and osteoclast surface were elevated above the normal range before transplantation and remained increased approximately 35 days afterward. However, osteoid and osteoblast surfaces, which also were increased before transplantation, were significantly decreased approximately 35 days after transplantation.³⁴⁷ Both bone formation and mineralization were also inhibited after transplantation. An important observation was that although none of the pretransplantation biopsy specimens showed evidence of apoptosis, 45% of post-transplantation specimens showed significant apoptosis after an average of only 35 days. Thus early post-transplantation apoptosis and a decrease in osteoblast number and osteoblast surface play a role in the pathogenesis of post-transplantation bone disease that may be related directly to the use of glucocorticoids. The few evaluations that examined the longer-term effect of transplantation on bone histology generally demonstrate low bone turnover and histology consistent with adynamic bone disease.^{348,349} Lehmann and colleagues³⁴⁹ performed a retrospective study of 57 patients after approximately 54 months after kidney transplantation. Seven histologic subgroups were identified: normal, osteitis fibrosa cystica, mild osteitis fibrosa, mixed uremic bone disease, osteomalacia, adynamic bone disease, and osteoporosis. Given the limited number of patients in the study, no significant correlations between the histologic subgroups and PTH or alkaline phosphatase levels were observed. However, this study did show a variable degree of histologic abnormalities following transplantation that could not have been predicted by biochemical testing and confirmed that preexisting bone disease lesions persist in the post-transplantation period. Protein expression profiles in pediatric solid organ transplant recipients demonstrate elevated FGF23 and sclerostin levels relative to nontransplant CKD.³⁵⁰ Paired bone biopsy specimens obtained before transplant and 12 months after transplant show increased bone sclerostin and decreased or unchanged FGF23.³⁵¹

The effects of other immunosuppressive agents on bone histology have also been examined. Bone biopsies performed approximately 10 years after transplantation in patients whose treatment included cyclosporine monotherapy, azathioprine and prednisone, or triple therapy revealed no differences among the immunosuppressant regimens.^{352,353} A subgroup analysis of 21 patients with normal serum PTH concentrations showed that cyclosporine monotherapy was associated with a more pronounced decrease in bone thickness than the other regimens.³⁵³ In addition, the cyclosporine group showed a lower trabecular appositional rate than the azathioprine and prednisone group. Multiple regression analysis showed that sex and time after transplantation were the most significant factors predicting bone volume and mineralizing surface. Predictive factors

for eroded bone surface and osteoclast number included age and time on dialysis before transplantation.^{352,353}

Initial studies examining BMD by DXA showed severe and rapid bone loss after kidney transplantation.³⁴⁶ However, later studies have not found the dramatic decrease initially described, perhaps because of the current practice of using lower steroid dosages or steroid-free immunosuppression regimens.^{334,354} The use of corticosteroids is the major determinant of low BMD because these agents impair calcium absorption from the gastrointestinal tract and inhibit bone cell recruitment and function. Cueto-Manzano and associates^{352,353} also evaluated BMD as a function of immunosuppressive therapy. In comparison with age- and sex-matched controls, none of the treated groups showed a significant reduction in BMD in the lumbar spine or femoral neck. However, in comparison with young normal controls, osteopenia was detected in the femoral neck, except in premenopausal women. There was no significant difference in BMD related to immunosuppressant regimen, sex, or menopausal status.

Although reductions in BMD have been associated with an increased fracture rate in postmenopausal women and in men treated with glucocorticoids, as well as in heart or liver transplant recipients, little data support its role in predicting fractures after kidney transplantation. In the one study that demonstrated low BMD to be predictive of fracture risk, 283 kidney transplant recipients with varying degrees of kidney function underwent 670 examinations of bone density at the hip. An absolute BMD value <0.9 g/cm² was associated with an increased risk of fracture.³⁵⁵ A study evaluating patients with early steroid withdrawal after kidney transplantation found preservation of BMD by DXA but continued loss at the ultradistal radius. The latter finding was confirmed by HRpQCT to be predominantly cortical loss, which was associated with persistent hyperparathyroidism.³⁵⁶ As indicated earlier, abnormal bone turnover may alter the predictive value of BMD testing in patients with CKD, and thus preexisting rate of bone turnover likely affects the assessment and outcomes of bone disease in transplant recipients.

VASCULAR CALCIFICATION CHANGES AFTER TRANSPLANTATION

Most transplant recipients enter their post-transplant course with a preexisting burden of vascular calcification; as in patients with CKD, vascular calcification strongly predicts cardiovascular events and all-cause mortality. Unfortunately, there are few studies on the nature and consequences of vascular calcification in kidney transplant recipients. Although preexisting calcifications progress after transplantation, they appear to do so at a much slower rate than before transplantation.^{357,358} A review of 13 clinical studies of the role of vascular calcifications in kidney transplant recipients shows that as in the CKD population, the associations of traditional and nontraditional risk factors with outcomes are variable in transplant recipients.³⁵⁹ There is a strong association between baseline coronary artery calcification score and progression of coronary artery calcification. A significant improvement in secondary hyperparathyroidism after transplantation is associated with slowing the progression of coronary artery calcification.

Independent risk factors of coronary artery calcification in kidney transplant recipients include low serum concentrations of calcidiol, fetuin-A, and MGP. Although diabetes is a risk factor for the presence of coronary artery calcification in transplant recipients, it has not been independently associated with progression of coronary artery calcification. Data on the effects of immunosuppressive drugs as factors in progression are few and inconclusive; however, it does appear that mycophenolate mofetil may have a beneficial effect because its antiproliferative action may inhibit smooth muscle cell proliferation, thus having a favorable effect on endothelial cell activity.³⁵⁹ In the general population and in patients with ESKD, there is an inverse relation between coronary artery calcification and bone density. This association has not been shown in kidney or kidney-pancreas transplant recipients.³⁶⁰

SUMMARY

In summary, some abnormalities of CKD-MBD persist after kidney transplantation and other conditions unique to transplantation frequently develop. With changing immunosuppressive protocols and persistence of CKD, treatment of these conditions can be particularly challenging. Studies have highlighted the importance of management of CKD-MBD before transplantation. Additional studies are required to determine the safety and efficacy of established and novel therapies in patients with CKD and ESKD and after kidney transplantation.^{336,337,361}

KEY REFERENCES

- Lee GJ, Marks J. Intestinal phosphate transport: a therapeutic target in chronic kidney disease and beyond? *Pediatr Nephrol*. 2015;30(3):363–371.
- King AJ, Siegel M, He Y, et al. Inhibition of sodium/hydrogen exchanger 3 in the gastrointestinal tract by tenapanor reduces paracellular phosphate permeability. *Sci Transl Med*. 2018;10(456):eaam6474.
- Zhou W, Simic P, Zhou Y, Caravan P, et al. Kidney glycolysis serves as a mammalian phosphate sensor that maintains phosphate homeostasis. *J Clin Invest*. 2023;133:133–146.
- Gauci C, Moranne O, Fouqueray B, et al. Pitfalls of measuring total blood calcium in patients with CKD. *J Am Soc Nephrol*. 2008;19(8):1592–1598.
- Spiegel DM, Brady K. Calcium balance in normal individuals and in patients with chronic kidney disease on low- and high-calcium diets. *Kidney Int*. 2012;81(11):1116–1122.
- Evenepoel P, Jørgensen HS, Bover J, Davenport A, et al. Recommended calcium intake in adults and children with chronic kidney disease—a European consensus statement. *Nephrol Dial Transplant*. 2024;39(2):341–366.
- Blaine J, Chonchol M, Levi M. Renal control of calcium, phosphate, and magnesium homeostasis. *Clin J Am Soc Nephrol*. 2015;10:1257–1272.
- Hou J, Renigunta A, Konrad M, et al. Claudin-16 and claudin-19 interact and form a cation-selective tight junction complex. *J Clin Invest*. 2008;118(2):619–628.
- Black DM, Bilezikian JP, Ensrud KE, et al. One year of alendronate after one year of parathyroid hormone (1-84) for osteoporosis. *N Engl J Med*. 2005;353(6):555–565.
- Hu MC, Shi M, Zhang J, et al. Renal production, uptake, and handling of circulating alphaKlotho. *J Am Soc Nephrol*. 2016;27(1):79–90.
- Hanudel MR, Laster M, Salusky IB. Non-renal-Related mechanisms of FGF23 pathophysiology. *Curr Osteoporos Rep*. 2018;16(6):724–729.
- Wolf M, Koch TA, Bregman DB. Effects of iron deficiency anemia and its treatment on fibroblast growth factor 23 and phosphate homeostasis in women. *J Bone Miner Res*. 2013;28:1793–1803.

85. Edmonston D, Wolf M. FGF23 at the crossroads of phosphate, iron economy and erythropoiesis. *Nat Rev Nephrol.* 2020;16:7–19.
86. Bożentowicz-Wikarek M, Owczarek A, Kocelak P, Olszanecka-Glinianowicz M, et al. C-Terminal to intact fibroblast growth factor 23 ratio in relation to estimated glomerular filtration rate in elderly population. *Kidney Blood Press Res.* 2016;41(5):519–526.
87. Singh S, Grabner A, Yanucil C, et al. Fibroblast growth factor 23 directly targets hepatocytes to promote inflammation in chronic kidney disease. *Kidney Int.* 2016;90(5):985–996.
88. Agoro R, Montagna A, Goetz R, et al. Inhibition of fibroblast growth factor 23 (FGF23) signaling rescues renal anemia. *FASEB J.* 2018;32(7):3752–3764.
89. Faul C, Amaral AP, Oskouei B, et al. FGF23 induces left ventricular hypertrophy. *J Clin Invest.* 2011;121(11):4393–4408.
91. Pastor-Arroyo EM, Gehring N, Krudewig C, et al. The elevation of circulating fibroblast growth factor 23 without kidney disease does not increase cardiovascular disease risk. *Kidney Int.* 2018;94(1):49–59.
107. McColm J, Hu L, Womack T, Tang CC, Chiang AY. Single- and multiple-dose randomized studies of blosozumab, a monoclonal antibody against sclerostin, in healthy postmenopausal women. *J Bone Miner Res.* 2014;29(4):935–943.
108. McClung MR, Grauer A, Boonen S, et al. Romosozumab in postmenopausal women with low bone mineral density. *N Engl J Med.* 2014;370(5):412–420.
109. Cejka D, Herberth J, Branscum AJ, et al. Sclerostin and Dickkopf-1 in renal osteodystrophy. *Clin J Am Soc Nephrol.* 2011;6(4):877–882.
110. Sabbagh Y, Gracioli FG, O'Brien S, et al. Repression of osteocyte Wnt/beta-catenin signaling is an early event in the progression of renal osteodystrophy. *J Bone Miner Res.* 2012;27(8):1757–1772.
114. Miller PD, Adachi JD, Albergaria BH, et al. Efficacy and safety of romosozumab among postmenopausal women with osteoporosis and mild-to-moderate chronic kidney disease. *J Bone Miner Res.* 2022;37:1437–1445.
115. Moe SM, Chen NX, Newman CL, et al. Anti-sclerostin antibody treatment in a rat model of progressive renal osteodystrophy. *J Bone Miner Res.* 2015;30(3):499–509.
131. Amaya-Garrido A, Brunet M, Buffin-Meyer B, et al. Calprotectin is a contributor to and potential therapeutic target for vascular calcification in chronic kidney disease. *Sci Transl Med.* 2023;15(712):eabn5939.
162. Mohamed ON, Mohamed MRM, Hassan IG, Alakkad AF, et al. The relationship of fetuin-A with coronary calcification, carotid atherosclerosis, and mortality risk in non-dialysis chronic kidney disease. *J Lipid Atheroscler.* 2024;13(2):194–211.
170. Nigwekar SU, Bloch DB, Nazarian RM, et al. Vitamin K-dependent carboxylation of matrix gla protein influences the risk of calciphylaxis. *J Am Soc Nephrol.* 2017;28(6):1717–1722.
173. Saritas T, Reinartz S, Krüger T, Ketteler M, et al. Vitamin K1 and progression of cardiovascular calcifications in hemodialysis patients: the VitaVasK randomized controlled trial. *Clin Kidney J.* 2022;15(12):2300–2311.
174. Ketteler M, Schanz M, Shricker S. Vitamin K: should we supplement to protect the kidneys and the heart? *Nephrol Dial Transplant.* 2021;36:2196–2198.
178. O'Neill WC, Sigrist MK, McIntyre CW. Plasma pyrophosphate and vascular calcification in chronic kidney disease. *Nephrol Dial Transplant.* 2010;25(1):187–191.
179. de Oliveira RB, Louvet L, Riser BL, et al. Peritoneal delivery of sodium pyrophosphate blocks the progression of pre-existing vascular calcification in uremic apolipoprotein-E knockout mice. *Calcif Tissue Int.* 2015;97(2):179–192.
189. Sprague SM, Bellorin-Font E, Jorgetti V, et al. Diagnostic accuracy of bone turnover markers and bone histology in patients with CKD treated by dialysis. *Am J Kidney Dis.* 2016;67(4):559–566.
196. Leaf DE, Jacob KA, Srivastava A, et al. Fibroblast growth factor 23 levels associate with AKI and death in critical illness. *J Am Soc Nephrol.* 2017;28(6):1877–1885.
197. van Ballegooijen AJ, Rhee EP, Elmariah S, de Boer IH, Kestenbaum B. Renal clearance of mineral metabolism biomarkers. *J Am Soc Nephrol.* 2016;27(2):392–397.
214. Bird ST, Smith ER, Gelperin K, Jung TH, et al. Severe hypocalcemia with denosumab among older female dialysis-dependent patients. *JAMA.* 2024;331(6):491–499.
234. Shevraja E, Reginster JY, Lamy O, Al-Daghri N, et al. Update on the clinical use of trabecular bone score (TBS) in the management of osteoporosis: results of an expert group meeting organized by the European society for clinical and economic aspects of osteoporosis, osteoarthritis and musculoskeletal diseases (ESCEO), and the international osteoporosis foundation (IOF) under the auspices of WHO collaborating center for epidemiology of musculoskeletal health and aging. *Osteoporos Int.* 2023;34(9):1501–1529.
235. Naylor KL, Lix L, Hans D, Garg AX, et al. Trabecular bone score in kidney transplant recipients. *Osteoporos Int.* 2016;27(3):1115–1121.
236. Yun HJ, Ryoo SR, Kim JE, Choi YJ, et al. Trabecular bone score may indicate chronic kidney disease-mineral and bone disorder (CKD-MBD) phenotypes in hemodialysis patients: a prospective observational study. *BMC Nephrol.* 2020;21(1):299.
237. Ketteler M, B G, Evenepoel P, et al. Executive summary of the 2017 KDIGO chronic kidney disease-mineral and bone disorder (CKD-MBD) guideline update: what's changed and why it matters. *Kidney Int.* 2017;92(6):26–36.
244. Levy RV, McMahon DJ, Agarwal S, et al. Comprehensive associations between acidosis and the skeleton in patients with kidney disease. *J Am Soc Nephrol.* 2023;34:668–681.
274. Moe SM, Chertow GM, Parfrey PS, et al. Cinacalcet, fibroblast growth factor-23, and cardiovascular disease in hemodialysis: the evaluation of cinacalcet HCl therapy to lower cardiovascular events (EVOLVE) trial. *Circulation.* 2015;132(1):27–39.
284. Ennis JL, Worcester EM, Coe FL, Sprague SM. Current recommended 25-hydroxyvitamin D targets for chronic kidney disease management may be too low. *J Nephrol.* 2016;29(1):63–70.
285. Dhillon-Jhattu S, McGill RL, Ennis JL, et al. Vitamin D and parathyroid hormone levels in CKD. *Am J Kidney Dis.* 2023;81:122–124.279.
312. Kim SM, Liu S, Long J, Montez-Rath ME, Leonard MB, Chertow GM. Declining rates of hip fracture in end-stage renal disease: analysis from the 2003–2011 nationwide inpatient sample. *J Bone Miner Res.* 2017;32(11):2297–2303.
316. Nickolas TL, Stein EM, Dworakowski E, et al. Rapid cortical bone loss in patients with chronic kidney disease. *J Bone Miner Res.* 2013;28(8):1811–1820.
317. Moe SM, Abdalla S, Chertow GM, et al. Effects of cinacalcet on fracture events in patients receiving hemodialysis: the EVOLVE trial. *J Am Soc Nephrol.* 2015;26(6):1466–1475.
318. Stehman-Breen CO, Sherrard DJ, Alem AM, et al. Risk factors for hip fracture among patients with end-stage renal disease. *Kidney Int.* 2000;58(5):2200–2205.
345. Naylor KL, Jamal SA, Zou G, et al. Fracture incidence in adult kidney transplant recipients. *Transplantation.* 2016;100(1):167–175.
351. Araújo MJCLN, Bacelar Marques ID, Gracioli FG, et al. Comparison of serum levels with bone content and gene expression indicate a contradictory effect of kidney transplantation on sclerostin. *Kidney Int.* 2019;96:1100–1104.
353. Cueto-Manzano AM, Konel S, Hutchison AJ, et al. Bone loss in long-term renal transplantation: histopathology and densitometry analysis. *Kidney Int.* 1999;55(5):2021–2029.

REFERENCES

- Moe S, Drueke T, Cunningham J, et al. Definition, evaluation, and classification of renal osteodystrophy: a position statement from Kidney Disease: Improving Global Outcomes (KDIGO). *Kidney Int*. 2006;69(11):1945–1953.
- Portale AA, Halloran BP, Morris Jr RC. Dietary intake of phosphorus modulates the circadian rhythm in serum concentration of phosphorus. Implications for the renal production of 1,25-dihydroxyvitamin D. *J Clin Invest*. 1987;80(4):1147–1154.
- Moe SM, Zidehsarai MP, Chambers MA, et al. Vegetarian compared with meat dietary protein source and phosphorus homeostasis in chronic kidney disease. *Clin J Am Soc Nephrol*. 2011;6(2):257–264.
- Sherman RA, Mehta O. Phosphorus and potassium content of enhanced meat and poultry products: implications for patients who receive dialysis. *Clin J Am Soc Nephrol*. 2009;4(8):1370–1373.
- Leon JB, Sullivan CM, Sehgal AR. The prevalence of phosphorus-containing food additives in top-selling foods in grocery stores. *J Ren Nutr*. 2013;23(4):265–270.
- Calvo MS, Moshfegh AJ, Tucker KL. Assessing the health impact of phosphorus in the food supply: issues and considerations. *Adv Nutr*. 2014;5(1):104–113.
- Hill KM, Martin BR, Wastney ME, et al. Oral calcium carbonate affects calcium but not phosphorus balance in stage 3–4 chronic kidney disease. *Kidney Int*. 2013;83(5):959–966.
- Lee GJ, Marks J. Intestinal phosphate transport: a therapeutic target in chronic kidney disease and beyond? *Pediatr Nephrol*. 2015;30(3):363–371.
- Amasheh S, Milatz S, Krug SM, et al. Tight junction proteins as channel formers and barrier builders. *Ann N Y Acad Sci*. 2009;1165:211–219.
- Sabbagh Y, O'Brien SP, Song W, et al. Intestinal npt2b plays a major role in phosphate absorption and homeostasis. *J Am Soc Nephrol*. 2009;20(11):2348–2358.
- Schiavi SC, Tang W, Bracken C, et al. Npt2b deletion attenuates hyperphosphatemia associated with CKD. *J Am Soc Nephrol*. 2012;23(10):1691–1700.
- Marks J, Churchill LJ, Debnam ES, Unwin RJ. Matrix extracellular phosphoglycoprotein inhibits phosphate transport. *J Am Soc Nephrol*. 2008;19(12):2313–2320.
- Miyamoto K, Ito M, Kuwahata M, Kato S, Segawa H. Inhibition of intestinal sodium-dependent inorganic phosphate transport by fibroblast growth factor 23. *Ther Apher Dial*. 2005;9(4):331–335.
- King AJ, Siegel M, He Y, et al. Inhibition of sodium/hydrogen exchanger 3 in the gastrointestinal tract by tenapanor reduces paracellular phosphate permeability. *Sci Transl Med*. 2018;10(456):eaam6474.
- Zhou W, Simic P, Zhou IY, Caravan P, et al. Kidney glycolysis serves as a mammalian phosphate sensor that maintains phosphate homeostasis. *J Clin Invest*. 2023;133:173–183.
- Gauci C, Moranne O, Fouqueray B, et al. Pitfalls of measuring total blood calcium in patients with CKD. *J Am Soc Nephrol*. 2008;19(8):1592–1598.
- Glendenning P. It is time to start ordering ionized calcium more frequently: preanalytical factors can be controlled and postanalytical data justify measurement. *Ann Clin Biochem*. 2013;50(Pt 3):191–193.
- Spiegel DM, Brady K. Calcium balance in normal individuals and in patients with chronic kidney disease on low- and high-calcium diets. *Kidney Int*. 2012;81(11):1116–1122.
- Evenepoel P, Jørgensen HS, Bover J, Davenport A, et al. Recommended calcium intake in adults and children with chronic kidney disease—a European consensus statement. *Nephrol Dial Transplant*. 2024;39(2):341–366.
- Woudenberg-Vrenken TE, Lameris AL, Weissgerber P, et al. Functional TRPV6 channels are crucial for transepithelial Ca²⁺ absorption. *Am J Physiol Gastrointest Liver Physiol*. 2012;303(7):G879–G885.
- Bronner F. Recent developments in intestinal calcium absorption. *Nutr Rev*. 2009;67(2):109–113.
- Masuyama R. Role of local vitamin D signaling and cellular calcium transport system in bone homeostasis. *J Bone Miner Metabol*. 2014;32(1):1–9.
- Liu N, Nguyen L, Chun RF, et al. Altered endocrine and autocrine metabolism of vitamin D in a mouse model of gastrointestinal inflammation. *Endocrinology*. 2008;149(10):4799–4808.
- Blaine J, Chonchol M, Levi M. Renal control of calcium, phosphate, and magnesium homeostasis. *Clin J Am Soc Nephrol*. 2015;10:1257–1272.
- Hou J, Renigunta A, Konrad M, et al. Claudin-16 and claudin-19 interact and form a cation-selective tight junction complex. *J Clin Invest*. 2008;118(2):619–628.
- Hoorn EJ, Zietse R. Disorders of calcium and magnesium balance: a physiology-based approach. *Pediatr Nephrol*. 2013;28(8):1195–1206.
- Bindels RJ, 2009 Homer W. Smith Award: minerals in motion: from new ion transporters to new concepts. *J Am Soc Nephrol*. 2010;21(8):1263–1269.
- de Groot T, Kovalevskaya NV, Verkaar S, et al. Molecular mechanisms of calmodulin action on TRPV5 and modulation by parathyroid hormone. *Mol Cell Biol*. 2011;31(14):2845–2853.
- Leunissen EH, Nair AV, Bull C, et al. The epithelial calcium channel TRPV5 is regulated differentially by klotho and sialidase. *J Biol Chem*. 2013;288(41):29238–29246.
- Brown EM, Gamba G, Riccardi D, et al. Cloning and characterization of an extracellular Ca²⁺-sensing receptor from bovine parathyroid. *Nature*. 1993;366(6455):575–580.
- Brennan SC, Thiem U, Roth S, et al. Calcium sensing receptor signalling in physiology and cancer. *Biochim Biophys Acta*. 2013;1833(7):1732–1744.
- Conigrave AD, Ward DT. Calcium-sensing receptor (CaSR): pharmacological properties and signaling pathways. *Best Pract Res Clin Endocrinol Metabol*. 2013;27(3):315–331.
- Brennan SC, Davies TS, Schepelmann M, Riccardi D. Emerging roles of the extracellular calcium-sensing receptor in nutrient sensing: control of taste modulation and intestinal hormone secretion. *Br J Nutr*. 2014;111(suppl 1):S16–S22.
- Kos CH, Karaplis AC, Peng JB, et al. The calcium-sensing receptor is required for normal calcium homeostasis independent of parathyroid hormone. *J Clin Invest*. 2003;111(7):1021–1028.
- Egbuna O, Quinn S, Kantham L, et al. The full-length calcium-sensing receptor dampens the calcemic response to 1,25(OH)₂ vitamin D₃ in vivo independently of parathyroid hormone. *Am J Physiol Ren Physiol*. 2009;297(3):F720–F728.
- Kantham L, Quinn SJ, Egbuna OI, et al. The calcium-sensing receptor (CaSR) defends against hypercalcemia independently of its regulation of parathyroid hormone secretion. *Am J Physiol Endocrinol Metab*. 2009;297(4):E915–E923.
- Rodriguez-Ortiz ME, Canalejo A, Herencia C, et al. Magnesium modulates parathyroid hormone secretion and upregulates parathyroid receptor expression at moderately low calcium concentration. *Nephrol Dial Transplant*. 2014;29(2):282–289.
- Latus J, Lehmann R, Roesel M, et al. Analysis of alpha-klotho, fibroblast growth factor-, vitamin-D and calcium-sensing receptor in 70 patients with secondary hyperparathyroidism. *Kidney Blood Press Res*. 2013;37(1):84–94.
- Sumida K, Nakamura M, Ubara Y, et al. Cinacalcet upregulates calcium-sensing receptors of parathyroid glands in hemodialysis patients. *Am J Nephrol*. 2013;37(5):405–412.
- Tyler MR. Control of renal calcium, phosphate, electrolyte, and water excretion by the calcium-sensing receptor. *Best Pract Res Clin Endocrinol Metabol*. 2013;27(3):345–358.
- Renkema KY, Velic A, Dijkman HB, et al. The calcium-sensing receptor promotes urinary acidification to prevent nephrolithiasis. *J Am Soc Nephrol*. 2009;20(8):1705–1713.
- Riccardi D, Brennan SC, Chang W. The extracellular calcium-sensing receptor, CaSR, in fetal development. *Best Pract Res Clin Endocrinol Metabol*. 2013;27(3):443–453.
- Xue Y, Xiao Y, Liu J, et al. The calcium-sensing receptor complements parathyroid hormone-induced bone turnover in discrete skeletal compartments in mice. *Am J Physiol Endocrinol Metab*. 2012;302(7):E841–E851.
- Gonzalez-Vazquez A, Planell JA, Engel E. Extracellular calcium and CaSR drive osteoinduction in mesenchymal stromal cells. *Acta Biomater*. 2014;10(6):2824–2833.
- Sun X, Kishore V, Fites K, Akkus O. Osteoblasts detect pericellular calcium concentration increase via neomycin-sensitive voltage-gated calcium channels. *Bone*. 2012;51(5):860–867.
- David V, Dai B, Martin A, Huang J, Han X, Quarles LD. Calcium regulates FGF-23 expression in bone. *Endocrinology*. 2013;154(12):4469–4482.

47. Alam MU, Kirton JP, Wilkinson FL, et al. Calcification is associated with loss of functional calcium-sensing receptor in vascular smooth muscle cells. *Cardiovasc Res*. 2009;81(2):260–268.
48. Henaut L, Boudot C, Massy ZA, et al. Calcimimetics increase CaSR expression and reduce mineralization in vascular smooth muscle cells: mechanisms of action. *Cardiovasc Res*. 2014;101(2):256–265.
49. Lopez I, Aguilera-Tejero E, Mendoza FJ, et al. Calcimimetic R-568 decreases extraosseous calcifications in uremic rats treated with calcitriol. *J Am Soc Nephrol*. 2006;17(3):795–804.
50. Koleganova N, Piecha G, Ritz E, Schmitt CP, Gross ML. A calcimimetic (R-568), but not calcitriol, prevents vascular remodeling in uremia. *Kidney Int*. 2009;75(1):60–71.
51. Ivanovski O, Nikolov IG, Joki N, et al. The calcimimetic R-568 retards uremia-enhanced vascular calcification and atherosclerosis in apolipoprotein E deficient (apoE^{-/-}) mice. *Atherosclerosis*. 2009;205(1):55–62.
52. Moe SM, Chen NX, Seifert MF, et al. A rat model of chronic kidney disease-mineral bone disorder. *Kidney Int*. 2009;75(2):176–184.
53. Ciceri P, Elli F, Brenna I, Volpi E, Brancaccio D, Cozzolino M. The calcimimetic calindol prevents high phosphate-induced vascular calcification by upregulating matrix GLA protein. *Nephron Exp Nephrol*. 2012;122(3–4):75–82.
54. Raggi P, Chertow GM, Torres PU, et al. The ADVANCE study: a randomized study to evaluate the effects of cinacalcet plus low-dose vitamin D on vascular calcification in patients on hemodialysis. *Nephrol Dial Transplant*. 2011;26(4):1327–1339.
55. Divieti P, John MR, Juppner H, Bringham FR. Human PTH-(7-84) inhibits bone resorption in vitro via actions independent of the type 1 PTH/PTHrP receptor. *Endocrinology*. 2002;143(1):171–176.
56. Vilardaga JP, Romero G, Friedman PA, Gardella TJ. Molecular basis of parathyroid hormone receptor signaling and trafficking: a family B GPCR paradigm. *Cell Mol Life Sci*. 2011;68(1):1–13.
57. Nechama M, Ben-Dov IZ, Silver J, Naveh-Man T. Regulation of PTH mRNA stability by the calcimimetic R568 and the phosphorus binder lanthanum carbonate in CKD. *Am J Physiol Ren Physiol*. 2009;296(4):F795–F800.
58. Olauson H, Lindberg K, Amin R, et al. Parathyroid-specific deletion of Klotho unravels a novel calcineurin-dependent FGF23 signaling pathway that regulates PTH secretion. *PLoS Genet*. 2013;9(12):e1003975.
59. Ritter CS, Brown AJ. Direct suppression of Pth gene expression by the vitamin D prohormones doxercalciferol and calcitriol requires the vitamin D receptor. *J Mol Endocrinol*. 2011;46(2):63–66.
60. Sprague SM, Moe SM. The case for routine parathyroid hormone monitoring. *Clin J Am Soc Nephrol*. 2013;8(2):313–318.
61. Black DM, Bilezikian JP, Ensrud KE, et al. One year of alendronate after one year of parathyroid hormone (1-84) for osteoporosis. *N Engl J Med*. 2005;353(6):555–565.
62. Zella LA, Shevde NK, Hollis BW, Cooke NE, Pike JW. Vitamin D-binding protein influences total circulating levels of 1,25-dihydroxyvitamin D3 but does not directly modulate the bioactive levels of the hormone in vivo. *Endocrinology*. 2008;149(7):3656–3667.
63. Dusso AS. Update on the biologic role of the vitamin D endocrine system. *Curr Vasc Pharmacol*. 2014;12(2):272–277.
64. Jones G. Extrarenal vitamin D activation and interactions between vitamin D(2), vitamin D(3), and vitamin D analogs. *Annu Rev Nutr*. 2013;33:23–44.
65. Chanakul A, Zhang MY, Louw A, et al. FGF-23 regulates CYP27B1 transcription in the kidney and in extra-renal tissues. *PLoS One*. 2013;8(9):e72816.
66. Hendy GN, Hruska KA, Mathew S, Goltzman D. New insights into mineral and skeletal regulation by active forms of vitamin D. *Kidney Int*. 2006;69(2):218–223.
67. Nigwekar SU, Tamez H, Thadhani RI. Vitamin D and chronic kidney disease-mineral bone disease (CKD-MBD). *BoneKEy Rep*. 2014;3:498.
68. Demay MB. Physiological insights from the vitamin D receptor knockout mouse. *Calcif Tissue Int*. 2013;92(2):99–105.
69. Feng JQ, Clinkenberg EL, Yuan B, White KE, Drezner MK. Osteocyte regulation of phosphate homeostasis and bone mineralization underlies the pathophysiology of the heritable disorders of rickets and osteomalacia. *Bone*. 2013;54(2):213–221.
70. Shimada T, Mizutani S, Muto T, et al. Cloning and characterization of FGF23 as a causative factor of tumor-induced osteomalacia. *Proc Natl Acad Sci U S A*. 2001;98(11):6500–6505.
71. Bellido T, Saini V, Pajevic PD. Effects of PTH on osteocyte function. *Bone*. 2013;54(2):250–257.
72. Kolek OI, Hines ER, Jones MD, et al. 1 α ,25-Dihydroxyvitamin D3 upregulates FGF23 gene expression in bone: the final link in a renal-gastrointestinal-skeletal axis that controls phosphate transport. *Am J Physiol Gastrointest Liver Physiol*. 2005;289(6):G1036–G1042.
73. Quarles LD. Role of FGF23 in vitamin D and phosphate metabolism: implications in chronic kidney disease. *Exp Cell Res*. 2012;318(9):1040–1048.
74. Belov AA, Mohammadi M. Molecular mechanisms of fibroblast growth factor signaling in physiology and pathology. *Cold Spring Harbor Perspect Biol*. 2013;5(6):a015958.
75. Hu MC, Shi M, Cho HJ, et al. Klotho and phosphate are modulators of pathologic uremic cardiac remodeling. *J Am Soc Nephrol*. 2015;26(6):1290–1302.
76. Hu MC, Shi M, Zhang J, et al. Renal production, uptake, and handling of circulating alphaKlotho. *J Am Soc Nephrol*. 2016;27(1):79–90.
77. Isakova T, Wahl P, Vargas GS, et al. Fibroblast growth factor 23 is elevated before parathyroid hormone and phosphate in chronic kidney disease. *Kidney Int*. 2011;79(12):1370–1378.
78. Hu MC, Kuro-o M, Moe OW. Renal and extrarenal actions of Klotho. *Semin Nephrol*. 2013;33(2):118–129.
79. Chang Q, Hoefs S, van der Kemp AW, Topala CN, Bindels RJ, Henderop JG. The beta-glucuronidase klotho hydrolyzes and activates the TRPV5 channel. *Science*. 2005;310(5747):490–493.
80. Cha SK, Hu MC, Kurosu H, Kuro-o M, Moe O, Huang CL. Regulation of renal outer medullary potassium channel and renal K(+) excretion by Klotho. *Mol Pharmacol*. 2009;76(1):38–46.
81. Satoh M, Nagasu H, Morita Y, Yamaguchi TP, Kanwar YS, Kashihara N. Klotho protects against mouse renal fibrosis by inhibiting Wnt signaling. *Am J Physiol Ren Physiol*. 2012;303(12):F1641–F1651.
82. Imel EA, Peacock M, Gray AK, Padgett LR, Hui SL, Econs MJ. Iron modifies plasma FGF23 differently in autosomal dominant hypophosphatemic rickets and healthy humans. *J Clin Endocrinol Metab*. 2011;96(11):3541–3549.
83. Hanudel MR, Laster M, Salusky IB. Non-renal-Related mechanisms of FGF23 pathophysiology. *Curr Osteoporos Rep*. 2018;16(6):724–729.
84. Wolf M, Koch TA, Bregman DB. Effects of iron deficiency anemia and its treatment on fibroblast growth factor 23 and phosphate homeostasis in women. *J Bone Miner Res*. 2013;28:1793–1803.
85. Edmonston D, Wolf M. FGF23 at the crossroads of phosphate, iron economy and erythropoiesis. *Nat Rev Nephrol*. 2020;16:7–19.
86. Bożentowicz-Wikarek M, Owczarek A, Kocelak P, Olszanecka-Glinianowicz M, et al. C-Terminal to intact fibroblast growth factor 23 in relation to estimated glomerular filtration rate in elderly population. *Kidney Blood Press Res*. 2016;41(5):519–526.
87. Singh S, Grabner A, Yanucil C, Schramm K, Czaja B, Krick S, Czaja MJ, Bartz R, Abraham R, Di Marco GS, et al. Fibroblast growth factor 23 directly targets hepatocytes to promote inflammation in chronic kidney disease. *Kidney Int*. 2016;90(5):985–996.
88. Agoro R, Montagna A, Goetz R, Aligbe O, Singh G, Coe LM, Mohammadi M, Rivella S, Sitara D. Inhibition of fibroblast growth factor 23 (FGF23) signaling rescues renal anemia. *FASEB J*. 2018;32(7):3752–3764.
89. Faul C, Amaral AP, Oskouei B, et al. FGF23 induces left ventricular hypertrophy. *J Clin Invest*. 2011;121(11):4393–4408.
90. Touchberry CD, Green TM, Tchikrizov V, et al. FGF23 is a novel regulator of intracellular calcium and cardiac contractility in addition to cardiac hypertrophy. *Am J Physiol Endocrinol Metab*. 2013;304(8):E863–E873.
91. Pastor-Arroyo EM, Gehring N, Krudewig C, Costantino S, Bettoni C, Knopfel T, Sabrautzki S, Lorenz-Depierreux B, Pastor J, Strom TM, et al. The elevation of circulating fibroblast growth factor 23 without kidney disease does not increase cardiovascular disease risk. *Kidney Int*. 2018;94(1):49–59.
92. Takeshita K, Fujimori T, Kurotaki Y, et al. Sinoatrial node dysfunction and early unexpected death of mice with a defect of klotho gene expression. *Circulation*. 2004;109(14):1776–1782.

93. Song S, Gao P, Xiao H, Xu Y, Si LY. Klotho suppresses cardiomyocyte apoptosis in mice with stress-induced cardiac injury via downregulation of endoplasmic reticulum stress. *PLoS One*. 2013;8(12):e82968.
94. Xie J, Cha SK, An SW, Kuro OM, Birnbaumer L, Huang CL. Cardioprotection by Klotho through downregulation of TRPC6 channels in the mouse heart. *Nat Commun*. 2012;3:1238.
95. Donate-Correa J, Mora-Fernandez C, Martinez-Sanz R, et al. Expression of FGF23/KLOTHO system in human vascular tissue. *Int J Cardiol*. 2013;165(1):179–183.
96. Lim K, Lu TS, Molostvov G, et al. Vascular Klotho deficiency potentiates the development of human artery calcification and mediates resistance to fibroblast growth factor 23. *Circulation*. 2012;125(18):2243–2255.
97. Moe SM. Klotho: a master regulator of cardiovascular disease? *Circulation*. 2012;125(18):2181–2183.
98. de Borst MH, Vervloet MG, ter Wee PM, Navis G. Cross talk between the renin-angiotensin-aldosterone system and vitamin D-FGF-23-klotho in chronic kidney disease. *J Am Soc Nephrol*. 2011;22(9):1603–1609.
99. Bonewald LF. The amazing osteocyte. *J Bone Miner Res*. 2011;26(2):229–238.
100. Parfitt AM. Targeted and nontargeted bone remodeling: relationship to basic multicellular unit origination and progression. *Bone*. 2002;30(1):5–7.
101. Kearns AE, Khosla S, Kostenuik PJ. Receptor activator of nuclear factor kappaB ligand and osteoprotegerin regulation of bone remodeling in health and disease. *Endocr Rev*. 2008;29(2):155–192.
102. Jamal SA, Ljunggren O, Stehman-Breen C, et al. Effects of denosumab on fracture and bone mineral density by level of kidney function. *J Bone Miner Res*. 2011;26(8):1829–1835.
103. Doumouchtsis KK, Kostakis AI, Doumouchtsis SK, et al. Associations between osteoprotegerin and femoral neck BMD in hemodialysis patients. *J Bone Miner Metabol*. 2008;26(1):66–72.
104. Padagas J, Colloton M, Shalhoub V, et al. The receptor activator of nuclear factor-kappaB ligand inhibitor osteoprotegerin is a bone-protective agent in a rat model of chronic renal insufficiency and hyperparathyroidism. *Calcif Tissue Int*. 2006;78(1):35–44.
105. Block GA, Bone HG, Fang L, Lee E, Padhi D. A single-dose study of denosumab in patients with various degrees of renal impairment. *J Bone Miner Res*. 2012;27(7):1471–1479.
106. Baron R, Kneissel M. WNT signaling in bone homeostasis and disease: from human mutations to treatments. *Nat Med*. 2013;19(2):179–192.
107. McColm J, Hu L, Womack T, Tang CC, Chiang AY. Single- and multiple-dose randomized studies of blosozumab, a monoclonal antibody against sclerostin, in healthy postmenopausal women. *J Bone Miner Res*. 2014;29(4):935–943.
108. McClung MR, Grauer A, Boonen S, et al. Romosozumab in postmenopausal women with low bone mineral density. *N Engl J Med*. 2014;370(5):412–420.
109. Cejka D, Herberth J, Branscum AJ, et al. Sclerostin and Dickkopf-1 in renal osteodystrophy. *Clin J Am Soc Nephrol*. 2011;6(4):877–882.
110. Sabbagh Y, Gracioli FG, O'Brien S, et al. Repression of osteocyte Wnt/beta-catenin signaling is an early event in the progression of renal osteodystrophy. *J Bone Miner Res*. 2012;27(8):1757–1772.
111. Kanbay M, Siritopol D, Saglam M, et al. Serum sclerostin and adverse outcomes in nondialyzed chronic kidney disease patients. *J Clin Endocrinol Metab*. 2014;99:E1854–E1861.
112. Gonçalves FLC, Elias RM, dos Reis LM, et al. Serum sclerostin is an independent predictor of mortality in hemodialysis patients. *BMC Nephrol*. 2014;15:190.
113. Viaene L, Behets GJ, Claes K, et al. Sclerostin: another bone-related protein related to all-cause mortality in haemodialysis? *Nephrol Dial Transplant*. 2013;28:3024–3030.
114. Miller PD, Adachi JD, Albergaria BH, et al. Efficacy and safety of romosozumab among postmenopausal women with osteoporosis and mild-to-moderate chronic kidney disease. *J Bone Miner Res*. 2022;37:1437–1445.
115. Moe SM, Chen NX, Newman CL, et al. Anti-sclerostin antibody treatment in a rat model of progressive renal osteodystrophy. *J Bone Miner Res*. 2015;30(3):499–509.
116. Fang Y, Ginsberg C, Seifert M, et al. CKD-induced wingless/integration1 inhibitors and phosphorus cause the CKD-mineral and bone disorder. *J Am Soc Nephrol*. 2014;25(8):1760–1773.
117. Guo J, Liu M, Yang D, et al. Suppression of Wnt signaling by Dkk1 attenuates PTH-mediated stromal cell response and new bone formation. *Cell Metab*. 2010;11(2):161–171.
118. Jilka RL, O'Brien CA, Bartell SM, Weinstein RS, Manolagas SC. Continuous elevation of PTH increases the number of osteoblasts via both osteoclast-dependent and -independent mechanisms. *J Bone Miner Res*. 2010;25(11):2427–2437.
119. Rhee Y, Lee EY, Lezcano V, et al. Resorption controls bone anabolism driven by parathyroid hormone (PTH) receptor signaling in osteocytes. *J Biol Chem*. 2013;288(41):29809–29820.
120. de Oliveira RB, Gracioli FG, dos Reis LM, et al. Disturbances of Wnt/beta-catenin pathway and energy metabolism in early CKD: effect of phosphate binders. *Nephrol Dial Transplant*. 2013;28(10):2510–2517.
121. Ibanez B, Badimon JJ, Garcia MJ. Diagnosis of atherosclerosis by imaging. *Am J Med*. 2009;122(1 Suppl):S15–S25.
122. Moe SM, Duan D, Doehle BP, O'Neill KD, Chen NX. Uremia induces the osteoblast differentiation factor Cbfa1 in human blood vessels. *Kidney Int*. 2003;63(3):1003–1011.
123. Shroff RC, McNair R, Figg N, et al. Dialysis accelerates medial vascular calcification in part by triggering smooth muscle cell apoptosis. *Circulation*. 2008;118(17):1748–1757.
124. Ducy P, Zhang R, Geoffroy V, Ridall AL, Karsenty G. Osf2/Cbfa1: a transcriptional activator of osteoblast differentiation. *Cell*. 1997;89(5):747–754.
125. Speer MY, Yang HY, Brabb T, et al. Smooth muscle cells give rise to osteochondrogenic precursors and chondrocytes in calcifying arteries. *Circ Res*. 2009;104(6):733–741.
126. Heath JM, Sun Y, Yuan K, et al. Activation of AKT by O-linked N-acetylglucosamine induces vascular calcification in diabetes mellitus. *Circ Res*. 2014;114(7):1094–1102.
127. Crouthamel MH, Lau WL, Leaf EM, et al. Sodium-dependent phosphate cotransporters and phosphate-induced calcification of vascular smooth muscle cells: redundant roles for PiT-1 and PiT-2. *Arterioscler Thromb Vasc Biol*. 2013;33(11):2625–2632.
128. Chen NX, Duan D, O'Neill KD, et al. The mechanisms of uremic serum-induced expression of bone matrix proteins in bovine vascular smooth muscle cells. *Kidney Int*. 2006;70(6):1046–1053.
129. Shanahan CM, Crouthamel MH, Kapustin A, Giachelli CM. Arterial calcification in chronic kidney disease: key roles for calcium and phosphate. *Circ Res*. 2011;109(6):697–711.
130. Jimbo R, Kawakami-Mori F, Mu S, et al. Fibroblast growth factor 23 accelerates phosphate-induced vascular calcification in the absence of Klotho deficiency. *Kidney Int*. 2014;85(5):1103–1111.
131. Amaya-Garrido A, Brunet M, Buffin-Meyer B, et al. Calprotectin is a contributor to and potential therapeutic target for vascular calcification in chronic kidney disease. *Sci Transl Med*. 2023;15(712):eabn5939.
132. Katsumata K, Kusano K, Hirata M, et al. Sevelamer hydrochloride prevents ectopic calcification and renal osteodystrophy in chronic renal failure rats. *Kidney Int*. 2003;64(2):441–450.
133. Cozzolino M, Dusso AS, Liapis H, et al. The effects of sevelamer hydrochloride and calcium carbonate on kidney calcification in uremic rats. *J Am Soc Nephrol*. 2002;13(9):2299–2308.
134. Neves KR, Gracioli FG, dos Reis LM, et al. Vascular calcification: contribution of parathyroid hormone in renal failure. *Kidney Int*. 2007;71(12):1262–1270.
135. Gracioli FG, Neves KR, dos Reis LM, et al. Phosphorus overload and PTH induce aortic expression of Runx2 in experimental uraemia. *Nephrol Dial Transplant*. 2009;24(5):1416–1421.
136. Price PA, June HH, Buckley JR, Williamson MK. SB 242784, a selective inhibitor of the osteoclastic V-H+ATPase, inhibits arterial calcification in the rat. *Circ Res*. 2002;91(6):547–552.
137. Lomashvili KA, Monier-Faugere MC, Wang X, Malluche HH, O'Neill WC. Effect of bisphosphonates on vascular calcification and bone metabolism in experimental renal failure. *Kidney Int*. 2009;75(6):617–625.
138. Moe SM, Seifert MF, Chen NX, et al. R-568 reduces ectopic calcification in a rat model of chronic kidney disease-mineral bone disorder (CKD-MBD). *Nephrol Dial Transplant*. 2009;24(8):2371–2377.
139. Allen MR, Chen NX, Gattone 2nd VH, et al. Skeletal effects of zoledronic acid in an animal model of chronic kidney disease. *Osteoporos Int*. 2013;24(4):1471–1481.

140. Davies MR, Lund RJ, Mathew S, Hruska KA. Low turnover osteodystrophy and vascular calcification are amenable to skeletal anabolism in an animal model of chronic kidney disease and the metabolic syndrome. *J Am Soc Nephrol*. 2005;16(4):917–928.
141. Fang Y, Ginsberg C, Sugatani T, Monier-Faugere MC, Malluche H, Hruska KA. Early chronic kidney disease-mineral bone disorder stimulates vascular calcification. *Kidney Int*. 2014;85(1):142–150.
142. Lomashvili KA, Wang X, O'Neill WC. Role of local versus systemic vitamin D receptors in vascular calcification. *Arterioscler Thromb Vasc Biol*. 2014;34(1):146–151.
143. Reynolds JL, Joannides AJ, Skepper JN, et al. Human vascular smooth muscle cells undergo vesicle-mediated calcification in response to changes in extracellular calcium and phosphate concentrations: a potential mechanism for accelerated vascular calcification in ESRD. *J Am Soc Nephrol*. 2004;15(11):2857–2867.
144. Chen NX, O'Neill KD, Chen X, Moe SM. Annexin-mediated matrix vesicle calcification in vascular smooth muscle cells. *J Bone Miner Res*. 2008;23(11):1798–1805.
145. Goettsch C, Hutcheson JD, Aikawa E. MicroRNA in cardiovascular calcification: focus on targets and extracellular vesicle delivery mechanisms. *Circ Res*. 2013;112(7):1073–1084.
146. Chen NX, Kiattisunthorn K, O'Neill KD, et al. Decreased microRNA is involved in the vascular remodeling abnormalities in chronic kidney disease (CKD). *PLoS One*. 2013;8(5):e64558.
147. Mackenzie NC, Staines KA, Zhu D, Genever P, Macrae VE. miRNA-221 and miRNA-222 synergistically function to promote vascular calcification. *Cell Biochem Funct*. 2014;32(2):209–216.
148. Ewence AE, Bootman M, Roderick HL, et al. Calcium phosphate crystals induce cell death in human vascular smooth muscle cells: a potential mechanism in atherosclerotic plaque destabilization. *Circ Res*. 2008;103(5):e28–e34.
149. Liu Y, Drozdov I, Shroff R, Beltran LE, Shanahan CM. Prelamin A accelerates vascular calcification via activation of the DNA damage response and senescence-associated secretory phenotype in vascular smooth muscle cells. *Circ Res*. 2013;112(10):e99–e109.
150. Dai XY, Zhao MM, Cai Y, et al. Phosphate-induced autophagy counteracts vascular calcification by reducing matrix vesicle release. *Kidney Int*. 2013;83(6):1042–1051.
151. Liu D, Cui W, Liu B, et al. Atorvastatin protects vascular smooth muscle cells from TGF-beta1-stimulated calcification by inducing autophagy via suppression of the beta-catenin pathway. *Cell Physiol Biochem*. 2014;33(1):129–141.
152. Giachelli CM. The emerging role of phosphate in vascular calcification. *Kidney Int*. 2009;75(9):890–897.
153. Murshed M, Harmey D, Millan JL, McKee MD, Karsenty G. Unique coexpression in osteoblasts of broadly expressed genes accounts for the spatial restriction of ECM mineralization to bone. *Genes Dev*. 2005;19(9):1093–1104.
154. Schinke T, Amendt C, Trindl A, Poschke O, Muller-Esterl W, Jahnhen-Dechent W. The serum protein alpha2-HS glycoprotein/fetuin inhibits apatite formation in vitro and in mineralizing calvaria cells. A possible role in mineralization and calcium homeostasis. *J Biol Chem*. 1996;271(34):20789–20796.
155. Reynolds JL, Skepper JN, McNair R, et al. Multifunctional roles for serum protein fetuin-A in inhibition of human vascular smooth muscle cell calcification. *J Am Soc Nephrol*. 2005;16(10):2920–2930.
156. Herrmann M, Kinkeldey A, Jahnhen-Dechent W. Fetuin-A function in systemic mineral metabolism. *Trends Cardiovasc Med*. 2012;22(8):197–201.
157. Smith ER, Ford ML, Tomlinson LA, et al. Serum calcification propensity predicts all-cause mortality in predialysis CKD. *J Am Soc Nephrol*. 2014;25(2):339–348.
158. Schafer C, Heiss A, Schwarz A, et al. The serum protein alpha 2-Heremans-Schmid glycoprotein/fetuin-A is a systemically acting inhibitor of ectopic calcification. *J Clin Invest*. 2003;112(3):357–366.
159. Westenfeld R, Schafer C, Kruger T, et al. Fetuin-A protects against atherosclerotic calcification in CKD. *J Am Soc Nephrol*. 2009;20(6):1264–1274.
160. Ketteler M, Bongartz P, Westenfeld R, et al. Association of low fetuin-A (AHSG) concentrations in serum with cardiovascular mortality in patients on dialysis: a cross-sectional study. *Lancet*. 2003;361(9360):827–833.
161. Scialla JJ, Kao WH, Crainiceanu C, et al. Biomarkers of vascular calcification and mortality in patients with ESRD. *Clin J Am Soc Nephrol*. 2014;9(4):745–755.
162. Mohamed ON, Mohamed MRM, Hassan IG, Alakkad AF, et al. The relationship of fetuin-A with coronary calcification, carotid atherosclerosis, and mortality risk in non-dialysis chronic kidney disease. *J Lipid Atheroscler*. 2024;13(2):194–211.
163. Moe SM, Reslerova M, Ketteler M, et al. Role of calcification inhibitors in the pathogenesis of vascular calcification in chronic kidney disease (CKD). *Kidney Int*. 2005;67(6):2295–2304.
164. Stenvinkel P, Wang K, Qureshi AR, et al. Low fetuin-A levels are associated with cardiovascular death: impact of variations in the gene encoding fetuin. *Kidney Int*. 2005;67(6):2383–2392.
165. Luo G, Ducy P, McKee MD, et al. Spontaneous calcification of arteries and cartilage in mice lacking matrix GLA protein. *Nature*. 1997;386(6620):78–81.
166. Beazley KE, Lima F, Borras T, Nurminskaya M. Attenuation of chondrogenic transformation in vascular smooth muscle by dietary quercetin in the MGP-deficient mouse model. *PLoS One*. 2013;8(9):e76210.
167. Ketteler M, Rothe H, Brandenburg VM, Westenfeld R. The K-factor in chronic kidney disease: biomarkers of calcification inhibition and beyond. *Nephrol Dial Transplant*. 2014;29(7):1267–1270.
168. McCabe KM, Booth SL, Fu X, et al. Dietary vitamin K and therapeutic warfarin alter the susceptibility to vascular calcification in experimental chronic kidney disease. *Kidney Int*. 2013;83(5):835–844.
169. Cranenburg EC, Brandenburg VM, Vermeer C, et al. Uncarboxylated matrix Gla protein (ucMGP) is associated with coronary artery calcification in haemodialysis patients. *Thromb Haemost*. 2009;101(2):359–366.
170. Nigwekar SU, Bloch DB, Nazarian RM, et al. Vitamin K-dependent carboxylation of matrix gla protein influences the risk of calciphylaxis. *J Am Soc Nephrol*. 2017;28(6):1717–1722.
171. Schurgers LJ, Joosen IA, Laufer EM, et al. Vitamin K-antagonists accelerate atherosclerotic calcification and induce a vulnerable plaque phenotype. *PLoS One*. 2012;7(8):e43229.
172. Westenfeld R, Krueger T, Schlieper G, et al. Effect of vitamin K2 supplementation on functional vitamin K deficiency in hemodialysis patients: a randomized trial. *Am J Kidney Dis*. 2012;59(2):186–195.
173. Saritas T, Reinartz S, Krüger T, Ketteler M, et al. Vitamin K1 and progression of cardiovascular calcifications in hemodialysis patients: the VitaVasK randomized controlled trial. *Clin Kidney J*. 2022;15(12):2300–2311.
174. Ketteler M, Schanz M, Shricker S. Vitamin K: should we supplement to protect the kidneys and the heart? *Nephrol Dial Transplant*. 2021;36:2196–2198.
175. Lomashvili KA, Garg P, Narisawa S, Millan JL, O'Neill WC. Up-regulation of alkaline phosphatase and pyrophosphate hydrolysis: potential mechanism for uremic vascular calcification. *Kidney Int*. 2008;73(9):1024–1030.
176. Rutsch F, Ruf N, Vaingankar S, et al. Mutations in ENPP1 are associated with 'idiopathic' infantile arterial calcification. *Nat Genet*. 2003;34(4):379–381.
177. Lomashvili KA, Khawandi W, O'Neill WC. Reduced plasma pyrophosphate levels in hemodialysis patients. *J Am Soc Nephrol*. 2005;16(8):2495–2500.
178. O'Neill WC, Sigrist MK, McIntyre CW. Plasma pyrophosphate and vascular calcification in chronic kidney disease. *Nephrol Dial Transplant*. 2010;25(1):187–191.
179. de Oliveira RB, Louvet L, Riser BL, et al. Peritoneal delivery of sodium pyrophosphate blocks the progression of pre-existing vascular calcification in uremic apolipoprotein-E knockout mice. *Calcif Tissue Int*. 2015;97(2):179–192.
180. Min H, Morony S, Sarosi I, et al. Osteoprotegerin reverses osteoporosis by inhibiting endosteal osteoclasts and prevents vascular calcification by blocking a process resembling osteoclastogenesis. *J Exp Med*. 2000;192(4):463–474.
181. Morony S, Tintut Y, Zhang Z, et al. Osteoprotegerin inhibits vascular calcification without affecting atherosclerosis in *ldlr*^{-/-} mice. *Circulation*. 2008;117(3):411–420.
182. Callegari A, Coons ML, Ricks JL, et al. Bone marrow- or vessel wall-derived osteoprotegerin is sufficient to reduce atherosclerotic lesion size and vascular calcification. *Arterioscler Thromb Vasc Biol*. 2013;33(11):2491–2500.
183. Callegari A, Coons ML, Ricks JL, Rosenfeld ME, Scatena M. Increased calcification in osteoprotegerin-deficient smooth muscle cells: dependence on receptor activator of NF-kappaB ligand and interleukin 6. *J Vasc Res*. 2014;51(2):118–131.

184. Deuell KA, Callegari A, Giachelli CM, Rosenfeld ME, Scatena M. RANKL enhances macrophage paracrine pro-calcific activity in high phosphate-treated smooth muscle cells: dependence on IL-6 and TNF- α . *J Vasc Res*. 2012;49(6):510–521.
185. Ben-Dov IZ, Galitzer H, Lavi-Moshayoff V, et al. The parathyroid is a target organ for FGF23 in rats. *J Clin Invest*. 2007;117(12):4003–4008.
186. Lavi-Moshayoff V, Wasserman G, Meir T, Silver J, Naveh-Many T. PTH increases FGF23 gene expression and mediates the high-FGF23 levels of experimental kidney failure: a bone parathyroid feedback loop. *Am J Physiol Ren Physiol*. 2010;299(4):F882–F889.
187. Rodriguez-Ortiz ME, Lopez I, Munoz-Castaneda JR, et al. Calcium deficiency reduces circulating levels of FGF23. *J Am Soc Nephrol*. 2012;23(7):1190–1197.
188. Moe SM, Chen NX, Newman CL, et al. A comparison of calcium to zoledronic acid for improvement of cortical bone in an animal model of CKD. *J Bone Miner Res*. 2014;29(4):902–910.
189. Sprague SM, Bellorin-Font E, Jorgetti V, et al. Diagnostic accuracy of bone turnover markers and bone histology in patients with CKD treated by dialysis. *Am J Kidney Dis*. 2016;67(4):559–566.
190. d'Amour P. Circulating PTH molecular forms: what we know and what we don't. *Kidney Int*. 2006;70:S29–S33.
191. Sturgeon CM, Sprague SM, Metcalfe W. Variation in parathyroid hormone immunoassay results—a critical governance issue in the management of chronic kidney disease. *Nephrol Dial Transplant*. 2011;26(11):3440–3445.
192. Group KDIGO-MW. KDIGO clinical practice guideline for the diagnosis, evaluation, prevention, and treatment of Chronic Kidney Disease-Mineral and Bone Disorder (CKD-MBD). *Kidney Int Suppl*. 2009;76(113):S1–S130.
193. Hollis BW. Measuring 25-hydroxyvitamin D in a clinical environment: challenges and needs. *Am J Clin Nutr*. 2008;88(2):507S–510S.
194. Sempos CT, Vesper HW, Phinney KW, Thienpont LM, Coates PM, Vitamin DSP. Vitamin D status as an international issue: national surveys and the problem of standardization. *Scand J Clin Lab Invest Suppl*. 2012;243:32–40.
195. Smith ER, McMahon LP, Holt SG. Fibroblast growth factor 23. *Ann Clin Biochem*. 2014;51(Pt 2):203–227.
196. Leaf DE, Jacob KA, Srivastava A, et al. Fibroblast growth factor 23 levels associate with AKI and death in critical illness. *J Am Soc Nephrol*. 2017;28(6):1877–1885.
197. van Ballegooijen AJ, Rhee EP, Elmariah S, de Boer IH, Kestenbaum B. Renal clearance of mineral metabolism biomarkers. *J Am Soc Nephrol*. 2016;27(2):392–397.
198. Kim HR, Nam BY, Kim DW, et al. Circulating alpha-klotho levels in CKD and relationship to progression. *Am J Kidney Dis*. 2013;61(6):899–909.
199. Seiler S, Wen M, Roth HJ, et al. Plasma Klotho is not related to kidney function and does not predict adverse outcome in patients with chronic kidney disease. *Kidney Int*. 2013;83(1):121–128.
200. Akimoto T, Shiizaki K, Sugase T, et al. The relationship between the soluble Klotho protein and the residual renal function among peritoneal dialysis patients. *Clin Exp Nephrol*. 2012;16(3):442–447.
201. Heijboer AC, Blankenstein MA, Hoenderop J, de Borst MH, Vervloet MG, consortium N. Laboratory aspects of circulating alpha-Klotho. *Nephrol Dial Transplant*. 2013;28(9):2283–2287.
202. Thambiah S, Roplekar R, Manghat P, et al. Circulating sclerostin and Dickkopf-1 (DKK1) in predialysis chronic kidney disease (CKD): relationship with bone density and arterial stiffness. *Calcif Tissue Int*. 2012;90(6):473–480.
203. Cejka D, Marculescu R, Kozakowski N, et al. Renal elimination of sclerostin increases with declining kidney function. *J Clin Endocrinol Metab*. 2014;99(1):248–255.
204. Brandenburg VM, Kramann R, Koos R, et al. Relationship between sclerostin and cardiovascular calcification in hemodialysis patients: a cross-sectional study. *BMC Nephrol*. 2013;14:219.
205. Cejka D, Jager-Lansky A, Kieweg H, et al. Sclerostin serum levels correlate positively with bone mineral density and microarchitecture in haemodialysis patients. *Nephrol Dial Transplant*. 2012;27(1):226–230.
206. Bakkaloglu SA, Wesseling-Perry K, Pereira RC, et al. Value of the new bone classification system in pediatric renal osteodystrophy. *Clin J Am Soc Nephrol*. 2010;5(10):1860–1866.
207. Moorthi RN, Moe SM. Recent advances in the noninvasive diagnosis of renal osteodystrophy. *Kidney Int*. 2013;84(5):886–894.
208. Nickolas TL, Cremers S, Zhang A, et al. Discriminants of prevalent fractures in chronic kidney disease. *J Am Soc Nephrol*. 2011;22(8):1560–1572.
209. Yamada S, Inaba M, Kurajoh M, et al. Utility of serum tartrate-resistant acid phosphatase (TRACP5b) as a bone resorption marker in patients with chronic kidney disease: independence from renal dysfunction. *Clin Endocrinol*. 2008;69(2):189–196.
210. Jørgensen HS, Behets G, Viaene L, et al. Diagnostic accuracy of noninvasive bone turnover markers in renal osteodystrophy. *Am J Kidney Dis*. 2022;79:667–676.
211. Dempster DW, Compston JE, Drezner MK, et al. Standardized nomenclature, symbols, and units for bone histomorphometry: a 2012 update of the report of the ASBMR Histomorphometry Nomenclature Committee. *J Bone Miner Res*. 2013;28(1):2–17.
212. Sherrard DJ, Hercz G, Pei Y, et al. The spectrum of bone disease in end-stage renal failure—an evolving disorder. *Kidney Int*. 1993;43(2):436–442.
213. Malluche HH, Mawad HW, Monier-Faugere MC. Renal osteodystrophy in the first decade of the new millennium: analysis of 630 bone biopsies in black and white patients. *J Bone Miner Res*. 2011;26(6):1368–1376.
214. Bird ST, Smith ER, Gelperin K, Jung TH, et al. Severe hypocalcemia with denosumab among older female dialysis-dependent patients. *JAMA*. 2024;331(6):491–499.
215. Alfrey AC. Aluminum and renal disease. *Contrib Nephrol*. 1993;102:110–124.
216. National Kidney F. K/DOQI clinical practice guidelines for bone metabolism and disease in chronic kidney disease. *Am J Kidney Dis*. 2003;42(4 suppl 3):S1–S201.
217. Jorgetti V. Review article: bone biopsy in chronic kidney disease: patient level end-point or just another test? *Nephrology*. 2009;14(4):404–407.
218. Andress DL. Adynamic bone in patients with chronic kidney disease. *Kidney Int*. 2008;73(12):1345–1354.
219. Moester MJ, Papapoulos SE, Lowik CW, van Bezooijen RL. Sclerostin: current knowledge and future perspectives. *Calcif Tissue Int*. 2010;87(2):99–107.
220. Ferreira JC, Ferrari GO, Neves KR, et al. Effects of dietary phosphate on adynamic bone disease in rats with chronic kidney disease—role of sclerostin? *PLoS One*. 2013;8(11):e79721.
221. Slatopolsky E, Finch J, Clay P, et al. A novel mechanism for skeletal resistance in uremia. *Kidney Int*. 2000;58(2):753–761.
222. Ott SM. Bone disease in CKD. *Curr Opin Nephrol Hypertens*. 2012;21(4):376–381.
223. Lund RJ, Davies MR, Brown AJ, Hruska KA. Successful treatment of an adynamic bone disorder with bone morphogenetic protein-7 in a renal ablation model. *J Am Soc Nephrol*. 2004;15(2):359–369.
224. Coco M, Rush H. Increased incidence of hip fractures in dialysis patients with low serum parathyroid hormone. *Am J Kidney Dis*. 2000;36(6):1115–1121.
225. Ott SM. Bone histomorphometry in renal osteodystrophy. *Semin Nephrol*. 2009;29(2):122–132.
226. Kurz P, Monier-Faugere MC, Bogner B, et al. Evidence for abnormal calcium homeostasis in patients with adynamic bone disease. *Kidney Int*. 1994;46(3):855–861.
227. London GM, Marchais SJ, Guerin AP, Boutouyrie P, Metivier F, de Vernejoul MC. Association of bone activity, calcium load, aortic stiffness, and calcifications in ESRD. *J Am Soc Nephrol*. 2008;19(9):1827–1835.
228. Barreto FC, Barreto DV, Moyses RM, et al. Osteoporosis in hemodialysis patients revisited by bone histomorphometry: a new insight into an old problem. *Kidney Int*. 2006;69(10):1852–1857.
229. Barreto DV, Barreto FC, Carvalho AB, et al. Association of changes in bone remodeling and coronary calcification in hemodialysis patients: a prospective study. *Am J Kidney Dis*. 2008;52(6):1139–1150.
230. Hui SL, Slemenda CW, Johnston Jr CC. Age and bone mass as predictors of fracture in a prospective study. *J Clin Invest*. 1988;81(6):1804–1809.
231. Jamal SA, Hayden JA, Beyene J. Low bone mineral density and fractures in long-term hemodialysis patients: a meta-analysis. *Am J Kidney Dis*. 2007;49(5):674–681.

232. Ambrus C, Almasi C, Berta K, et al. Vitamin D insufficiency and bone fractures in patients on maintenance hemodialysis. *Int Urol Nephrol*. 2011;43(2):475–482.
233. Yenchek RH, Ix JH, Shlipak MG, et al. Bone mineral density and fracture risk in older individuals with CKD. *Clin J Am Soc Nephrol*. 2012;7(7):1130–1136.
234. Shevroja E, Reginster JY, Lamy O, Al-Daghri N, et al. Update on the clinical use of trabecular bone score (TBS) in the management of osteoporosis: results of an expert group meeting organized by the European society for clinical and economic aspects of osteoporosis, osteoarthritis and musculoskeletal diseases (ESCEO), and the international osteoporosis foundation (IOF) under the auspices of WHO collaborating center for epidemiology of musculoskeletal health and aging. *Osteoporos Int*. 2023;34(9):1501–1529.
235. Naylor KL, Lix L, Hans D, Garg AX, et al. Trabecular bone score in kidney transplant recipients. *Osteoporos Int*. 2016;27(3):1115–1121.
236. Yun HJ, Ryoo SR, Kim JE, Choi YJ, et al. Trabecular bone score may indicate chronic kidney disease-mineral and bone disorder (CKD-MBD) phenotypes in hemodialysis patients: a prospective observational study. *BMC Nephrol*. 2020;21(1):299.
237. Ketteler MBG, Evenepoel P, et al. Executive summary of the 2017 KDIGO chronic kidney disease-mineral and bone disorder (CKD-MBD) guideline update: what's changed and why it matters. *Kidney Int*. 2017;92(6):26–36.
238. Torres A, Lorenzo V, Gonzalez-Posada JM. Comparison of histomorphometry and computerized tomography of the spine in quantitating trabecular bone in renal osteodystrophy. *Nephron*. 1986;44(4):282–287.
239. Jamal SA, Gilbert J, Gordon C, Bauer DC. Cortical pQCT measures are associated with fractures in dialysis patients. *J Bone Miner Res*. 2006;21(4):543–548.
240. Cejka D, Patsch JM, Weber M, et al. Bone microarchitecture in hemodialysis patients assessed by HR-pQCT. *Clin J Am Soc Nephrol*. 2011;6(9):2264–2271.
241. West SL, Jamal SA. Determination of bone architecture and strength in men and women with stage 5 chronic kidney disease. *Semin Dial*. 2012;25(4):397–402.
242. Nickolas TL, Stein E, Cohen A, et al. Bone mass and microarchitecture in CKD patients with fracture. *J Am Soc Nephrol*. 2010;21(8):1371–1380.
243. Jamal S, Cheung AM, West S, Lok C. Bone mineral density by DXA and HR pQCT can discriminate fracture status in men and women with stages 3 to 5 chronic kidney disease. *Osteoporos Int*. 2012;23(12):2805–2813.
244. Levy RV, McMahon DJ, Agarwal S, et al. Comprehensive associations between acidosis and the skeleton in patients with kidney disease. *J Am Soc Nephrol*. 2023;34:668–681.
245. Hopper TA, Wehrli FW, Saha PK, et al. Quantitative microcomputed tomography assessment of intratrabecular, intertrabecular, and cortical bone architecture in a rat model of severe renal osteodystrophy. *J Comput Assist Tomogr*. 2007;31(2):320–328.
246. Wehrli FW, Leonard MB, Saha PK, Gomberg BR. Quantitative high-resolution magnetic resonance imaging reveals structural implications of renal osteodystrophy on trabecular and cortical bone. *J Magn Reson Imag*. 2004;20(1):83–89.
247. Kauppila LI, Polak JF, Cupples LA, Hannan MT, Kiel DP, Wilson PW. New indices to classify location, severity and progression of calcific lesions in the abdominal aorta: a 25-year follow-up study. *Atherosclerosis*. 1997;132(2):245–250.
248. London GM, Guerin AP, Marchais SJ, Metivier F, Pannier B, Adda H. Arterial media calcification in end-stage renal disease: impact on all-cause and cardiovascular mortality. *Nephrol Dial Transplant*. 2003;18(9):1731–1740.
249. Joshi SB, Okabe T, Roswell RO, et al. Accuracy of computed tomographic angiography for stenosis quantification using quantitative coronary angiography or intravascular ultrasound as the gold standard. *Am J Cardiol*. 2009;104(8):1047–1051.
250. Bellasi A, Ferramosca E, Muntner P, et al. Correlation of simple imaging tests and coronary artery calcium measured by computed tomography in hemodialysis patients. *Kidney Int*. 2006;70(9):1623–1628.
251. Verbeke F, Van Biesen W, Honkanen E, et al. Prognostic value of aortic stiffness and calcification for cardiovascular events and mortality in dialysis patients: outcome of the calcification outcome in renal disease (CORD) study. *Clin J Am Soc Nephrol*. 2011;6(1):153–159.
252. Bellasi A, Raggi P. Vascular imaging in chronic kidney disease. *Curr Opin Nephrol Hypertens*. 2012;21(4):382–388.
253. Bellasi A, Block GA, Ferramosca E, Ratti C, Raggi P. Integration of clinical and imaging data to predict death in hemodialysis patients. *Hemodial Int*. 2013;17(1):12–18.
254. Kestenbaum B, Sampson JN, Rudser KD, et al. Serum phosphate levels and mortality risk among people with chronic kidney disease. *J Am Soc Nephrol*. 2005;16(2):520–528.
255. Menon V, Greene T, Pereira AA, et al. Relationship of phosphorus and calcium-phosphorus product with mortality in CKD. *Am J Kidney Dis*. 2005;46(3):455–463.
256. Mehrotra R, Peralta CA, Chen SC, et al. No independent association of serum phosphorus with risk for death or progression to end-stage renal disease in a large screen for chronic kidney disease. *Kidney Int*. 2013;84(5):989–997.
257. Isakova T, Gutierrez OM, Chang Y, et al. Phosphorus binders and survival on hemodialysis. *J Am Soc Nephrol*. 2009;20(2):388–396.
258. Lopes AA, Tong L, Thumma J, et al. Phosphate binder use and mortality among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study (DOPPS): evaluation of possible confounding by nutritional status. *Am J Kidney Dis*. 2012;60(1):90–101.
259. Cannata-Andia JB, Fernandez-Martin JL, Locatelli F, et al. Use of phosphate-binding agents is associated with a lower risk of mortality. *Kidney Int*. 2013;84(5):998–1008.
260. Palmer SC, Hayen A, Macaskill P, et al. Serum levels of phosphorus, parathyroid hormone, and calcium and risks of death and cardiovascular disease in individuals with chronic kidney disease: a systematic review and meta-analysis. *JAMA*. 2011;305(11):1119–1127.
261. Stevens LA, Djurdjev O, Cardew S, Cameron EC, Levin A. Calcium, phosphate, and parathyroid hormone levels in combination and as a function of dialysis duration predict mortality: evidence for the complexity of the association between mineral metabolism and outcomes. *J Am Soc Nephrol*. 2004;15(3):770–779.
262. Tentori F, Blayney MJ, Albert JM, et al. Mortality risk for dialysis patients with different levels of serum calcium, phosphorus, and PTH: the Dialysis Outcomes and Practice Patterns Study (DOPPS). *Am J Kidney Dis*. 2008;52(3):519–530.
263. Block GA, Kilpatrick RD, Lowe KA, Wang W, Danese MD. CKD-mineral and bone disorder and risk of death and cardiovascular hospitalization in patients on hemodialysis. *Clin J Am Soc Nephrol*. 2013;8(12):2132–2140.
264. Isakova T, Xie H, Yang W, et al. Fibroblast growth factor 23 and risks of mortality and end-stage renal disease in patients with chronic kidney disease. *JAMA*. 2011;305(23):2432–2439.
265. Fliser D, Kollerits B, Neyer U, et al. Fibroblast growth factor 23 (FGF23) predicts progression of chronic kidney disease: the Mild to Moderate Kidney Disease (MMKD) Study. *J Am Soc Nephrol*. 2007;18(9):2600–2608.
266. Gutierrez OM, Januzzi JL, Isakova T, et al. Fibroblast growth factor 23 and left ventricular hypertrophy in chronic kidney disease. *Circulation*. 2009;119(19):2545–2552.
267. Smith K, deFilippi C, Isakova T, et al. Fibroblast growth factor 23, high-sensitivity cardiac troponin, and left ventricular hypertrophy in CKD. *Am J Kidney Dis*. 2013;61(1):67–73.
268. Seiler S, Reichart B, Roth D, Seibert E, Fliser D, Heine GH. FGF-23 and future cardiovascular events in patients with chronic kidney disease before initiation of dialysis treatment. *Nephrol Dial Transplant*. 2010;25(12):3983–3989.
269. Nakano C, Hamano T, Fujii N, et al. Intact fibroblast growth factor 23 levels predict incident cardiovascular event before but not after the start of dialysis. *Bone*. 2012;50(6):1266–1274.
270. Gutierrez OM, Mannstadt M, Isakova T, et al. Fibroblast growth factor 23 and mortality among patients undergoing hemodialysis. *N Engl J Med*. 2008;359(6):584–592.
271. Olason H, Qureshi AR, Miyamoto T, et al. Relation between serum fibroblast growth factor-23 level and mortality in incident dialysis patients: are gender and cardiovascular disease confounding the relationship? *Nephrol Dial Transplant*. 2010;25(9):3033–3038.
272. Hsu HJ, Wu MS. Fibroblast growth factor 23: a possible cause of left ventricular hypertrophy in hemodialysis patients. *Am J Med Sci*. 2009;337(2):116–122.

273. Jean G, Terrat JC, Vanel T, et al. High levels of serum fibroblast growth factor (FGF)-23 are associated with increased mortality in long haemodialysis patients. *Nephrol Dial Transplant*. 2009;24(9):2792–2796.
274. Moe SM, Chertow GM, Parfrey PS, et al. Cinacalcet, fibroblast growth factor-23, and cardiovascular disease in hemodialysis: the evaluation of cinacalcet HCl therapy to lower cardiovascular events (EVOLVE) trial. *Circulation*. 2015;132(1):27–39.
275. Dawson-Hughes B, Heaney RP, Holick MF, Lips P, Meunier PJ, Vieth R. Estimates of optimal vitamin D status. *Osteoporos Int*. 2005;16(7):713–716.
276. Holick MF. Vitamin D status: measurement, interpretation, and clinical application. *Ann Epidemiol*. 2009;19(2):73–78.
277. Ross AC, Manson JE, Abrams SA, et al. The 2011 report on dietary reference intakes for calcium and vitamin D from the Institute of Medicine: what clinicians need to know. *J Clin Endocrinol Metab*. 2011;96(1):53–58.
278. Holick MF, Binkley NC, Bischoff-Ferrari HA, et al. Evaluation, treatment, and prevention of vitamin D deficiency: an Endocrine Society clinical practice guideline. *J Clin Endocrinol Metab*. 2011;96(7):1911–1930.
279. Hossein-Nezhad A, Holick MF. Vitamin D for health: a global perspective. *Mayo Clin Proc*. 2013;88(7):720–755.
280. LaClair RE, Hellman RN, Karp SL, et al. Prevalence of calcidiol deficiency in CKD: a cross-sectional study across latitudes in the United States. *Am J Kidney Dis*. 2005;45(6):1026–1033.
281. Elder GJ, Mackun K. 25-Hydroxyvitamin D deficiency and diabetes predict reduced BMD in patients with chronic kidney disease. *J Bone Miner Res*. 2006;21(11):1778–1784.
282. Gonzalez EA, Sachdeva A, Oliver DA, Martin KJ. Vitamin D insufficiency and deficiency in chronic kidney disease. A single center observational study. *Am J Nephrol*. 2004;24(5):503–510.
283. Levin A, Bakris GL, Molitch M, et al. Prevalence of abnormal serum vitamin D, PTH, calcium, and phosphorus in patients with chronic kidney disease: results of the study to evaluate early kidney disease. *Kidney Int*. 2007;71(1):31–38.
284. Ennis JL, Worcester EM, Coe FL, Sprague SM. Current recommended 25-hydroxyvitamin D targets for chronic kidney disease management may be too low. *J Nephrol*. 2016;29(1):63–70.
285. Dhillon-Jhattu S, McGill RL, Ennis JL, et al. Vitamin D and parathyroid hormone levels in CKD. *Am J Kidney Dis*. 2023;81:122–124.279.
286. Wolf M, Shah A, Gutierrez O, et al. Vitamin D levels and early mortality among incident hemodialysis patients. *Kidney Int*. 2007;72(8):1004–1013.
287. Wang AY, Lam CW, Sanderson JE, et al. Serum 25-hydroxyvitamin D status and cardiovascular outcomes in chronic peritoneal dialysis patients: a 3-y prospective cohort study. *Am J Clin Nutr*. 2008;87(6):1631–1638.
288. Shoben AB, Rudser KD, de Boer IH, Young B, Kestenbaum B. Association of oral calcitriol with improved survival in nondialyzed CKD. *J Am Soc Nephrol*. 2008;19(8):1613–1619.
289. Teng M, Wolf M, Ofsthun MN, et al. Activated injectable vitamin D and hemodialysis survival: a historical cohort study. *J Am Soc Nephrol*. 2005;16(4):1115–1125.
290. Melamed ML, Eustace JA, Plantinga L, et al. Changes in serum calcium, phosphate, and PTH and the risk of death in incident dialysis patients: a longitudinal study. *Kidney Int*. 2006;70(2):351–357.
291. Zheng Z, Shi H, Jia J, Li D, Lin S. Vitamin D supplementation and mortality risk in chronic kidney disease: a meta-analysis of 20 observational studies. *BMC Nephrol*. 2013;14:199.
292. Pilz S, Iodice S, Zittermann A, Grant WB, Gandini S. Vitamin D status and mortality risk in CKD: a meta-analysis of prospective studies. *Am J Kidney Dis*. 2011;58(3):374–382.
293. Li YC, Kong J, Wei M, Chen ZF, Liu SQ, Cao LP. 1,25-Dihydroxyvitamin D(3) is a negative endocrine regulator of the renin-angiotensin system. *J Clin Invest*. 2002;110(2):229–238.
294. Bucharles S, Barberato SH, Stingham AE, et al. Hypovitaminosis D is associated with systemic inflammation and concentric myocardial geometric pattern in hemodialysis patients with low iPTH levels. *Nephron Clin Pract*. 2011;118(4):c384–c391.
295. Xiang W, Kong J, Chen S, et al. Cardiac hypertrophy in vitamin D receptor knockout mice: role of the systemic and cardiac renin-angiotensin systems. *Am J Physiol Endocrinol Metab*. 2005;288(1):E125–E132.
296. de Boer IH, Ioannou GN, Kestenbaum B, Brunzell JD, Weiss NS. 25-Hydroxyvitamin D levels and albuminuria in the third national health and nutrition examination survey (NHANES III). *Am J Kidney Dis*. 2007;50(1):69–77.
297. Isakova T, Gutierrez OM, Patel NM, Andress DL, Wolf M, Levin A. Vitamin D deficiency, inflammation, and albuminuria in chronic kidney disease: complex interactions. *J Ren Nutr*. 2011;21(4):295–302.
298. Freundlich M, Quiroz Y, Zhang Z, et al. Suppression of renin-angiotensin gene expression in the kidney by paricalcitol. *Kidney Int*. 2008;74(11):1394–1402.
299. Heart Protection Study Collaborative G. MRC/BHF Heart Protection Study of cholesterol lowering with simvastatin in 20,536 high-risk individuals: a randomised placebo-controlled trial. *Lancet*. 2002;360(9326):7–22.
300. Agarwal R, Acharya M, Tian J, et al. Antiproteinuric effect of oral paricalcitol in chronic kidney disease. *Kidney Int*. 2005;68(6):2823–2828.
301. Fishbane S, Chittineni H, Packman M, Dutka P, Ali N, Durie N. Oral paricalcitol in the treatment of patients with CKD and proteinuria: a randomized trial. *Am J Kidney Dis*. 2009;54(4):647–652.
302. de Zeeuw D, Agarwal R, Amdahl M, et al. Selective vitamin D receptor activation with paricalcitol for reduction of albuminuria in patients with type 2 diabetes (VITAL study): a randomised controlled trial. *Lancet*. 2010;376(9752):1543–1551.
303. Chitalia N, Recio-Mayoral A, Kaski JC, Banerjee D. Vitamin D deficiency and endothelial dysfunction in non-dialysis chronic kidney disease patients. *Atherosclerosis*. 2012;220(1):265–268.
304. Bucharles S, Barberato SH, Stingham AEM, et al. Impact of cholecalciferol treatment on biomarkers of inflammation and myocardial structure in hemodialysis patients without hyperparathyroidism. *J Ren Nutr*. 2012;22(2):284–291.
305. Matias PJ, Jorge C, Ferreira C, et al. Cholecalciferol supplementation in hemodialysis patients: effects on mineral metabolism, inflammation, and cardiac dimension parameters. *Clin J Am Soc Nephrol*. 2010;5(5):905–911.
306. Thadhani R, Appelbaum E, Pritchett Y, et al. Vitamin D therapy and cardiac structure and function in patients with chronic kidney disease: the PRIMO randomized controlled trial. *JAMA*. 2012;307(7):674–684.
307. Canziani ME, Tomiyama C, Higa A, Draibe SA, Carvalho AB. Fibroblast growth factor 23 in chronic kidney disease: bridging the gap between bone mineral metabolism and left ventricular hypertrophy. *Blood Purif*. 2011;31(1–3):26–32.
308. Nickolas TL, McMahon DJ, Shane E. Relationship between moderate to severe kidney disease and hip fracture in the United States. *J Am Soc Nephrol*. 2006;17(11):3223–3232.
309. Leonard MB. A structural approach to skeletal fragility in chronic kidney disease. *Semin Nephrol*. 2009;29(2):133–143.
310. Nitsch D, Mylne A, Roderick PJ, Smeeth L, Hubbard R, Fletcher A. Chronic kidney disease and hip fracture-related mortality in older people in the UK. *Nephrol Dial Transplant*. 2009;24(5):1539–1544.
311. Elliott MJ, James MT, Quinn RR, et al. Estimated GFR and fracture risk: a population-based study. *Clin J Am Soc Nephrol*. 2013;8(8):1367–1376.
312. Kim SM, Liu S, Long J, Montez-Rath ME, Leonard MB, Chertow GM. Declining rates of hip fracture in end-stage renal disease: analysis from the 2003–2011 nationwide inpatient sample. *J Bone Miner Res*. 2017;32(11):2297–2303.
313. Alem AM, Sherrard DJ, Gillen DL, et al. Increased risk of hip fracture among patients with end-stage renal disease. *Kidney Int*. 2000;58(1):396–399.
314. Chen YJ, Kung PT, Wang YH, et al. Greater risk of hip fracture in hemodialysis than in peritoneal dialysis. *Osteoporos Int*. 2014;25(5):1513–1518.
315. Danese MD, Kim J, Doan QV, Dylan M, Griffiths R, Chertow GM. PTH and the risks for hip, vertebral, and pelvic fractures among patients on dialysis. *Am J Kidney Dis*. 2006;47(1):149–156.
316. Nickolas TL, Stein EM, Dworakowski E, et al. Rapid cortical bone loss in patients with chronic kidney disease. *J Bone Miner Res*. 2013;28(8):1811–1820.
317. Moe SM, Abdalla S, Chertow GM, et al. Effects of cinacalcet on fracture events in patients receiving hemodialysis: the EVOLVE trial. *J Am Soc Nephrol*. 2015;26(6):1466–1475.

318. Stehman-Breen CO, Sherrard DJ, Alem AM, et al. Risk factors for hip fracture among patients with end-stage renal disease. *Kidney Int.* 2000;58(5):2200–2205.
319. Jadoul M, Albert JM, Akiba T, et al. Incidence and risk factors for hip or other bone fractures among hemodialysis patients in the Dialysis Outcomes and Practice Patterns Study. *Kidney Int.* 2006;70(7):1358–1366.
320. Kaji H, Suzuki M, Yano S, et al. Risk factors for hip fracture in hemodialysis patients. *Am J Nephrol.* 2002;22(4):325–331.
321. West SL, Jamal SA, Lok CE. Tests of neuromuscular function are associated with fractures in patients with chronic kidney disease. *Nephrol Dial Transplant.* 2012;27(6):2384–2388.
322. Ibels LS, Alfrey AC, Huffer WE, Craswell PW, Anderson JT, Weil 3rd R. Arterial calcification and pathology in uremic patients undergoing dialysis. *Am J Med.* 1979;66(5):790–796.
323. Schwarz U, Buzello M, Ritz E, et al. Morphology of coronary atherosclerotic lesions in patients with end-stage renal failure. *Nephrol Dial Transplant.* 2000;15(2):218–223.
324. Gross ML, Meyer HP, Ziebart H, et al. Calcification of coronary intima and media: immunohistochemistry, backscatter imaging, and x-ray analysis in renal and nonrenal patients. *Clin J Am Soc Nephrol.* 2007;2(1):121–134.
325. Moe SM, O'Neill KD, Duan D, et al. Medial artery calcification in ESRD patients is associated with deposition of bone matrix proteins. *Kidney Int.* 2002;61(2):638–647.
326. Braun J, Oldendorf M, Moshage W, Heidler R, Zeitler E, Luft FC. Electron beam computed tomography in the evaluation of cardiac calcification in chronic dialysis patients. *Am J Kidney Dis.* 1996;27(3):394–401.
327. Block GA, Spiegel DM, Ehrlich J, et al. Effects of sevelamer and calcium on coronary artery calcification in patients new to hemodialysis. *Kidney Int.* 2005;68(4):1815–1824.
328. Mehrotra R, Budoff M, Christenson P, et al. Determinants of coronary artery calcification in diabetics with and without nephropathy. *Kidney Int.* 2004;66(5):2022–2031.
329. Block GA, Raggi P, Bellasi A, Kooienga L, Spiegel DM. Mortality effect of coronary calcification and phosphate binder choice in incident hemodialysis patients. *Kidney Int.* 2007;71(5):438–441.
330. Bellasi A, Kooienga L, Block GA, Veledar E, Spiegel DM, Raggi P. How long is the warranty period for nil or low coronary artery calcium in patients new to hemodialysis? *J Nephrol.* 2009;22(2):255–262.
331. Hutchison AJ, Whitehouse RW, Boulton HF, et al. Correlation of bone histology with parathyroid hormone, vitamin D3, and radiology in end-stage renal disease. *Kidney Int.* 1993;44(5):1071–1077.
332. London GM, Marty C, Marchais SJ, Guerin AP, Metivier F, de Vernejoul MC. Arterial calcifications and bone histomorphometry in end-stage renal disease. *J Am Soc Nephrol.* 2004;15(7):1943–1951.
333. Cejka D, Weber M, Diarra D, Reiter T, Kainberger F, Haas M. Inverse association between bone microarchitecture assessed by HR-pQCT and coronary artery calcification in patients with end-stage renal disease. *Bone.* 2014;64:33–38.
334. Sprague SM, Josephson MA. Bone disease after kidney transplantation. *Semin Nephrol.* 2004;24(1):82–90.
335. Alshayeb HM, Josephson MA, Sprague SM. CKD-mineral and bone disorder management in kidney transplant recipients. *Am J Kidney Dis.* 2013;61(2):310–325.
336. Haffner D, Schuler U. Metabolic bone disease after renal transplantation. *Curr Opin Pediatr.* 2014;26(2):198–206.
337. Molnar MZ, Naser MS, Rhee CM, Kalantar-Zadeh K, Bunnapradist S. Bone and mineral disorders after kidney transplantation: therapeutic strategies. *Transplant Rev.* 2014;28(2):56–62.
338. Sprague SM, Belozeroff V, Danese MD, Martin LP, Olgaard K. Abnormal bone and mineral metabolism in kidney transplant patients—a review. *Am J Nephrol.* 2008;28(2):246–253.
339. Trombetti A, Richert L, Hadaya K, et al. Early post-transplantation hypophosphatemia is associated with elevated FGF-23 levels. *Eur J Endocrinol.* 2011;164(5):839–847.
340. Ramsey-Goldman R, Dunn JE, Dunlop DD, et al. Increased risk of fracture in patients receiving solid organ transplants. *J Bone Miner Res.* 1999;14(3):456–463.
341. Chiu MY, Sprague SM, Bruce DS, Woodle ES, Thistlethwaite Jr JR, Josephson MA. Analysis of fracture prevalence in kidney-pancreas allograft recipients. *J Am Soc Nephrol.* 1998;9(4):677–683.
342. Nikkel LE, Hollenbeak CS, Fox EJ, Uemura T, Ghahramani N. Risk of fractures after renal transplantation in the United States. *Transplantation.* 2009;87(12):1846–1851.
343. Nair SS, Lenihan CR, Montez-Rath ME, Lowenberg DW, Chertow GM, Winkelmayer WC. Temporal trends in the incidence, treatment and outcomes of hip fracture after first kidney transplantation in the United States. *Am J Transplant.* 2014;14(4):943–951.
344. Edwards BJ, Desai A, Tsai J, et al. Elevated incidence of fractures in solid-organ transplant recipients on glucocorticoid-sparing immunosuppressive regimens. *J Osteoporos.* 2011;2011:591793.
345. Naylor KL, Jamal SA, Zou G, et al. Fracture incidence in adult kidney transplant recipients. *Transplantation.* 2016;100(1):167–175.
346. Julian BA, Laskow DA, Dubovsky J, Dubovsky EV, Curtis JJ, Quarles LD. Rapid loss of vertebral mineral density after renal transplantation. *N Engl J Med.* 1991;325(8):544–550.
347. Rojas E, Carlini RG, Clesca P, et al. The pathogenesis of osteodystrophy after renal transplantation as detected by early alterations in bone remodeling. *Kidney Int.* 2003;63(5):1915–1923.
348. Monier-Faugere MC, Mawad H, Qi Q, Friedler RM, Malluche HH. High prevalence of low bone turnover and occurrence of osteomalacia after kidney transplantation. *J Am Soc Nephrol.* 2000;11(6):1093–1099.
349. Lehmann G, Ott U, Stein G, Steiner T, Wolf G. Renal osteodystrophy after successful renal transplantation: a histomorphometric analysis in 57 patients. *Transplant Proc.* 2007;39(10):3153–3158.
350. Pereira RC, Delany AM, Khouzam NM, et al. Primary osteoblast-like cells from patients with end-stage kidney disease reflect gene expression, proliferation, and mineralization characteristics ex vivo. *Kidney Int.* 2015;87:593–601.
351. Araújo MJCLN, Bacelar Marques ID, Gracioli FG, et al. Comparison of serum levels with bone content and gene expression indicate a contradictory effect of kidney transplantation on sclerostin. *Kidney Int.* 2019;96:1100–1104.
352. Cueto-Manzano AM, Konel S, Crowley V, et al. Bone histopathology and densitometry comparison between cyclosporine a monotherapy and prednisolone plus azathioprine dual immunosuppression in renal transplant patients. *Transplantation.* 2003;75(12):2053–2058.
353. Cueto-Manzano AM, Konel S, Hutchison AJ, et al. Bone loss in long-term renal transplantation: histopathology and densitometry analysis. *Kidney Int.* 1999;55(5):2021–2029.
354. Josephson MA, Schumm LP, Chiu MY, Marshall C, et al. Calcium and calcitriol prophylaxis attenuates posttransplant bone loss. *Transplantation.* 2004;78(8):1233–1236.
355. Akaberi S, Simonsen O, Lindergard B, Nyberg G. Can DXA predict fractures in renal transplant patients? *Am J Transplant.* 2008;8(12):2647–2651.
356. Iyer SP, Nikkel LE, Nishiyama KK, et al. Kidney transplantation with early corticosteroid withdrawal: paradoxical effects at the central and peripheral skeleton. *J Am Soc Nephrol.* 2014;25(6):1331–1341.
357. Moe SM, O'Neill KD, Reslerova M, Fineberg N, Persohn S, Meyer CA. Natural history of vascular calcification in dialysis and transplant patients. *Nephrol Dial Transplant.* 2004;19(9):2387–2393.
358. Oschatz E, Benesch T, Kodras K, Hoffmann U, Haas M. Changes of coronary calcification after kidney transplantation. *Am J Kidney Dis.* 2006;48(2):307–313.
359. Cianciolo G, Capelli I, Angelini ML, et al. Importance of vascular calcification in kidney transplant recipients. *Am J Nephrol.* 2014;39(5):418–426.
360. Chau K, Martinez G, Elder GJ. Vascular calcification in patients undergoing kidney and simultaneous pancreas-kidney transplantation. *Nephrology.* 2014;19(5):275–281.
361. Sgambat K, Moudgil A. Optimization of bone health in children before and after renal transplantation: current perspectives and future directions. *Front Pediatr.* 2014;2:13.

QUESTIONS

1. A 56-year-old woman with ESRD on hemodialysis is evaluated for painful nodules on the abdomen, which have been present for the past month. Medications include warfarin, metoprolol, atorvastatin, renal vitamin, and calcium acetate. She is receiving doxercalciferol and darbopoietin intravenously with dialysis. Physical examination reveals an overweight woman with several hard, subcutaneous nodules approximately 1 cm in diameter in the lower abdomen; there is one open wound 1.5 cm in diameter with surrounding erythema. Laboratory studies reveal the following results: calcium 9.7 mg/dL; phosphate 6.0 mg/dL; PTH 550 pg/mL. She undergoes a punch biopsy of the wound, which shows findings consistent with calcific uremic arteriolopathy. Which of the following therapeutic approaches would you recommend?
 - a. discontinue warfarin
 - b. start sodium thiosulfate with dialysis
 - c. start vitamin K supplementation
 - d. a and b
 - e. a, b, and c

Answer: d

Rationale: Calciphylaxis or calcific uremic arteriolopathy is a devastating complication in CKD and ESRD, with mortality approaching 50%. Epidemiologic studies have linked warfarin use with calciphylaxis, and a potential mechanism is related to the inhibition by warfarin of the calcification inhibitor matrix Gla protein (MGP). MGP is gamma-carboxylated similar to factors in the coagulation cascade, and higher amounts of undercarboxylated MGP are correlated with vascular calcification. While vitamin K supplementation in theory should rescue the defect in MGP, no definitive data are yet available from ongoing studies on the efficacy of vitamin K supplementation in reducing or reversing vascular calcification, or specifically in outcomes in calciphylaxis. In epidemiologic studies and case reports, sodium thiosulfate was shown to be at times useful in treatment of calciphylaxis, although the exact mechanism by which it reduces extraosseous calcification is not known.

2. A 60-year-old female patient 6 months post kidney transplant is evaluated in routine follow-up. Physical examination is unremarkable. Laboratory studies reveal the following: creatinine 1.0 g/dL, calcium 10.8 mg/dL; phosphate 2.8 mg/dL; PTH 150 pg/mL. Which of the following statements about her condition is true?
 - a. Hyperparathyroidism is uncommon following a successful kidney transplant
 - b. She should be referred to an endocrine surgeon for parathyroidectomy

- c. Cinacalcet may offer effective therapy for her hyperparathyroidism
- d. Post-transplant hypercalcemia is not associated with increased mortality or graft loss
- e. She should have FGF23 levels checked given borderline low phosphate levels

Answer: c

Rationale: Postrenal transplant hyperparathyroidism is commonly observed, although it usually resolves by 6 to 12 months post transplant. The associated hypercalcemia has been associated with increased mortality and graft loss in epidemiologic studies, and thus control of the hypercalcemia is indicated. While parathyroidectomy can be used for refractory cases or those that fail medical management, cinacalcet is currently a reasonable option, especially early in the post-transplant period. FGF23 levels may be increased in post-transplant patients and contribute to the hypophosphatemia, although the primary endocrine abnormality in this case is hyperparathyroidism, not hyperphosphatemia.

3. Which one of the following statements about phosphate control in patients with CKD not on dialysis is correct?
 - a. Phosphate binders effectively lower phosphate levels in CKD patients not on dialysis.
 - b. Phosphate binders have been shown to decrease vascular calcifications and mortality in placebo-controlled trials.
 - c. Phosphate levels drawn during an office visit accurately reflect total body phosphate stores.
 - d. Vegetarian protein sources have less bioavailable phosphate than animal protein sources.
 - e. The optimal phosphate level of a patient with CKD is <4 mg/dL.

Answer: d

Rationale: Whereas epidemiologic studies have shown that higher phosphate levels are associated with increased mortality, no prospective placebo-controlled study has shown that lowering phosphate levels to a particular target lowers clinical outcomes in the CKD or ESRD populations. Few studies have been done in patients with CKD, and they have not shown consistent decreases in serum phosphate concentrations, partly because circulating phosphate concentrations do not adequately reflect total body stores. Phosphate in plant sources is less bioavailable because some of it is in phytate, which humans cannot digest, and small studies using vegetarian protein sources compared with animal product protein sources have shown an effect on not only phosphate levels but also FGF23.

4. Which of the following statements are true concerning the integrated regulation of calcium and phosphorus?
- a. Both PTH and FGF 23 stimulate the activation of 25 hydroxycholecalciferol to 1,25 dihydroxy cholecalciferol to enhance gastrointestinal absorption of calcium and phosphate.
 - b. Both hypocalcemia and PTH stimulate the release of FGF23.
 - c. Both FGF23 and 1,25 dihydroxy cholecalciferol increase the production of PTH.
 - d. Increased phosphate load in CKD results in increases of both PTH and FGF23, which downregulate the renal sodium phosphate transporters.
 - e. b and d

Answer: d

Rationale: Hyperphosphatemia, or more specifically increased phosphate loading, causes an increase in both PTH and FGF-23, the latter from bone. Both the elevated PTH and FGF-23 increase urinary phosphate excretion through downregulation of NaPi transporters. The effect of FGF-23 in the kidney is klotho dependent. PTH also causes an increase in calcium reabsorption, thus decreasing urinary calcium. PTH stimulates the conversion of 25 hydroxycholecalciferol to 1,25 dihydroxy cholecalciferol, by stimulating CYP27B1 activity, whereas FGF23 inhibits CYP27B1 and stimulates CYP24, thereby decreasing 1,25 dihydroxycholecalciferol levels. Hypocalcemia is a major stimulus for PTH release; however, it has an inhibitory effect on FGF23, whereas hypercalcemia will stimulate FGF23. Both FGF23 and 1,25 dihydroxy cholecalciferol inhibit the release of PTH.