

Tip Lesion Variant of FSGS

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. The localized sclerotic lesion that only involves the proximal tubular pole of the glomerulus has been classified as the tip lesion, with some evidence suggesting a better prognosis than typical FSGS. As in minimal change disease and FSGS, there are no immune complexes, and electron microscopy shows extensive foot process effacement (not shown, see minimal change disease). (Jones' silver stain, $\times 200$).




Fig 2. The sclerotic lesion here is sharply localized to the proximal tubular pole, with adhesion and continuity of the capillary loop to Bowman's capsule, and intercapillary foam cells localized to this area, characteristic of the so-called tip lesion variant of FSGS. (Jones' silver stain, $\times 400$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 38(6):E30.

© 2001 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90047-0](http://dx.doi.org/10.1053/S0272-6386(13)90047-0)