

Immunotactoid Glomerulopathy

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Immunotactoid glomerulopathy typically shows mesangial proliferation or a membranoproliferative pattern by light microscopy, as in this case. There is widespread splitting of the glomerular basement membrane and increased mesangial matrix, with thickened Bowman's capsule and adhesions of the tuft to Bowman's capsule (Jones' silver stain, $\times 400$).




Fig 2. Immunotactoid glomerulopathy shows positivity for IgG and rarely other immunoglobulins, along with complement. The positivity is in a segmental capillary loop distribution, along with mesangial positivity, mirroring the distribution of deposits by light microscopy. Deposits more typically are polyclonal, although there may be predominance of one light chain, as in this case, where there was predominant IgG kappa staining, but with concomitant lambda staining (anti-kappa antibody, $\times 400$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman.


Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 41(1):E23-E24.

© 2003 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.


0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90048-2](http://dx.doi.org/10.1053/S0272-6386(13)90048-2)



Copyright © 2003 by the National Kidney Foundation

Fig 3. Immunotactoid glomerulopathy is defined, in our view, by the presence of organized deposits by electron microscopy, which show stacking in parallel arrays and/or microtubular substructure. This morphology may be seen in immunotactoid glomerulopathy, or be associated with paraproteins such as cryoglobulin (see Cryoglobulinemic Glomerulonephritis section). In this case there are massive mesangial and subendothelial deposits with deposits organized in parallel arrays, some cut longitudinally and some cut in cross-section. The microtubules are quite large, close to 50 nm in diameter (transmission electron microscopy, $\times 4,000$).



Copyright © 2003 by the National Kidney Foundation

Fig 4. Immunotactoid glomerulopathy deposits show organized structure by electron microscopy, either organized in parallel arrays, and/or with microtubular substructure. Some deposits are cut in cross-section, clearly revealing the microtubular substructure, whereas others, stacked in adjacent parallel arrays, are cut longitudinally. These deposits permeate the glomerular basement membrane; there is complete overlying effacement of foot processes on the left (transmission electron microscopy, $\times 9,500$).