

Light Chain Cast Nephropathy

Agnes Fogo, MD

The AJKD Atlas of Renal Pathology presents a compilation of figures on a specific pathologic entity. You may download the figures to create your own personal, non-commercial library of images or to create slides for teaching purposes.




Fig 1. Tubular casts with surrounding syncytial giant cell reaction and chronic tubulointerstitial nephritis and fibrosis, diagnostic of myeloma cast nephropathy. The glomerulus on the right shows no lesions, although occasionally glomeruli may be involved with a paraprotein process such as light-chain deposition disease or amyloid in cases of myeloma cast nephropathy (hematoxylin & eosin; original magnification $\times 100$).




Fig 2. High power view of intratubular casts that appear fractured and very refractile with surrounding syncytial giant cell reaction. These findings are diagnostic of myeloma kidney. There is moderate interstitial fibrosis and lymphocytic infiltrate (PAS stain; original magnification $\times 400$).

From the Department of Pathology, Vanderbilt University Medical Center, Nashville, TN.

Medical Photographer: Brent Weedman; with assistance from Kim Solez, MD, of the National Kidney Foundation's cyberNephrology™ Team.

Address author queries to Agnes Fogo, MD, Department of Pathology, Vanderbilt University Medical Center, MCN C-3310, Nashville, TN 37232. E-mail: Agnes.Fogo@vanderbilt.edu

Am J Kidney Dis. 32(4):E1-E2.

© 1998 by the National Kidney Foundation, Inc. Published by Elsevier Inc. All rights reserved.

0272-6386/\$36.00

[http://dx.doi.org/10.1053/S0272-6386\(13\)90009-3](http://dx.doi.org/10.1053/S0272-6386(13)90009-3)




Fig 3. Close-up of intratubular refractile casts with surrounding syncytial giant cell reaction with chronic tubulointerstitial nephritis and fibrosis, characteristic of myeloma cast nephropathy (PAS stain; original magnification $\times 400$).




Fig 4. Myeloma intratubular casts show a slightly wormy substructure with fractured ends, which may also be visualized by light microscopy (Fig 2) (transmission electron microscopy; original magnification $\times 10,000$).